AQA GCSE Chemistry | 8462

SPECIFICATION POINT 4.1.1.7

Electronic structure

Mark schemes and examiner comments

Tier	**Foundation**
Question groups	**32**
Available paper years	**2008–2025**

Using this booklet

1. Booklet answer numbers match the question booklet.
2. Use the original question number within marking tables.
3. Examiner comments are included where available.

Name: _____ **Date:** _____

Where a precise marking extract is unavailable, the full source document is included in an appendix. Any missing mark scheme is explicitly identified.

4.1.1.7 Electronic structure

Answer 01 · Mark scheme

Question 3

question	answers	extra information	mark
(a)	gives out (heat)		1
(b)	D		1
(c)	L		1
(d)	magnesium chloride		1
total			4

4.1.1.7 Electronic structure

Answer 01 · Mark scheme

January 2008 | CHY2F | Original Q3 | Legacy

CHY2F

4.1.1.7 Electronic structure

Answer 02 · Mark scheme

Question 1

question	answers	extra information	mark
(a)(i)	B		1
(a)(ii)	A		1
(a)(iii)	E		1
(a)(iv)	D		1
(b)(i)	Mendeleev and Newlands		1
(b)(ii)	atomic weight		1
(b)(iii)	chemical reactions		1
(b)(iv)	electrons		1
total			8

4.1.1.7 Electronic structure

Answer 02 · Mark scheme January 2008 | CHY3F | Original Q1 | Legacy

CHY3F

Answer 03 · Mark scheme June 2008 | CHY2F | Original Q7 | Legacy

question	answers	extra information	mark
7(a)	157	correct answer with **or** without working (2 ×19 + 119) for **1** mark only allow (119 + 19 =) 138 for **1** mark only ignore units	2
7(b)	24.2	accept answers in the range 24 to 24.2038..... ignore incorrect rounding after correct answer 25 only without working gains **1** mark **or** 38/157 × 100 gains **1** mark **or** (19/157 × 100 =) 12 to 12.1 gains **1** mark allow error carried forward from part(a) 38/(a) × 100 gains **2** marks if calculated correctly (19/138 ×100 =) 13.8 gains **1** mark	2
7(c)	0.29	accept answers in the range 0.28 to 0.3 allow error carried forward from part (b) (b)/100 × 1.2 correctly calculated ignore units	1

Question 7 continued on the next page…

COMPONENT NUMBER: CHY2F

COMPONENT NAME: Additional Science / Chemistry

STATUS: Final

DATE: June 2008

Question 7 continued…

question	answers	extra information	mark
7(d)	an <u>electron</u>	allow electrons allow electron shared / lost for **1** mark apply list principle for additional particles	1
	is gained owtte	must be linked to electron accept can hold / take in if in correct context eg it can hold another electron (in its outer shell) = **2** marks it can take an electron (from another atom) = **2** marks ignore reference to fluoride ions incorrect number of electrons gained does **not** gain the second mark	1
Total			**7**

4.1.1.7 Electronic structure

Answer 03 · Mark scheme

June 2008 | CHY2F | Original Q7 | Legacy

COMPONENT NUMBER: CHY2F

COMPONENT NAME: Additional Science / Chemistry

STATUS: Final

DATE: June 2008

4.1.1.7 Electronic structure

Answer 04 · Mark scheme

June 2009 | CHY2F | Original Q6 | Legacy

Question 6

question	answers	extra information	mark
6(a)(i)	gives out a large amount of energy only water produced / product is non polluting (owtte)	 allow it does not harm the environment	1 1
6(a)(ii)	does not explode / burst into flames owtte	ignore will not react	1
6(a)(iii)	hydrogen absorbed and released much faster	allow more efficient allow can store a larger amount	1
6(b)(i)	B		1
6(b)(ii)	a lithium atom loses an electron		1
6(b)(iii)	C		1
6(c)	reversible		1
6(d)(i)	much smaller		1
6(d)(ii)	surface area		1
Total			**10**

4.1.1.7 Electronic structure

Answer 04 · Mark scheme


June 2009 | CHY2F | Original Q6 | Legacy

CHY2F

4.1.1.7 Electronic structure

Answer 05 · Mark scheme

Question 7

question	answers	extra information	mark
7(a)		accept dots / crosses / e must be drawn on diagram electrons do not need to be paired ignore brackets or + or −charges ignore 2,8,7	1
7(b)	(one) electron lost / given away / transferred from sodium / transferred to chlorine owtte	recognition that electrons are involved must be linked to electrons accept loses electron(s) for **2** marks NB loses 2 or more electrons gains **1** mark reference to sharing / covalent max **1** mark ignore charges on ions formed	1 1
7(c)(i)	any **one** from: • ions / atoms / they are / it is negatively charged / anions • opposite (charges) attract	 accept they are negative accept they are attracted or it is oppositely charged ignore opposite forces attract	1
7(c)(ii)	hydrogen	accept H_2 ignore H or H^+	1

Question 7 continues on next page

CHY2F

Question 7 continued

question	answers	extra information	mark
7(d)(i)	poisons released into environment (owtte)	accept any sensible idea of harm / harmful / poisons / poisonous / pollution / damaging do **not** accept answers such as global warming / ozone layer etc. ignore safety unless qualified	1
7(d)(ii)	any **one** sensible idea eg • loss of work / unemployment **or** company goes out of business • any adverse effect on local economy (owtte) • any adverse effect on paper production / cost of paper / cost of water (treatment) • chlorine (compounds) have been used (for many years) without causing harm owtte • only a tiny amount of chlorine is released so it would not cause harm	eg shops / house prices etc. allow less expensive to use chlorine or converse ignore uses of chlorine to treat drinking water unless qualified	1
7(d)(iii)	ideas related to bias	accept more reliable or valid or fair ignore more accurate / fair test	1
Total			**8**

4.1.1.7 Electronic structure

Answer 05 · Mark scheme

January 2010 | CHY2F | Original Q7 | Legacy

CHY2F

4.1.1.7 Electronic structure

Answer 06 · Mark scheme

June 2010 | CHY3F | Original Q3 | Legacy

Question 3

question	answers	extra information	mark
3(a)(i)	**A**		1
3(a)(ii)	**F**		1
3(a)(iii)	**E**		1
3(a)(iv)	**C**		1
3(a)(v)	**A or B**		1
3(b)(i)	Rb K Na	allow rubidium, potassium, sodium do **not** accept RB or NA	1
3(b)(ii)	decrease **or** become lower / smaller / less	allow from 180° C to 27° C	1
3(c)	They are harder than Group 1 metals. They have higher melting points than Group 1 metals. They often form coloured compounds but Group 1 compounds are usually white.		1 1 1
Total			**10**

4.1.1.7 Electronic structure

Answer 06 · Mark scheme June 2010 | CHY3F | Original Q3 | Legacy

CHY3F

4.1.1.7 Electronic structure

Answer 07 · Mark scheme

Question 1

question	answers	extra information	mark
1(a)	less dense than air		1
1(b)(i)	water	accept hydrogen oxide do **not** accept hydrogen dioxide / hydro oxide	1
1(b)(ii)	unreactive		1
1(c)	atoms		1
1(d)	electron(s) nucleus		1 1
Total			**6**

4.1.1.7 Electronic structure

Answer 07 · Mark scheme

January 2011 | CHY1F | Original Q1 | Legacy

CHY1F

4.1.1.7 Electronic structure

Answer 08 · Mark scheme

Question 7

question	answers	extra information	mark
7(a)(i)	65	correct answer with or without working = **2** marks if answer incorrect evidence of (81 – 16) for **1** mark ignore units	2
7(a)(ii)	zinc	accept error carried forward from (a)(i) allow correct symbol answer given should be element / metal closest to their answer do **not** allow compounds	1
7(b)(i)	• it loses electrons • three electrons	sharing / covalency = max **1** mark	1 1
7(b)(ii)	8 electrons shown in second shell.	accept dots / crosses / mixture of dots and crosses / e electrons do not need to be paired do **not** allow extra electrons in first shell	1
Total			**6**

4.1.1.7 Electronic structure

Answer 09 · Mark scheme

Question 1

question	answers	extra information	mark
1(a)(i)	B		1
1(a)(ii)	E		1
1(a)(iii)	F		1
1(a)(iv)	D		1
1(a)(v)	C		1
1(b)(i)	Br	do not accept BR or br or bR ignore numbers allow written in table if answer blank	1
1(b)(ii)	I Br Cl	allow iodine, bromine, chlorine allow I,B,C allow capitals or lower case allow 184, 58, –34 ignore numbers	1
1(c)	they are halogens they become less reactive down Group 7		1 1
Total			9

4.1.1.7 Electronic structure

Answer 09 · Mark scheme

January 2011 | CHY3F | Original Q1 | Legacy

CHY3F

4.1.1.7 Electronic structure

Answer 10 · Mark scheme

June 2011 | CHY2F | Original Q5 | Legacy

Question 5

question	answers	extra information	mark
5(a)	gives out / releases / transfers to surroundings heat / energy	ignore light / burns ignore the wire gets hot	1
5(b)	activation energy		1
5(c)	(aluminium +) oxygen (→) aluminium oxide	accept correct formulae	1
5(d)	C		1
5(e)(i)	a negative		1
5(e)(ii)	loses		1
5(e)(iii)	gains two		1 1
Total			**8**

4.1.1.7 Electronic structure

Answer 10 · Mark scheme

June 2011 | CHY2F | Original Q5 | Legacy

CHY2F

4.1.1.7 Electronic structure

Answer 11 · Mark scheme

June 2012 | CH1FP | Original Q1 | Legacy

Question 1

question	answers	extra information	mark
1(a)(i)	nucleus		1
1(a)(ii)	protons		1
1(b)	protons / + / positive electrons / – / negative	both words needed in any order for **1** mark	1
1(c)	nitrogen	allow N or N_2	1
1(d)	B **and** C (both) have one electron **or** same number of electrons in the outer energy level / shell	both letters needed in any order for **1** mark allow Li **and** Na allow both are in Group **1** allow both are alkali metals allow both can lose only one electron **or** become +1 ions allow this mark if no letters given in boxes	1 1
Total			**6**

4.1.1.7 Electronic structure


Answer 11 · Mark scheme

June 2012 | CH1FP | Original Q1 | Legacy

CH1FP

4.1.1.7 Electronic structure

Answer 12 · Mark scheme

question	answers	extra information	mark
1(a)(i)	nucleus		1
1(a)(ii)	an energy level (shell)		1
1(b)			1 1 1
1(c)	2/two(%)		1
1(d)(i)	10/ten		1
1(d)(ii)	(group) 0	accept noble gases ignore (group) 8	1
Total			**8**

4.1.1.7 Electronic structure

Answer 12 · Mark scheme

January 2013 | CH1FP | Original Q1 | Legacy

CH1FP

4.1.1.7 Electronic structure

Answer 13 · Mark scheme

January 2013 | CH2FP | Original Q8 | Legacy

question	answers	extra information	mark
8(a)(i)	because they are positively charged	accept they are positive / H^+ accept oppositely charged **or** opposites attract ignore they are attracted	1
8(a)(ii)	gains one / an electron	accept $H^+ + e^- \rightarrow H$ or multiples allow gains electrons	1
8(b)	3 bonding pairs 1 lone pair	 accept 2 non-bonding electrons on outer shell of nitrogen	1 1
8(c)(i)	hydroxide / OH^-	do **not** accept sodium hydroxide	1
8(c)(ii)	$H^+ + OH^- \rightarrow H_2O$	ignore state symbols ignore word equation	1

January 2013

CH2FP

question	answers	extra information	mark
8(d)	Marks awarded for this answer will be determined by the Quality of Written Communication (QWC) as well as the standard of the scientific response. Examiners should also refer to the information on page 5.		6

0 marks	Level 1 (1-2 marks)	Level 2 (3-4 marks)	Level 3 (5-6 marks)
No relevant content.	There are basic descriptions of advantages or disadvantages of the electrolysis cells.	There are clear descriptions of environmental or economic advantages or disadvantages of the electrolysis cells. Comparisons may be implied.	There are detailed descriptions of environmental and economic advantages and disadvantages, comparing the electrolysis cells.

examples of chemistry points made in the response:

Accept converse where appropriate.

- mercury cell is more expensive to construct
- mercury is recycled but membranes must be replaced
- mercury is toxic but membrane / polymer is not
- removing traces of mercury from waste is expensive
- mercury cell uses more electricity
- mercury cell produces chlorine that is purer
- mercury cell produces higher concentration / better quality of sodium hydroxide (solution)

Total			12

UMS conversion calculator www.aqa.org.uk/umsconversion

4.1.1.7 Electronic structure

Answer 14 · Mark scheme

June 2014 | CH2FP | Original Q4 | Legacy

Question	Answers	Extra information	Mark	AO / Spec. Ref.	ID
4(a)(i)	14		1	2 / 2.3.1c	G
4(a)(ii)	isotope		1	1 / 2.3.1d	G
4(a)(iii)	(very) small	*accept smaller / tiny / (very) little*	1	1 / 2.3.1b	G
4(b)(i)	C		1	1 / 2.1	A
4(b)(ii)	NH_3		1	1 / 2.1.1g	A
4(c)(i)	nitric (acid)		1	1 / 2.6.2b	G
4(c)(ii)	indicator		1	1 / 2.6.1b	E
4(c)(iii)	crystallisation **or** evaporation	allow by heating **or** cooling **or** leave (on windowsill) *do not accept freezing*	1	1 / 2.6.1c	E
4(c)(iv)	any **one** from: • grass grows faster • grass grows taller **or** thicker	*allow grass grows better / greener*	1	3 / 2.6.2c	E
4(d)	potassium (atom) loses (an electron) chlorine (atom) gains (an electron) 1 (electron) electron	reference to incorrect bonding or particle = max **3** ignore references to full outer shells	1 1 1 1	1 + 2 / 2.1.1c/d/e	E
Total			**13**		

4.1.1.7 Electronic structure

Answer 14 · Mark scheme

June 2014 | CH2FP | Original Q4 | Legacy



4.1.1.7 Electronic structure

Answer 15 · Mark scheme

June 2015 | CH1FP | Original Q1 | Legacy

Question	Answers	Extra information	Mark	AO / Spec. Ref.
1(a)	gold		1	1 / 1.3.1c
1(b)	atom (s)		1	1 / 1.1.1a
1(c)(i)	protons neutrons	any order allow proton allow neutron	1 1	1 / 1.1.1c/h
1(c)(ii)	3 / three		1	2 / 1.1.1e/g / 1.1.2a
1(d)(i)	Al	ignore any numbers / charges	1	2 / 1.1.1b
1(d)(ii)	any **two** from: • limited resource • expensive in terms of energy / mining • effects on the environment, such as, landfill, atmospheric pollution, quarrying	allow uses a lot of energy to extract	2	1 / 1.3.1e/i/j
1(e)	resistant to corrosion does not react (with water or food)	allow **one** mark for low density with a suitable reason given	1 1	2+3 / 1.3.3c
Total			10	

4.1.1.7 Electronic structure

Answer 15 · Mark scheme

June 2015 | CH1FP | Original Q1 | Legacy



4.1.1.7 Electronic structure

Answer 16 · Mark scheme


June 2015 | CH2FP | Original Q1 | Legacy

Question	Answers	Extra information	Mark	AO / Spec. Ref.
1(a)	1 very small	must be in this order accept negligible, 1/2000 allow zero	1 1	**AO1** 2.3.1b
1(b)	The mass number		1	**AO1** 2.3.1c
1(c)	C		1	**AO2** 2.3.1a
1(d)(i)	2		1	**AO2** 2.1.1a
1(d)(ii)	3		1	**AO2** 2.1.1a
1(e)(i)	28		1	**AO2** 2.3.1f
1(e)(ii)	42.9	accept ecf from (e)(i) accept 42 - 43	1	**AO2** 2.3.3a
1(f)(i)	0.9		1	**AO2** 2.3
1(f)(ii)	any **one** from: • accurate • sensitive • rapid • small sample		1	**AO1** 2.3.2a
Total			**10**	

4.1.1.7 Electronic structure

Answer 16 · Mark scheme

June 2015 | CH2FP | Original Q1 | Legacy



4.1.1.7 Electronic structure

Answer 17 · Mark scheme


June 2015 | CH2FP | Original Q4 | Legacy

Question	Answers	Extra information	Mark	AO / Spec. Ref.
4(a)	any **one** from: • protection/improve lifespan • improve appearance		1	**AO2** 2.7.1d
4(b)(i)	Bleach		1	**AO1** 2.7.1i
4(b)(ii)	Hydrogen is less reactive than sodium.		1	**AO2** 2.7.1f
4(b)(iii)	1 bonding pair of electrons 6 unbonded electrons on Cl	accept dot, cross or e or – or any combination	1	**AO1** 2.7.1e
4(b)(iv)	Covalent		1	**AO1** 2.7.1f
4(b)(v)	Hydrogen chloride has a low boiling point. Hydrogen chloride is made of simple molecules.		1 1	**AO1** 2.2.1a 2.1.1g
4(c)(i)	oxygen	accept carbon dioxide	1	**AO1** 2.7.1h
4(c)(ii)	aluminium ions are positive so are attracted (to the negative electrode)	allow opposites attract	1 1	**AO1/AO2** 2.7.1h 2.2.2b
4(c)(iii)	Reduction		1	**AO1** 2.7.1e
4(c)(iv)	slide	allow move	1	**AO1** 2.2.4b
4(d)(i)	C		1	**AO1** 2.2.3a,c
4(d)(ii)	strong covalent bonds		1	**AO1** 2.2.3a
Total			**14**	

4.1.1.7 Electronic structure

Answer 17 · Mark scheme

June 2015 | CH2FP | Original Q4 | Legacy



4.1.1.7 Electronic structure

Answer 18 · Mark scheme


June 2015 | CH3FP | Original Q2 | Legacy

Question	Answers	Extra information	Mark	AO / Spec. Ref.
2(a)(i)	Na	allow sodium	1	**AO1** 3.1.3.a
2(a)(ii)	Cu	allow copper	1	**AO2** 3.1.3.c
2(a)(iii)	C	allow carbon	1	**AO2** 3.1.2.b
2(a)(iv)	He	allow helium	1	**AO1** 3.1.2.b
2(b)	H	allow hydrogen do **not** allow H_2	1	**AO2** 3.1.2.b
Total			**5**	

4.1.1.7 Electronic structure

Answer 18 · Mark scheme

June 2015 | CH3FP | Original Q2 | Legacy



4.1.1.7 Electronic structure

Answer 19 · Mark scheme

June 2016 | CH1FP | Original Q1 | Legacy

Question 1

Question	Answers	Extra information	Mark	AO / Spec. Ref.
1(a)(i)	7		1	AO2 1.1.1g
1(a)(ii)	-1		1	AO1 1.1.1d
1(a)(iii)	neutrons		1	AO1 1.1.1c
1(b)	number of protons		1	AO1 1.1.1f
1(c)	atom **Y**		1	AO2 1.1.1 h; 1.1.2a
1(d)(i)	Ne	allow neon	1	AO2 1.1.2a/b
1(d)(ii)	has a full outer shell **or** full outer energy level **or** has 8 electrons in its outer shell	allow in Group 0 allow a noble gas allow the shells are full ignore in Group 8	1	AO2 1.1.2b; 1.1.3a
Total			**7**	

4.1.1.7 Electronic structure


Answer 20 · Mark scheme June 2016 | CH2FP | Original Q1 | Legacy

Question	Answers	Extra information	Mark	AO / SpecRef
1(a)(i)	7 / seven		1	AO1 2.1.1e
1(a)(ii)	1 Electron	do **not** accept -1	1 1	AO1 2.3.1b
1(a)(iii)	isotopes		1	AO1 2.3.1d
1(b)(i)	(sodium +) fluorine → sodium fluoride		1	AO2 2.1.1d,e
1(b)(ii)	compounds		1	AO1 2.1.1a
1(b)(iii)	mole		1	AO1 2.3.1g
1(b)(iv)	sodium (atom) loses fluorine (atom) gains one electron ions formed	**max 3** if reference to incorrect particle / bonding allow sodium forms positive (ion) **or** fluorine forms negative (ion) allow form ionic bond allow to gain a full outer shell of electrons allow forms noble gas structure	 1 1 1 1	AO1 / AO2 2.1.1b,c,d,e,f
1(b)(v)	Dissolve in water High melting point		1 1	AO1 2.2.2a,b
Total			**13**	

4.1.1.7 Electronic structure



Answer 20 · Mark scheme

June 2016 | CH2FP | Original Q1 | Legacy



4.1.1.7 Electronic structure

Answer 21 · Mark scheme June 2017 | CH1FP | Original Q6 | Legacy

Question	Answers	Extra information	Mark	AO / Spec. Ref.
6(a)(i)	electronic structure drawn	allow any representation of an electron allow 2,4	1	**AO2** 1.1.1h
6(a)(ii)	six / 6 protons (protons) are positively charged	 do **not** allow electrons in nucleus otherwise ignore electrons do **not** allow nucleus is neutral allow (protons are) + / +1 / 1+ ignore statements about mass if no other mark awarded allow one mark for nucleus	1 1 1	**AO1** **AO2** 1.1.1c;d;e
6(b)	(a hydrocarbon is) made up of hydrogen and carbon (atoms) only	do **not** allow mixture	1	**AO1** 1.4.1a;c
6(c)(i)	 	ethane correctly drawn with another hydrogen on each carbon and another three single bonds ethene correctly drawn with a double bond between the carbon atoms	1 1	**AO1** 1.4.2a 1.5.1c
6(c)(ii)	as the number of carbon atoms increases the boiling point increases	allow converse ignore alkene	1	**AO3** 1.4.2c

4.1.1.7 Electronic structure


Answer 21 · Mark scheme

Question	Answers	Extra information	Mark	AO / Spec. Ref.
6(c)(iii)	the boiling points of alkanes are higher than the boiling points of alkenes	allow the boiling points of alkenes are lower than the boiling points of alkanes	1	AO2 1.4.2b;c
	that has the same number of carbon atoms	consequential on first marking point	1	
	or state a correct example	e.g. hexane has a higher boiling point than hexene **or** hexane boils at 69°C hexene boils at 64°C if no other mark awarded allow both alkanes and alkenes have the same pattern of the more carbon atoms the higher the boiling point for one mark		
6(d)	(alkane molecules are) cracked	allow break down / up **or** decompose **or** split up	1	AO1 AO2 AO3
	to produce small(er) molecules / alkanes / hydrocarbons	allow short chain for small	1	1.5.1a;b;d
	and alkenes **or** a named alkene	allow molecules that have a double bond **or** are unsaturated if no other mark awarded allow evaporate **or** boil for one mark ignore turns to gas	1	
Total			13	

4.1.1.7 Electronic structure

Answer 21 · Mark scheme

June 2017 | CH1FP | Original Q6 | Legacy



4.1.1.7 Electronic structure


Answer 22 · Mark scheme

Question	Answers	Extra information	Mark	AO / Spec. Ref.
3(a)(i)	tick (✓) by: covalent		1	AO1 2.1.1g
3(a)(ii)	ticks (✓) by: oxygen consists of small molecules oxygen has a low boiling point		1 1	AO1 2.1.1g
3(b)(i)	magnesium (atom) loses (electrons) oxygen (atom) gains (electrons) two (electrons) ions formed	max 3 if reference to incorrect particle or incorrect bonding accept ionic bonding or full outer shells/noble gas structure is formed allow magnesium forms positive (ion) **or** oxygen forms negative (ion)	1 1 1 1	AO1 AO2 2.1.1a,b,c, d,e,f
3(b)(ii)	ticks (✓) by: nanoparticles contain a few hundred atoms nanoparticles have a high surface area to volume ratio		1 1	AO1 2.2.6a
3(c)	giant covalent		1	AO1 2.2.3a
Total			10	

4.1.1.7 Electronic structure

Answer 22 · Mark scheme

June 2017 | CH2FP | Original Q3 | Legacy



4.1.1.7 Electronic structure

Answer 23 · Mark scheme

June 2017 | CH2FP | Original Q4 | Legacy

Question	Answers	Extra information	Mark	AO / Spec. Ref.
4(a)	58.5		1	AO2 2.5.1b
4(b)	water		1	AO2 2.3.3a
4(c)(i)	prevents loss of thermal energy	allow prevents heat loss	1	AO1 2.6.1
4(c)(ii)	13/thirteen		1	AO3 2.5.1
4(c)(iii)	temperature changes/results are all similar	accept no anomalous results allow they are all similar do **not** accept temperature	1	AO2 2.5.1
4(c)(iv)	exothermic	allow giving out energy	1	AO3 2.5.1
4(d)(i)	ions move	allow they move ignore free ions do not accept movement of electrons	1	AO1 2.7.1a 2.2.2b
4(d)(ii)	(chloride ions are) negative	accept opposites attract do **not** accept chlorine is negative	1	AO1 2.7.1c
4(d)(iii)	Tick (✓) by: soap		1	AO1 2.7.1i
4(d)(iv)	1 bonding pair of electrons 6 unbonded electrons on Cl	accept dot, cross, e or – or any combination	1 1	AO2 2.1

4.1.1.7 Electronic structure


Answer 23 · Mark scheme

4(d)(v)	Tick (✓) by: $\frac{1}{36.5}$ x 100		1	**AO2** 2.3.3a
Total			**12**	

4.1.1.7 Electronic structure

Answer 23 · Mark scheme


June 2017 | CH2FP | Original Q4 | Legacy



4.1.1.7 Electronic structure

Answer 24 · Mark scheme


June 2017 | CH3FP | Original Q1 | Legacy

Question	Answers	Extra information	Mark	AO / Spec. Ref.
1 (a)(i)	numbers	allow number allow numbers circled in the box	1	AO1 3.1.2b
1 (a)(ii)	electrons	allow electron allow electrons circled in the box	1	AO1 3.1.2b
1 (b)(i)	Cl/chlorine	allow CL / cL / cl	1	AO1 3.1.3f
1 (b)(ii)	Cu/copper	allow CU / cU / cu	1	AO2 3.1.3c
1 (b)(iii)	7/seven		1	AO1 3.1.3e
1(c)(i)	tick (✓) by: displacement		1	AO1 3.1.3g
1(c)(ii)	tick (✓) by: Fluorine		1	AO3 3.1.3g
1(d)	tick (✓) by: yellow		1	AO1 3.4.1e
1(e)(i)	circle around: 		1	AO2 3.6.2a
1(e)(ii)	carbon dioxide	allow carbon dioxide circled in the box	1	AO2 3.6.2b
1(e)(iii)	esters	allow esters circled in the box allow ethers	1	AO2 3.6.2b
Total			11	

4.1.1.7 Electronic structure

Answer 24 · Mark scheme

June 2017 | CH3FP | Original Q1 | Legacy



Answer 25 · Mark scheme

June 2018 | 8462/1F | Original Q2

The full source mark scheme begins on page 66. Follow original question 2 in that document.

4.1.1.7 Electronic structure

Answer 26 · Mark scheme

June 2019 | 8462/1F | Original Q1

Question 1


Question	Answers	Extra information	Mark	AO / Spec. Ref.
01.1	**A** nucleus **B** electron		1 1	AO1 4.1.1.4 4.1.1.7
01.2	electron		1	AO1 4.1.1.5
01.3	3 / three		1	AO2 4.1.2.1
01.4	(atomic number) 5 (mass number) 11		1 1	AO2 4.1.1.5
01.5	isotope		1	AO1 4.1.1.5
01.6	there are the same number of $^{79}_{35}Br$ atoms and $^{81}_{35}Br$ atoms		1	AO2 4.1.1.6
Total			**8**	

4.1.1.7 Electronic structure

Answer 27 · Mark scheme

June 2019 | 8462/1F | Original Q2

Question 2

Question	Answers	Extra information	Mark	AO / Spec. Ref.
02.1	H_2O_2		1	AO2 4.1.1.1 4.2.1.4
02.2	covalent		1	AO1 4.2.1.4
02.3	transition metals		1	AO1 4.1.3.2
02.4	B		1	AO1 4.5.1.2
02.5	A		1	AO2 4.5.1.2
02.6	exothermic		1	AO1 4.5.1.1
02.7	 1 bonding pair of electrons in the right hand overlap 4 non-bonding electrons on oxygen	scores **2** marks allow dots, crosses, circles or $e^{(-)}$ for electrons do **not** accept any change to the number of electrons in the left hand overlap do **not** accept non-bonding electrons on hydrogen ignore inner shell electrons drawn on oxygen	 1 1	AO1 4.2.1.4
Total			**8**	

4.1.1.7 Electronic structure

Answer 28 · Mark scheme

November 2021 | 8462/1F | Original Q3

Question 3

Question	Answers	Extra information	Mark	AO / Spec. Ref.
03.1	2,8,8,1		1	AO2 4.1.1.7

Question	Answers	Extra information	Mark	AO / Spec. Ref.
03.2	they have the same number of outer shell electrons		1	AO1 4.1.2.1 4.1.2.5

Question	Answers	Extra information	Mark	AO / Spec. Ref.
03.3	metallic		1	AO1 4.2.1.5 1-3

Question	Answers	Extra information	Mark	AO / Spec. Ref.
03.4	any **two** from: • bubbles (very) quickly • melts (into a ball) • floats • moves (very) quickly	allow flame	2	AO1 4.1.2.5 4.4.1.2

Question	Answers	Extra information	Mark	AO / Spec. Ref.
03.5	(reactivity) increases (down the group)		1	AO1 4.1.2.5

Question 3 continued

Question	Answers	Extra information	Mark	AO / Spec. Ref.
03.6	any **two** from: • increasing speed of movement • increasing rate of bubble production • doesn't melt → melts • no flame → flame **or** flame → explosion		2	AO3 4.1.2.5 4.4.1.2

Question	Answers	Extra information	Mark	AO / Spec. Ref.
03.7	hydrogen		1	AO1 4.1.2.5 4.4.1.2

Question 3 continued

Question	Answers	Extra information	Mark	AO / Spec. Ref.
03.8	sodium ion structure 2,8 fluoride ion structure 2,8 + charge on sodium ion **and** – charge on fluoride ion	allow any combination of circles, dots, crosses or $e^{(-)}$	1 1 1	AO2 4.2.1.2
	an answer of  sodium ion fluoride ion scores **3** marks			

Total		12

4.1.1.7 Electronic structure

Answer 29 · Mark scheme

June 2022 | 8462/1F | Original Q9

Question 9

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.1	**D**		1	AO3 4.1.2.3


Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.2	**B**		1	AO3 4.1.3.2

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.3	any **two** from: (Group 1 elements) • have lower melting / boiling points • have lower densities • are less strong • are softer	allow converse statements for transition elements allow (Group 1 elements are) more malleable / ductile allow (Group 1 elements) are not useful as catalysts ignore transition elements form coloured compounds ignore transition elements form ions with different charges ignore references to chemical properties	2	AO1 4.1.3.1 4.1.3.2

4.1.1.7 Electronic structure

Answer 29 · Mark scheme

June 2022 | 8462/1F | Original Q9

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.4		allow any combination of x, •, o, $e^{(-)}$ for electrons	1	AO2 4.1.1.7

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.5	delocalised electrons (the electrons) carry (electrical) charge (the electrons move) through the metal / aluminium / structure	allow free electrons ignore current / electricity for charge ignore throughout for through	1 1 1	AO1 4.2.1.5 4.2.2.8

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.6	ionic		1	AO1 4.2.1.1

4.1.1.7 Electronic structure

Answer 29 · Mark scheme

June 2022 | 8462/1F | Original Q9

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.7	magnesium (atom) loses electrons		1	AO2 4.2.1.1 4.2.1.2
	oxygen (atom) gains electrons		1	
	two electrons (are transferred)		1	
	magnesium ions **and** oxide ions are formed	allow Mg^{2+} (ions) **and** O^{2-} (ions) are formed allow magnesium forms positive ions and oxygen forms negative ions allow (both) form a complete outer shell	1	

Total Question 9		**13**

4.1.1.7 Electronic structure

Answer 30 · Mark scheme

June 2023 | 8462/1F | Original Q9

Question 9

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.1	a ball of positive charge with (negative) electrons embedded	do **not** accept references to protons, nuclei, neutrons	1 1	AO1 4.1.1.3

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.2	(earliest) electrons protons (latest) neutrons		1	AO1 4.1.1.3

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.3	(number of outer shell electrons) 7 (reason) (tennessine is in) Group 7	 allow the number of outer electrons is the same as the group number allow tennessine is a halogen MP2 is dependent on MP1 being awarded	1 1	AO2 4.1.2.1 4.1.2.6

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.4	(time needed for) peer review	allow the idea that other scientists had to check the results	1	AO3 4.1.2.2

4.1.1.7 Electronic structure

Answer 30 · Mark scheme

June 2023 | 8462/1F | Original Q9

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.5	(A_r =) $\frac{(6 \times 7.6) + (7 \times 92.4)}{100}$ = 6.924 = 6.9	 allow $\frac{45.6 + 646.8}{100}$ allow (6 × 0.076) + (7 × 0.924) allow 0.456 + 6.468 allow an answer correctly rounded to 1 decimal place from an incorrect calculation which uses all the data in the table	 1 1 1	AO2 4.1.1.6

Total Question 9	9

4.1.1.7 Electronic structure

Answer 31 · Mark scheme

June 2024 | 8462/1F | Original Q9

Question 9

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.1		allow energy level for shell		AO1 4.1.2.1
	(similarity) both have one outer (shell) electron	allow same number of outer (shell) electrons	1	
	(difference) sodium has 3 shells but potassium has 4 shells	allow potassium has more shells allow (different) number of shells	1	

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.2	any **two** from: • effervescence / bubbles / fizzing • (potassium) floats • (potassium) moves around • (potassium) becomes smaller • (potassium) melts • flame • explosion	ignore gas produced allow (potassium) forms a ball ignore colour of flame	2	AO1 4.1.2.5 4.4.1.2

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.3	blue / violet / purple		1	AO2
	(the solution is) alkaline	allow (the solution) contains OH^- (ions) allow (the solution) contains hydroxide ions allow the solution is basic	1	AO1 4.1.2.5 4.4.2.4

4.1.1.7 Electronic structure

Answer 31 · Mark scheme

June 2024 | 8462/1F | Original Q9

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.4	all five points correctly plotted	allow a tolerance of ± ½ a small square allow **1** mark for three or four points correctly plotted	2	AO2 4.1.2.4

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.5	3.4 (mg/cm^3)	allow a value in the range 3.0 to 3.8 (mg/cm^3)	1	AO3 4.1.2.4

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.6	chlorine and potassium bromide		1	AO2 4.1.2.6

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.7	relative molecular mass increases and boiling point increases		1	AO1 4.1.2.6

Total Question 9	**11**


Answer 32 · Mark scheme

June 2025 | 8462/1F | Original Q8

Question 8

Question	Answers	Extra information	Mark	AO / Spec. Ref.
08.1	towards the right **and** the top	allow a description of the position in the periodic table allow in Groups (3) 4 to 0	1	AO1 4.1.2.3

Question	Answers	Extra information	Mark	AO / Spec. Ref.
08.2	a melting point below 100 °C		1	AO3 4.1.2.3 4.2.2.7 4.2.2.8

Question	Answers	Extra information	Mark	AO / Spec. Ref.
08.3		allow any combination of ×, ●, o, $e^{(-)}$ for electrons	1	AO1 4.1.1.7

4.1.1.7 Electronic structure

Answer 32 · Mark scheme

Question	Answers	Extra information	Mark	AO / Spec. Ref.
08.4	any **one** from: • so elements / tellurium / iodine were in groups with similar properties • iodine has similar properties to Group 7 elements • iodine has different properties to Group 6 elements	ignore reference to atomic structure allow named elements for Group 6 / 7 elements allow converse arguments in terms of tellurium	1	AO1 4.1.2.2

Question	Answers	Extra information	Mark	AO / Spec. Ref.
08.5	argon has a full outer shell (of electrons)	allow energy level for shell allow argon has eight outer shell electrons ignore argon has no outer shell electrons	1	AO1 4.1.2.4
	(so argon) has a stable arrangement of electrons **or** (so argon) has a stable electronic structure	allow (so argon) does not gain / lose / share electrons	1	

4.1.1.7 Electronic structure

Answer 32 · Mark scheme

Question	Answers	Extra information	Mark	AO / Spec. Ref.
08.6	atoms with the same number of protons (but with) different numbers of neutrons	allow atoms of the same element allow atoms with the same atomic number ignore atoms with the same number of electrons	1 1	AO1 4.1.1.5

Question	Answers	Extra information	Mark	AO / Spec. Ref.
08.7	~115		1	AO2 4.1.1.6

Total Question 8	**9**

AQA

GCSE
CHEMISTRY

8462/1F - PAPER 1 FOUNDATION TIER

Mark scheme

8462

June 2018

Version/Stage: 1.0 Final

4.1.1.7 Electronic structure

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Assessment Writer.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from aqa.org.uk

Copyright © 2018 AQA and its licensors. All rights reserved.
AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Information to Examiners

1. General

The mark scheme for each question shows:

- the marks available for each part of the question
- the total marks available for the question
- the typical answer or answers which are expected
- extra information to help the Examiner make his or her judgement
- the Assessment Objectives and specification content that each question is intended to cover.

The extra information is aligned to the appropriate answer in the left-hand part of the mark scheme and should only be applied to that item in the mark scheme.

At the beginning of a part of a question a reminder may be given, for example: where consequential marking needs to be considered in a calculation; or the answer may be on the diagram or at a different place on the script.

In general the right-hand side of the mark scheme is there to provide those extra details which confuse the main part of the mark scheme yet may be helpful in ensuring that marking is straightforward and consistent.

2. Emboldening and underlining

2.1 In a list of acceptable answers where more than one mark is available 'any **two** from' is used, with the number of marks emboldened. Each of the following bullet points is a potential mark.

2.2 A bold **and** is used to indicate that both parts of the answer are required to award the mark.

2.3 Alternative answers acceptable for a mark are indicated by the use of **or**. Different terms in the mark scheme are shown by a / ; eg allow smooth / free movement.

2.4 Any wording that is underlined is essential for the marking point to be awarded.

3. Marking points

3.1 Marking of lists

This applies to questions requiring a set number of responses, but for which students have provided extra responses. The general principle to be followed in such a situation is that 'right + wrong = wrong'.

Each error / contradiction negates each correct response. So, if the number of error / contradictions equals or exceeds the number of marks available for the question, no marks can be awarded.

However, responses considered to be neutral (indicated as * in example 1) are not penalised.

Example 1: What is the pH of an acidic solution?

[1 mark]

Student	Response	Marks awarded
1	green, 5	0
2	red*, 5	1
3	red*, 8	0

Example 2: Name two planets in the solar system.

[2 marks]

Student	Response	Marks awarded
1	Neptune, Mars, Moon	1
2	Neptune, Sun, Mars, Moon	0

3.2 Use of chemical symbols / formulae

If a student writes a chemical symbol / formula instead of a required chemical name, full credit can be given if the symbol / formula is correct and if, in the context of the question, such action is appropriate.

3.3 Marking procedure for calculations

Marks should be awarded for each stage of the calculation completed correctly, as students are instructed to show their working. Full marks can, however, be given for a correct numerical answer, without any working shown.

3.4 Interpretation of 'it'

Answers using the word 'it' should be given credit only if it is clear that the 'it' refers to the correct subject.

3.5 Errors carried forward

Any error in the answers to a structured question should be penalised once only.

Papers should be constructed in such a way that the number of times errors can be carried forward is kept to a minimum. Allowances for errors carried forward are most likely to be restricted to calculation questions and should be shown by the abbreviation ecf in the marking scheme.

3.6 Phonetic spelling

The phonetic spelling of correct scientific terminology should be credited **unless** there is a possible confusion with another technical term.

3.7 Brackets

(…..) are used to indicate information which is not essential for the mark to be awarded but is included to help the examiner identify the sense of the answer required.

3.8 Allow

In the mark scheme additional information, 'allow' is used to indicate creditworthy alternative answers.

3.9 Ignore

Ignore is used when the information given is irrelevant to the question or not enough to gain the marking point. Any further correct amplification could gain the marking point.

3.10 Do not accept

Do **not** accept means that this is a wrong answer which, even if the correct answer is given as well, will still mean that the mark is not awarded.

4. Level of response marking instructions

Extended response questions are marked on level of response mark schemes.

- Level of response mark schemes are broken down into levels, each of which has a descriptor.
- The descriptor for the level shows the average performance for the level.
- There are two marks in each level.

Before you apply the mark scheme to a student's answer, read through the answer and annotate it (as instructed) to show the qualities that are being looked for. You can then apply the mark scheme.

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Step 1: Determine a level

Start at the lowest level of the mark scheme and use it as a ladder to see whether the answer meets the descriptor for that level. The descriptor for the level indicates the different qualities that might be seen in the student's answer for that level. If it meets the lowest level then go to the next one and decide if it meets this level, and so on, until you have a match between the level descriptor and the answer.

When assigning a level you should look at the overall quality of the answer. Do **not** look to penalise small and specific parts of the answer where the student has not performed quite as well as the rest. If the answer covers different aspects of different levels of the mark scheme you should use a best fit approach for defining the level.

Use the variability of the response to help decide the mark within the level, ie if the response is predominantly level 2 with a small amount of level 3 material it would be placed in level 2 but be awarded a mark near the top of the level because of the level 3 content.

Step 2: Determine a mark

Once you have assigned a level you need to decide on the mark. The descriptors on how to allocate marks can help with this.

The exemplar materials used during standardisation will help. There will be an answer in the standardising materials which will correspond with each level of the mark scheme. This answer will have been awarded a mark by the Lead Examiner. You can compare the student's answer with the example to determine if it is the same standard, better or worse than the example. You can then use this to allocate a mark for the answer based on the Lead Examiner's mark on the example.

You may well need to read back through the answer as you apply the mark scheme to clarify points and assure yourself that the level and the mark are appropriate.


Indicative content in the mark scheme is provided as a guide for examiners. It is not intended to be exhaustive and you must credit other valid points. Students do **not** have to cover all of the points mentioned in the indicative content to reach the highest level of the mark scheme.

You should ignore any irrelevant points made. However, full marks can be awarded only if there are no incorrect statements that contradict a correct response.

An answer which contains nothing of relevance to the question must be awarded no marks.

4.1.1.7 Electronic structure

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Question	Answers	Extra information	Mark	AO / Spec. Ref.
01.1			1 1 1	AO1 4.1.1.2
01.2	include a (filter) funnel	allow funnel drawn on the diagram ignore clamp stand	1	AO3 4.1.1.2
01.3	evaporate condense	must be this order	1 1	AO1 4.1.1.2
01.4	$\frac{2}{20} \times 100$ = 10 (%)	an answer of 10 (%) scores **2** marks an answer of 11.1(%) **or** 90 (%) scores **1** mark	1 1	AO2 4.2.2.7
01.5	an alloy		1	AO1 4.2.2.7
01.6	the layers in the mixture are distorted		1	AO1 4.2.2.7

4.1.1.7 Electronic structure

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Question	Answers	Extra information	Mark	AO / Spec. Ref.
01.7	8000 nm^3		1	AO2 4.2.4.1
Total			**11**	

4.1.1.7 Electronic structure

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Question	Answers	Extra information	Mark	AO / Spec. Ref.
02.1	7		1	AO1 4.1.2.6
02.2	small molecule		1	AO2 4.1.2.6 4.2.2.4
02.3	F_2		1	AO1 4.1.1.1 4.1.2.6
02.4	the reactivity decreases (going down Group 7) (because) chlorine displaces bromine and iodine (and) bromine displaces iodine **or** iodine does not react	allow the reactivity decreases from chlorine to iodine allow (because) chlorine has two reactions allow (because) neither bromine nor iodine can displace chlorine allow (and) bromine has one reaction **or** iodine has no reactions allow (and) iodine cannot displace bromine	1 1 1	AO1 4.1.2.6 AO3 4.1.2.6 AO3 4.1.2.6
02.5	80		1	AO2 4.3.1.2

4.1.1.7 Electronic structure

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Question	Answers	Extra information	Mark	AO / Spec. Ref.
02.6	(1.2 kg =) 1200 (g) **or** (900 g =) 0.9 (kg)	an answer of 75 (%) scores **2** marks	1	AO2 4.3.3.1
	$(\frac{900}{1200} \times 100) = 75(\%)$ **or** $(\frac{0.9}{1.2} \times 100) = 75(\%)$	allow an answer correctly calculated from: $(\frac{900}{\text{incorrect attempt at conversion of 1.2}} \times 100)$ **or** $(\frac{\text{incorrect attempt at conversion of 900}}{1.2} \times 100)$	1	
Total			**9**	

4.1.1.7 Electronic structure

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Question	Answers	Extra information	Mark	AO / Spec. Ref.
03.1	nucleus neutron neutron electron proton	must be in this order	1 1 1 1 1	AO1 4.1.1.3 4.1.1.4 4.1.1.5
03.2	$(A_r)\ \frac{(63 \times 70) + (65 \times 30)}{100}$ = 63.6	an answer of 63.6 scores **2** marks	1 1	AO2 4.1.1.6
03.3	copper / Cu	allow ecf from answer to question **03.2**	1	AO2 4.1.1.1
03.4	$\frac{1.2 \times 10^{-10}}{10\,000}$ **or** $1.2 \times 10^{-10} \times 1 \times 10^{-4}$ $= 1.2 \times 10^{-14}$ (m)	an answer of 1.2×10^{-14} (m) scores **2** marks a correct answer not in standard form scores **1** mark	1 1	AO2 4.1.1.5
Total			**10**	

4.1.1.7 Electronic structure

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.1	3.6 (cm^3)		1	AO2 4.4.3.4
04.2	hydrogen line only		1	AO3 4.4.3.4
04.3	both lines		1	AO3 4.4.3.4
04.4	graphite has delocalised electrons		1	AO1 4.2.3.2
04.5	**cathode** zinc (1) **anode** chlorine (1) hydrogen (1) bromine (1)	do **not** accept chloride allow **1** mark if chlorine and zinc the wrong way around do **not** accept bromide allow **1** mark if bromine and hydrogen the wrong way around	1+1 1+1	AO2 4.4.3.2 4.4.3.4
Total			**8**	

4.1.1.7 Electronic structure

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Question	Answers	Extra information	Mark	AO / Spec. Ref.
05.1	gas		1	AO1 4.2.2.2
05.2	the gas escapes	allow carbon dioxide escapes do **not** accept references to evaporation	1	AO1 4.3.1.3
05.3	5.12 (g)		1	AO2 4.3.1.3
05.4	4.00 (g) trial 1	allow 2.89 written in either space, or ringed in the table, unless contradicted by mass of copper carbonate or trial number	1	AO3 4.3.1.3
05.5	reheat (and reweigh) until constant mass	 an answer of heat to constant mass scores **2** marks if no other mark scored allow for **1** mark heat for longer **or** (heat at a) higher temperature **alternative approach**: (1) continue heating and pass gas through limewater (1) until the (lime)water stops bubbling **or** until the limewater no longer turns cloudy	1 1	AO3 4.3.1.3

4.1.1.7 Electronic structure


MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Question	Answers	Extra information	Mark	AO / Spec. Ref.
05.6	straight line of best fit	must touch at least 5 of the 6 plots	1	AO2 4.3.1.3
05.7	correct value read from line of best fit in Figure 8	allow tolerance of ± ½ small square	1	AO2 4.3.1.3
05.8	(mass =) $168 \times \frac{\text{answer from question } \mathbf{05.7}}{8.4}$	a correctly calculated value from their answer to question **05.7** scores **2** marks allow (mass =) answer from question **05.7** × 20	1	AO2 4.3.1.3
	correctly calculated value (g)		1	
Total			**10**	

4.1.1.7 Electronic structure

PMT

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Question	Answers	Extra information	Mark	AO / Spec. Ref.
06.1	 allow **one** mark if answers are reversed		1 1	AO1 4.5.1.1
06.2	polystyrene is a better insulator		1	AO3 4.5.1.1
06.3	both bars labelled both bars correctly plotted	 allow tolerance of ± ½ small square ignore width and spacing of bars if no other mark scored, allow **1** mark for any one bar correctly plotted and labelled	1 1	AO2 4.5.1.1

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Question	Answers	Extra information	Mark	AO / Spec. Ref.
06.4	temperature increases **or** temperature does not decrease	ignore because it is exothermic allow (because) energy / 'heat' is transferred to the surroundings energy / 'heat' is not taken in from the surroundings allow the energy of the products is less than the energy of the reactants	1	AO3 4.5.1.1
06.5	(most reactive) magnesium (zinc) nickel	this order only	1	AO3 4.5.1.1 4.4.1.2
06.6	suitable method described the observations / measurements required to place in order an indication of how results would be used to place the unknown metal in the reactivity series **approaches that could be used:** **approach 1:** add the unknown metal to copper sulfate solution (1) measure temperature change (1) place the metals in order of temperature change (1)		1 1 1	AO3 4.4.1.2 4.5.1.1

Question	Answers	Extra information	Mark	AO / Spec. Ref.
06.6 cont.	**approach 2:** add the metal to salt solutions of the other metals **or** heat the metal with oxides of the other metals (1) measure temperature change (only if salt solutions used) **or** observe whether a chemical change occurs (1) compare temperature change or whether there is a reaction to place in correct order (1) **approach 3:** add all of the metals to an acid (1) measure temperature change or means of comparing rate of reaction (1) place the metals in order of temperature change or rate of reaction (1) **approach 4:** set up electrochemical cells with the unknown metal as one electrode and each of the other metals as the other electrode (1) measure the voltage of the cell (1) place the metals in order of voltage (1)			
06.7	D		1	AO1 4.5.1.2
06.8	C		1	AO1 4.5.1.2
Total			**12**	

4.1.1.7 Electronic structure

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Question	Answers	Extra information	Mark	AO / Spec. Ref.
07.1	sodium oxide	allow Na_2O	1	AO2 4.1.1.1 4.1.2.5 4.4.1.1
07.2	oxidation		1	AO1 4.4.1.1
07.3	13		1	AO1 4.4.2.4
07.4	sodium hydroxide		1	AO2 4.1.1.1
07.5	OH^-		1	AO1 4.4.2.4
07.6	(volume =) $\frac{250}{1000}$ or ¼ or 0.25 (dm^3) **or** (mass per cm^3 =) $\frac{40}{1000}$ (g) or 0.04 (g) ($\frac{250}{1000}$ × 40 =) 10 (g)	an answer of 10 (g) scores **2** marks	1 1	AO2 4.3.2.5

4.1.1.7 Electronic structure

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Question	Answers	Extra information	Mark	AO / Spec. Ref.
07.7	all points correct	allow a tolerance of ± ½ a small square allow **1** mark for 3 points correct ignore any attempt at a line of best fit	2	AO2 4.1.2.5
07.8	39 (°C)	allow any value from 34 to 46 (°C)	1	AO2 4.1.2.5
Total			**10**	

4.1.1.7 Electronic structure

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Question	Answers	Extra information	Mark	AO / Spec. Ref.
08.1	any **one** from: • metal • (metal) hydroxide • (metal) carbonate • alkali	allow named example allow correct formula ignore base allow ammonium hydroxide allow ammonium carbonate allow soluble base allow ammonia	1	AO1 4.4.2.1 4.4.2.2 4.4.2.3
08.2	$Ca(NO_3)_2$	allow $Ca^{2+}(NO_3^-)_2$	1	AO2 4.4.2.2

4.1.1.7 Electronic structure

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Question	Answers	Mark	AO / Spec. Ref.
01.3	**Level 3:** The method would lead to the production of a valid outcome. All key steps are identified and logically sequenced.	5–6	AO1 4.4.2.3
	Level 2: The method would not necessarily lead to a valid outcome. Most steps are identified, but the method is not fully logically sequenced.	3–4	
	Level 1: The method would not lead to a valid outcome. Some relevant steps are identified, but links are not made clear.	1–2	
	No relevant content	0	
	Indicative content • use magnesium oxide and sulfuric acid • add sulfuric acid to a beaker • warm sulfuric acid • add magnesium oxide • stir • continue adding until magnesium oxide is in excess • filter • using a filter paper and funnel • to remove excess magnesium oxide • heat solution in an evaporating basin • to crystallisation point • leave to crystallise • pat dry with filter paper credit may be given for diagrams		
Total			**8**

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.1	FeS_2	do **not** accept equations	1	AO2 4.2.1.3
09.2	26 30 26	must be this order	1 1 1	AO2 4.1.1.4 4.1.1.5
09.3	any **two** from: • iron has a high(er) melting / boiling point • iron is dense(r) • iron is hard(er) • iron is strong(er) • iron is less reactive • iron has ions with different charges • iron forms coloured compounds • iron can be a catalyst	allow the converse statements for sodium allow transition metal for iron allow Group 1 metal for sodium ignore references to atomic structure ignore iron rusts allow iron is less malleable / ductile allow specific reactions showing difference in reactivity allow iron is magnetic	2	AO1 4.1.2.5 4.1.3.1 4.1.3.2

4.1.1.7 Electronic structure

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.4	carbon is more reactive (than nickel)	allow converse	1	AO1 4.4.1.2 4.4.1.3
	(so) carbon will displace / replace nickel (from nickel oxide) **or** (so) carbon will remove oxygen (from nickel oxide)	allow (so) nickel ions gain electrons allow (so) carbon transfers electrons to nickel (ions)	1	
09.5		an answer of 67.8 (%) scores **3** marks an answer of 67.8160919 (%) or correctly rounded answer to 2, 4 or more sig figs scores **2** marks an incorrect answer for one step does **not** prevent allocation of marks for subsequent steps		AO2 4.3.3.2
	(total M_r of reactants =) 87		1	
	(percentage atom economy) $= \frac{59}{87} \times 100$	allow (percentage atom economy) $= \frac{59}{\text{incorrectly calculated } M_r} \times 100$	1	
	= 67.8 (%)	allow an answer from an incorrect calculation to 3 sig figs	1	
Total			**11**	

4.1.1.7 Electronic structure

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

Question	Answers	Extra information	Mark	AO / Spec. Ref.
10.1	copper, zinc, sodium chloride solution		1	AO2 4.5.2.1
10.2	a reactant is used up	allow the reaction stops allow electrolyte / electrode / ions / metal / metal hydroxide / alkali for reactant	1	AO1 4.5.2.1
10.3	the reaction is not reversible		1	AO1 4.5.2.1
10.4	$2H_2 + O_2 \rightarrow 2H_2O$	allow fractions / multiples allow **1** mark for O_2	2	AO1 AO2 4.1.1.1 4.5.2.2

4.1.1.7 Electronic structure

MARK SCHEME – GCSE CHEMISTRY – 8462/1F – JUNE 2018

<table>
<tr><th>Question</th><th>Answers</th><th>Mark</th><th>AO / Spec. Ref.</th></tr>
<tr><td rowspan="5">10.5</td><td>Level 3: A judgement, strongly linked and logically supported by a sufficient range of correct reasons, is given.</td><td>5–6</td><td>AO3</td></tr>
<tr><td>Level 2: Some logically linked reasons are given. There may also be a simple judgement.</td><td>3–4</td><td>AO3</td></tr>
<tr><td>Level 1: Relevant points are made. This is not logically linked.</td><td>1–2</td><td>AO2</td></tr>
<tr><td>No relevant content</td><td>0</td><td></td></tr>
<tr><td>Indicative content

reasons why fuel cells could be judged as better
<table>
<tr><th>from the table</th><th>from other knowledge</th></tr>
<tr><td>• time for refuelling a fuel cell is faster than recharging
or
a fuel cell does not need to be recharged
• a fuel cell has a greater range</td><td>• hydrogen can be renewable if made by electrolysis using renewable energy
• lithium-ion batteries can catch fire
• produces only water
or
no pollutants produced
• lithium-ion batteries may release toxic chemicals on disposal
• lithium-ion batteries (eventually cannot be recharged so) have a finite life</td></tr>
</table>
reasons why the lithium-ion battery could be judged as better
<table>
<tr><th>from the table</th><th>from other knowledge</th></tr>
<tr><td>• lithium-ion uses energy more efficiently
• cost of lithium-ion car much less
• cost of recharging much less than refuelling with hydrogen</td><td>• hydrogen is often made from fossil fuels so is not renewable
• charging points are more widely available than hydrogen filling stations
• hydrogen takes up a lot of space
or
is difficult to store
• hydrogen can be highly flammable / explosive
• no emissions produced
• (catalyst in the hydrogen fuel-cell eventually becomes poisoned so) have a finite life</td></tr>
</table></td><td></td><td>4.4.3.4
4.5.2.1
4.5.2.2</td></tr>
<tr><td>Total</td><td></td><td></td><td>11</td></tr>
</table>

Answer 25 · Examiner comments

June 2018 | 8462/1F | Original Q2

Question 2 (low and standard demand)

02.1 42% of students did not realise that the number of outer shell electrons would be the same as the Group number.

02.2 19% of students were able to deduce that hydrogen bromide exists as small molecules. The majority of answers were fairly evenly split between giant covalent and ionic lattice.

02.3 59% of students were able to select F_2 as the correct formula for fluorine, with most of the others opting for F.

02.4 26% of students scored two or more marks, usually for giving the correct trend in reactivity and stating that iodine had no reactions or no changes.

Very few students could articulate their reasoning in terms of displacement. Several tried to answer in terms of reactivity of the potassium halides rather than the halogens, or were misusing the term ‘halide’ instead of ‘halogen’, making it difficult to gain any credit. Others thought that darkness of colour was an indication of reactivity, whilst some referred to melting or boiling points, which were not relevant.

Some students tried to answer in terms of ease of gaining an electron, rather than by using the information in the table as intended. Those who did so were given credit, with the most common mark being for a description of the iodine atom being larger than the chlorine atom.

02.5 74% of students were able to calculate the M_r of TiO_2 correctly. Of those who didn’t gain credit, often they didn’t recognise that there are two oxygen atoms, giving an answer of 64

02.6 This question discriminated well with 32% of students achieving both marks. Many students could convert 1.2 kg to 1200 g. However, many used the mass of reactant, 500 g, as either the numerator or denominator in the percentage yield expression. Others inverted the % yield expression or tried to subtract values.

4.1.1.7 Electronic structure

Answer 26 · Examiner comments

June 2019 | 8462/1F | Original Q1

Question 1 (low demand)

01.1 Approximately 80% of students identified both the nucleus and the electron in the diagram of an atom.

01.2 Approximately 60% of students identified the electron as having the lowest mass.

01.3 Approximately three-quarters of students correctly applied their knowledge and understanding of the link between electronic structure and group number.

01.4 Almost 60% of students scored both marks. Here, they applied their knowledge and understanding of the links between atomic structure and both atomic number and mass number.

01.5 Approximately 70% of students knew the term ‘isotope’ for the type of atom described.

01.6 Students found this multiple choice question about the link between isotope ratios and relative atomic mass difficult.

The first incorrect option was chosen more often than the correct one. This suggests that, in many cases, students did not understand how the relative atomic mass is determined from the average mass of all isotope atoms and were perhaps drawn towards the word ‘fewer’ alongside the smaller number ‘79’.

4.1.1.7 Electronic structure

Answer 27 · Examiner comments

June 2019 | 8462/1F | Original Q2

Question 2 (low demand)

02.1 Over 85% of students identified the correct formula of hydrogen peroxide.

02.2 Approximately 70% of students identified covalent bonding as the bonding in a hydrogen peroxide molecule.

02.3 Students found this question about elements as catalysts difficult. The 3 options were chosen in similar proportions.

02.4 Approximately three-quarters of students correctly identified the activation energy on the reaction profile diagram.

02.5 This question proved more difficult with approximately 45% of students being able to identify the energy of hydrogen peroxide on the reaction profile diagram.

02.6 Approximately 80% of students identified the reaction as exothermic, having been given a description.

02.7 Approximately 55% of students were able to score both marks for completing the dot and cross diagram.

The mark for the bonding electrons was scored more often than the mark for the non-bonding electrons.

4.1.1.7 Electronic structure

Answer 29 · Examiner comments — June 2022 | 8462/1F | Original Q9

Question 9 (Standard demand)

09.1 Fewer than a quarter of students recognised that element **Q** must be a non-metal because of the lack of conductivity, and therefore in section **D**. **B** and **C** were both seen more often than the correct answer, though responses were fairly evenly spread across all options.

09.2 This was slightly better answered than the previous question. However, again responses were spread across all options.

09.3 Most students described a property of either a Group 1 element or a transition element without reference to the other; this did not gain credit as a comparison was needed. Of those students who did answer in comparative terms, most knew that Group 1 metals are softer than transition metals, but there was confusion about which had lower melting points. Some students thought that one group or the other would not conduct electricity. Very few students were awarded both marks; about a sixth scored one mark.

09.4 Most students could draw the electronic structure correctly. The most common error was to draw only one electron on the outer shell.

09.5 Most students referred to electrons in their answer as required, but fewer than a third described those electrons as delocalised. Some also referred to the electrons carrying charge. However, very few articulated the idea that the electrons themselves move through the metal in a particular direction. Of those who referred to electrons moving, many implied that the movement was random. Some students incorrectly referred to electrons bumping into other atoms or electrons.

Answer 29 · Examiner comments — June 2022 | 8462/1F | Original Q9

09.6 Just over half of students correctly identified the type of bonding. The most common incorrect response was covalent.

09.7 Many students gave good descriptions of the transfer of two electrons from a magnesium atom to an oxygen atom to produce magnesium ions and oxide ions. However, some students subsequently referred to electrons being shared. About 40% of students were awarded at least one mark, with about a sixth gaining all four marks.

4.1.1.7 Electronic structure

Answer 30 · Examiner comments

June 2023 | 8462/1F | Original Q9

Question 9 (standard demand)

09.1 The description of the plum pudding model in the specification (a ball of positive charge with negative electrons embedded in it) is not well known. Very few students scored both marks and fewer than a quarter scored one. Many students believed that the positive

Answer 30 · Examiner comments

June 2023 | 8462/1F | Original Q9

charge represented a proton, or that the atom as a whole is positively charged. Some referred to a ‘sea’ or ‘cloud’ of positive charge, but these terms do not convey the shape.

09.2 Fewer than a third of students knew the correct order of discovery of the particles.

09.3 About a third of students scored both marks. Most missed the relevance of the instruction to use the period table. Many tried to work out the electron configuration, instead of simply locating element 117 in Group 7 and deducing that there must be 7 outer shell electrons.

09.4 This question was very poorly answered and did not discriminate well. Students seemed unfamiliar with the concept of the need for peer review. Most answered in terms of the rarity of the element, but this was also true for the original scientists and does not explain the time lag before acceptance.

09.5 Whilst one in ten students scored all three marks, overall, few remembered how to do this standard calculation. The four numbers in the table were manipulated in various random ways. Some students did not even use all the data. Most did not realise that the answer must lie somewhere between 6 and 7, the two mass numbers. One in five students made no attempt to answer.

Answer 31 · Examiner comments

June 2024 | 8462/1F | Original Q9

Question 9 (standard demand)

09.1 This question discriminated well, with one in six students scoring both marks and more than two fifths scoring at least one. Many students did not read the question fully and gave similarities and differences in properties rather than electronic structures. These responses gained no credit. Of those who did reference the electronic structure, most stated correctly that both have one electron on the outer shell, but were less successful in giving a correct difference. Sometimes the wrong number of shells was given for the two elements: usually 2 shells for sodium and 3 for potassium. Some stated that potassium has more outer shells than sodium. This is not correct as the phrase 'outer shell' is always taken to mean the single outermost shell.

09.2 More than a third of students could give two correct observations, and more than four fifths could give one, with bubbling, floating and a flame being the most popular. 'Gas produced' did not gain credit as it is not an observation but a deduction from the observation of bubbling. 'Fizzing' is taken to mean effervescence, so fizzing and bubbling are the same thing rather than two different observations. Credit was not given for any observation requiring extra equipment such as a thermometer or an indicator.

09.3 Only two fifths of students could give a correct colour and even fewer could express the reason in terms of the solution being alkaline; many thought it would be neutral. Some students gave purple as the colour, but gave potassium burning with a purple flame as their reason.

09.4 Nearly three fifths of students scored both marks for plotting the data, with the vast majority scoring at least one. The most commonly mis-plotted point was (86, 9.1) which was often plotted at (92, 9.1). There was no expectation that a line of best fit would be drawn.

09.5 Students were good at estimating the missing value, with nearly three quarters scoring the mark. Sometimes incorrect and inappropriate methods were seen which produced values in the allowed range, such as averaging all five values, or finding the difference between the points adjacent to the missing one. These were not given credit.

09.6 Over a third of students identified the reagents. There was an even distribution of incorrect responses.

© 2024 AQA and its licensors. All rights reserved.

Answer 31 · Examiner comments

June 2024 | 8462/1F | Original Q9

09.7 Two fifths of students identified the correct pair of trends. The most popular error was to identify that the boiling point decreases down the group.

Answer 32 · Examiner comments

June 2025 | 8462/1F | Original Q8

Question 8 (standard demand)

08.1 Only one student in six could give a correct description of where the non-metals are found, despite there being a clear statement in the specification. Many referred to the right of the table, but few referred to the top. Those who referred to specific groups were often correct but some omitted Group 0. Some referred to a 'staircase' but did not specify what it means; it should be noted that the periodic table provided does not show a staircase.

08.2 About half correctly chose the low melting point, with the three distractors all being popular. Several ticked more than one box, which automatically scores zero when the instruction is to tick **one** box.

08.3 Three fifths of students drew the structure correctly. A few tried to put in 23 electrons and a few 12. One in seven students did not attempt to answer; this is a common occurrence when a question is not accompanied by answer lines. Students do need to read all the words, so they are aware there is a task to do.

08.4 This was answered less successfully. Many stated that Mendeleev left spaces for other elements to be discovered which, while true, does not answer the question. Others answered in terms of atomic number, which was not known in Mendeleev's time. Even those who tried to answer in terms of placing elements in the 'correct' group rarely mentioned properties.

08.5 Whilst about two fifths knew that argon has a full outer shell, not many were able to go on to say that this is a stable electronic structure, or that electrons are not transferred or shared. Many simply repeated information in the question with different words, such as 'argon is a noble gas' or 'argon is stable'.

08.6 Whilst about a quarter of students knew that a different number of neutrons was involved, few stated that isotopes are <u>atoms</u> with the same number of protons or <u>atoms</u> of the same element. Some referred to 'elements' implying more than one element, which contradicts the idea of atoms of the same element. Some students implied incorrectly that an isotope is a single atom in which the number of neutrons is different to the number of protons and electrons.

08.7 Fewer than half of students could correctly estimate that the A_r must be closest to 115 as there was such a high proportion of the 115 isotopes. The most popular answer was ~114, taking a simple mean rather than a weighted mean of the mass numbers.