AQA GCSE Chemistry | 8462

SPECIFICATION POINT 4.6.2.5

The effect of changing concentration (HT only)

Past-paper questions

Tier	**Higher**
Question groups	**3**
Available paper years	**2020–2023**

Using this booklet

1. Answer the questions in order and show your working.
2. Original question numbers and marks are retained.
3. Print at actual size (100%) for measuring questions.
4. Use the companion answer booklet to check your work.

Name: _____ **Date:** _____

Whole questions are retained for context, so some parts assess other specification points. Current and legacy sources are labelled. Continue on extra paper if needed.

4.6.2.5 The effect of changing concentration (HT only)

Question 01

November 2020 | 8462/2H | Original Q10

1 0 This question is about a reversible reaction.

The reaction between solutions of iron(III) ions (Fe^{3+}) and thiocyanate ions (SCN^-) is reversible.

The ionic equation for the reaction is:

	$Fe^{3+}(aq)$	+	$SCN^-(aq)$	$\rightleftharpoons$	$FeSCN^{2+}(aq)$
Colour of solution:	yellow		colourless		red

The colour of the equilibrium mixture is orange at room temperature.

1 0 . 1 Give the name of the solvent used to dissolve the ions in this reaction.

[1 mark]

1 0 . 2 A few drops of a colourless solution containing a high concentration of thiocyanate ions (SCN^-) are added to the orange equilibrium mixture.

Explain the colour change observed.

[3 marks]

4.6.2.5 The effect of changing concentration (HT only)

Question 01 · continued

November 2020 | 8462/2H | Original Q10

1 0 . 3 A water bath is set up at a temperature above room temperature.

When a test tube containing the orange equilibrium mixture is placed in the water bath, the mixture becomes more yellow.

Explain what this shows about the energy change for the forward reaction.

[3 marks]

1 0 . 4 Explain why a change in pressure does **not** affect the colour of the equilibrium mixture.

[2 marks]

1 0 . 5 Other metal ions form coloured equilibrium mixtures with thiocyanate ions.

Which metal ion could form a coloured equilibrium mixture with thiocyanate ions?

[1 mark]

Tick (✓) **one** box.

Al^{3+} ☐

Co^{2+} ☐

Mg^{2+} ☐

Na^{+} ☐

10

4.6.2.5 The effect of changing concentration (HT only)

Question 02

June 2022 | 8462/2H | Original Q7

0 7 Ammonia is produced in the Haber process.

The raw materials for the Haber process are nitrogen and hydrogen.

The equation for the reaction is:

$$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$$

0 7 . 1 Give the sources of the nitrogen and of the hydrogen used in the Haber process.

[2 marks]

Nitrogen _____

Hydrogen _____

0 7 . 2 How does the equation for the reaction show that the atom economy of the forward reaction is 100%?

[1 mark]

_____


_____

4.6.2.5 The effect of changing concentration (HT only)

Question 02 · continued

0 7 . 3 **Figure 4** represents the Haber process.

Figure 4



Explain how the ammonia produced is separated from the unreacted nitrogen and hydrogen in **X**.

[2 marks]

Question 7 continues on the next page

4.6.2.5 The effect of changing concentration (HT only)

Question 02 · continued

The Haber process uses a temperature of 450 °C and a pressure of 200 atmospheres.

Table 6 shows the percentage yield of ammonia produced at 450 °C using different pressures.

Table 6

Pressure in atmospheres	Percentage (%) yield of ammonia
60	9
120	18
180	25
240	31
300	36
360	40
420	43

4.6.2.5 The effect of changing concentration (HT only)

Question 02 · continued

0 7 . 4 Complete **Figure 5**.


The first two points have been plotted.

You should:

- use a suitable scale for the *x*-axis
- plot the remaining data from **Table 6**
- draw a line of best fit.

[4 marks]

Figure 5



0 7 . 5 Determine the percentage yield of ammonia at 450 °C and 500 atmospheres.

Show your working on **Figure 5**.

[2 marks]

Percentage yield = _____ %

4.6.2.5 The effect of changing concentration (HT only)

Question 02 · continued

0 7 . 6 The equation for the production of ammonia in the Haber process is:

$$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$$

The forward reaction is exothermic.

The conditions used are:

- a temperature of 450 °C
- a pressure of 200 atmospheres
- the presence of an iron catalyst.

Explain why these conditions are chosen for economical production of ammonia in the Haber process.

You should include references to the rate of reaction and the position of equilibrium.

[6 marks]

17

4.6.2.5 The effect of changing concentration (HT only)

Question 03

June 2023 | 8462/2H | Original Q9

0 9 This question is about reversible reactions.

When 4.68 g of hydrated copper sulfate changes into anhydrous copper sulfate:

- 2.99 g of anhydrous copper sulfate is produced
- 1.47 kJ of energy is taken in from the surroundings.

The equation for the reversible reaction is:

hydrated copper sulfate ⇌ anhydrous copper sulfate + water

0 9 . 1 Calculate the maximum mass of water that can be produced from 11.7 g of hydrated copper sulfate.

[3 marks]

Mass = _____ g

0 9 . 2 15.0 g of anhydrous copper sulfate completely changes into hydrated copper sulfate when water is added.

Calculate the amount of energy transferred to the surroundings.

[2 marks]

Energy = _____ kJ

4.6.2.5 The effect of changing concentration (HT only)

Question 03 · continued June 2023 | 8462/2H | Original Q9

The gases nitrogen dioxide and dinitrogen tetroxide reach dynamic equilibrium in a sealed container.

The equation for the reaction is:

$$2\ NO_2(g) \rightleftharpoons N_2O_4(g)$$

nitrogen dioxide (brown) — dinitrogen tetroxide (colourless)

The forward reaction is exothermic.

0 9 . 3 What happens to the position of the equilibrium in this reaction if the temperature is increased?

[1 mark]


Tick (✓) **one** box.

- Shifts to the left ☐
- Stays the same ☐
- Shifts to the right ☐

0 9 . 4 A teacher seals a brown-coloured mixture of nitrogen dioxide and dinitrogen tetroxide in a gas syringe.

Figure 8 shows the sealed gas syringe.

Figure 8



The teacher pushes the syringe piston in.

This increases the pressure in the gas syringe.

What is the colour of the mixture when a new equilibrium position is reached?

[1 mark]

Tick (✓) **one** box.

The mixture is a darker shade of brown. ☐

The mixture is the same shade of brown. ☐

The mixture is a lighter shade of brown. ☐

Question 9 continues on the next page

4.6.2.5 The effect of changing concentration (HT only)

Question 03 · continued June 2023 | 8462/2H | Original Q9

Hydrogen iodide gas decomposes into hydrogen gas and iodine gas at high temperatures.

The equation for the reaction is:

$$2\,HI(g) \rightleftharpoons H_2(g) + I_2(g)$$

0 9 . 5 Explain the effect of increasing the pressure on the equilibrium position of this reaction.

[2 marks]

0 9 . 6 Suggest the effect of adding a catalyst on the equilibrium position of this reaction.

[1 mark]

4.6.2.5 The effect of changing concentration (HT only)

Question 03 · continued

Copper forms coloured compounds.

Hydrochloric acid is added to an aqueous solution of copper compound **A**.

The word equation for the reaction is:

copper compound **A** (blue) + hydrochloric acid ⇌ copper compound **B** (yellow) + water

0 9 . 7 The reaction mixture is green when both copper compounds are present in a solution at equilibrium.

How can the equilibrium position be shifted to make the reaction mixture more yellow?

[1 mark]

Tick (✓) **one** box.

- Add more hydrochloric acid ☐
- Add more water ☐
- Leave the reaction mixture for 30 minutes ☐

0 9 . 8 The concentrations of the substances in this reaction do **not** change at dynamic equilibrium.

Explain why.

[2 marks]

13