AQA GCSE Combined Science - Trilogy | 8464

SPECIFICATION POINT 5.4.2.5

Strong and weak acids (HT only)

Mark schemes and examiner comments

Tier	**Higher**
Question groups	**4**
Available paper years	**2020–2025**

Using this booklet

1. Booklet answer numbers match the question booklet.
2. Use the original question number within marking tables.
3. Examiner comments are included where available.

Name: _____ **Date:** _____

Where a precise marking extract is unavailable, the full source document is included in an appendix. Any missing mark scheme is explicitly identified.

5.4.2.5 Strong and weak acids (HT only)

Answer 01 · Mark scheme November 2020 | 8464/C/1H | Original Q4

04.1	ZnO (**s**) + HCl (**aq**) → $ZnCl_2$ (**aq**) + H_2O (**l**)	allow **1** mark for 2/3 correct state symbols	2	AO2 5.2.2.2 5.4.2.3 RPA8
04.2	any **one** from: • warm / heat the mixture • increase the concentration of the (hydrochloric) acid	ignore add a catalyst ignore stir ignore powder ignore add more zinc oxide do **not** accept volume / amount of (hydrochloric) acid do **not** accept increase the surface area	1	AO1 5.4.2.2 5.4.2.3 RPA8
04.3	zinc oxide remains **or** solid remains	ignore colour allow zinc oxide is added until in excess	1	AO1 5.3.2.4 5.4.2.2 5.4.2.3 RPA8
04.4	filtration / filter		1	AO1 5.4.2.2 5.4.2.3 RPA8
04.5	heat leave to crystallise / cool	do **not** accept heat to dryness allow leave to evaporate some water	1 1	AO1 5.4.2.2 5.4.2.3 RPA8

5.4.2.5 Strong and weak acids (HT only)

Answer 01 · Mark scheme

04.6	(at start) value in range 12–14 (at end) value in range 0–3	must be in this order	1 1	AO1 AO2.2 5.4.2.2 5.4.2.4
04.7	2 NaOH + H_2SO_4 → Na_2SO_4 + 2 H_2O	allow 1 mark for Na_2SO_4 **and** H_2O	2	AO2 5.1.1.1 5.4.2.2
04.8	0.10 mol/dm^3		1	AO3 5.4.2.2 5.4.2.4
Total			**12**	

5.4.2.5 Strong and weak acids (HT only)

Answer 01 · Mark scheme

November 2020 | 8464/C/1H | Original Q4

Question	Answers	Extra information	Mark	AO / Spec. Ref.

5.4.2.5 Strong and weak acids (HT only)

Answer 02 · Mark scheme

June 2022 | 8464/C/1H | Original Q6

Question 6

Question	Answers	Extra information	Mark	AO / Spec. Ref.
06.1	one shared pair in overlap	allow any combination of circles, dots, crosses or $e^{(-)}$	1	AO1 5.1.2.6 5.2.1.4
	6 non-bonding electrons in outer shell of chlorine	do **not** accept extra electron(s) on outer shell of hydrogen ignore any inner shell electrons	1	
		an answer of  scores **2** marks		

Question	Answers	Extra information	Mark	AO / Spec. Ref.
06.2	completely ionises in aqueous solution	allow completely dissociates in aqueous solution	1	AO1 5.4.2.5

5.4.2.5 Strong and weak acids (HT only)

Answer 02 · Mark scheme

June 2022 | 8464/C/1H | Original Q6

Question	Answers	Extra information	Mark	AO / Spec. Ref.
06.3	fizzing / effervescence **or** magnesium disappears		1	AO3 5.4.2.5
	at a greater rate with a strong acid	allow converse with weak acid allow for **2** marks strong acid has a greater temperature increase	1	

Question	Answers	Extra information	Mark	AO / Spec. Ref.
06.4	(pH) decreases	allow pH is lower	1	AO1
	by (a unit of) 2		1	AO2 5.4.2.5

5.4.2.5 Strong and weak acids (HT only)

Answer 02 · Mark scheme

June 2022 | 8464/C/1H | Original Q6

Question	Answers	Extra information	Mark	AO / Spec. Ref.
06.5	(bonds broken = (4 × 413) + C=C + 431 =) 2083 + **C=C**		1	AO2 5.5.1.3
	(bonds made = (346 + 339 + (5 × 413) =) 2750		1	
	(energy released = bonds made – bonds broken =) 56 = 2750 – [2083 + C=C]	allow correct use of incorrect value(s) from step 1 and / or step 2	1	
	(C=C) = 611 (kJ/mol)		1	

Total Question 6		11

5.4.2.5 Strong and weak acids (HT only)

Answer 03 · Mark scheme

Question 4

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.1	completely ionised in aqueous solution	allow completely ionised (when dissolved) in water allow fully dissociated in aqueous solution	1	AO1 5.4.2.5

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.2	low concentration of (nitric) acid dissolved in water	allow low concentration of solute / solid / gas dissolved in water	1	AO1 5.4.2.5

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.3	decreases by a factor of 100		1	AO2 5.4.2.5

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.4	$H^+ + OH^- \rightarrow H_2O$	ignore state symbols	1	AO1 5.4.2.4

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.5	1 : 2		1	AO2 5.3.2.3

5.4.2.5 Strong and weak acids (HT only)

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.6	red		1	AO3 5.3.2.4 5.4.2.2 RPA10

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.7	(for each mass of magnesium carbonate) repeat the experiment (discard anomalous results) **and** calculate the mean		1	AO3 5.5.1.1 RPA10
	measure the highest temperature		1	

Total Question 4	**8**

5.4.2.5 Strong and weak acids (HT only)

Answer 04 · Mark scheme

June 2025 | 8464/C/1H | Original Q5

Question 5

Question	Answers	Extra information	Mark	AO / Spec. Ref.
05.1	(hydrogen chloride) g / (g) **and** (water) l / (l) **and** (hydrochloric acid) aq / (aq)		1	AO2 5.2.2.2

Question	Answers	Extra information	Mark	AO / Spec. Ref.
05.2	3.16×10^{-4} (mol/dm^3)		1	AO2 5.4.2.5

Question	Answers	Extra information	Mark	AO / Spec. Ref.
05.3		allow any combination of x, O, $e^{(-)}$, ● for electrons		AO2 5.2.1.1 5.2.1.4
	1 pair of electrons in both overlaps		1	
	4 non-bonding electrons on oxygen atom **and** 6 non-bonding electrons on chlorine atom	do **not** accept non-bonding electrons on hydrogen	1	
		an answer of  scores **2** marks		

5.4.2.5 Strong and weak acids (HT only)

Answer 04 · Mark scheme

Question	Answers	Extra information	Mark	AO / Spec. Ref.
05.4	a weak acid partially ionises in an aqueous solution (so) produces a lower concentration of hydrogen ions		1 1	AO1 5.4.2.4 5.4.2.5

Question	Answers	Extra information	Mark	AO / Spec. Ref.
05.5	(bonds broken = (4 × 413) + 243 =) 1895 (bonds made = (3 × 413) + 346 + 432 =) 2017 (energy change = 1895 − 2017 =) (−) 122 (kJ/mol) **alternative approach** (bonds broken = 413 + 243 =) 656 (1) (bonds made = 346 + 432 =) 778 (1) (energy change = 656 − 778 =) (−) 122 (kJ/mol) (1)	allow correct use of an incorrectly determined value for bonds broken and / or bonds made allow correct use of an incorrectly determined value for bonds broken and / or bonds made	1 1 1	AO2 5.5.1.3

Total Question 5	9

5.4.2.5 Strong and weak acids (HT only)

Answer 02 · Examiner comments

June 2022 | 8464/C/1H | Original Q6

Question 6 (Standard, standard/high and high demand)

06.1 Two third of students correctly drew a dot and cross diagram for hydrogen chloride. The most common mark was for the two shared electrons in the overlap between the atoms. Many did not show that there are six unshared electrons.

06.2 Less than 10% were able to state what is meant by the term strong acid. Many omitted ‘in aqueous solution’.

06.3 40% of students gained 1 mark usually for referring to an observation when magnesium reacts with a strong acid and a weak acid. The second mark was for a comparison of the rates of reaction between magnesium and the two acids, with few students being able to develop their response to give this.

06.4 Nearly 40% of students gained 1 mark usually for stating that the pH decreases when the concentration was increased. However very few recognised that pH was related to the power of 10, so when the concentration of hydrochloric acid is increased by a factor of 100 then the pH will decrease by 2.

06.5 Many students were good at calculating the energy changes for bonds broken and bonds made, however then went on to simply subtract these two numbers without taking into account the C=C bond energy. Many students kept their working to a minimum which often did not help them score. This question discriminated well with better responses showing clear working where the energy released was equated to the difference between the bonds made and the bonds broken.

5.4.2.5 Strong and weak acids (HT only)

Answer 03 · Examiner comments

Question 4 (standard & standard/high demand)

04.1 Many students stated that strong acids completely ionise but omitted to include that this happens in water or in aqueous solution. 10% of students gave the correct definition.

04.2 Less than 5% of students gave a correct explanation for 'dilute aqueous solution'. Very few students made reference to being dissolved in water or used the term (low) concentration.

Common incorrect responses made reference to reducing the concentration of the acid or stating that there would be a small volume of acid per unit volume or small volume of acid in aqueous solution. A number of students stated that a high concentration of substance/acid had been dissolved.

04.3 This question proved challenging. To deduce the change in the hydrogen ion concentration students need to know that as the pH decreases by one unit, the hydrogen ion concentration of the solution increases by a factor of 10. A third of students gained the mark with approximately equal numbers choosing the alternative responses.

04.4 Nearly one-fifth of students were able to recall an ionic equation to represent the neutralisation between an acid and an alkali. This question had the highest not attempted rate at 12.5%, which was nearly double that of any other question. A common response was 'acid + alkali → salt' suggesting that students were not familiar with the ions present in acid or alkaline solutions or that this involved neutralisation to produce water.

04.5 Most students were able to deduce the correct ratio of moles from the equation. The most common distractor was 2:1 where the correct ratio was reversed.

04.6 Just over half of students recognised that universal indicator would turn red in the acidic solution produced when magnesium carbonate and excess nitric acid were reacted. Green was the most common distractor which was not correct as excess acid was used in the reaction.

© 2024 AQA and its licensors. All rights reserved.

5.4.2.5 Strong and weak acids (HT only)

Answer 03 · Examiner comments June 2024 | 8464/C/1H | Original Q4

04.7 This question required improvements to the method, so references to changing the volume or concentration of solution were insufficient. Many students wrote that the experiment should be repeated but to be creditworthy they needed to include that the mean should be calculated. 'Measuring until the temperature stopped increasing' was taken as equivalent to 'measure the highest temperature'. However, 'measuring until the temperature is constant' is insufficient as this may not be the highest temperature or the mixture may have cooled to room temperature.

Just under a quarter of students gained 1 mark and 1% gained 2 marks.


0 4 . 7 Give **two** improvements to the **method** to produce more accurate results.

Do **not** refer to improvements to the apparatus in your answer.

[2 marks]

1 Repeat more than 5 times

2 Have more nitric acid. ~~Measure the temperature of the acid before~~

This student did not gain any credit: repeating the experiment on its own is insufficient as the mean is not calculated (after anomalous results are discarded).

© 2024 AQA and its licensors. All rights reserved.

Answer 04 · Examiner comments

Question 5 (Standard/High and High demand)

05.1 The state symbols for hydrogen chloride gas and water were well known but aqueous for hydrochloric acid was not. Some students mixed up l and aq. Most students correctly use lower case letters.

05.2 Few students gained this high demand mark. Those students who gained the mark often wrote down a few words next to the values given in the question, e.g. 'add 1, divide by 10' or 'concentration increases by factor of 10, pH decreases by 1'. Numerous students multiplied by 10 rather than dividing and many struggled to evaluate data in standard form.

05.3 More than half of students gained two marks for completing the dot and cross diagram. When 1 mark was gained it was usually for 1 pair of electrons in both overlaps with extra non-bonding electrons on hydrogen or incorrect numbers of non-bonding electrons on the oxygen atom and/or the chlorine atom.

05.4 Very few gained both marks. Students had to explain why a weak acid has a higher pH than a strong acid of the same concentration. Most answers were in terms of a pH comparison. Lots of students did not achieve MP1 by not using the full definition including in aqueous solution. For MP2 many students said there were less H^+ ions omitting reference to concentration or number per unit volume.

05.5 Many students were able to calculate the energy changes for bonds broken and for bonds made and then subtracted the latter value from the former. The value of bonds broken was less than bonds made so the overall energy change was a negative number. Those students who miscounted the bonds broken or made, could not gain MP1 and/or MP2 but were still able to access MP3 by evaluating their own figures correctly.

© 2024 AQA and its licensors. All rights reserved.

5.4.2.5 Strong and weak acids (HT only)

Answer 04 · Examiner comments June 2025 | 8464/C/1H | Original Q5

0 5 . 5 Hydrogen chloride is produced in the reaction between methane and chlorine.

Figure 6 represents the equation for the reaction.

Figure 6



Table 4 shows the bond energies.

Table 4

Bond	C — H	Cl — Cl	C — Cl	H — Cl
Bond energy in kJ/mol	413	243	346	432

Calculate the overall energy change for the reaction.

[3 marks]

reactants	Products
413 x 4 = 1652	1 x 346 = 346
243 x 2 = 486	413 x 3 = 1239
total = 1652 + 486 =	432 x 1 = 432
2138 − 2017 = 121	total = 2346 + 1239 + 432 = 2017

Overall energy change = 121 kJ/mol

This student has broken 2× Cl—Cl bonds (not 1×) for the reactants so is not awarded MP1. They then correctly subtract the correct value for bond making (2017) so MP2 and MP3 are credited.

© 2024 AQA and its licensors. All rights reserved.