Chhetri Academy

chhetriacademy.com

GCSE MATHS

Number

Edexcel 1MA1 | Subtopic collection Topic 03

Fractions and mixed numbers

Higher tier

Answers and examiner comments

48	2007-2025	Higher
question groups	available papers	tier

Using this booklet

1. Answer numbers match the companion question booklet.
2. Follow the original question number on marking tables.
3. Examiner comments are included where available.

Name _____ **Date** _____

Some entries use full source marking material in an appendix. Any unavailable mark schemes are identified beside the relevant answer.

Original question 18

The full source mark scheme begins on page 70. Use the original paper code, tier and question number shown above.

Original question 1

No matching mark scheme was available in the supplied archive for this question.

Question		Working	Answer	Mark	Notes
23		$x = 0.363636\ldots$ $100x = 36.363636\ldots$ $99x = 36$ $x = \frac{36}{99} = \frac{4}{11}$ or $10000x = 3636.36\ldots$ $9999x = 3636$ $x = \frac{3636}{9999} = \frac{4}{11}$ or $9900x = 3600$ etc	Proof	3	M1 for $100x = 36.363636\ldots$ or $10000x = 3636.3636..$ M1 (dep) for subtraction of both sides A1 for $\frac{36}{99} = \frac{4}{11}$ from correct proof. OR M1 starts long/short division of 11 into 4, set out correctly, with 0.36 seen on the top of the bus stop (oe) with a remainder of 7 M1(dep) Remainder of 4 after the remainder of 7 seen in correct place A1 At least 2 remainders of 4 and one of 7 seen in the correct place and with a statement that the decimal will recur with a cycle length 2 because the remainders have a cycle length 2.

2	(a)		28.38461538	2	B2 for 28.3846…. (B1 for 107.01 or 3.77 or 28.38(…) or $28\frac{5}{13}$ oe seen)
	(b)		30	1	B1ft for 30 or for answer >1sf in (a) rounded to 1 sf

6	(a)	$\frac{2}{3.95}$	0.5063(29113...)	2	B2 for 0.5063 or better. (B1 for 0.5 or 0.50 or 0.506 or 0.51 or 3.95 or the fraction $\frac{40}{79}$ seen)
	(b)		0.51	1	B1 ft for 0.51 from their answer to part (a) which is written to two or more decimal places.

Question	Working	Answer	Mark	Notes
9		2.42927(0474)	2	B2 for 2.42927 or better (B1 for 19.56 or 8.0518 seen or 2.43 or 2.429 or 2.4292 or 2.4293 or digits 242927... or $\frac{97800}{40259}$ seen)

Question		Working	Answer	Mark	Notes
16	(a)	$\frac{2}{3} \times \frac{6}{5}$	$\frac{4}{5}$	3	M1 for $\frac{2}{3} \times \frac{6}{5}$ M1 for $\frac{2\times6}{3\times5}$ or 12/15 oe A1 cao
	(b)	$(2-1) + \frac{5}{15} - \frac{6}{15}$ or $\frac{35}{15} - \frac{21}{15}$ Or $\times$ | 1 | 3 2 | | 6 5 | 5 | 15 $\times$ | 7 | 3 7 | | 21 5 | 35 | 15	$\frac{14}{15}$	3	M1 for attempt to find a common denominator or sight of $\frac{5}{15}$ or $\frac{6}{15}$ or $\frac{35}{15}$ or $\frac{21}{15}$ oe or fully correct table A1 for sight of $\frac{5}{15} - \frac{6}{15}$ or $\frac{35}{15} - \frac{21}{15}$ oe A1 for $\frac{14}{15}$ oe Alternative M1 for 0.33(3…) or 0.4 OR 2.33(3…) or 1.4 A1 for 0.33(3…) – 0.4 OR 2.33(3…) –1.4 A1 for 0.93(recurring)

13	(a)	$\sqrt{\frac{920-364.5661765}{0.046}}$ $= \sqrt{12074.64834...}$	109.88(47047)	2	B2 for 109.88(….. (B1 for 12074 seen …. or 555.43seen …or 109 to 110)
	(b)		110	1	B1 f.t. provided answer to (a) is more than 3 s.f.

1380_4H

Question		Working	Answer	Mark	Notes
14		. 0.25 = 0.2555… 0.25555…×10 = 2.5555… 0.25555…×100 = 25.5555… 25.5555…– 2.5555… = 23 **Alternative Method** x = 0.25555 … 10x = 2.5555 … 100x = 25.5555… 90x = 23	$\frac{23}{90}$	3	M1 for 0.25555... or 0.2 + 0.05555… M1 for two correct recurring decimals that, when subtracted, would result a terminating decimal, and attempting the subtraction or $\frac{2.3}{9}$ or $\frac{253}{990}$ A1 cao **Alternative Method** M1 for x = 0.25555… M1 for two correct recurring decimals that, when subtracted, would result a terminating decimal, e.g. 10x = 2.5555..., 100x = 25.5555… and attempting the subtraction or 9x = 2.3 or 90x = 23 or 900x = 230 or $\frac{2.3}{9}$ or $\frac{253}{990}$ A1 cao SC: B1 for fraction with denominator 90 if M0

Original question 24

The full source mark scheme begins on page 122. Use the original paper code, tier and question number shown above.

2	(a)	$\frac{546.7}{12.5} =$	43.736	2	B2 for 43.736 (B1 for 546.7 **or** $\frac{5467}{10}$ **or** $\frac{5467}{125}$ **or** 12.5 **or** $\frac{25}{2}$ or 43.7 **or** 43.8 **or** 43.73 **or** 43.74 **or** 40 **or** 44)
	(b)		40	1	B1 for 40 **or** ft from their answer to (a) provided (a) is written to 2 or more significant figures

1MA0_2H

Question		Working	Answer	Mark	Notes
1		$\frac{\sqrt{20.4}}{6.2\times0.48}=\frac{4.5166359}{2.976}$	1.5176(868)	2	B2 for 1.5176… (B1 for sight of 4.51(66359..) **or** 4.52 **or** 2.97(6) **or** 2.98 **or** 1.51 **or** 1.52 **or** 1.518 **or** 1.517 **or** 1.5177**or** $\frac{\sqrt{510}}{5}$)

Question		Working	Answer	Mark	Notes
17		$\sqrt{\frac{2.73\ldots}{0.732\ldots}}$	1.931851…	2	M1 for 2.73… or 0.732…or 3.73…or 1.931 or 1.932 or 1.93 or $(1+\sqrt{3})$ or $(\sqrt{3}-1)$ or $(2+\sqrt{3})$ **or** 1.65... **or** 0.855... A1 for 1.9318(5…) SC: B1 for 2.5127(17...)

15		$x = 0.7505050...$ $10x = 7.505050...$ $1000x = 750.505050...$ $990x = 743$ OR $100x = 75.0505050...$ $99x = 74.3$	$\frac{743}{990}$	3	M1 for 0.75050(50….) or 0.7 + 0.050(5050….) M1 (dep) for two recurring decimals that, when subtracted, leave a terminating decimal A1 for $\frac{743}{990}$

PAPER: 5MB2H_01

3			1.3852..	2	M1 for 6.4 or 4.62 or $\frac{320}{231}$ A1 for 1.3852(81385)
*4			4 rolls	4	M1 for $\pi \times 2.4$ M1 for $(\pi \times 2.4) \div 2$ or 7.5 to 7.541 M1 for or 3.75 or 3.76... or 3.77... or (2, 4,) 6 , 8 C1 for a clear statement that 4 (rolls) are needed

PAPER: 5MB3H_01

PAPER: 1MA0_2H

Question	Working	Answer	Mark	Notes
1 (a)		18.75	2	M1 for 84 **or** 4.48 **or** $\frac{112}{25}$ **or** 18.7 **or** 18.8 **or** 19 **or** 20 **or** $\frac{75}{4}$ A1 cao
(b)		20	1	B1 for 20 **or** ft from their answer to (a) provided (a) is written to 2 or more significant figures

2		1263	4	M1 for a correct method to find 65% (= 5473) of the customers M1 for a correct method to find $\frac{1}{5}$ (= 1684) of the customers M1 (dep on M2) for a correct method to find the remaining number of customers A1 cao OR M1 for a correct method of adding 65% and $\frac{1}{5}$ when both correctly written as percentages (= 85%) or decimals (= 0.85) or fractions (= $\frac{85}{100}$ oe) M1 ft for a correct method to find the percentage or decimal or fraction of the customers (= 7157) M1 (dep on M2) for a correct method to find the remaining number of customers A1 cao OR M1 for a correct method of adding 65% and $\frac{1}{5}$ when both correctly percentages (= 85%) or decimals (= 0.85) or fractions (= $\frac{85}{100}$ oe) M1 ft for a correct method to find the remaining percentage (= 15%) or

PAPER: 5MB3H_01				
Question	**Working**	**Answer**	**Mark**	**Notes**
				decimal (= 0.15) or fraction ($\frac{15}{100}$ oe) of the customers M1 (dep on M2) for a correct method to find the remaining number of customers A1 cao

2		1263	4	M1 for a correct method to find 65% (= 5473) of the customers M1 for a correct method to find $\frac{1}{5}$ (= 1684) of the customers M1 (dep on M2) for a correct method to find the remaining number of customers A1 cao OR M1 for a correct method of adding 65% and $\frac{1}{5}$ when both correctly written as percentages (= 85%) or decimals (= 0.85) or fractions (= $\frac{85}{100}$ oe) M1 ft for a correct method to find the percentage or decimal or fraction of the customers (= 7157) M1 (dep on M2) for a correct method to find the remaining number of customers A1 cao OR M1 for a correct method of adding 65% and $\frac{1}{5}$ when both correctly percentages (= 85%) or decimals (= 0.85) or fractions (= $\frac{85}{100}$ oe) M1 ft for a correct method to find the remaining percentage (= 15%) or

PAPER: 5MB3H_01				
Question	**Working**	**Answer**	**Mark**	**Notes**
				decimal (= 0.15) or fraction ($\frac{15}{100}$ oe) of the customers M1 (dep on M2) for a correct method to find the remaining number of customers A1 cao

PAPER: 5MB2H_01

Question		Working	Answer	Mark	Notes
15		$x = 0.15555\ldots$ $10x = 1.5555\ldots$ $9x = 1.4$ $x = \frac{1.4}{9} = \frac{14}{90}$ OR $x = 0.1 + y$ where $y = 0.0555\ldots$ $10y = 0.5555\ldots$ $100y = 5.5555..$ $90y = 5$ so $y = 5/90$ $x = 0.1+5/90= 1/10+ 5/90$	$\frac{7}{45}$	3	M1 for $0.155(5\ldots)$ or $0.1+ 0.055(5\ldots)$; This can be implied in subsequent working. M1 for 2 correct recurring decimals which when subtracted will leave an integer or a terminating decimal number with a correct fraction for their 2 recurring decimals A1 for $\frac{7}{45}$ [**SC**: B1 for an answer of $\frac{15}{99}$ oe, with or without working]

8			$14\frac{2}{3}$	3	M1 for method to write fractions as improper fractions with one correct M1 (dep on M1) for multiplying numerators and denominators A1 cao SC: B2 for $\frac{220}{15}$ oe

Question		Working	Answer	Mark	Notes
23			$\frac{14}{3}$	5	M1 for correct substitution into a volume formula for a cylinder or a cone, eg. $\frac{1}{3}\times\pi\times3^2\times4$ (= 12π) or $\pi\times3^2\times(6-4)$ (= 18π) or $\pi\times3^2\times h$ (= $9\pi h$) or $\pi\times3^2\times(h-2)$ M1 for method to find volume after 5 hours, eg. "12π" + "18π" (= 30π) M1 (dep on M1) for use of a correct ratio, eg. "30π" $\times\frac{9}{5}$ (= 54π) or "30π" $\times\frac{4}{5}$ (= 24π) M1 for deriving an equation in h, eg. "54π" = "$9\pi h$" + "12π" A1 for $\frac{14}{3}$ or equivalent fraction

PAPER: 1MA0_2H

Question		Working	Answer	Mark	Notes
3	(a)		4.58006(9567)	2	M1 for 1.83 or 8.381(527307) or 4.6 or 4.58 or 4.580 or 4.5801 A1 for 4.58006(9567)
	(b)		4.5801	1	B1 ft provided at least 5 decimal places in (a)

Original question 10

The full source mark scheme begins on page 142. Use the original paper code, tier and question number shown above.

18	(a)		$2\frac{4}{5}$	3	M1 for writing as improper fractions eg $\frac{6}{5}$ or $\frac{7}{3}$ M1 (dep) for multiplying improper fractions eg $\frac{6\times7}{5\times3}$ or $\frac{14}{5}$ oe A1 cao
	(b)		$\frac{4}{5}$	3	M1 for finding two correct fractions with a common denominator eg $\frac{7}{15}-\frac{10}{15}$ or $\frac{21-30}{45}$ M1 (dep) for complete and correct method eg $1-\frac{3}{15}$ or $\frac{37}{15}-\frac{25}{15}$ or $\frac{111-75}{45}$ oe A1 for $\frac{4}{5}$ oe

PAPER: 1MA0_1H

13			$11\frac{2}{3}$	3	M1 for writing as improper fractions eg $\frac{25}{3}$ or $\frac{7}{5}$ M1 (dep) for multiplying improper fractions eg $\frac{25\times7}{3\times5}$ $(=\frac{175}{15})$ or $\frac{5\times7}{3\times1}$ $(=\frac{35}{3})$ A1 cao

PAPER: 5MB2H/01

16		$\frac{\sqrt{3}}{5}+\frac{2}{\sqrt{3}}\times\frac{\sqrt{3}}{\sqrt{3}}$ $=\frac{3\sqrt{3}}{15}+\frac{10\sqrt{3}}{15}$; $\frac{\sqrt{3}\sqrt{3}+10}{3\sqrt{3}}=\frac{13\sqrt{3}}{5\sqrt{3}\sqrt{3}}$	$\frac{13}{15}$	3	M1 for rationalising a denominator M1 for finding same denominator (dep M1 or with √3) A1 oe Accept $\frac{13}{15}\sqrt{3}$

5		1.4091(…)	2	B2 for 1.4091(…) (B1 for 2.1025 or 1.492 or 2.397…or 2.398)

20		$\frac{59}{330}$	3	M1 for $100x = 17.87878787\ldots$ or $1000x = 178.7878787\ldots$ **and** $10x = 1.7878787$ M1 (dep) for subtraction, $100x - x$ or $1000x - 10x$ or $\frac{17.7}{99}$ or $\frac{177}{990}$ seen A1 working leading to given fraction

Question	Working	Answer	Mark	Notes
14		$\frac{1}{3}$	P1	process to solve the problem e.g. $\frac{3}{10}\times\frac{4}{9}(=\frac{12}{90}=\frac{2}{15})$ OR finds the number of white circles for their chosen number OR for 9 : 21 (or a multiple of 9 : 21)
			P1	second step of the process e.g. $\frac{7}{10}\times\frac{2}{7}(=\frac{14}{70}=\frac{2}{10}=\frac{1}{5})$ OR finds the number of black circles for their chosen number OR for a multiple of 2 : 5 where the ratio parts sum to "21"
			P1	for complete process e.g. $"\frac{2}{15}"+"\frac{1}{5}"\left(=\frac{4}{30}+\frac{6}{30}\right)$ OR finds the total number of circles for their chosen number OR for 3 ratios that could be used to solve the problem eg 9 : 21 with 4 : 5 with 6 : 15
			A1	for $\frac{1}{3}$ oe

Question	Working	Answer	Mark	Notes
4		Daisy is wrong	P1	for process to find area of any relevant circle ie $\pi\times4^2$ (=16π), $\pi\times7^2$ (=49π), $\pi\times10^2$ (=100π) **or** 7^2 and 4^2
		(supported)	P1	for completed method to find shaded area eg "$\pi\times7^2$" – "$\pi\times4^2$" (=33π) **or** use of radii eg $7^2 - 4^2$ (=33)
			A1	for 2 comparable figures, eg 33π and 100π **or** 33 and 100 **or** 103 to 103.7 and 314 to 314.2 **or** 103 to 103.7 and 104.6 to 104.8
			C1	statement eg No because it should be $\frac{33}{100}$ and their accurate figures Allow use of π = 3 or better

Question	Working	Answer	Mark	Notes
17 (a)	1.5, 6, 10.2, 7.2, 1.2	Histogram drawn	C1	for 2 correct bars of different widths or at least 3 correct frequency densities.
			C1	for all bars in correct proportions or 4 correct bars with axes scaled and labelled.
			C1	for fully correct histogram with axes scaled and labelled.
(b)		$\frac{123}{150}$	M1	for a method to find number of students in interval, eg $30 + 51 + 36 + \frac{1}{3} \times 18$ (= 123) or $150 - 15 - \frac{2}{3} \times 18$ (= 123)
			A1	for $\frac{123}{150}$ oe or 0.82 or 82%

5	(a)		0.625	B1	cao
	(b)		$9.75 \leq x < 9.85$	B2	for $9.75 \leq x < 9.85$
				[B1	for 9.75 or 9.85 (or $9.84\dot{9}$)]

Question	Answer	Mark	Mark scheme	Additional guidance
1 (a)	$\frac{95}{28}$	M1	for a method to add using common denominators with at least one fraction correct (matching numerator with common denominator) eg $\frac{60}{28}+\frac{35}{28}$ **or** $(2)\frac{4}{28}+(1)\frac{7}{28}$	Use of decimals gets no credit unless it leads to a correct fraction
		A1	$\frac{95}{28}$ oe eg $3\frac{11}{28}$	
(b)	$1\frac{3}{5}$	M1	for $\frac{6}{5}\times\frac{4}{3}$ **or** $\frac{24}{20}\div\frac{15}{20}$ **or** $\frac{8}{5}$ oe eg $1\frac{9}{15}$	Use of decimals gets no credit unless it leads to a correct fraction
		A1	cao	

9	$5\frac{3}{5}$	M1 M1 A1	for writing as improper fractions with at least one correct, eg $\frac{7}{2}\times\frac{8}{5}$ oe (dep) for multiplying improper fractions, eg $\frac{"56"}{"10"}$ **or** $5\frac{6}{10}$ **or** $\frac{28}{5}$ oe cao	

3	$2\frac{1}{3}$	M1 M1 A1	for either $\frac{7}{4}$ oe or $\frac{4}{3}$ oe for method to find the product, eg. $\frac{7\times4}{4\times3}$ or $\frac{21\times16}{12\times12}$ oe or for $\frac{28}{12}$ or $\frac{7}{3}$ oe for $2\frac{1}{3}$ or an equivalent mixed number	

Question	Answer	Mark	Mark scheme	Additional guidance
15	$\frac{414}{990}$	M1	for (x =) 0.41818… or (10x =) $4.\dot{1}\dot{8}$ or 4.1818… or (100x =) $41.\dot{8}\dot{1}$ or 41.818… or (1000x =) $418.\dot{1}\dot{8}$ or 418.18…	
		M1	for using two recurring decimals with a terminating decimal difference, eg. (1000x – 10x =) $418.\dot{1}\dot{8} - 4.\dot{1}\dot{8}$ or 418.18… − 4.1818… (= 414)	Accept (100x – x =) $41.\dot{8}\dot{1} - 0.4\dot{1}\dot{8}$ or 41.818... – 0.41818... (= 41.4)
		A1	for $\frac{414}{990}$ oe, eg $\frac{23}{55}$	$\frac{41.4}{99}$ must be simplified to gain the accuracy mark

Paper: 1MA1/3H

Question	Answer	Mark	Mark scheme	Additional guidance
14	$\frac{13}{20}$	P1	for finding the fraction who chose either soup ($\frac{2}{5}$ oe) or chose prawns ($\frac{3}{5}$ oe) **or** for process to share any number in the ratio 2 : 3 eg 100 ÷ (2 + 3) × 2 (=40)	Starting number 100 Soup : Prawn 40:60
		P1	for a process that could lead to the proportion who chose lasagne or curry for either starter, eg sharing 40% (soup) in the ratio 5 : 3 **or** sharing 60% (prawns) in the ratio 1 : 5 **or** $\frac{2}{5} \times \frac{5}{8}$ **or** $\frac{2}{5} \times \frac{3}{8}$ **or** $\frac{3}{5} \times \frac{1}{6}$ **or** $\frac{3}{5} \times \frac{5}{6}$ **or** for continuing the process with their starting number to find the number who chose lasagne or curry for either starter	L:C L:C 25:15 10:50
		P1	for a complete process to find the proportion who chose curry for **both** starters, eg($\frac{2}{5} \times \frac{3}{8}$) + ($\frac{3}{5} \times \frac{5}{6}$) **or** to find the number who chose curry for **both** starter for their starting number	15 + 50 =65 and $\frac{15+50}{100}$
		A1	$\frac{13}{20}$ or equivalent fraction	

Paper: 1MA1/3H

2	Shown	M1 M1 C1	for conversion to improper fractions eg. $\frac{7}{3}$ **or** $\frac{15}{4}$ (dep) for method to multiply fractions, eg. $\frac{7\times15}{3\times4}\left(=\frac{105}{12}\right)$ **or** $\frac{28\times45}{12\times12}\left(=\frac{1260}{144}\right)$ oe for complete working showing each stage as far as $\frac{35}{4}$ or $8\frac{9}{12}$	Need not be shown with operators

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
23	$\frac{27}{56}$	P1	for $\frac{3}{8}$ **and** $\frac{7}{9}$ **OR** uses a total of 72 cards and shows a process to find the number of cards with a black shape **or** the number of cards with a triangle, eg $72 \div 8 \times 3$ (=27) **or** $72 \div 9 \times 7$ (=56)	72 or any multiple of 72 Could be seen in a ratio, eg 27 : 45 or 16 : 56
		P1	for process shown to divide fractions $\frac{3}{8} \div \frac{7}{9}$ **or** $\frac{3}{8} \times \frac{9}{7}$ **OR** for $\frac{3}{8} \times \frac{9}{9}$ $(= \frac{27}{72})$ **and** $\frac{7}{9} \times \frac{8}{8} (= \frac{56}{72})$ **OR** uses a total of 72 cards and shows a process to find the number of cards with a black shape **and** the number of cards with a triangle, eg $72 \div 8 \times 3$ (=27) **and** $72 \div 9 \times 7$ (=56)	Accept the division shown as $\frac{\frac{3}{8}}{\frac{7}{9}}$ Could be seen in ratios, eg 27 : 45 and 16 : 56
		A1	for $\frac{27}{56}$ or any other equivalent fraction	Answer of 27 : 56 gets P2A0

Modifications to the mark scheme for Modified Large Print (MLP) papers: 1MA1 1H

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 1MA1/1H			
Question		**Modification**	**Mark scheme notes**
3		Wording added "Look at the diagram for Question 3 in the Diagram Book." The wording "The diagram shows four graphs" removed and replaced by "It shows four graphs labelled graph A, graph B, graph C and graph D." Diagrams enlarged. Graph lines made thicker. Open headed arrows. Headings moved above the graph.	Standard mark scheme
4		Wording added "Look at the diagram for Question 4 in the Diagram Book." The wording "The diagram shows four triangles" removed and replaced by "It shows four triangles." Diagram enlarged. Headings moved above the diagrams. Angles moved outside of the angle arcs and the angle arcs made smaller. Triangles straightened up so a 10 cm side lies horizontally. Braille only: Description added of the triangles.	Standard mark scheme
6		The wording "ADC is a triangle" removed. Diagram enlarged. Wording added "Look at the diagram for Question 6 in the Diagram Book. It shows the triangle ADC." Angles moved outside of the angles arcs and the angle arcs made smaller.	Standard mark scheme

PAPER: 1MA1/1H			
Question		**Modification**	**Mark scheme notes**
7		Wording added "Look at the table and the diagram for Question 7 in the Diagram Book." Wording "This stem and leaf diagram shows…" removed and replaced with "The stem and leaf diagram shows…" The wording "…of a group of Year 9 girls" removed and replaced by "…of a group of girls in Year 9". The wording "…of a group of 15 Year 9 boys" removed and replaced by "…of a group of 15 boys in Year 9". Table enlarged and kept on the same page as the stem and leaf diagram. Diagram enlarged and a tracking line added. Key moved above the diagram.	Standard mark scheme
8		Model provided for all candidates with a base added to represent the horizontal floor. Wording added "Look at the diagram for Question 8 in the Diagram Book. You may be provided with a model." The wording "The diagram shows a prism…" removed and replaced by "The diagram and the model show a prism…". Diagram enlarged. Dashed lines to be made thicker and longer. The pressure formula to be kept in the Question Paper and moved to the left of the diagram in the Diagram Book.	Standard mark scheme
9		The wording "Write these numbers in order of size" removed and replaced by "Write these four numbers in order of size."	Standard mark scheme
10		The letter a changed to w. The letter b changed to x. The letter c changed to y.	Standard mark scheme except for the letter changes indicated.

PAPER: 1MA1/1H				
Question		**Modification**	**Mark scheme notes**	
12	(a)	The value of the 350-400 interval changed from '20' to '25'. The value of the 450-500 interval changed from '10' to '5'. Wording added "Look at the table for Question 12(a) in the Diagram Book." Table enlarged. The wording "The table gives information…" removed and replaced by "It gives information…"	Standard mark scheme	
	(b)	Wording added "Look at the diagram for Question 12(b) in the Diagram Book." The wording "On the grid opposite…" removed and replaced by "On the grid,…" Diagram enlarged. Right axis labelled. Small squares removed. The axes labels moved to the top of the vertical axis and to the left of the horizontal axis.	Standard mark scheme	
	(c)	The wording "60% of this group of people have a weekly wage of £360 or less" removed and replaced by "75% of this group of people have a weekly wage of £375 or less" to make the question accessible.	M1 for 75 ÷ 100 × 80 (=60) oe	M1 reading value from graph at wage = 375 (=48)
			M1 reading value from graph at cf = 60 (=400)	M1 for "48" ÷ 80 × 100 (=60(%)) **or** for 75 ÷ 100 × 80 (=60)
			Standard mark scheme (NO) with 60 and "400" **or** with "48" and 60(%) **or** with "48" and 60	
17		The letter *f* changed to *p*, *d* changed to *m*.	Standard mark scheme but note the changes to the letters.	
21		The wording "DEF is a triangle" replaced by "Look at the diagram for Question 21 in the Diagram Book. It shows the triangle DEF." Diagram enlarged.	Standard mark scheme	

PAPER: 1MA1/1H			
Question		**Modification**	**Mark scheme notes**
22		Wording added "Look at the diagram for Question 22 in the Diagram Book." The wording "The diagram shows…" removed and replaced by "It shows…" Diagram enlarged. Open headed arrows. Shading changed to dotty shading. Angle moved outside of the angle arc the angle arc made smaller. The shapes labelled 'shape A' and 'shape B.' The labels "(3x-1) cm" and "2x cm" added to the bottom of the diagram. The label "(5x-1) cm" added to the top of the diagram.	Standard mark scheme
23		Wording added: "Look at the information for Question 23 in the Diagram Book. It shows the four types of cards in a game." Diagram enlarged. Black cards changed to dotty shading. All reference of 'black' changed to 'shaded'. Headings added above the cards e.g. "shaded circle".	Standard mark scheme

3	$1\frac{8}{15}$	M2 (M1 A1	for a complete method, eg $4-2+\frac{3}{15}-\frac{10}{15}$ condoning error with one numerator **or** for $\frac{21}{5}-\frac{8}{3}=\frac{63}{15}-\frac{40}{15}\left(=\frac{23}{15}\right)$ with no more than one error for finding two fractions with a correct common denominator, with at least one correct corresponding numerator, eg $\frac{3}{15}, \frac{10}{15}$ **or** for converting both to improper fractions, eg $\frac{21}{5}, \frac{8}{3}$) $1\frac{8}{15}$ oe	At least one improper fraction must be correct Any equivalents must be a mixed number

Paper: 1MA1/1H

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
13	Explanation	C1	explanation **Acceptable examples** he should have used 100(x) rather than 10(x) he should have used 1000x and 10x Ted's working does not eliminate the recurring decimals the recurring numbers after the decimal point have to be in the same sequence he should have multiplied by 100 to subtract easier after the decimal point he should have multiplied by 100 because two numbers are recurring **Not acceptable examples** it is not correct the method is not complete he should have used 1000x he should have multiplied by 100 he should have multiplied by 100 and then done 100x – 10x to give 43/90	

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
12	$\frac{116}{990}$	M1	for ($x =$) 0.11717... **or** ($10x =$) $1.\dot{1}\dot{7}$ or 1.1717... **or** ($100x =$) $11.\dot{7}\dot{1}$ or 11.7171... **or** ($1000x =$) $117.\dot{1}\dot{7}$ or 117.1717...	
		M1	(dep M1) for a method using two recurring decimals that leads to a terminating decimal difference, using correct multiples of x eg ($1000x - 10x =$) $117.\dot{1}\dot{7} - 1.\dot{1}\dot{7}$ ($= 116$) or 117.1717... – 1.1717... ($= 116$)	Accept ($100x - x =$) $11.\dot{7}\dot{1} - 0.1\dot{1}\dot{7}$ or 11.7171... – 0.11717... ($= 11.6$)
		A1	for $\frac{116}{990}$ oe, eg $\frac{58}{495}$	$\frac{11.6}{99}$ must be written in the form $\frac{a}{b}$ where a and b are integers to gain the accuracy mark

Paper: 1MA1/1H

2	(a)	$3\frac{17}{20}$	M1	for finding two fractions with a correct common denominator (multiple of 20), with at least one correct corresponding numerator, eg $\frac{12}{20}$, $\frac{5}{20}$ or $\frac{32}{20}$, $\frac{45}{20}$	May be from $\frac{3}{5}$ and $\frac{1}{4}$ or from $\frac{8}{5}$ and $\frac{9}{4}$
			A1	for $3\frac{17}{20}$ or an equivalent mixed number SC B1 for an answer of 3.85 if M0 scored	
	(b)	shown	M1	for $\frac{8}{3}\times\frac{1}{6}$ oe or $\frac{4}{9}\times\frac{6}{1}$ oe or $\frac{8}{3}\times\frac{9}{4}$ oe	
			A1	for unsimplified fraction which could lead to $\frac{4}{9}$, eg $\frac{8}{18}$ or for $\frac{4}{3}\times\frac{1}{3}$ or $\frac{24}{9}\div 6$ **or** for unsimplified fraction which could lead to $2\frac{2}{3}$, eg $\frac{24}{9}$ **or** for unsimplified fraction which could lead to 6, eg $\frac{72}{12}$	

Paper: 1MA1/1H

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
6 (a)(i)	2 : 6 : 5	P1	for process to compare ratios, eg $a : b = 2 : 6$ or $b : c = 3 : 2.5$	Could use 3 or any common multiple of 3 and 6
		A1	for 2 : 6 : 5 oe	
(ii)	$\frac{2}{13}$	M1	for process to find fraction, eg $\frac{[2]}{[2+6+5]}$ or for $\frac{a}{a+b+c}$	
		A1	for $\frac{2}{13}$ oe or ft (a)(i)	
(b)	1 : 10	P1	for process to express all numbers in terms of one number, eg $p = 5 \times 2m$ (= $10m$) or $m = \frac{n}{2}$ **or** for $2m = \frac{p}{5}$ **or** for assigning values in the ratio given, eg $m = 1$, $n = 2$, $p = 10$ **or** for $n : m : p = 2 : 1 : 10$ oe **or** 10 : 1 oe	
		A1	for 1 : 10 oe	

Paper: 1MA1/1H

2	$4\frac{7}{8}$	M2 (M1 A1	for a complete method, eg $7-2+\frac{3}{8}-\frac{4}{8}$ condoning error with one numerator **or** for $\frac{59}{8}-\frac{5}{2}=\frac{59}{8}-\frac{20}{8}$ $(=\frac{39}{8})$ with no more than one error **OR** for an answer of 4.875 for finding two fractions with a correct common denominator, with at least one correct corresponding numerator, eg $\frac{3}{8}, \frac{4}{8}$ **or** for converting both to improper fractions, eg $\frac{59}{8}, \frac{5}{2}$ **OR** for 7.375 – 2.5) for $4\frac{7}{8}$ oe eg $4\frac{14}{16}$	At least one improper fraction must be correct Both decimals must be correct Any equivalents must be a mixed number

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
13 (a)	Histogram drawn	B3	for fully correct histogram with axes scaled and labelled	Relative heights 2, 3.5, 4.4, 2, 0.8
		(B2	for 4 correct bars **or** for frequency ÷ class width for all 5 frequencies **and** 2 correct bars of different widths)	
		(B1	for 2 correct bars of different widths **or** for frequency ÷ class width for at least 3 frequencies)	
(b)	$\frac{67}{150}$	M1	for a method to find number of people in interval eg $\frac{1}{2} \times 70 + 22 + \frac{1}{3} \times 30$ (= 67) **or** $150 - 20 - 8 - \frac{1}{2} \times 70 - \frac{2}{3} \times 30$ (= 67) **or** 10 × [3.5] + 5 × [4.4] + 5 × [2] ft their histogram	Where [3.5], [4.4] and [2] are the heights of the bars of their histogram
		A1	for $\frac{67}{150}$ or ft their histogram	

Question	Answer	Mark	Mark scheme	Additional guidance
3 (a)	$2\frac{1}{3}$	M1	for a method to subtract by writing both fractions with a common denominator with at least one correct numerator, eg $3\frac{3}{6}-1\frac{1}{6}$ **or** $\frac{3}{6}-\frac{1}{6}\left(=\frac{2}{6}\right)$ **or** $\frac{21}{6}-\frac{7}{6}\left(=\frac{14}{6}\right)$ **or** $\frac{42}{12}-\frac{14}{12}\left(=\frac{28}{12}\right)$	
		A1	for $2\frac{1}{3}$ or an equivalent mixed number	Do not isw incorrect further work from correct equivalent mixed number
(b)	Shown	M1	for conversion to improper fractions, eg $\frac{21}{4}$ **or** $\frac{7}{3}$ **or** $\frac{9}{4}$	
		M1	(dep) for method to divide by a fraction, eg $\frac{21}{4}\times\frac{3}{7}$ or $\frac{63}{12}\div\frac{28}{12}$	
		C1	for complete work showing each stage as far as $\frac{9}{4}$ **or** $2\frac{7}{28}$	Must see an intermediate step, eg $\frac{63}{28}$ must be seen and then cancelled or correct cancelling seen before the multiplication

Paper: 1MA1/1H

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
13	$\frac{5}{33}$	M1	for $10x = 2.\dot{2}$ or $2.22\ldots$ or $(10x - x =)\ 2.\dot{2} - 0.\dot{2}\ (= 2)$ or $2.22\ldots - 0.22\ldots\ (= 2)$ or $\frac{2}{9}$ oe fraction	eg $\frac{20}{90}$, $\frac{22}{99}$
		M1	for a method using two recurring decimals that leads to a terminating decimal difference, using correct multiples of y eg $(1000y - 10y =)\ 681.\dot{8}\dot{1} - 6.\dot{8}\dot{1}\ (= 675)$ or $681.81\ldots - 6.81\ldots\ (= 675)$ or $\frac{675}{990}$ or $(100y - y) = 68.\dot{1}\dot{8} - 0.6\dot{8}\dot{1}\ (= 67.5)$ or $68.181\ldots - 0.681\ldots\ (= 67.5)$ or $\frac{67.5}{99}$	
		A1	for $(x =)\frac{2}{9}$ oe **and** $(y =)\frac{675}{990}$ oe	Accept $(y =)\frac{67.5}{99}$
		M1	for "$\frac{2}{9}$" × "$\frac{675}{990}$"	Award 4 marks for an answer equivalent to $\frac{5}{33}$, eg $\frac{15}{99}$, $\frac{135}{891}$, $\frac{1350}{8910}$ unless from incorrect working
		A1	cao	

Paper: 1MA1/1H

edexcel

Edexcel GCSE

Mathematics A 1387

June 2007

advancing learning, changing lives

Mark Scheme

Edexcel GCSE

Mathematics A 1387

Table Of Contents

Notes on Marking Principles	3 - 4
5521 / 01 Mark Scheme	5 - 11
5521 / 02 Mark Scheme	12 – 17
5523 / 03 Mark Scheme	18 – 27
5523 / 04 Mark Scheme	28 – 35
5525 / 05 Mark Scheme	36 – 43
5525 / 06 Mark Scheme	44 - 52

NOTES ON MARKING PRINCIPLES

1 **Types of mark**

M marks: method marks
A marks: accuracy marks
B marks: unconditional accuracy marks (independent of M marks)

2 Abbreviations

cao – correct answer only
ft – follow through
isw – ignore subsequent working
SC: special case
oe – or equivalent (and appropriate)
dep – dependent
indep - independent

3 No working

If no working is shown then correct answers normally score full marks
If no working is shown then incorrect (even though nearly correct) answers score no marks.

4 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.
If there is no answer on the answer line then check the working for an obvious answer.

5 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

6 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

7 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

8 Linear equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.



9 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

Remember: if you are having difficulty making a decision on how you should mark a candidate response contact your Team Leader for advice, or send the item to review.


Paper 5521_01					
No		Working	Answer	Mark	Notes
1		Cat |||| /||| 8 Dog |||| | 6 Fish || 2 Hamster |||| 4	8, 6, 2, 4	3	M1 for attempt to tally or one frequency correct in either column A1 for 1 frequency correct or all tallies correct in correct column A1 for all frequencies correct (accept if /20)
2			27, 35, 42, 67, 118	1	B1 cao
3	(a)		Diameter drawn	1	B1 for a diameter
	(b)		Right angle marked	1	B1 *R* marked correctly
	(c)		Rectangle drawn	1	B1 for a rectangle
4	(a)		40	1	B1 cao
	(b)		50	1	B1 cao
	(c)		3 full loaves 4 full loaves + 1 half loaf	2	B1 for 3 full loaves B1 for 4 full loaves + 1 half loaf
5	(a)		7252	1	B1 cao
	(b)		Three thousand and eighty six	1	B1 accept 3 thousand and eighty six (condone 0 hundred)
	(c)		4600	1	B1 accept 4600
	(d)		200	1	B1 for 200 or 2 hundred or 100 or hundred
6	(i)		Cube	2	B1 for 'cube' (accept 'cuboid') ignore spelling
	(ii)		Cylinder		B1 for 'cylinder' ignore spelling
7		5×100	500	2	B2 for 490 or 500 or 510 (B1 for either 5 or 5.0 or 100 seen)

Paper 5521_01					
No		**Working**	**Answer**	**Mark**	**Notes**
8	(a)		8.4 cm or 84 mm	2	B2 allow ±2mm (B1 for 8.4 or 84, B1 for appropriate unit)
	(b)		37°	1	B1 allow ±2°
9	(a)		Carbon black	1	B1 accept 'black carbon' accept 26%
	(b)		0.1(0)	1	B1 cao
	(c)		0.04	1	B1 cao
	(d)	$\frac{26}{100}$	$\frac{13}{50}$	2	M1 for $\frac{26}{100}$ A1 cao
10	(a)		97	1	B1 cao
	(b)		London Reading	1	B1 cao
	(c)	41 + 57 + 58	156	3	M1 for two of 41, 57, 58 M1(dep) for '41' + '57' + '58' A1 cao
11	(a)		3	1	B1 cao allow ± 0.2
	(b)		–5	1	B1 cao allow ±0.2

Paper 5521_01					
No		**Working**	**Answer**	**Mark**	**Notes**
12	(a)		Centimetres (cm) miles litres (*l*)	3	B3 (B1 for each correct answer) accept abbreviations
	(b)		300	1	B1 cao
	(c)	'1500>1400' or '1.5>1.4'	Reason	1	B1 for No and '1500>1400' or '1.5>1.4'
13	(a)	Height of bars 12, 8, 6 lines drawn between points	Bars drawn	2	B2 for 3 bars correctly drawn (B1 for 2 bars correct)
	(b)		July and August	1	B1 oe
	(c)	24 – 4	20	1	B1 (Accept '4 to 24' oe)
	(d)	The temperatures are rising	Temp's rising oe	1	B1 for reason
14	(a) (i)		$(-4, 3)$	2	B1 cao
	(ii)		$(2, -1)$		B1 cao
	(b) (i)	*D* marked at $(-4, -1)$	Point marked on grid	2	B1 for point marked at $(-4, -1)$ cao
	(ii)		$(-4, -1)$		B1 ft
15	(a)			1	B1 cao
	(b)			1	B1 cao

Paper 5521_01				
No	**Working**	**Answer**	**Mark**	**Notes**
16 (a)		(Pat +) reason	1	B1 correct comment (Pat may be implied)
(b)	21 ÷ 3	7	1	B1 cao
17 (i)		S marked at 1	3	B1 for S within ½ cm of 1
(ii)		P marked at 0		B1 for P marked at 0 cao
(iii)		Q marked at "1/3"		B1 for Q marked at 1/3 ± 1 cm use overlay
18	6 + 6 + 3 or $2\frac{1}{2}\times 6$	15	3	M1 for realizing 6 glasses in one bottle M1 for realizing 3 glasses in ½ a bottle A1 cao (M2 for attempt to find $2\frac{1}{2}\times 6$) oe
19	15 and 16 parts shaded Alternative 1 $\frac{3}{4}=0.75,\ \frac{4}{5}=0.8$ Alternative 2 $\frac{3}{4}=\frac{15}{20},\ \frac{4}{5}=\frac{16}{20}$	$\frac{4}{5}$ + reason	3	M1 for correctly shading 15 parts for 3/4 M1 for correctly shading 16 parts for 4/5 A1 (dependent on M2) for selection of 4/5 Alternative 1 M1 for $\frac{3}{4}=0.75$ M1 for $\frac{4}{5}=0.8$ A1 (dep on M2) for selection of 0.8 or $\frac{4}{5}$ or $\frac{16}{20}$ Alternative 2 M1 for $\frac{3}{4}=\frac{15}{20}$ M1 for $\frac{4}{5}=\frac{16}{20}$ A1(dep on M2) for selection of $\frac{4}{5}$ or $\frac{16}{20}$

Paper 5521_01				
No	**Working**	**Answer**	**Mark**	**Notes**
20 (a)		−5 (−3) (−1) 1 3 5	2	B2 cao (B1 for any 2 or 3 correct)
(b)	Points plotted	Line	2	B2 for line from (−1, −5) to (4, 5) (B1 ft for plotting at least 5 "points")
21	Shapes shaded on grid	6 tessellating shapes	2	B2 for fully correct with 5 or more additional shapes, no gaps (B1 for 4 shapes tessellating with at least one shape inverted, with or without the given shape ignore extras)
22 (a)		$\frac{7}{20}$	1	B1 for $\frac{7}{20}$ oe
(b)	7 + 2 (or 20 – 11) are not lime flavour	$\frac{9}{20}$	1	B1 for $\frac{9}{20}$
(c)		0	1	B1 for 0, zero or nought ($\frac{0}{20}$ gets B0)
23 (a)		$4a$	1	B1 accept $4\times a$, $a \times 4$, $a4$
(b)		$12b$	1	B1 accept $12\times b$, $b \times 12$, $b12$
(c)		$2a + 6b$	2	B2 cao (B1 for $2a$ or $6b$ seen)
(d)		$x(x-6)$	2	B2 cao (B1 for $x(ax+b)$ where a, b are numbers not equal to 0 or $x - 6$ seen on its own, or as part of an expression)
24 (a)		40	1	B1 cao
(b)		45	1	B1 for 42 – 48 accept ¾ hour
(c)	40×2 or $\frac{40}{30}\times 60$ or $40\div\frac{1}{2}$	80	2	M1 for 40×2 or $\frac{40}{30}$ or $40\div\frac{1}{2}$ A1 cao NB $\frac{40}{45}\times 60$ gets M0 A0

Paper 5521_01					
No		**Working**	**Answer**	**Mark**	**Notes**
25	(a)		4560	1	B1 cao
	(b)		45.6	1	B1 cao
	(c)		2.4	1	B1 cao
26		$-7 - 3 = -10$ $2 \times -10 = -20$ $-20 \div 4$	-5	3	M1 for substitution of 2 and -7 into $p(q - 3)$ or sight of -20 or $-14 - 6$ M1dep for '-20' $\div 4$ A1 cao B1 SC for sight of -10 if M0 awarded
27	(a)		Reflection in y-axis	1	B1 cao
	(b)		Rotation by half turn about (0,0)	2	B2 cao (B1 for half turn not about (0,0).)
	(c)		Enlargement Scale factor 3 Centre (0,0)	3	B1 for 'enlargement' B1 for "scale factor 3" or 3 seen B1 for 'centre (0,0)
28	(a)		Reason	1	B1 for 'The frequencies are nearly equal' oe
	(b)	$1\times26+2\times26+3\times23+4\times25 = 247$ $247/100$	2.47	3	M1 for fx (attempting at least 2 relevant products) M1 for $\sum fx \div 100$ A1 2.47 cao

Paper 5521_01				
No	**Working**	**Answer**	**Mark**	**Notes**
29	5 × 5 × 6	150	4	M1 for attempt at 1 division (e.g. 40 ÷ 8), may be implied by marks or number on one edge of diagram or by 5 or 6 seen M1 for attempt at 3 divisions (40÷8, 40÷8, 60÷10), may be implied by marks or numbers on diagram or by 5,5 and 6 seen. M1 (dep on 1st M1) for "5" × "5" × "6" A1 cao Alternatively M1 for 40 × 40 × 60 **or** 8 × 8 × 10 **or** 96000 or 640 seen M1 for 40 × 40 × 60 **and** 8 × 8 × 10 or 96000 **and** 640 seen M1 (dep on 1st M1) for "(40 × 40 × 60)" ÷ "(8 × 8 × 10)" A1 cao SC:B1 for dividing area of one carton face by area of corresponding box face if M0

Paper 5521_02					
No		Working	Answer	Mark	Notes
1	(a)	Draw diagram.	Diagram	1	B1 cao
	(b)		13,16	1	B1 cao
	(c)		31	1	B1 cao
2	(a)		8	1	B1 cao
	(b)		14	2	B2 for 14 (B1 for 13 or 15)
	(c)		16	2	B2 for 16 (B1 for 15, 17 or 8)
3	(a)		4,7 drawn	2	B2 for car height 4 and bus height 7, (B1 for one correct)
	(b)		6	1	B1 cao
	(c)		Walk	1	B1
	(d)		27	1	B1 cao
4	(a)		40	1	B1 for 40–41 inclusive
	(b)		12	1	B1 for 11.5 – 12.5 inclusive
5	(a)		Row complete	2	B2 for 1+..+11; 36 (B1 for one of the 2 cells complete)
	(b)		Square	1	B1 "square"
6		One line of symmetry		1	B1 within 2mm of centre of base / 2mm of vertex

Paper 5521_02				
No	**Working**	**Answer**	**Mark**	**Notes**
7 (a) (i)		09 06	2	B1 (accept 9 06 oe)
(ii)		39		B1 cao
(b)		06 55	1	B1 (accept 6 55 oe)
(c)		2h 6min	1	B1 cao
(d)		15 min	2	M1 for 0906 – 0645 – "(c)" or 0906 – 0645 − 2hr 6min or 2hr 21min − "(c)" or 2hr 21min − 2hr 6 min or 141 − 126 or 20 − 5 A1 cao SC: B1 for 55 or 75 or 93 seen
8 (i)		8,10,12,20 or 30	3	B1 at least one of 8, 10, 12, 20, 30 (no extras)
(ii)		8,12 or 20		B1 at least one of 8,12, 20 (no extras)
(iii)		3 or 5		B1 3 or 5 or both (no extras)
9 (a)		C or G	1	B1 at least one of C or G (no extras)
(b)		A and F	1	B1 cao
(c)		2	1	B1 (accept –2)
10 (a)		$\frac{7}{10}$	1	B1 7/10 oe
(b)(i)		4 squares	2	B1 4 squares shaded
(ii)		80%		B1 80% or ft from unshaded part (no ft from 0% or 100%)
(c)(i)		2.5	2	B1 2.4–2.6 inclusive
(ii)		1.7		B1 1.6–1.8 inclusive
11 (a)		2	1	B1 for 2 or –2
(b)		14	1	B1 for 14 or –14

Paper 5521_02					
No		**Working**	**Answer**	**Mark**	**Notes**
12		2 × 8.50 = 17.00 3 × 4.50 = 13.50 Total = 30.50 50.00 – 30.50	19.5(0)(*p*)	3	M1 for adding 5 correct values or 2 × 8.50 + 3 × 4.50 (ignore units) or 30.5(0) or 3050 seen M1 dep for 50 − "30.50" (ignore units) (OR M1for adding at least 1 adult ticket and at least 1 child ticket **and** subtracting from 50) A1 cao SC: B1 for 24 or 37 or 2400 or 3700 seen
13	(a)		Hexagon	1	B1
	(b)(i)		120	2	B1 cao
	(ii)		Str line		B1 reference to a (straight) line and 180°
	(c)		Obtuse	1	B1 Accept "interior"
14	(a)	2, 2, 3, 3, 3, 4, 4, 4, 5, 6	3.5	2	M1 ordering the numbers (condone 1 error or omission) A1 cao
	(b)	36 ÷ 10	3.6	2	M1 sum of numbers ÷ 10 A1 cao SC B1 for 3r 6
	(c)	6 – 2	4	1	B1 cao
15	(a)		Paul	1	B1 cao
	(b)	36 ÷ 2 oe	18	1	B1 cao
	(c)	60/360 =	1/6	2	M1 60/360 oe A1 cao

Paper 5521_02					
No		**Working**	**Answer**	**Mark**	**Notes**
16		4.7 ÷ 5.9 = 0.796610169	0.7966..	2	B2 for 0.7966 or better (B1 for 0.8, 0.80, 0.79, 0.796, 0.797 or digits 59 seen
17		$6x - 7 + 7 = 38 + 7$ $6x = 45$	7.5	2	M1 $6x = 45$ or +7 both sides A1 7.5 oe; accept 45/6 oe
18		**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	3	B3 all six entries correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
19	(a)	900 × 1.70 =	1530	2	M1 900 × 1.7(0) or digits 153(0) seen A1 cao
	(b)	160 ÷ 1.70 =	£94.12 or £94.11	2	M1 160 ÷ 1.7(0) or digits 941(....) seen A1 cao
20	(a)(i)	180 – 54 (=126) "126" ÷ 2	63	3	M1 for (180–54) ÷ 2 A1 cao
	(ii)		Reason		B1 (indep) angles in triangle add to 180 OR equal angles in isosceles triangle OR equal angles and 2 sides the same (B0 if any incorrect reasoning given eg parallel, equilateral triangle)
	(b)	180 – "x"	117	1	B1 117 or ft 180 – "x" if $x < 90$
21	(a)	3 × 35 + 50	155	2	M1 for 3 × 35 + 50 or digits 155 seen A1 cao
	(b)	260 – 50 = 210 210 ÷ 35 =	6	3	M1 for 260–50 or 210 seen. M1 for "260-50" ÷ 35 or 210÷35 A1 cao SC B1 for starting at a number between 100 and 170 and adding at least two 35's and showing a total between 230 and 290 OR For adding at least three 35's, perhaps with other numbers, and showing a total between 180 and 240 (or between 230 and 290 if 50 is included in the sum)

Paper 5521_02				
No	**Working**	**Answer**	**Mark**	**Notes**
(c)		P=35h + 50	3	B3 for P=35h+50 or P=35×h+50 oe (B2 for correct RHS or P=h + 50 ×35 or P=35h+k where k is numerical oe) (B1 for P = some other linear expression in h, OR h + 50 ×35 OR 35h seen) NB: P=h scores no marks; ignore £ signs. SC B2 for $h = \frac{P-50}{35}$
22 (a)		Elevation	2	B2 for 4 vertical squares. Accept 4 by 1 rectangle. (B1 for 4 vertical squares with one square added or one parallelogram added at the top, or 3 vertical squares, or 4 horizontal squares)
(b)		Plan	2	B2 for 2 adjacent squares, vertical or horizontal. Accept 2 by 1 rectangle. (B1 for 3 adjacent horizontal or vertical squares or a rectangle with sides in the ratio 2:1)
23 (i) (ii) (iii)		5 9 6	3	B1 cao B1 cao B1 cao
24	$45.00 + 45.00 \times \frac{15}{100} =$ $45.00 + 6.75 =$	51.75	3	M2 for $45.00 + 45.00 \times \frac{15}{100}$ oe or 45.00×1.15 oe OR $45.00 + 6.75$ OR complete method or 5175 seen. (M1 for $45.00 \times \frac{15}{100}$ oe OR 6.75 seen OR 675 seen OR correct method for calculating 15% of 45) A1 cao SC Award B2 for an answer of 38.25

Paper 5521_02				
No	**Working**	**Answer**	**Mark**	**Notes**
25 (a)		Points	1	B1 all three points ±1 full square
(b)		Negative	1	B1 Negative (ignore additional descriptors unless contradictory)
(c)		lobf	1	B1 A single straight line drawn to cross between (5,30), (5,40) and (40,0), (40,15); accept freehand if considered to be straight.
(d)(i)		18–25	2	B1 18g–25g inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient.
(ii)		30–40		B1 30–40 min inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient
26		300 3 75 150	3	B3 for 4 correct answers (B2 for 2 or 3 correct answers) (B1 for 1 correct answer)
27	π × 0.65	2.04–2.05	2	M1 for π × 0.65 or 3.14 x 0.65 or 3.142 x 0.65 A1 for 2.04–2.05 SC Award B1 for 2.0 seen (not 2)
28	5 miles = 8 km 70mph÷5×8 = 112 km/h OR 120km/h ÷ 8 × 5 = 75 mph Faster than 70 mph	70mph (Great Britain) (112 km)	3	M1 5 miles = 8 km; OR 70 mph is about 100 km/h OR 1km=0.6(25) miles OR 1mile=1.6km oe M1 70 ÷ 5 × 8 (=112) or 120 ÷ 8 × 5 (=75) A1 (dep on at least M1) GB or 70 mph Refer to both answer line and working. NB GB or 70 mph without working scores 0 marks

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
1		Cuboid drawn	2	B2 for correct isometric drawing in any orientation (ignore points 'behind', mark 7 vertices only); accept lines drawn near to dots as long as there is no ambiguity. (B1 for one of the three faces drawn correctly **or** for an isometric drawing of any cuboid)
2	Different makes of car Tally Frequency	Make of car Tally Frequency	3	B1 for make of car or list of at least 3 different makes B1 for tally or tally marks B1 for frequency or totals
3		6 tessellating shapes	2	B2 for fully correct with 5 or more additional shapes, no gaps (B1 for 4 or more shapes tessellating, with at least one shape inverted, with or without the given shape, ignore extras)
4	24.90 ÷ 3 or 8.30 24.90 – '8.30' or 2 × 8.30	16.6(0)	3	M1 for 24.90 ÷ 3 or 8.30 M1 (dep) for 24.90 – “8.30” or 2 × “8.30” A1 for 16.60 or 16.6

Paper 5523_03



No	Working	Answer	Mark	Notes
5	315 × 24 24 × 315 1260 6300 7560 7200 7560 **3 1 5** 0 6 | 0 2 | 1 0 **2** 1 2 | 0 4 | 2 0 **4** 7 5 6 0 **300 10 5** 6000 | 200 | 100 **20** 1200 | 40 | 20 **4** 6000+200+100+1200+40+20 = 7560 **3 0.1 0.05** 60 | 2 | 1 **20** 12 | 0.4 | 0.2 **4** 60 + 2 + 1 + 12 + 0.4 + 0.2 = 75.6	75.6(0)	3	M1 for a complete method with relative place value correct. Condone 1 multiplication error, addition not necessary. **OR** M1 for a complete grid with not more than 1 multiplication error, addition not necessary. **OR** M1 for sight of a complete partitioning method, condone 1 multiplication error, final addition not necessary. A1 for 7560 or digits 756(0) A1 (dep on M1, but not previous A1) for correct placement of decimal point.

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
6	15 and 16 parts shaded Alternative 1 $\frac{3}{4} = 0.75$ or 75%, $\frac{4}{5} = 0.8$ or 80% Alternative 2 $\frac{3}{4} = \frac{15}{20}$, $\frac{4}{5} = \frac{16}{20}$	$\frac{4}{5}$ + reason	3	M1 for shading 15 parts for $\frac{3}{4}$ M1 for shading 16 parts for $\frac{4}{5}$ A1 (dep on M2) for selection of $\frac{4}{5}$ with correct shading Alternative 1 M1 for $\frac{3}{4} = 0.75$ or 75% M1 for $\frac{4}{5} = 0.8$ or 80% A1 (dep on M2) for selection of 0.8 or 80% or $\frac{4}{5}$ with correct decimals or percentages Alternative 2 M1 for $\frac{3}{4} = \frac{15}{20}$ oe M1 for $\frac{4}{5} = \frac{16}{20}$ oe A1 (dep on M2) for selection of $\frac{4}{5}$ or $\frac{16}{20}$ with equivalent fractions

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
7		$5 \times 5 \times 6$	150	4	M1 for attempt at 1 division (e.g. 40 ÷ 8), may be implied by marks or number on one edge of diagram or by 5 or 6 seen M1 for attempt at 3 divisions (40÷8, 40÷8, 60÷10), may be implied by marks or numbers on diagram or by 5,5 and 6 seen. M1 (dep on 1st M1) for "5" × "5" × "6" A1 cao Alternatively M1 for 40 × 40 × 60 **or** 8 × 8 × 10 **or** 96000 or 640 seen M1 for 40 × 40 × 60 **and** 8 × 8 × 10 or 96000 **and** 640 seen M1 (dep on 1st M1) for "(40 × 40 × 60)" ÷ "(8 × 8 × 10)" A1 cao SC:B1 for dividing area of one carton face by area of corresponding box face if M0
8	(a)		$\frac{7}{20}$	1	B1 for $\frac{7}{20}$ oe
	(b)	7 + 2 (or 20 – 11) are not lime flavour	$\frac{9}{20}$	1	B1 for $\frac{9}{20}$ oe
	(c)		0	1	B1 for 0, zero or nought ($\frac{0}{20}$ gets B0)
9	(a)		$80x$	1	B1 for $80x$ (accept $80 \times x$, $x80$, $x \times 80$) seen
	(b)		$95y$	1	B1 for $95y$ (accept $95 \times y$, $y95$, $y \times 95$) seen
	(c)		$80x + 95y$	2	M1ft for adding "$80x$" and "$95y$" (algebraic expressions only) A1 for $80x + 95y$

Paper 5523_03

No	Working	Answer	Mark	Notes
10 (a)		40	1	B1 cao
(b)		45	1	B1 for 42 to 48 (accept 3/4 hour)
(c)	40×2 or $\frac{40}{30}\times60$ or $40\div\frac{1}{2}$	80	2	M1 for 40×2 or $\frac{40}{30}$ or $40\div\frac{1}{2}$ A1 cao NB $\frac{40}{45}\times 60$ gets M0 A0
11 (a)	$3\times3-4\times2$ or $9-8$	1	2	M1 for substitution of 3 and 2 into expression or 9 and 8 seen A1 cao
(b)	$-7-3=-10$ $2\times-10=-20$ $-20\div4$	-5	3	M1 for substitution of 2 and -7 into $p(q-3)$ or sight of -20 or $-14-6$ M1 (dep) for "-20" $\div 4$ A1 cao SC: B1 for -10 seen if M0
12 (a)	6 | 8 9 7 7 | 8 5 9 6 3 8 | 1 3 1 7 1 9 | 0 1	6 | 7 8 9 7 | 3 5 6 8 9 8 | 1 1 1 3 7 9 | 0 1	3	M1 for unordered diagram (condone one error) A1 cao B1 for key (eg 6 | 7 = 67)
(b)(i)		Explanation	2	B1 for '(order numbers and) select middle value' oe
(ii)		79		B1 cao

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
13	(a)		Reflection in y-axis	1	B1 for triangle with vertices at (-1, 1) (-3, 1) and (-1,4)
	(b)		Rotation by half turn about (0, 0)	2	B2 for triangle with vertices (-1, -1) (-3, -1) and (-1, -4) (B1 for half turn not about (0,0))
	(c)		Enlargement Scale factor 3 Centre (0, 0)	3	B1 for 'enlargement' B1 for 'scale factor 3' or 3 seen B1 for 'centre (0,0)' B0 for any combination of transformations
14	(a)		4560	1	B1 cao
	(b)		45.6	1	B1 cao
	(c)		2.4	1	B1 cao
15	(a)	$4a-2a+5b+b$	$2a+6b$	2	B2 cao (B1 for $2a$ or $6b$ seen)
	(b)		$x(x-6)$	2	B2 cao (B1 for $x(ax+b)$ where a, b are numbers not equal to zero or $x-6$ seen on its own, or part of an expression)
	(c)		$3x-2x^3$	2	B2 cao (B1 for $3x$ or $2x^3$)
	(d)		$4x(3y+x)$	2	B2 cao (B1 for $2(6xy+2x^2)$ or $4(3xy+x^2)$ or $x(12y+4x)$ or $2x(6y+2x)$ or $4x(\quad)$)

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
16	(a)	$1 - (0.2 + 0.3 + 0.1)$	0.4	2	M1 for $1 - (0.2 + 0.3 + 0.1)$ A1 for 0.4 oe, accept $\frac{0.4}{1}$
	(b)	0.2×200	40	2	M1 for 0.2×200 A1 cao NB $\frac{40}{200}$ is M1 A0, 40 out of 200 is M1 A1
17	(a) (i)	$180 - 2 \times 25$	130	3	M1 for $180 - 2 \times 25$ A1 cao
	(ii)		Reason		B1 for mentioning isosceles and equal (or base) angles or equal sides and equal (or base) angles
	(b)	$180 - 95$	85	1	B1 cao
18	(a) (i)		7^5	3	B1 cao
	(ii)		7^4		B2 cao (B1 for sight of 7^5 or 7^{2+3} or 7×7^3 or $7^1 \times 7^3$ or $7^2 \times 7^2$ or 7^{2+3-1})
	(b)		$\frac{1}{2}$	1	B1 for $\frac{1}{2}$ or 0.5 or 2^{-1}
19	(a)		3×10^7	1	B1 cao
	(b)		0.002	1	B1 cao

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
20		Box plot	2	3 aspects: 1st aspect – vertical line for median 2nd aspect – box using correct quartiles 3rd aspect – whiskers (could be single line) drawn with correct end points B2 for fully correct box plot (B1 for 1 aspect)
21 (a)	e.g. 2 | 126 3 | 63 3 | 21 7	$2\times3\times3\times7$	2	M1 for a systematic method of at least 2 correct divisions by a prime number oe factor trees; can be implied by digits 2, 3, 3, 7 on answer line. A1 for $2\times3^2\times7$ or $2\times3\times3\times7$
(b)	$2\times3\times7$	42	2	B2 cao (B1 for 6, 14, 21 or $2\times3\times7$)
22	$\frac{8}{3}\times\frac{5}{4}=\frac{8\times5}{3\times4}=\frac{40}{12}$	$3\frac{1}{3}$	3	B1 for $\frac{8}{3}$ oe improper fraction or $\frac{5}{4}$ oe improper fraction M1 (dep on B1) for multiplying numerator and denominator of "$\frac{8}{3}$" and "$\frac{5}{4}$" A1 for $3\frac{1}{3}$ oe mixed number or $\frac{10}{3}$ OR B1 for 1.25 and 2.67 or 2.66(…) M1 (dep on B1) for correct method of multiplication A1 for $3.\dot{3}$

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
23			2	M1 for a relevant pair of intersecting arcs A1 for line drawn within guidelines, at least 3cm in length, accept broken line [SC: B1 for line drawn within guidelines if M0]
24 (a)		$-1, 0, 1, 2, 3$	2	B2 cao (-1 each error or omission)
(b)(i)		$x \geq \frac{7}{2}$	3	M1 for $2x \geq 7$, condone use of = sign or wrong equality A1 for $x \geq \frac{7}{2}$ oe as final answer
(ii)		4		SC:B1 for 3.5 or $\frac{7}{2}$ seen if M0 B1 ft from $x \geq$ "$\frac{7}{2}$"
25	$4x + 2y = 8$ $4x - 10y = 20$ $12y = -12$ $y = -1$ $4x + 2(-1) = 8$ $x = 2.5$	$x = 2.5$ $y = -1$	3	M1 for correct process to eliminate either x or y (condone one arithmetical error) M1 (dep) for substituting found value into either equation A1 for $x = 2.5$, $y = -1$ [SC: B1 for $x = 2.5$ or $y = -1$ if M0]

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
26		Interior angle of hexagon = 180 – (360 ÷ 6) = 120 360 – (90 + 120)	150	4	Alternative 1 M1 for 360 ÷ 6 A1 for 60 M1 (dep on M1) for "60" + 90 A1 cao Alternative 2 M1 for 360 ÷ 6 A1 for 60 M1 (dep on M1) for 360 – (2 × "60" + 90) A1 cao Alternative 3 M1 for (6 – 2) × 180 ÷ 6 A1 for 120 M1 (dep on M1) for 360 – (90 + "120") A1 cao
27	(a)		(16), 50, 82, 96, 100	1	B1 cao
	(b)	Cumulative freq. diag. curve/ segments	Cum. freq graph	2	B1 for 4 or 5 points plotted correctly ± 1 full (2mm) square depending on sensible table (condone 1 addition error) B1 (dep) for points joined by curve or line segments provided no gradient is negative - ignore any part of graph outside range of their points. (SC:B1 if 4 or 5 points plotted not at end but consistent within each interval and joined)
	(c)	100 – 42	58	2	M1 (ft dep on graph being cf) for reading from graph at 18 or 19, can be implied by answer in range 40 to 46 A1 for answer in range 56 to 60 or ft for 100 – '42' ±1 full (2mm) square

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
1	(a)	900 × 1.70 =	1530	2	M1 900 × 1.7(0) or digits 153(0) seen A1 cao
	(b)	160 ÷ 1.70 =	£94.12 or £94.11	2	M1 160 ÷ 1.7(0) or digits 941(....) seen A1 cao
2	(a)	4.7 ÷ 5.9 = 0.796610169	0.7966..	2	B2 for 0.7966 or better (B1 for 0.8, 0.80, 0.79, 0.796, 0.797 or digits 59 seen)
	(b)	0.82, 0.8, 0.85, 0.66, 0.875 0.66, 0.8, 0.82, 0.85, 0.875 2/3, 4/5, 0.82, 85%, 7/8	2/3, 4/5, 0.82, 85%, 7/8	2	B2 correct order (oe decimals in order) (B1 correct order reversed, or one error in ordered listing) with or without decimal equivalents. NB Accept 0.67 or 0.66
3		**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	3	B3 all 6 correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
4	(a)	–3 (–1) (1) 3 5 7	**–3**, 3, **5**, **7**	2	B2 all correct (B1 2,3 correct)
	(b)	(-2,-3),(-1,-1),(0,1),(1,3),(2,5),(3,7)	line	2	B2 cao for line from $x = -2$ to $x = 3$ (B1 plotting at least 5 points correctly or single line passing through (0, 1) or single line of gradient 2) The six possible points are: (-2,-3), (-1,-1), (0,1), (1,3), (2,5), (3,7)
	(c)(i)		–2	2	B1 $y = -2$ or ft from line segment
	(ii)		2.5		B1 $x = 2.5$ or ft from line segment

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
5	(a) (i)	180 – 54 (=126) "126" ÷ 2	63	3	M1 for (180–54) ÷ 2 A1 cao
	(ii)		Reason		B1 (indep) angles in triangle add to 180 OR equal angles in isosceles triangle OR equal angles and 2 sides the same (B0 if any incorrect reasoning given eg parallel, equilateral triangle)
	(b)	180 – "x"	117	1	B1 117 or ft 180 – "x" if $x < 90$
6		15/120×360 = 45 Plain 40/120×360 = 120 Cheese & Onion 55/120×360 = 165 Salt & Vinegar 10/120 × 360 = 30 Beef	45°, 120°, 165°, 30°	4	M1 evidence of method for at least one angle (could be implied by one correct angle of four on pie chart or in the table) A2 All four angles drawn ±2° tolerance, any order (A1 at least 2 angles correctly drawn ±2°, or all 4 angles in the table) B1 (dep on at least 1 angle drawn correctly, and exactly 4 sectors) for labels (flavour or frequency; initials will do) NB: Ignore the table if the pie chart provides the marks.
7	(a)	260 – 50 = 210 210 ÷ 35 =	6	3	M1 for 260–50 or 210 seen. M1 for "260-50" ÷ 35 or 210÷35 A1 cao
	(b)		P=35h + 50	3	B3 for P=35h+50 or P=35×h+50 oe (B2 for correct RHS or P=h + 50 ×35 or P=35h+k where k is numerical oe) (B1 for P = some other linear expression in h, OR h + 50 ×35 OR 35h seen) NB: P=h scores no marks; ignore £ signs. SC B2 for $h = \frac{P-50}{35}$
8	(a)		Elevation	2	B2 for 4 vertical squares. Accept 4 by 1 rectangle. (B1 for 4 vertical squares with one square added or one parallelogram added at the top, or 3 vertical squares, or 4 horizontal squares)

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
(b)		Plan	2	B2 for 2 adjacent squares, vertical or horizontal. Accept 2 by 1 rectangle. (B1 for 3 adjacent horizontal or vertical squares or a rectangle with sides in the ratio 2:1)
9 (a)		Points	1	B1 all three points ±1 full square.
(b)		Negative	1	B1 Negative. Ignore other descriptors unless contradictory.
(c)		lobf	1	B1 A single straight line drawn to cross between (5,30), (5,40) and (40,0), (40,15); accept freehand if considered to be straight.
(d)(i) (ii)		18–25 30–40	2	B1 18g–25g inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient. B1 30–40 min inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient.
10		300, 3, 75, 150	3	B3 for 4 correct (B2 for 2 or 3 correct) (B1 for 1 correct)
11 (a)	$6x - 7 + 7 = 38 + 7$ $6x = 45$	7.5	2	M1 $6x = 45$ or +7 both sides A1 7.5 oe; accept 45/6
(b)	$5y - 2 = 10$ or $20y - 8 = 40$ $5y = 12$ $20y = 48$	$2\frac{2}{5}$	3	M1 $20y–8(=40)$ or $\frac{4(5y-2)}{4} = \frac{40}{4}$ or $5y–2=10$ M1 (indep) for correct rearrangement into the form $ay=b+c$ or better (eg $20y=40+8$ or $5y=10+2$, using own terms) A1 for $2\frac{2}{5}$, 2.4 oe
12	5 miles = 8 km 70mph÷5×8 = 122 km/h OR 120km/h ÷ 8 × 8 = 75 mph Faster than 70 mph	70mph (Great Britain) (112 km)	3	M1 5 miles = 8 km; OR 70 mph is about 100 km/h OR 1km=0.6(25) miles OR 1mile=1.6km oe M1 70 ÷ 5 × 8 (=112) or 120 ÷ 8 × 5 (=75) oe A1 (dep on at least M1) GB or 70 mph Refer to both answer line and working. NB GB or 70 mph without working scores 0 marks.

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
13 (a)	$\pi \times 0.65$	2.04–2.05	2	M1 for $\pi \times 0.65$ or 3.14×0.65 or 3.142×0.65 oe A1 2.04–2.05 SC Award B1 for 2.0 seen (not 2)
(b)	1000 ÷ "(a)"	487–491	2	M1 for 1000 (or 100) ÷ "(a)" A1 for 487–491
14	$45.00 + 45.00 \times \frac{15}{100} =$ $45.00 + 6.75 =$	51.75	3	M2 for $45.00 + 45.00 \times \frac{15}{100}$ oe or 45.00×1.15 oe OR $45.00 + 6.75$ OR complete method or 5175 seen. (M1 for $45.00 \times \frac{15}{100}$ oe OR 6.75 seen OR 675 seen OR correct method for calculating 15% of 45) A1 cao SC Award B2 for an answer of 38.25
15		273	2	M1 for 728 ÷ 8 or 728 ÷ "3+5" or 91 A1 cao SC B1 for 455, or for 273:455
16	3 24 3.7 46.9(53) 4 60 3.8 51.0(72) 3.1 26.6(91) 3.9 55.4(19) 3.2 29.5(68) 3.21 29.8(66…) 3.3 32.6(37) 3.22 30.1(66…) 3.4 35.9(04) 3.23 30.4(68..) 3.5 39.3(75) 3.24 30.7(72...) 3.6 43.0(56) 3.25 31.0(78...) or 31	3.2	4	B2 for trial between 3.2 and 3.3 inclusive (B1 for trial between 3 and 4 inclusive) B1 for different trial between 3.21 and 3.25 inclusive B1 (dep on at least one previous B1) cao for 3.2 as final answer. NB: embedded answers: –B1; award Bs for evaluations rounded or truncated to at least 1 dp or for 31

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
17	$45^2 + 34^2 =$ $2025 + 1156 = 3181$ $\sqrt{3181} = 56.4$	56	4	M1 for $45^2 + 34^2$ M1 (dep) for $\sqrt{(2025 + 1156)}$ A1 for 56.4 … B1 for rounding their diagonal to the nearest integer (dep on evidence from decimal) NB 56 as the final answer gets full marks. NB Scale drawings result in 0 marks.
18	$2000 \times (1.055)^3$ Interest = 2348.48 – 2000 =	348.48	3	M1 for 5.5/100 × 2000 (oe) or 330 or 2330 or 110 or 2110 M1 (dep) for 5.5/100 × (2000 + "110" + "116.05") or 122.4… seen A1 cao (accept only 348.48 or 348.49) **OR** M2 for $2000 \times (1.055)^3$ or 2348.48(…) or 2348.49 seen (M1 for $2000 \times (1.055)^n$, $n \neq 3$) A1 for 348.48 or 348.49 [SC: B2 for 2348.48 - 2348.49]
19		Line	2	B2 line fully within tramlines, crossing AB and CD (B1 a straight line which crosses AB within the tramline, and also crosses CD) NB: Accept dotted or dashed lines, but not curves; accept freehand if considered to be straight. SC B1 for a perpendicular bisector of AB that is at least half way from AB to CD within the tramlines
20 (a)		$90<t\leq100$	1	B1 for $90<t\leq100$; accept 90-100.
(b)	65×12=780 75×22=1650 85×23=1955 95×24=2280 105×19=1995 8660/100=	86.6	4	M1 for use of fx with x consistent within intervals (including end points). Allow one slip. M1 (dep) for use of midpoints M1 (dep on 1st M1) for use of $\sum fx/100$ or $\sum fx/\sum f$ A1 86.6

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
21	(a)	$8 \times \frac{10}{4}$	20	2	M1 $\frac{10}{4}$ or $\frac{4}{10}$ or 0.4 or 2.5 oe seen A1 cao NB ratios get M0 unless of the form 1:*n* OR M1 $\frac{8}{4}, \frac{4}{8}$ oe seen A1 cao
	(b)	$15 \times \frac{4}{10}$	6	2	M1 $15 \times \frac{4}{10}$ oe A1 cao
22	(a)	$x^2 - 4x + 3x - 12 = x^2 - x - 12$	$x^2 - x - 12$	2	M1 for exactly 4 terms correct ignoring signs (eg x^2, 4*x*, 3*x*, 12) or 3 correct terms out of 4 terms with correct signs (eg 3 out of 4 of x^2, –4*x*, +3*x*,–12) A1 cao
	(b)	$(x+2)(x+5)$	$(x+2)(x+5)$	2	B2 cao (B1 for exactly one of $(x+2)$, $(x+5)$)
	(c)	$6=15+4q-20$ $6-15=4(q-5)$ $p-3t=4q-4t$ $6-3\times5=4(q-5)$	2 ¾	3	M1 for correct substitution of *p* and *t*. M1 for correct expansion of $4(q-t)$ oe (eg $4q-20$, $4q-4t$) A1 11/4 or 2 ¾ or 2.75 OR M1 for correct substitution of *p* and *t*. M1 for $\frac{p-3t}{4} = q - t$ oe A1 11/4 or 2 ¾ or 2.75

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
23	(a)	$9.9\times10^8 - 6.0\times10^7$	9.3×10^8	2	M1 for $99\times10^7 - 6\times10^7$ or $9.9\times10^8 - 0.6\times10^8$ or conversion of either to an ordinary number, or 930000000 or 93×10^7 or 9.3×10^n where n is any positive integer A1 cao
	(b)	$\frac{6.0-4.5}{4.5}\times100 = \frac{1.5}{4.5}\times100 =$ or $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}\times100$	33.3%	3	M2 $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}\times100$ oe (M1 for $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}$ or $\frac{6.0\times10^7 - 4.5\times10^7}{6.0\times10^7}\times100$ oe A1 cao OR M2 $\frac{6.0\times10^7}{4.5\times10^7}\times100$ -100 = (M1 $\frac{6.0\times10^7}{4.5\times10^7}\times100$ or 133.33(%)) A1 cao NB Accept any of the above expressions without any reference to 10^7.
24		LCM (40,24) = 120 Bread buns 120÷40 Burgers 120÷24 OR Bread buns: 40 is $2\times2\times2$ (×5) Burgers: 24 is $2\times2\times2$ (×3)	Bread buns 3 Burgers 5	3	M1 attempt to find LCM by eg lists of multiples, or summing of 40 and 24. A1 identify 120 (as LCM) A1 cao (both) OR M1 expansion of either number into its prime factors in a factor tree or 8×5 or 8×3 A1 both expansions correct A1 cao (both) SC B2 if answers given the wrong way around

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
25	8 × 41 = 328 2 × 29 = 58 328 – 58 = 270 270 ÷ 6 = 45	45	3	M1 for either 8 × 41 (=328) or 2 × 29 (=58) M1 (dep) "328" – "58" (=270) A1 cao NB 328 and/or 58 on the answer line gets M1 (implied); 270 on the answer line gets M2 (implied)

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
1 (a)	$1 - (0.2 + 0.3 + 0.1)$	0.4	2	M1 for $1 - (0.2 + 0.3 + 0.1)$ A1 for 0.4 oe, accept $\frac{0.4}{1}$
(b)	0.2×200	40	2	M1 for 0.2×200 A1 cao NB $\frac{40}{200}$ is M1 A0, 40 out of 200 is M1 A1
2	$650 - 430 = 220$ 1 choc ice costs 110p $650 - 5 \times 110 = 100$p	50	3	M1 for $650 - 430$ or 220 or 110 oe seen M1 for $650 - 5 \times \frac{'220'}{2}$ or $430 - 3 \times \frac{'220'}{2}$ oe A1 for 50p or £0.50 or £0.5 *Alternative scheme* $2x + 5y = 650$ $2x + 3y = 430$ oe M1 for subtracting two simultaneous equations to eliminate x (lollies)(2 or 3 terms correct) M1 for $650 - 5 \times 'y'$ or $430 - 3 \times 'y'$ oe A1 for 50p or £0.50 or £0.5 *Alternative scheme* M1 for $3\times (2x + 5y = 650)$ evaluated and $5 \times (2x + 3y = 430)$ evaluated oe (5 or 6 terms correct) M1 for subtraction of equations to eliminate y (choc ices)(2 or 3 terms correct ft) A1 for 50p or £0.50 or £0.5

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
3		question + response boxes oe	2	1st aspect: One question (eg 'how long does it take you to travel to school?' or 'What time did you leave home to get to school?'); ignore other questions. 2nd aspect: Response list (at least two), not overlapping. 3rd aspect: Some mention of units (eg minutes) in either question or responses B2 for all three aspects, or B1 for just one aspects.
4	2 [(3×1) + (4×1)] + (3×6) + (1×6) + (2×6) + (4×6) + (1×6) + (5×6)	110 cm^2	4	M1 for attempt to find the area of one face M1 for at least 6 faces with intention to add A1 cao B1 (indep) for cm^2 (with or without numerical answer)
5	$\left(\frac{-6+4}{2}\right), \left(\frac{5+3}{2}\right)$	(–1, 4)	2	B2 cao [B1 for (–1,*a*) or (b,4) or (4,-1)]
6		Box plot	2	3 aspects: 1st aspect – vertical line for median 2nd aspect – box using correct quartiles 3rd aspect – whiskers (could be single line) drawn with correct end points B2 for fully correct box plot (B1 for 1 aspect)
7			2	M1 for a relevant pair of intersecting arcs A1 for line drawn within guidelines, at least 3cm in length, accept broken line [SC: B1 for line drawn within guidelines if M0]

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
8	(a)	e.g. 2 | 126 3 | 63 3 | 21 7	$2\times3\times3\times7$	2	M1 for a systematic method of at least 2 correct divisions by a prime number oe factor trees; can be implied by digits 2, 3, 3, 7 on answer line. A1 for $2\times3^2\times7$ or $2\times3\times3\times7$
	(b)	$2\times3\times7$	42	2	B2 cao (B1 for 6, 14, 21 or $2\times3\times7$)
9	(a)		$-1,0,1,2,3$	2	B2 cao (-1 each error or omission)
	(b)(i)		$x\geq\frac{7}{2}$	3	M1 for $2x\geq7$, condone use of = sign or wrong equality A1 for $x\geq\frac{7}{2}$ oe as final answer
	(ii)		4		SC:B1 for 3.5 or $\frac{7}{2}$ seen if M0 B1 ft from $x\geq$"$\frac{7}{2}$" or $x>$"$\frac{7}{2}$"
10	(a)(i)		7^5	3	B1 cao
	(ii)		7^4		B2 cao (B1 for sight of 7^5 or 7^{2+3} or 7×7^3 or $7^1\times7^3$ or $7^2\times7^2$ or 7^{2+3-1})
	(b)		$\frac{1}{2}$	1	B1 for $\frac{1}{2}$ or 0.5 or 2^{-1}
11	(a)	$5n=m+21$	$n=\frac{m+21}{5}$	2	M1 for $5n=m+21$ or for attempt to divide three terms by 5 A1 $n=\frac{m+21}{5}$ oe

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
(b)	$4p-8q=3p+2$ $p-8q=2$ $p=8q+2$	$p=8q+2$	3	M1 for $4p-aq$ or $\frac{3}{4}p+b$ where a is an integer and b is a number M1 (dep) for taking one term correctly to LHS or RHS of expression A1 $p=8q+2$ oe
12		$y=\frac{1}{2}x+3$	2	B2 for $y = ½x + 3$ oe (B1 for $y = ½x + c$, $c \neq 7$ or $y = mx + 3$ oe or $½x + 3$ or $M = ½x + 3$)
13	$\frac{8}{3}\times\frac{5}{4}=\frac{8\times5}{3\times4}=\frac{40}{12}$	$3\frac{1}{3}$	3	B1 for $\frac{8}{3}$ oe or $\frac{5}{4}$ oe M1 (dep on B1) for multiplying numerator and denominator of "$\frac{8}{3}$" and "$\frac{5}{4}$" A1 for $3\frac{1}{3}$ oe mixed number or $\frac{10}{3}$ *OR* B1 for 1.25 and 2.67 or 2.66(…) M1(dep on B1) for correct method of multiplication A1 for $3.\dot{3}$
14	$4x+2y=8$ $4x-10y=20$ $12y=-12$ $y=-1$ $4x+2(-1)=8$ $x=2.5$	$x=2.5$ $y=-1$	3	M1 for correct process to eliminate either x or y (condone one arithmetical error) M1 (dep) for substituting found value into either equation A1 for $x = 2.5$, $y = -1$ [SC: B1 for $x = 2.5$ or $y = -1$ if M0]

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
15		Interior angle of hexagon = 180 – (360 ÷ 6) = 120 360 – (90 + 120)	150	4	Alternative 1 M1 for 360 ÷ 6 A1 for 60 M1 (dep on M1) for "60" + 90 A1 cao Alternative 2 M1 for 360 ÷ 6 A1 for 60 M1(dep on M1) for 360 – (2 × "60" + 90) A1 cao Alternative 3 M1 for (6 – 2) × 180 ÷ 6 A1 for 120 M1(dep on M1) for 360 – (90 + "120") A1 cao
16	(a)		(16), 50, 82, 96, 100	1	B1 cao
	(b)	Cumulative freq. diag. curve/ segments	Cum. freq graph	2	B1 ft for 4 or 5 points plotted correctly ± 1 full (2mm) square depending on sensible table (condone 1 addition error) B1 (dep) for points joined by curve or line segments provided no gradient is negative - ignore any part of graph outside range of their points. (SC:B1 if 4 or 5 points plotted not at end but consistent within each interval and joined)

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
(c)	$100 - 42$	58	2	M1 (ft dep on graph being cf) for reading from graph at 18 or 19, can be implied by answer in range 40 to 46 A1 for answer in range 56 to 60 or ft for 100 –'42' ±1 full (2mm) square
17 (a)		0.6 and 0.7,0.3,0.7	2	B1 for 0.6 on LH branch B1 for 0.7, 0.3 and 0.7 on RH branches
(b)	0.4×0.3	0.12	2	M1 for 0.4×0.3 A1 0.12 oe
(c)	$0.4\times0.7+0.6\times0.3$	0.46	3	M1 for '0.4×0.7' or '0.6×0.3' M1 for addition of two products from correct branches A1 0.46 oe Alternative M2 for an attempt to evaluate 1-(0.3×0.4 +'0.6×0.7') A1 cao
18	$x = 0.4545...$ so $100x = 45.4545...$ $99x = 45$ $x = \frac{45}{99} = \frac{15}{33}$	proof	3	M1 for $100x = 45.45\ldots$ or $10000x = 4545.45\ldots$ M1 (dep) for subtraction of both sides A1 for $\frac{15}{33}$ from correct proof
19	$\left(\sqrt{3}\right)^2 + \sqrt{3}\sqrt{2} - \sqrt{3}\sqrt{2} - \left(\sqrt{2}\right)^2$ $= 3 - 2$	1	2	B2 cao (B1 for $\sqrt{3}\sqrt{3} + \sqrt{3}\sqrt{2} - \sqrt{3}\sqrt{2} - \sqrt{2}\sqrt{2}$ oe, $\sqrt{3}\sqrt{3} - \sqrt{2}\sqrt{2}$ oe, or for 2,3, $\sqrt{4}$ $\sqrt{6}$ $\sqrt{9}$ seen)
20 (a)	$\frac{96}{24}$ or 4 $\sqrt{4}$ or 2	8	3	M1 for $\frac{96}{24}$ or $\frac{24}{96}$ or 4 or $\frac{1}{4}$ oe M1 for $\sqrt{\frac{96}{24}}$ or $\sqrt{\frac{24}{96}}$ or $\sqrt{'4'}$ or $\frac{1}{\sqrt{'4'}}$ or 2 or $\frac{1}{2}$ oe A1 cao

Paper 5525_05					
No		Working	Answer	Mark	Notes
	(b)	12×2^3	96	2	M1 for '2'^{3} or 8 A1 cao
21	(a)	$x\times3-x\times2x^2$	$3x-2x^3$	2	B2 cao (B1 for a two term expression with either $3x$ or $2x^3$)
	(b)		$4x(3y+x)$	2	M1 for taking out a factor of x, $2x$, 2, 4 or $4x$ A1 cao
	(c)	$\frac{5a}{b^2}$	$\frac{5a}{b^2}$	2	B2 for $\frac{5a}{b^2}$ or $5ab^{-2}$ (accept $\frac{5a}{1b^2}$) (B1 for either dealing with the numbers or dealing with the powers of a)
	(d)	$\frac{\cancel{x-3}}{\cancel{(x-3)}(x+3)}$	$\frac{1}{x+3}$	2	M1 for $(x-3)(x+3)$ A1 cao
22	(a)		$\mathbf{a}+\mathbf{b}$	1	B1 $\mathbf{a}+\mathbf{b}$ oe
	(b)	$ED=\mathbf{a}$ $DX=-2\mathbf{b}+2\ AC=2\mathbf{a}$ (So, $DX=2ED$)	$2\mathbf{a}$	3	M1 for $(DX=)\ DA+AX$ A1 for $(DX=)\ -2\boldsymbol{b}+2$ ("$\boldsymbol{a}+\boldsymbol{b}$") A1 $2\boldsymbol{a}$ from fully correct proof
23		$\pi(2x)^2h=\frac{4}{3}\pi(3x)^3$ $h=\frac{\frac{4}{3}\pi(3x)^3}{\pi(2x)^2}$	$9x$	3	M1 for $\pi(2x)^2h=\frac{4}{3}\pi(3x)^3$ (condone absence of brackets) M1 (dep) for valid algebra that gets to $h=ax$ (condone one error in powers of numerical constants) A1 cao

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
24	(i)	$n^2+(n+1)^2=2(n^2+n)+1$	$2n^2+2n+1$	4	M1 for at least 3 terms correct from $n^2+n+n+1$ A1 for $2n^2+2n+1$ oe
	(ii)		$2(n^2+n)$ is always even so $2(n^2+n)+1$ is always odd		M1 for recognizing $2n^2$ is always even A1ft complete proof for their quadratic *Alternative method* M1 for recognizing that if n^2 is odd then $(n+1)^2$ is even or vice versa A1 for complete proof
25			42 318	2	B1 for answer in range 36 – 48 B1 for answer in range 312 – 324
26	(a)	$(x-3)^2-3^2+15$	$p=3, q=6$	2	B2 for $p=3$ and $q=6$ (B1 for $p=3$ OR $q=6$) SC: award B2 for $(x-3)^2+6$ if p and q are not identified
	(b)		Sketch	2	B1 for U shaped curve B1 ft for TP in first quadrant (ft if TP not in first quadrant)
27	(a)	Graph translated 2 units upwards through points (–4, 2), (–2, 4), (0,2) and (3,5)	Sketch	2	M1 for a vertical translation A1 curve through points (–4, 2), (–2, 4), (0,2) and (3,5) ± ½ square
	(b)	Graph reflected in x-axis through points (–4,0), (–2, –2), (0, 0) and (3, -3)	Sketch	2	M1 for reflection in x-axis or y-axis A1 curve through points (–4,0), (–2, –2), (0, 0) and (3, -3) ± ½ square

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
1		273	2	M1 for 728 ÷ 8 or 728 ÷ "3+5" or 91 A1 cao SC B1 for 455 or 273:455
2		$3n - 1$	2	B2 for $3n - 1$ oe (B1 for $3n + k$ where $k \neq -1$ but k could be 0)
3	3 24 3.7 46.9(53) 4 60 3.8 51.0(72) 3.1 26.6(91) 3.9 55.4(19) 3.2 29.5(68) 3.21 29.8(66…) 3.3 32.6(37) 3.22 30.1(66…) 3.4 35.9(04) 3.23 30.4(68..) 3.5 39.3(75) 3.24 30.7(72...) 3.6 43.0(56) 3.25 31.0(78...) or 31	3.2	4	B2 for trial between 3.2 and 3.3 inclusive (B1 for trial between 3 and 4 inclusive) B1 for different trial between 3.21 and 3.25 inclusive B1 (dep on at least one previous B1) cao for 3.2 NB: embedded answers: –B1; award Bs for evaluations rounded or truncated to at least 1 dp or for 31
4	2 is the only even prime number and the product of 2 odd numbers is odd	Yes	2	B2 for 'yes' and '2 is the only even prime number and the product of two odd numbers is odd' oe (B1 for 'yes' and either '2 is the only even prime number' oe **or** 'the product of two odd numbers is odd' oe)
5	$45^2 + 34^2 =$ $2025 + 1156 = 3181$ $\sqrt{3181} = 56.4$	56	4	M1 for $45^2 + 34^2$ M1 for $\sqrt{(2025 + 1156)}$ A1 for 56.4 … B1 for rounding their diagonal to the nearest integer (dep on evidence from decimal) NB 56 with no incorrect working as the final answer gets full marks. NB Scale drawings result in 0 marks.

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
6	$2000 \times (1.055)^3$ Interest = 2348.48 – 2000 =	£348.48	3	M1 for 5.5/100 × 2000 (oe) or 330 or 16.5/100×2000 or 2330 or 110 or 2110 M1 (dep) for 5.5/100 × (2000 + "110" + "116.05") or 122.4… seen A1 cao (accept only 348.48 or 348.49) **OR** M2 for $2000 \times (1.055)^3$ or 2348.48(…) or 2348.49 seen (M1 for $2000 \times (1.055)^n$, $n \neq 3$) A1 for 348.48 or 348.49 [SC: B2 for 2348.48 - 2348.49]
7 (a)		Line	2	B2 line fully within tramlines, crossing AB and CD (B1 a straight line which crosses AB within the tramline, and also crosses CD) NB: Accept dotted or dashed lines, but not curves; accept freehand if considered to be straight. SC B1 for the perpendicular bisector of *AB* reaching halfway or more from *AB*
(b)		Region		B2 correct arc ±2mm and shaded within. Allow dotty or continous arc. (B1 inaccurate arc and shaded or accurate arc unshaded)
8	65×12=780 75×22=1650 85×23=1955 95×24=2280 105×19=1995 8660/100=	86.6	4	M1 for use of fx with x consistent within intervals (including end points). Allow one slip even if outside interval M1 (dep) for use of midpoints M1 (dep on 1st M1) for use of $\sum fx/100$ or $\sum fx/\sum f$ A1 86.6

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
9		Distance ÷ time: 1400 ÷ 2 h 20 min 20 mins is $\frac{1}{3}$ hour 1400 × 3 ÷ 7 =	600 kph	3	B1 20 mins as $\frac{1}{3}$ hour or as 0.33……hour M1 for distance ÷ time eg 1400 ÷ "2h 20 min" A1 cao OR B1 2 hour 20min = 140 (min) M1 Speed = $\frac{1400}{140}$ = (10 km per minute) A1 cao
10	(a)	$(x+18)+2x+(2x+7)=180$	Equation	2	B2 for $(x+18)+2x+(2x+7)=180$ oe (B1 for $(x+18)+2x+(2x+7)$)
	(b)	$5x+25 = 180$ $5x = 155$	31	2	M1 for simplifying to at least $5x+25=180$ or 360 (may be earned in (a)) A1 for $x=31$
11	(a)	(–14) –4 (0) 4 14	–4, 4, 14	2	B2 for all 3 values correct (B1 for 1 or 2 values correct)
	(b)		curve	2	B1 for all 5 points plotted correctly ± ½ square (ft from table if at least B1 awarded in (a)) B1 (indep) ft for any smooth curve through their points

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
12	(a)	$8 \times \frac{10}{4}$	20	2	M1 $\frac{10}{4}$ or $\frac{4}{10}$ or 0.4 or 2.5 oe seen A1 cao NB ratios get M0 unless of the form 1:*n* OR M1 $\frac{8}{4}, \frac{4}{8}$ oe seen A1 cao
	(b)	$15 \times \frac{4}{10}$	6	2	M1 $15 \times \frac{4}{10}$ oe A1 cao
13	(a)	$9.9\times10^8 - 6.0\times10^7$	$\pm 9.3\times10^8$	2	M1 for $99\times10^7 - 6.0\times10^7$ or $9.9\times10^8 - 0.60\times10^8$ or correct conversion of either to an ordinary number or 930000000 or 9.3×10^n where *n* is any positive integer A1 cao
	(b)	$\frac{6.0-4.5}{4.5} \times 100 = \frac{1.5}{4.5} \times 100 =$ or $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7} \times 100$	33.3%	3	M2 $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7} \times 100$ oe (M1 for $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}$ or $\frac{6.0\times10^7 - 4.5\times10^7}{6.0\times10^7} \times 100$ oe A1 33.3 – 33.4 OR M2 $\frac{6.0\times10^7}{4.5\times10^7} \times 100$ -100 or 33.3% (M1 $\frac{6.0\times10^7}{4.5\times10^7} \times 100$ or 133.3%) A1 33.3-33.4 NB Accept any of the above expressions without any reference to 10^7.

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
14	LCM (40,24) = 120 Bread buns 120÷40 Burgers 120÷24 OR Bread buns: 40 is $2 \times 2 \times 2$ (×5) Burgers: 24 is $2 \times 2 \times 2$ (×3)	Bread Buns 3 Burgers 5	3	M1 attempt to find LCM by eg lists of multiples, or summing of 40s and summing of 24s, with at least 3 numbers in each list A1 identify 120 as LCM A1 cao (both) **OR** M1 expansion of either number into its prime factors in a factor tree or 8×5 or 8×3 A1 both expansions correct A1 cao (both) SC B2 if answers given the wrong way around
15	$\sin 32 = \frac{AB}{12}$ $AB = 12 \times \sin 32$ $AB = 6.35903\ldots$	6.36	3	M1 $\sin 32 = \frac{AB}{12}$ (accept $\text{Sin}\ \frac{AB}{12}$) M1 $12 \times \sin 32$ or $12 \times 0.5299..$ A1 accept 6.359 – 6.360 SC Gradians 5.78(1…) Radians 6.62 Get M1M1A0 **OR** Use of Sine Rule $\frac{\sin 32}{AB} = \frac{\sin 90}{12}$ or $\frac{AB}{\sin 32} = \frac{12}{\sin 90}$ M1 $AB = \frac{12 \times \sin 32}{\sin 90}$ M1 $AB =$ 6.359 – 6.36 A1 SC Gradians 5.85(…) Radians 7. 40(…) M1M1A0

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
16		$8 \times 41 = 328$ $2 \times 29 = 58$ $328 - 58 = 270$ $270 \div 6 = 45$	45	3	M1 for either 8×41 (=328) or 2×29 (=58) M1 (dep) "328" – "58" (=270) A1 cao NB 328 and /or 58 on the answer line gets M1 (implied); 270 on the answer line gets M2 (implied)
17	(a)	$x^2 - 4x + 3x - 12 = x^2 - x - 12$	$x^2 - x - 12$	2	M1 for exactly 4 terms correct ignoring signs (x^2, $4x$, $3x$, 12) or 3 out of 4 terms with correct signs (x^2, $-4x$, $+3x$, -12) A1 cao
	(b)	$6x^2 - 8x + 15x - 20 = 6x^2 + 7x - 20$	$6x^2 + 7x - 20$	2	M1 for exactly 4 terms correct ignoring signs ($6x^2$, $8x$, $15x$, 20) or 3 out of 4 terms with correct signs ($6x^2$, $-8x$, $+15x$, -20) A1 cao
	(c)	$(x+2)(x+5)$	$(x+2)(x+5)$	2	B2 cao (B1 for exactly one of $(x + 2)$, $(x + 5)$)
	(d)		$12p^8q^3$	2	B2 cao (B1 for any 2 out of 3 terms correct in a product or 3 terms correct in a sum or part product)
	(e)	$6 = 15 + 4(q - 5)$ $6 = 15 + 4q - 20$ $11 = 4q$	$2\frac{3}{4}$	3	M1 for correct substitution of p and t. M1 for correct expansion of $4(q\text{-}t)$ oe (eg $4q$-20, $4q$-$4t$) A1 11/4 or 2 ¾ or 2.75 OR M1 for correct substitution of p and t. M1 for $\frac{p - 3t}{4} = q - t$ oe A1 11/4 or 2 ¾ or 2.75
18	(a)	$T=kx$; $150=6k$; $k=25$	$T=25x$	3	M1 for $T=kx$, k algebraic M1 subs $T=150$ and $x=6$ into $T = kx^n$ $(n \neq 0)$ A1 for $T=25x$ oe SC B1 $T \propto 25x$ oe

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
	(b)	$T = 25 \times 15 =$	375	1	B1 ft on k , $k \neq 1$
	(c)	$600 = 25x$; $x = 600 \div 25 =$	24	1	B1 ft on k , $k \neq 1$
19		$\frac{120}{570} \times 50 = 10.5263$ $\frac{250}{570} \times 50 = 21.9298$ $\frac{200}{570} \times 50 = 17.5438$	10 22 18	3	M1 method shown eg $\frac{120}{120+250+200} \times 50$ or one of 10.5(263) 21.9(298), 17.5(438) **OR** 570 ÷ 50 = 11.4 and one of 120 ÷ 11.4 =10.526…) , 270 ÷ 11.4 (=21.9298…) , 200 ÷ 11.4 (=17.5438…) A1 for 10.5(263), 21.9(298), 17.5(438) or 11, 22, 18 B1 correction to add to 50: 10, 22, 18 or 11, 22, 17
20		$x(x+1) + 4(2x–3) = (2x–3)(x+1)$ $x^2 + x + 8x – 12 = 2x^2 + 2x – 3x – 3$ $x^2 – 10x + 9 = 0$ $(x – 9)(x – 1) = 0$	$x = 1, 9$	5	M1 for multiplying through by common denominator $(2x–3)(x+1)$ M1 (dep)for either $x^2 + x + 8x – 12$ or $2x^2 + 2x – 3x – 3$ oe A1 for correct quadratic (= 0) M1 for a correct method to solve 3 term quadratic A1 cao for both solutions
21		$AB^2 = 8^2 + 9^2 – 2\times8\times9\times\cos40$ $AB^2 = 64 + 81 – 144\times\cos40$ $AB^2 = 145 – 144 \times 0.766$ $AB^2 = 145 – 110.31… = 34.6896$ $AB = \sqrt{34.6796} = 5.8897877$	5.89	3	M1 Subs in Cos Rule: $8^2 + 9^2 – 2\times8\times9\times\cos40$ M1 correct order of evaluation of $8^2 + 9^2 – 2\times8\times9\times\cos40$ A1 cao 5.88 – 5.89 SC: Award B2 for one of $AB^2 = 241.03…$ or $AB = 15.525…$ (radians) $AB^2 = 28.50…$ or $AB = 5.33…$ (gradians)
22	(a)		24 30	2	B1 24 cao B1 30 cao
	(b)	Column $60 \leq w < 75$ at 10 Column $75 \leq w < 95$ at 4	10 4	2	B1 10 and correct width (tol $\pm\frac{1}{2}$ small square) B1 4 and correct width (tol $\pm\frac{1}{2}$ small square)

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
23 (a)		100.5	1	B1 cao
(b)		10.515	1	B1 cao
(c)	$\frac{100.5}{10.515}$	9.5577746	2	M1 for greatest distance divided by least time Where 100 < greatest distance ≤ 100.5, 10.51 ≤ least time < 10.52 A1 for 9.555 – 9.56
(d)	$\frac{99.5}{10.525}$	9.45368..	2	M1 for least distance divided by greatest time Where 99.5 ≤ least distance < 100, 10.52 < greatest time ≤10.53 A1 for 9.45 – 9.455
24	$\frac{4}{7}\times\frac{4}{7}\times\frac{3}{7} + \frac{3}{7}\times\frac{3}{7}\times\frac{4}{7} =$ $\frac{48+36}{343} = \frac{84}{343}$ But there are three ways this can be achieved: BBG, BGB, GBB So the probability is $\frac{84}{343}\times 3$ OR $1-\frac{64}{343}-\frac{27}{343}$ NB: 84/343 = 0.244897; 252/343 = 0.73469 $\frac{4}{7} = 0.57(142...)$, $\frac{3}{7} = 0.42(857...)$	$\frac{252}{343}$	3	M1 for $\frac{4}{7}\times\frac{4}{7}\times\frac{3}{7}$ or $\frac{3}{7}\times\frac{3}{7}\times\frac{4}{7}$ oe or $\left(\frac{4}{7}\right)^3$ oe or $\left(\frac{3}{7}\right)^3$ oe or $\frac{91}{343}$ or 0.10(49…) or 0.13(99….) M1 (indep) for identification of all 6 outcomes (M2 for $1-\left[\left(\frac{4}{7}\right)^3+\left(\frac{3}{7}\right)^3\right]$) oe A1 $\frac{252}{343}$, $\frac{36}{49}$, 0.73(469…) oe

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
25	(a)	$\frac{1}{3} \times \pi \times 5^2 \times 8 = \pi \times 25 \times 8 \div 3 =$ 209.4395	209–210	2	M1 for $\frac{1}{3} \times \pi \times 5^2 \times 8$ A1 for 209–210
	(b)	Base radius = $\frac{216}{360} \times 15 = 9$ Height = $\sqrt{(15^2 - 9^2)} =$	12	4	M1 for 216 ÷ 360 A1 for 9 M1 for $\sqrt{(15^2 - \text{“9”}^2)}$, where “9” < 15 A1 cao

PMT

edexcel

Mark Scheme (Results)

Summer 2012

GCSE Mathematics (Linear) 1MA0
Foundation (Non-Calculator) Paper 1F

PMT

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at www.edexcel.com.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2012
Publications Code UG032619
All the material in this publication is copyright
© Pearson Education Ltd 2012

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labeling conventions.

ii) *select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect canceling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.



12 Parts of questions
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers
Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

Guidance on the use of codes within this mark scheme
M1 – method mark A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working


1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
1	(a)		380	1	B1 cao
	(b)		6.2	1	B1 cao
	(c)		Arrow at 34	1	B1 cao
2	(a)		8	1	B1 for 8 ± 0.2
	(b)		35	1	B1 for 35 ± 2°
	(c)		Circle drawn	1	B1 for all parts within ± 2mm, (use overlay)
3	(a)		4, 7, 4, 3, 2	2	M1 for at least 3 correct tallies **or** at least 3 correct frequencies A1 for all frequencies correct
	(b)		7	1	B1 for 7 **or** ft from frequencies in (a) or tallies if no frequencies
	(c)		Diagram drawn	3	M1 for bar chart or other suitable chart with at least 3 correct heights for their scale (can f.t.) A1 for all 5 bars correctly labelled and vertical axis correctly scaled A1 for fully correct **or** ft frequencies in (a) **OR** M1 for pictogram with at least 3 correct rows (can f.t.) A1 for correct labels on all 5 rows and correctly key A1 for fully correct **or** ft frequencies in (a) **OR** M1 for pie chart with at least 3 correct sectors ± 2° (can f.t.) A1 for all 5 sectors correctly labelled A1 for fully correct **or** ft frequencies in (a)

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
4		£1.18 + 94p = £2.12 £5 – £2.12 – 30p = £2.58 £2.58 ÷ 2 =	1.29	3	M2 for (5 – 1.18 – 0.94 – 0.30) ÷ 2 oe **or** digits 129 (M1 for 1.18 + 0.94 **or** 2.12 seen **or** 1.18 + 0.94 + 0.30 oe **or** 2.42 seen **or** 5 – 1.18 – 0.94 oe **or** 2.88 seen **or** (5 – 1.18 – 0.94) ÷ 2 **or** 1.44 seen **or** 5 – 1.18 – 0.94 – 0.30 oe **or** 2.58 seen) A1 cao **NOTE**: Accept working in £ or pence
5	(a)(i) (ii) (b)		(2, 3) (–3, 1) Point plotted at (3, –4)	2 1	B1 cao B1 cao B1 cao
6	(a) (b) (c)		– 5 6 3	1 1 1	B1 cao B1 for 6 or – 6 B1 cao
7			(P, B), (P, S), (P, L) (M, B), (M, S), (M, L) (H, B), (H, S), (H, L)	2	M1 for any 3 combinations with no incorrect combinations A1 for all 9 combinations with no duplicates or extras
8	(a) (b)	 24 ÷ 4 =	Walk 6	1 2	B1 cao M1 for 24 ÷ 4 oe or ¼ oe seen A1 cao

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
9	(a)		Isosceles triangle	1	B1 for isosceles triangle
	(b)		Rectangle with area 12 cm^2	2	M1 for rectangle drawn A1 cao
10	(a)		A marked at 0	1	B1 for A marked at 0 (within overlay)
	(b)		B marked at 1/4	1	B1 for B marked at 1/4 (within overlay)
11	(a)		9	1	B1 cao
	(b)		33	2	M1 for 5×5 **or** 25 seen in the working **or** $2 \times 2 \times 2$ **or** 8 seen in the working A1 cao
12	(a)		20	2	M1 $3 \times 3 \times 3$ oe seen or drawn or 27 seen or use of 3 layers A1 cao
	(b)			2	B2 for correct view (B1 for  or )
13	(a)(i)		07 29	2	B1 for 07 29
	(ii)		36		B1 for 36 or ft difference between (i) and 06 53
	(b)		07 51	1	B1 cao
	(c)		09 55	1	B1 for 09 55 or 9 55 or five to ten

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
14		2 + 8 + 2 + 8 = 20 20 ÷ 4 =	5	4	M2 for 2 + 8 + 2 + 8 oe **or** 20 seen **or** (2 + 8) ÷ 2 oe (M1 for the sum of 3 sides of the rectangle) M1 (dep) for the sum of 3 or 4 sides of the rectangle ÷ 4 **or** an attempt to evaluate (2 + 8) ÷ 2 oe to get the length of one side A1 cao **SC**: B1 for an answer of 4 coming from $\sqrt{2\times 8}$ oe
15	(a)		4	1	B1 cao
	(b)	9.5 – 4.75 = **OR** 9.5 ÷ 2 =	4.75	2	M1 for 9.5 – 4.75 **or** 9.5 ÷ 2 **or** 4.75 – 9.5 A1 cao
	(c)		6	1	B1 cao
	(d)	12 × 4 =	48	2	M1 for × 4 seen **or** identifying + 0.5 for every 2 inches **or** 12 + 12 + 12 + 12 oe **or** build up method eg 12, 24, 36, 48 allow one error A1 cao
16	(a)		trapezium	1	B1 for trapezium or isosceles trapezium
	(b)			2	B2 for correct tessellation (at least 5 more shapes) (B1 for at least 4 shapes (including initial shape) correctly tessellating)

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
17*		S: 35 ÷ 100 × 40 = 14 W: 40 ÷ 8 × 3 = 15 **OR** D: 16 ÷ 40 (× 100) = 0.4 (40%) W: 3 ÷ 8 (× 100) = 0.375 (37.5%) **OR** D: $\frac{16}{40} = \frac{80}{200}$ S: $\frac{35}{100} = \frac{70}{200}$ W: $\frac{3}{8} = \frac{75}{200}$	Debbie and correct calculations	4	**Compares Marks out of 40 or fractions with denominator of 40** M1 for 35 ÷ 100 × 40 oe **or** 14 seen (or 14/40 seen) M1 for 40 ÷ 8 × 3 **or** 15 seen (or 15/40 seen) A1 for 14 and 15 **or** $\frac{14}{40}$ and $\frac{15}{40}$ C1 (dep on M1) for correct conclusion for their working QWC with 3 comparable marks: Decision and justification should be clear with working clearly presented and attributable. **OR Decimals (or Percentages)** M1 for 16 ÷ 40 (× 100) oe **or** 0.4 (or 40) seen M1 for 3 ÷ 8 (× 100) oe **or** 0.375 (or 37.5) seen A1 for 0.4 and 0.375 (**or** 40 and 37.5) C1 (dep on M1) for correct conclusion for their working QWC: with 3 comparable decimals (or percentages: Decision and justification should be clear with working clearly presented and attributable. **OR Compares Fractions with denominator other than 40** M1 for attempt to convert all to fractions with a common denominator other than 40 M1 for at least 1 correct A1 for $\frac{80}{200}$ and $\frac{70}{200}$ and $\frac{75}{200}$ oe C1 (dep on M1) for correct conclusion for their working QWC with 3 comparable fractions: Decision and justification should be clear with working clearly presented and attributable.

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
18*	(a)		10	1	B1 cao
	(b)	 **Miles** 0 10 20 30 40 50 **Ed** 0 15 30 45 60 75 **Bill** 10 20 30 40 50 60	Ed is cheaper up to 20 miles, Bill is cheaper for more than 20 miles	3	M1 for correct line for Ed intersecting at (20,30) ±1 sq tolerance **or** $10 + x = 1.5x$ oe C2 (dep on M1) for a correct full statement ft from graph eg. Ed cheaper up to 20 miles **and** Bill cheaper for more than 20 miles (C1 (dep on M1) for a correct conclusion ft from graph eg. cheaper at 10 miles with Ed ; eg. cheaper at 50 miles with Bill eg. same cost at 20 miles; eg for £5 go further with Bill **or** A general statement covering short **and** long distances eg. Ed is cheaper for shorter distances **and** Bill is cheaper for long distances) **OR** (continued on next page)

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
18 (contd)					**OR** M1 for correct method to work out Ed's delivery cost for at least 2 values of n miles where $0 < n \leq 50$ **or** for correct method to work out Ed and Bill's delivery cost for n miles where $0 < n \leq 50$ C2 (dep on M1) for 20 miles linked with £30 for Ed and Bill with correct full statement eg. Ed cheaper up to 20 miles **and** Bill cheaper for more than 20 miles (C1 (dep on M1) for a correct conclusion eg. cheaper at 10 miles with Ed; eg. cheaper at 50 miles with Bill eg. same cost at 20 miles; eg for £5 go further with Bill **or** A general statement covering short **and** long distances eg. Ed is cheaper for shorter distances **and** Bill is cheaper for long distances) SC: B1 for correct full statement seen with no working eg. Ed cheaper up to 20 miles **and** Bill cheaper for more than 20 miles QWC Decision and justification should be clear with working clearly presented and attributable

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
19		1,96 × 2.25 = 4.41 **OR** 4.23 ÷ 9 = 0.47 1.96 ÷ 4 = 0.49 **OR** 4.23 × 4 = 16.92 1.96 × 9 = 17.64 **OR** 4.23 ÷ 9 = 0.47 0.47 × 4 = 1.88 **OR** 1.96 ÷ 4 = 0.49 0.49 × 9 = 4.41 **OR** 9 ÷ 4.23 = 2.12 4 ÷ 1.96 = 2.04	Pack of 9	3	M2 for a fully correct method to enable a conclusion eg 1.96 × 2¼ **OR** M1 for 4.23 ÷ 9 or 423 ÷ 9 **or** 0.47 seen **or** 47 seen M1 for 1.96 ÷ 4 or 196 ÷ 4 or 0.49 seen or 49 seen **OR** M1 for 4.23 × 4 **or** 423 × 4 **or** 16.92 seen **or** 1692 seen M1 for 1.96 × 9 **or** 196 × 9 **or** 17.64 seen **or** 1764 seen **OR** M1 for 4.23 ÷ 9 **or** 423 ÷ 9 **or** 0.47 seen **or** 47 seen M1 for 0.47 × 4 **or** 47 × 4 **or** 1.88 seen **or** 188 seen **OR** M1 for 1.96 ÷ 4 **or** 196 ÷ 4 **or** 0.49 seen **or** 49 seen M1 for 0.49 × 9 **or** 49 × 9 **or** 4.41 seen **or** 441 seen **OR** M1 for 9 ÷ 4.23 **or** 2.12(...) seen **or** 2.13 seen M1 for 4 ÷ 1.96 **or** 2.04(...) seen A1 for Pack of 9 and fully correct calculations **NOTE:** B0 for an answer of 9 not supported by working.
20	(a)		6	1	B1 cao
	(b)		44	1	B1 cao
	(c)		31	2	M1 for 60 – 29 **or** 29 – 60 **or** any correct method that is attempting to find the difference between 29 and 60 (allow 1 arithmetic error) A1 cao

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
21*		Angle *DBC* = (180 – 50) ÷ 2 Base angles of isosceles triangle are equal Angle *ABD* = 180 – 65 Angles on a straight line add up to 180 *x* = 180 – 20 – 115 Angles in a triangle add up to 180 **OR** Angle *DBC* = (180 – 50) ÷ 2 Base angles of isosceles triangle are equal *x* = 65 – 20 Exterior angle of triangle is equal to sum of interior opposite angles **OR** Angle *DCB* = (180 – 50) ÷ 2 Base angles of isosceles triangle are equal *x* = 180 – 50 – 20 – 65 Angles in a triangle add up to 180	45 with reasons	4	M1 for (180 – 50) ÷ 2 oe **or** 65 seen M1 for 180 – 20 – (180 – "65") **or** "65" – 20 or 180 – 50 – 20 –'65' oe C2 for *x* identified as 45 with full reasons QWC: Reasons clearly laid out with correct geometrical language used (C1 (dep on M1) for one reason QWC: Reasons clearly laid out with correct geometrical language used) NOTE: *x* = 45 with no working or without any correct angles marked on the diagram cannot score.

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
22	(a)	360 ÷ 60 = 6 300 ÷ 60 = 5 6 × 5 =	Yes and 30	3	M1 for dividing side of patio by side of paving slab eg 360 ÷ 60 **or** 300 ÷ 60 **or** 3.6 ÷ 0.6 **or** 3 ÷ 0.6 **or** 6 and 5 seen **or** 6 divisions seen on length of diagram **or** 5 divisions seen on width of diagram M1 for correct method to find number of paving slabs eg (360 ÷ 60) × (300 ÷ 60) oe **or** 6 × 5 **or** 30 squares seen on diagram (units may not be consistent) A1 for Yes **and** 30 (or 2 extra) with correct calculations **OR** M1 for correct method to find area of patio **or** paving slab eg 360 × 300 **or** 108000 seen **or** 60 × 60 **or** 3600 seen **or** 3.6 × 3 **or** 10.8 seen **or** 0.6 × 0.6 **or** 0.36 seen M1 for dividing area of patio by area of a paving slab eg. (3.6 × 3) ÷ (0.6 × 0.6) oe (units may not be consistent) A1 for Yes **and** 30 (or 2 extra) with correct calculations **OR** M1 for method to find area of patio and area of 32 slabs eg. 60 × 60 × 32 **or** 360 × 300 M1 for method to find both areas eg. 60 × 60 × 32 **and** 360 × 300 (units may not be consistent) A1 for Yes **and** 115200 **and** 108000 **OR** Yes **and** 11.52 **and** 10.8 **NB :** Throughout the question, candidates could be working in metres or centimetres

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
22	(b)	1726 25890 27616 **8 6 3** **2** 2 4 1 8 9 **3** **7** 1 6 1 2 6 **2** 6 1 6 800 60 3 30 24000 1800 90 2 1600 120 6 24000 + 1800 + 90 + 1600 + 120 + 6 = 27616	276.16	3	M1 for complete correct method with relative place value correct. Condone 1 multiplication error, addition not necessary. **OR** M1 for a complete grid. Condone 1 multiplication error, addition not necessary. **OR** M1 for sight of a complete partitioning method, condone 1 multiplication error. Final addition not necessary. A1 for digits 27616 A1 ft (dep on M1) for correct placement of decimal point after addition (of appropriate values) (SC: B1 for attempting to add 32 lots of 8.63)
23	(a)		30	2	M1 for 25 ÷ 10 **or** 2.5 seen **or** 10 ÷ 25 **or** 0.4 seen **or** 12+12+6 oe **or** a complete method eg. 25×12÷10 oe A1 cao
	(b)	1000 ÷ 200 × 12	60	2	M1 for 500 ÷ 50 **or** 1000 ÷ 200 **or** 500 ÷ 10 **or** correct scale factor clearly linked with one ingredient eg 10 with sugar or 5 with butter or flour or 50 with milk **or** an answer of 120 or 600 A1 cao

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
24		Acton after 24, 48, 72, 96, .. Barton after 20, 40, 60, 80, .. LCM of 20 and 24 is 120 9:00 am + 120 minutes **OR** Acton after 24, 48, 1h 12 min... Barton after 20, 40, 1 h LCM is 2 hours 9:00 am + 2 hours **OR** Times from 9:00 am when each service leaves the bus station Acton at 9:24, 9:48, 10:12.. Barton at 9:20, 9:40, 10:00.. **OR** 20 = 2 × 2 × 5 24 = 2 × 2 × 2 × 3 2 × 2 × 2 × 3 × 5 = 120	11:00 am	3	M1 for listing multiples of 20 and 24 with at least 3 numbers in each list ; multiples could be given in minutes or in hours and minutes (condone one addition error in total in first 3 numbers in lists) A1 identify 120 (mins) **or** 2 (hours) as LCM A1 for 11:00 (am) **or** 11(am) **or** 11 o'clock **OR** M1 for listing times after 9am when each bus leaves the bus station, with at least 3 times in each list (condone one addition error in total in first 3 times after 9am in lists) A1 for correct times in each list up to and including 11:00 A1 for 11:00 (am) **or** 11(am) **or** 11 o'clock **OR** M1 for correct method to write 20 and 24 in terms of their prime factors 2, 2, 5 and 2, 2, 2, 3 (condone one error) A1 identify 120 as LCM A1 for 11:00 (am) **or** 11(am) **or** 11 o'clock

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
25	(a)		$6y - 15$	1	B1 cao
	(b)		$4x(2x + y)$	2	B2 cao (B1 for $x(8x + 4y)$ **or** $2x(4x + 2y)$ **or** $4(2x^2 + xy)$ **or** $4x(ax + by)$ where a, b are positive integers **or** $ax(2x + y)$ where a is a positive integer **or** $4x(2x \quad y)$)
	(c)	$10t = gh$ $h = \frac{10t}{g}$	$\frac{10t}{g}$	2	M1 for clear intention to multiply both sides of the equation by 10 (eg. ×10 seen on both sides of equation) **or** clear intention to divide both sides of the equation by g (eg. ÷ g seen on both sides of equation) **or** $10t = gh$ **or** $\frac{t}{g} = \frac{h}{10}$ **or** fully correct reverse flow diagram eg. ← ×10 ← ÷g ← A1 for $\frac{10t}{g}$ oe

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
26	(a)	$2 \times 5 \times 2 = 20$ $300 \div 20 =$	15	3	M2 for $300 \div (2 \times 5 \times 2)$ oe (M1 for $2 \times 5 \times 2$ **or** 20 seen **or** $300 \div (2 \times 5)$ **or** 30 seen A1 cao
	(b)	$c = \frac{30 \times 40}{150}$ =	8	2	M1 for $\frac{30 \times 40}{150}$ **or** 1200 seen A1 cao
27		$3x - 15 = 2x + 24$ $x = 39$ **OR** $2x + 3x - 15 + 2x + 2x + 24 = 360$ $9x + 9 = 360$ $9x = 351$ $x = 39$ **OR** $2x + 2x + 24 = 180$ $4x + 24 = 180$ $4x = 156$ $x = 39$ **OR** $2x + 3x - 15 = 180$ $5x - 15 = 180$ $5x = 195$ $x = 39$	39	3	M1 for forming an appropriate equation eg $3x - 15 = 2x + 24$ **or** $2x + 3x - 15 + 2x + 2x + 24 = 360$ oe **or** $2x + 2x + 24 = 180$ oe **or** $2x + 3x - 15 = 180$ oe **or** $2x + 3x - 15 = 2x + 2x + 24$ M1 (dep) for correct operation(s) to isolate x and non-x terms in an equation to get $ax = b$ A1 cao **OR** M2 for $\frac{351}{9}$ **or** $\frac{195}{5}$ **or** $\frac{156}{4}$ oe A1 cao

PMT

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481
Email publication.orders@edexcel.com

Order Code UG032619 Summer 2012

For more information on Edexcel qualifications, please visit our website
www.edexcel.com

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual
Llywodraeth Cynulliad Cymru
Welsh Assembly Government
Rewarding Learning

PMT

edexcel

Mark Scheme (Results)

November 2015

Pearson Edexcel GCSE
In Mathematics B (2MB01)
Higher (Non-Calculator) Unit 2

PMT

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

November 2015
Publications Code UG042910
All the material in this publication is copyright
© Pearson Education Ltd 2015

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Note that in some cases a correct answer alone will not score marks unless supported by working; these situations are made clear in the mark scheme. Examiners should be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will award marks for the quality of written communication (QWC).
The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labelling conventions.

ii*) select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Partial answers shown (usually indicated in the ms by brackets) can be awarded the method mark associated with it (implied).
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks; transcription errors may also gain some credit. Send any such responses to review for the Team Leader to consider.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

PMT

11 Linear equations
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded (embedded answers).

12 Parts of questions
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers
Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

14 The detailed notes in the mark scheme, and in practice/training material for examiners, should be taken as precedents over the above notes.

Guidance on the use of codes within this mark scheme
M1 – method mark for appropriate method in the context of the question A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

PMT

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
1.			6 525 900 450	3	M1 for × 6 ÷ 4 oe or $\frac{6}{4}$ or 1.5 or 1 correct answer A2 for all correct (A1 for 2 or 3 correct)
2.			$8n + 24$	3	M1 for generating at least 3 terms of sequence 32, 40, 48 or sequence 4, 5, 6 M1 for $8n \pm k$ or $8(n \pm m)$ where $k \neq 24$, $m \neq 3$ A1 for $8n + 24$ or $8(n + 3)$ oe
3.			105 cm^3	4	M1 for writing factor pairs of 35, 21 or 15 M1 for "3" × "5" × "7" using their lengths dep on one at least one correct factor pair A1 ft for 105 (dep on one correct factor pair) B1 for cm^3

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
4.		$y = \frac{1}{2}x + 3$ x : −2, −1, 0, 1, 2, 3, 4 y : 2, 2.5, 3, 3.5, 4, 4.5, 5	Correct line from (−2, 2) to (4, 5)	3	**(Table of values / calculation of values)** M1 for at least 2 correct attempts to find points by substituting values of x. M1 ft for plotting at least 2 of their points (any points plotted from their table must be correctly plotted) A1 for correct line between $x = -2$ and $x = 4$ **(No table of values)** M1 for at least 2 correct points with no more than 2 incorrect points plotted M1 for at least 2 correct points (and no incorrect points) plotted **OR** line segment of $y = \frac{1}{2}x + 3$ drawn A1 for correct line between $x = -2$ and $x = 4$ **(Use of $y = mx + c$)** M1 for line drawn with gradient of $\frac{1}{2}$ **OR** line drawn with a y intercept of 3 M1 for line drawn with gradient of $\frac{1}{2}$ **AND** line drawn with a y intercept of 3 A1 for correct line between $x = -2$ and $x = 4$ **SC** : B2 for correct line from $x = 0$ to $x = 4$

PMT

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
5.			68	5	B1 for correct conversion of units M1 for 250 ÷ 25 (= 10) or 300 ÷ 20 (= 15) M1 for "10" × "15" M1 (dep on M2) for number of tiles ÷ 20, round up, × 8.5(0) A1 for 68 OR B1 for correct conversion of units M1 for 250 ÷ 20 (= 12.5) or 300 ÷ 25 (= 12) M1 for "12.5" × "12" M1 (dep on M2) for number of tiles ÷ 20, round up, × 8.5(0) A1 for 68 OR B1 for correct conversion of units M1 for 3 × 2.5 (= 7 .5) or 20 × 25 (= 500) M1 for "3 × 2.5" ÷ "0.2 × 0.25" or "300 × 250" ÷ "20 × 25" M1 (dep on M2) for number of tiles ÷ 20, round up, × 8.5(0) A1 for 68
*6		1, 4, 7, 10, 13 8, 6, 4, 2, 0	Explanation	2	M1 for listing at least 3 terms of both sequences C1 for Yes and explanation from fully correct working that 4 is in both sequences; numbers in A are increasing; numbers in B are decreasing

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
7			7.21 (am)	3	M1 for listing multiples 9,18,27,36 and 12,24,36 (condone 1 arithmetic error) or method to find LCM M1 for identifying 36 as LCM A1 cao OR M1 for listing times 6.54, 7.03, 7.12, 7.21 or for listing times 6.57, 7.09, 7.21 (condone one arithmetic error) M1 for listing times 6.54, 7.03, 7.12, 7.21 **and** 6.57, 7.09, 7.21 (condone one arithmetic error) A1 cao
8	(a)		$4(3e+1)$	1	B1 cao
	(b)		$15c - 10d$	1	B1 cao
	(c)		$28a^4b^3$	2	B2 for $28a^4b^3$ (B1 for two of 28, a^4 , b^3 as a product)
	(d)		$(x + 7)(x - 7)$	1	B1 cao
	(e)	$2y^2 - 6y + 7y - 21$	$2y^2 + y - 21$	2	M1 for 3 out of no more than 4 terms correct with correct signs or the 4 terms $2y^2$, $6y$, $7y$ and 21 seen, ignoring signs A1 cao

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
9		360 – 64 140 + 40 + 296 + 40 = 516 540 − 516 = 24	24°	4	M1 for 1 correct relevant angle calculation A1 for 24 cao C2 for all reasons (C1 (dep M1) for reason relating to parallel lines or parallelogram which is relevant to their chosen method) eg angles on a straight line add to 180° angles at a point add to 360° angles in a triangle add to 180° angles in a quadrilateral add to 360° alternate angles are equal corresponding angles are equal allied angles / co-interior angles add up to 180° opposite angles in a parallelogram are equal
10			$4\frac{8}{35}$	3	M1 for converting both fractions to get a common denominator of a multiple of 35 with at least one correctly converted. M1 (dep on M1) for $3+\frac{"28"}{35}+\frac{"15"}{35}(=3\frac{43}{35})$ oe A1 for $4\frac{8}{35}$ cao

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
11	(a)		7.8×10^{-3}	1	B1 cao
	(b)		6 710 000	1	B1 cao
	(c)	$-9,\ 0.9,\ 1,\ 3$	$-9,\ 0.9,\ 9^0,\ 9^{\frac{1}{2}}$	2	M1 for either $9^{\frac{1}{2}}$ or 9^0 evaluated correctly A1 for fully correct list from correct working, accept original numbers or evaluated. (SC: B1 for 1 error in position or a correct list in reverse order)
12.	(a)		(0, 5, 5)	1	B1 cao
	(b)		$(2\frac{1}{2}, 5, 2\frac{1}{2})$	1	B1 for $(2\frac{1}{2}, 5, 2\frac{1}{2})$ oe
13.			120	3	M1 for $80\% = 480$ or $\frac{480}{80}(=6)$ or $\frac{480}{80} \times 100\ (=600)$ oe M1 for $600 - 480$ or $\frac{480}{80} \times 20$ oe A1 cao
14.			$A = 9x^2 + 19x - 6$	4	B1 for one of $5x - 2$ or x found M1 for correct method to find area of one relevant rectangle. M1 for complete method to find whole area or simplified expression $9x^2 + 19x - 6$ or correct but not simplified formula A1 for correct, simplified formula $A = 9x^2 + 19x - 6$

PMT

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
15.			$y = \frac{3}{2}x - \frac{5}{2}$	4	M1 for attempt to find gradient of AB M1 (dep) for attempt to find gradient of perpendicular line eg use of $-\frac{1}{m}$ M1 for substituting $x = 3, y = 2$ into $y =$ "m" $x + c$ A1 for complete correct equation $y = \frac{3}{2}x - \frac{5}{2}$ oe
16.			$2\sqrt{7}$	3	M1 for multiplying numerator and denominator by $\sqrt{7}$ M1 for correct method to expand $(4+\sqrt{2})(4-\sqrt{2})$ with 3 out of no more than 4 terms correct with correct signs or the 4 terms seen, ignoring signs A1 for $2\sqrt{7}$ (accept $\sqrt{28}$)
17.		$\frac{(x-5)(x+3)}{(x+3)(2x+1)}$	$\frac{x-5}{2x+1}$	3	M1 for correct factorisation of numerator M1 for correct factorisation of denominator A1 for $\frac{x-5}{2x+1}$

PMT

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom

Question 9

Many students gained at least two of the three marks for converting both mixed numbers into improper fractions (with at least one correct) and then multiplying correctly. The requirement for the answer to be given as a mixed number in its simplest form meant that the final accuracy mark was often not awarded. Answers of $5\frac{6}{10}$ and $\frac{28}{5}$ were common. Some answers were spoilt by arithmetic errors. Having converted both mixed numbers into improper fractions some students chose to use a common denominator of 10. Unfortunately $\frac{35}{10} \times \frac{16}{10}$ was often followed by $\frac{560}{10}$ or by $\frac{51}{10}$ rather than by $\frac{560}{100}$. A few students decided to use decimals and they were often successful. When no marks were scored this was usually because students had chosen to multiply the whole numbers and fractions separately, giving an answer of $3\frac{3}{10}$.

Question 3

Although a correct mixed fraction was found by many students, a large number still find great difficulty in working with fractions. The less able students, instead of working with "top heavy" fractions, split the fractions to multiply them and working of 1 (1×1) and $\frac{3}{12}$ $(3 \times \frac{1}{4})$ was sometimes seen. Other students tried to add the fractions with $\frac{21}{12} + \frac{16}{12} = \frac{37}{12}$ regularly seen. Many answers were left as 28/12, when the question clearly stated that a mixed number was required.

Question 15

Students are familiar with this type of question and at least one mark was usually gained for showing an understanding of the meaning of the recurring decimal notation, although 0.418418…was not uncommon. Many identified two decimals with a terminating decimal difference but were unable to accurately find the difference. Any subsequent incorrect cancelling of $\frac{414}{990}$ was ignored.

Question 14

This question was found by many to be challenging and some did not know how to start. A number of students simply added the numbers in all three ratios together to obtain 8/19, showing no understanding of what was expected.

One of the approaches which gained marks was by choosing a starting number and dividing it in the correct ratios. An alternative approach seen was a tree diagram, often with the first process mark being awarded for use of fractions for either soup (2/5) or Prawns (3/5), however, many students did not know what to do next after this point. The most common incorrect answer for this question, when working with fractions was seen, was to work only using the curry part of each ratio ($3/8 \times 5/6$) and ignoring the relationship to the choice of starter.

Question 2

This question was generally answered well. The question required students to show that the answer is $8\frac{3}{4}$ so it was necessary for each stage of working to be shown. Many students gained at least two of the three marks for converting both mixed numbers into improper fractions (with at least one correct) and then multiplying correctly. It was pleasing to see some students making the arithmetic more straightforward by simplifying $\frac{7}{3}\times\frac{15}{4}$ to $\frac{7}{1}\times\frac{5}{4}$. Those students that wrote $\frac{7}{3}\times\frac{15}{4}=\frac{105}{12}$ had to simplify $\frac{105}{12}$ to $\frac{35}{4}$ or write it as $8\frac{9}{12}$ in order to complete the working and gain the final mark.
Having converted both mixed numbers into improper fractions some students made the arithmetic unnecessarily complicated by choosing to use a common denominator of 12. Sometimes these students multiplied the numerators but left the denominator as 12. Some students wrote both mixed numbers as improper fractions and made no further progress. When no marks were scored this was often because students had attempted to multiply the whole numbers and the fractions separately.

Question 23

Many students made a successful start by showing the fractions $\frac{3}{8}$ and $\frac{7}{9}$ and gained the first mark. However, a very common incorrect next step was to multiply the two fractions and give an answer of $\frac{21}{72}$. Some students did divide $\frac{3}{8}$ by $\frac{7}{9}$ to get $\frac{27}{56}$ and scored full marks. A more common route to the correct answer was for students to recognise the significance of 72 and write $\frac{3}{8} = \frac{27}{72}$ and $\frac{7}{9} = \frac{56}{72}$ or use

the ratios 3 : 5 and 2 : 7 to get 27 : 45 and 16 : 56. Some students could not then make the final step to $\frac{27}{56}$. The final mark was lost if the answer was given as a ratio and not as a fraction.

Summary

Based on their performance on this paper, students should:

- practise finding missing angles using angles on parallel lines and giving correctly worded reasons
- practise their arithmetic skills, particularly division and operations with decimals
- take care when interpreting the scales on the axes of graphs
- present their working clearly and in an organised way on the page so that the order of the process of solution is clear and unambiguous.

Question 3

This question was generally answered quite well. The most common method seen was to convert both mixed numbers into improper fractions and then write both fractions over a common denominator. Many students gained at least two of the three marks for getting this far. The accuracy mark was often lost because students failed to write their answer as a mixed number or made an arithmetic error. Having written both mixed numbers as improper fractions some students made no further correct progress and gained one mark only. Far fewer students chose to subtract the whole numbers and deal with the fractional parts separately. The difficulty with this method was subtracting a larger fraction from a smaller fraction which meant that $4\frac{3}{15}-2\frac{10}{15}$ was often followed by an answer of $2\frac{7}{10}$ or $2\frac{-7}{10}$.

Question 13

Some good evaluations of Ted's method were seen. Successful students often explained that Ted should have used $100x$ and x or that he should have used $1000x$ and $10x$. Some students gave acceptable explanations that focused on the recurring numbers after the decimal points.
Any numerical examples given as part of an explanation needed to be correct for the mark to be awarded. This was often not the case. Some answers consisted only of a correct numerical method showing that the recurring decimal could be written as the fraction $\frac{43}{99}$ and these were accepted. Some answers, such as "he should have used $100x$" or "he needs to find $1000x$", did not go quite far enough. Many answers were simply incorrect. A common misconception was that Ted had made a multiplication error because the value he found for $10x$ was wrong. A significant number of answers consisted of a commentary of what Ted was doing but no evaluation of his method.

Question 12

This is a familiar type of question and most students were able to gain the first mark for showing an understanding of the recurring decimal notation. It was pleasing that many students were able to show a complete method leading to a correct fraction. Any subsequent incorrect cancelling of $\frac{116}{990}$ was ignored. Some students got as far as $\frac{11.6}{99}$ but were then unable to complete the method to arrive at a correct fraction. After finding two appropriate decimals to subtract some students spoilt their solution by making careless arithmetic errors such as $1000x - 10x = 900x$. Some students could recall the need to multiply the recurring decimal by powers of ten but were either unable to find the multiples needed to eliminate the recurring nature of the decimal or could not carry out the multiplications correctly. It was common, for example, to see $10x = 1.\dot{1}\dot{7}$ followed by $100x = 11.1717..$ or by $100x = 117.\dot{1}\dot{7}$. Some students did not understand the recurring notation, writing 0.117117... or 0.171717...

Question 2

Part (a) was generally answered well. The most common method was to convert both mixed numbers into improper fractions and then write both fractions over a common denominator. Many students gained at least one of the two marks for getting this far. The accuracy mark was often lost because students failed to write their answer as a mixed number or made an arithmetic error. Fewer students chose to add the whole numbers and deal with the fractional parts separately but those that did were often successful. Considering that this is a familiar question early in the paper, a surprising number of students were unable to achieve both marks.

A similar proportion of students gained full marks in part (b) as in part (a) but a greater number of students gained no marks. Students who showed a correct method usually went on

to get full marks. Most of the successful students started with $\frac{8}{3}\times\frac{1}{6}$ and evaluated this as $\frac{8}{18}$ or cancelled to get $\frac{4}{3}\times\frac{1}{3}$. Some students answered the question by showing that $2\frac{2}{3}\div\frac{4}{9}=6$ or $\frac{4}{9}\times 6=2\frac{2}{3}$ but these approaches were rare.

Question 6

In order to make progress in part (a)(i) students needed to associate corresponding parts from the two ratios. Those that wrote the ratio 1 : 3 as 2 : 6 were often able to go on and write down the ratio $a : b : c$ as 2 : 6 : 5 although 2 : 6 : 10 was quite common. Some used a common multiple of 3 and 6 to get a common element representing the value of b and 6 : 18 : 15 was a common correct answer. Many students did not understand how to combine the two ratios. A very common incorrect answer was 1 : 9 : 5 from students simply adding the two values of b.

In part (a)(ii) many students gained both marks for expressing a as a fraction of their three numbers a, b and c. When students gained one mark only this was usually because they made an arithmetic error when adding the values of a, b and c or they gave an answer such as $\frac{1}{6.5}$ where the denominator is not an integer. Students who had not given a correct ratio in (a)(i) could still gain both marks in this part for writing a correct fraction for their values of a, b and c.

Students found part (b) quite challenging. After writing $p = 10m$ or $m = \frac{n}{2}$ or $2m = \frac{p}{5}$ and gaining the first mark many students did not find a correct ratio. The statement $p = 10m$ was quite often followed by $m : p = 10 : 1$. By leaving their ratio as $\frac{n}{2} : 5n$ some students failed to achieve the final mark. A few students answered the question by assigning values in the ratio given and were often successful. A large proportion of the students were unable to make any meaningful progress with this question.

Question 2

This question was answered very well. The most common method was to convert both mixed numbers into improper fractions and then write the fractions over a common denominator. Many students gained at least two of the three marks for getting this far. The accuracy mark was often lost because students failed to write their answer as a mixed number or they made an arithmetic error. Many of the arithmetic errors occurred when converting the mixed numbers to improper fractions. Some students used 16 rather than 8 as the common denominator which increased the chances of them making an arithmetic error. Having written both mixed numbers as improper fractions some students made no further correct progress and gained one mark only. Far fewer students chose to subtract the whole numbers and deal with the fractional parts separately. The difficulty that some had with this method was subtracting a larger fraction from a smaller fraction which meant that $7\frac{3}{8}-2\frac{4}{8}$ was often not evaluated correctly.

Question 13

In part (a) the students who worked out the frequency densities and used these to draw a histogram often drew five correct bars. This alone was not sufficient for full marks because the two axes needed to be correctly scaled and labelled. Some arithmetic errors such as $22 \div 5 = 4.5$ were made when working out the frequency densities. Students should be encouraged to show the method they use to work out each frequency density so that examiners can see if incorrect values come from a correct method of dividing frequency by class width. Some students made an error drawing one of the bars but still gained two of the three marks. A few students with the correct frequency densities did not draw the bars in the correct intervals and gained one mark only. Weaker students often did not recognise the need to find frequency densities and drew a bar chart or a frequency polygon or a cumulative frequency graph.

A similar proportion of students gained full marks in part (b) for working out an estimate for the fraction of the 150 people who were in the shop for between 20 minutes and 40 minutes. As this can be done by using the table or by using the histogram some students who had gained no marks in part (a) were able to give a correct fraction in part (b) by using the table. The marks were also awarded in part (b) for answers which followed through correctly from incorrect histograms. A common mistake was to give the answer as 67, not as a fraction, and this gained one mark only.

Question 3

Part (a) was generally answered well. The most common method was to convert both mixed numbers into improper fractions and then write both fractions over a common denominator. Many learners gained at least one of the two marks for getting this far. The accuracy mark was sometimes lost because learners failed to write their answer as a mixed number or made an arithmetic error. Some chose to subtract the whole numbers and deal with the fractional parts separately and those that did were usually successful.

A slightly smaller proportion of learners gained full marks in part (b) compared to part (a) but most gained at least one mark. The question required learners to show that the answer to the division is $2\frac{1}{4}$ so it was necessary for each stage of the method to be shown. Many gained at least two of the three marks for converting both mixed numbers into improper fractions (with at least one correct) and then showing a method to divide by a fraction. It was pleasing to see some learners making the arithmetic more straightforward by simplifying $\frac{21}{4}\times\frac{3}{7}$ to $\frac{3}{4}\times\frac{3}{1}$ before multiplying. Those that wrote $\frac{21}{4}\times\frac{3}{7}=\frac{63}{28}$ had to simplify $\frac{63}{28}$ to $\frac{9}{4}$ or write it as $2\frac{7}{28}$ in order to complete the working and gain the final mark. Having converted both mixed numbers into improper fractions some learners made the arithmetic unnecessarily complicated by choosing to use a common denominator of 12. Some wrote both mixed numbers as improper fractions and made no further progress; some flipped the wrong fraction or flipped both fractions. When no marks were scored this was often because learners had attempted to multiply the whole numbers and the fractions separately

Question 13

Converting a recurring decimal to a fraction is a familiar type of question but here candidates had two recurring decimals to deal with. However, this was a well answered question with many responses gaining 4 or 5 marks. Many responses gained the first method mark for the start of a method to convert $0.\dot{2}$ to a fraction. Some knew that $0.\dot{2} = \frac{2}{9}$ without needing to show a method. A method to convert $0.6\dot{8}\dot{1}$ to a fraction was needed for the second method mark and many responses found two recurring decimals that lead to a terminating decimal difference using correct multiples of y. After finding two appropriate decimals to subtract some responses were spoilt by careless arithmetic errors such as $1000y - 10y = 900y$. Some responses could recall the need to multiply the recurring decimal by powers of ten but were either unable to find the multiples needed to eliminate the recurring nature of the decimal or could not carry out the multiplications correctly. Some did not interpret the recurring notation correctly and $y = 0.6\dot{8}\dot{1}$ was sometimes followed by $1000y = 681.681...$, for example. Many responses, however, did show a correct method to convert $0.6\dot{8}\dot{1}$ to a decimal and gained the second method mark. When correct fractions were found for both x and y the first of the two accuracy marks could be awarded. Candidates still had to multiply the two fractions, which earned the third method mark, and give the answer as a fraction in its simplest form. Some responses simplified $\frac{675}{990}$ before multiplying, others wrote the multiplication first and

simplified later. Arithmetic errors were often made when simplifying and these errors meant that many responses did not reach an answer of $\frac{5}{33}$ and so could not be awarded the final accuracy mark. In some responses, the fraction had been cancelled correctly but not completely. A few responses made it more difficult for themselves by putting the fractions over a common denominator before multiplying. Some responses attempted to add the fractions rather than multiply them.