Chhetri Academy

chhetriacademy.com

GCSE MATHS

Number

Edexcel 1MA1 | Subtopic collection Topic 05

Indices and roots

Higher tier

Answers and examiner comments

32	2003-2025	Higher
question groups	available papers	tier

Using this booklet

1. Answer numbers match the companion question booklet.
2. Follow the original question number on marking tables.
3. Examiner comments are included where available.

Name _____ **Date** _____

Some entries use full source marking material in an appendix. Any unavailable mark schemes are identified beside the relevant answer.

Original question 17

No matching mark scheme was available in the supplied archive for this question.

Original question 18

The full source mark scheme begins on page 50. Use the original paper code, tier and question number shown above.

Original question 21

No matching mark scheme was available in the supplied archive for this question.

Original question 10

The full source mark scheme begins on page 101. Use the original paper code, tier and question number shown above.

Original question 16

No matching mark scheme was available in the supplied archive for this question.

24	(i)		1	1	B1 cao
	(ii)		8	1	B1 for 8 or -8 or $\pm$ 8
	(iii)		$\frac{4}{9}$	2	M1 for $\left(\frac{8}{27}\right)^{\frac{2}{3}}$ oe or $\left(\frac{3}{2}\right)^{-2}$ oe or $\left(\frac{2}{3}\right)^{2}$ oe or $\left(\frac{1}{\sqrt[3]{27/8}}\right)^{2}$ or better or $\frac{9}{4}$ oe seen A1 cao

1380_3H
Summer 2010

22	(a)	p^{5+4}	p^9	1	B1 (accept p^{5+4})
	(b)	q^{5-2}	q^3	1	B1 (accept q^{5-2})
	(c)		$2u$	2	B2 (accept $2t^0u$, $2t^0u^1$oe) (B1 for 2 correct terms from 2, t^0 and u oe eg u^1)
	(d)		$3wy^3$	2	B2 cao (B1 for 2 correct terms from 3, w and y^3 oe) NB: accept w^1 for w.
	(e)		x^{-2} x^0 $x^{\frac{1}{2}}$ x x^2	2	B2 cao (B1 for any 4 in relative correct order, or all correct but in reverse order)

1380_4H

Question		Working	Answer	Mark	Notes
11	(a)		1	1	B1 cao
	(b)		3	1	B1 cao
	(c)		$\frac{1}{8}$	1	B1 for $\frac{1}{8}$ oe

9	(a)		a^9	1	B1 for a^{4+5} or a^9
	(b)		$9e^5f^6$	2	B2 cao (B1 for two of 9, e^{6-1}, f^{8-2} as a product)
	(c)		3	1	B1 (accept ± 3 but not just –3)

1MA0_1H					
Question		**Working**	**Answer**	**Mark**	**Notes**
*10		Angle *AED* = 38 alternate angles are equal Angle *ADE* = (180 – 38) ÷ 2 = 71 *x* = 180 – 71 base angles of an isosceles triangle are equal angles in a triangle add up to 180 angles on a straight line sum to 180 **OR** angle *AEF* = 142 allied angles/co-interior angles add up to 180 *ADE* = 142 ÷ 2 = 71 base angles of an isosceles triangle are equal exterior angle of a triangle is equal to the sum of the interior opposite angles , *x* = 180 – 71 angles in a straight line add to 180 **OR** Angle *AED* = 38 alternate angles are equal for angles *BAE* and *AED* **and** *BAD* and *ADC* (*x*) Angle *DAE*= (180 – 38) ÷ 2 = 71 base angles of an isosceles triangle are equal angles in a triangle add up to 180 **Or** Angle *AED* = 38 alternate angles are equal Angle *ADE* = (180 – 38) ÷ 2 = 71 base angles of an isosceles triangle are equal and angles in a triangle sum to 180 *x* = 38 + 71 alternate angles *BAD* and *ADC*(*x*) are equal	*x* = 109	4	B1 for angle *AED* = 38 or *AEF* = 142 M1 for a complete method to find one of the base angles of the isosceles triangle C2 (dep M1) for *x* = 109 with complete reasons (C1 (dep M1) for one reason correctly used and stated)

1MA0_1H

17	(a)		$\frac{1}{10}$	1	B1 for 0.1 or $\frac{1}{10}$ oe
	(b)		9	2	M1 for $(\sqrt[3]{27})^2$ or $\sqrt[3]{27^2}$ oe or $\sqrt[3]{27} = 3$ A1 cao
	(c)	$\sqrt{75} = \sqrt{25}\sqrt{3}$	$5\sqrt{3}$	2	M1 for $\sqrt{25 \times 3}$ or $\sqrt{25}\sqrt{3}$ oe A1 cao

2	(a)		5^{10}	1	B1 cao
	(b)		7^3	1	B1 cao

PAPER: 1MA0_1H

Question		Working	Answer	Mark	Notes
22	(a)		1	1	B1 cao
	(b)		$\frac{5y}{8x^3}$	2	M1 for correct square root or correct use of reciprocal eg $\frac{8x^3}{5y}$ **or** $\frac{25y^2}{64x^6}$ A1 for $\frac{5y}{8x^3}$ **or** $\frac{5}{8}yx^{-3}$ oe
	(c)		$\frac{x+27}{(x-3)(x+3)}$	3	M1 for denominator $(x-3)(x+3)$ **or** x^2-9 M1 for $\frac{5(x+3)}{(x-3)(x+3)}$ oe **or** $\frac{4(x-3)}{(x-3)(x+3)}$ oe (NB The denominator must be $(x-3)(x+3)$ or x^2-9 **or** another suitable common denominator) A1 for $\frac{x+27}{(x-3)(x+3)}$ **or** $\frac{x+27}{x^2-9}$

PAPER: 1MA0_1H

10	(a)		1	1	B1
	(b)		$\frac{1}{16}$	1	B1
	(c)		$2\sqrt{7}$	2	M1 for multiplying numerator and denominator by $\sqrt{7}$ A1 for $2\sqrt{7}$

PAPER: 5MB2H_01

22	(a)		$\frac{1}{8}$	1	B1 for $\frac{1}{8}$ oe
	(b)		1.5	1	B1 for 1.5 oe
	(c)		3	2	M1 for √12×√12 – √12×√3 – √3×√12 + √3×√3 or √144 – √36 – √36 + √9 oe. with no more than one sign error A1 cao OR M1 for writing (√12 – √3) as (2√3 – √3) (= √3) A1cao

PAPER: 1MA0_1H

10	(a)		a^7	1	B1 cao
	(b)		b^{14}	1	B1 cao
	(c)		1	1	B1 cao
	(d)		$\frac{1}{4}$	1	B1 for $\frac{1}{4}$ oe

Original question 20

The full source mark scheme begins on page 153. Use the original paper code, tier and question number shown above.

12	(a)		$\frac{1}{9}$	M1	for showing a method using either reciprocal or square root e.g. $\frac{1}{n}$ or 9 seen
				A1	cao Accept $\pm\frac{1}{9}$ or 0.1 recurring
	(b)			M1	for showing cube root of 64 as 4 and the cube root of 125 as 5 or $\frac{16}{n}$ $(n \neq 25)$ or $\frac{n}{25}$ $(n \neq 16)$ or an intention to find the cube root and square.
			$\frac{16}{25}$		
				A1	cao Accept 0.64

17		Completes proof	M1 C1	Expands both expressions eg $\frac{1}{2}(n^2 + n + n^2 + n + 2n + 2)$ **or** $n^2 + n$ and $n^2 + n + 2n + 2$ **or** factorises $\frac{1}{2}(n+1)(n+n+2)$ Completes proof with explanation and reference to $(n+1)^2$

Paper: 1MA1/1H

Question	Working	Answer	Mark	Notes
6 (a)		6	B1	cao
(b)		5	B1	cao
(c)		Shown	M1	for writing 100^a or 1000^b as a power of 10 ($=10^{2a}$ or 10^{3b}) or 10^{2a+3b} or $100 = 10^2$ and $1000 = 10^3$
			C1	for complete chain of reasoning leading to conclusion

Paper: 1MA1/2H

19	$\frac{3}{4}$ oe	P1 P1 A1	for a first step to converting to a common base with one correct conversion, eg. $9^{-\frac{1}{2}} = 3^{-1}$ or $\frac{1}{3}$ or $27^{\frac{1}{4}} = 3^{\frac{3}{4}}$ oe (dep) for $3^{-1} = 3^{\frac{3}{4}} \div 3^{x+1}$ oe cao	$9^{-\frac{1}{2}} = 3^{-1}$ (or $\frac{1}{3}$) oe or $27^{\frac{1}{4}} = 3^{\frac{3}{4}}$ oe seen alone gets the P1

Paper: 1MA1/1H

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
11 (a)	300	M1	for working out $\sqrt[4]{81}$ as 3 **or** $\sqrt[4]{10^8}$ as 10^2 or 100	Mark may be awarded if operations are attempted on 8100000000 eg 300000000
		A1	for 300 **or** 3×10^2 or 3×100	
(b)	$\frac{1}{8}$	M1	for showing a square root of 64 as 8 **or** recognition of the reciprocal eg $\frac{1}{n}$ **or** shows expressions that show an understanding of the ½ index and the minus index eg $\frac{1}{\sqrt{64}}$ or other equivalent forms	
		A1	oe	Accept $\pm\frac{1}{8}$ oe
(c)	3^{2-n}	M1	for $3^{2(n-1)}$ **or** 3^{2n-2} **or** $\left(3^2\right)^{n-1}$	
		A1	for 3^{2-n} oe eg $3^{n-2(n-1)}$	

Paper: 1MA1/1H

17	$y = \frac{40}{\sqrt{x^3}}$	P1	for setting up an equation with a constant term, eg $y = k\sqrt{t}$ **or** $t = \frac{K}{x^3}$	Condone the use of 'α' instead of '=' for the first two P marks Equation can be implied by correct substitution
		P1	for a process to substitute values in one equation, eg $15 = k\sqrt{9}$ or $k = 5$ **or** $8 = \frac{K}{2^3}$ or $K = 64$	
		P1	(dep P2) for combining the two equations ft their values of k and K, eg $y = 5\sqrt{\frac{64}{x^3}}$ **OR** for $y = 5\sqrt{t}$ **and** $t = \frac{64}{x^3}$	
		A1	oe	Formula must include 40 Accept other forms for the power of x but must be a single term in x

Paper: 1MA1/1H


Paper: 1MA1/3H

Question	Answer	Mark	Mark scheme	Additional guidance
9 (a)	10^{60}	M1	for a correct first step using one of the rules of indices, eg. $10^{150} \times 10^{90} = 10^{240}$ or $10^{360} \div 10^{150} = 10^{210}$ or $10^{360} \div 10^{90} = 10^{270}$ or $\sqrt{10^{360}} = 10^{180}$ or $\sqrt{10^{150}} = 10^{75}$ or $\sqrt{10^{90}} = 10^{45}$	
		M1	for correct use of rules of indices leading as far as $\sqrt{10^{120}}$ or $\frac{10^{180}}{10^{120}}$	
		A1	cao	
(b)	reason	C1	for correct reasoning **Acceptable examples** eg should do 50×2 (not 50^2) because $(12^{50})^2 = 12^{100}$ because when you have a power inside and outside the bracket you times them because $(a^b)^c = a^{bc}$ (not a^{b^c}) **Not acceptable examples** because you need to multiply everything in the brackets by 2 because he should have squared 12 as well you add the powers instead of timesing	

Paper: 1MA1/3H

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
3	2^6	M1	for the start of a method of simplification, eg 2^{-5+8} (= 2^3) or $2^{-5\times2}$ (= 2^{-10}) or $2^{8\times2}$ (= 2^{16})	
		A1	cao	
			SC B1 for answer of 64 or 8^2 or 4^3 if M0 scored.	

Question	Answer	Mark	Mark scheme	Additional guidance
5 (a)	Venn diagram	B3	for a fully correct Venn diagram	Ignore all entries except the region you are marking for each method mark
		(B2	for two or three of the four regions correct)	
		(B1	for just one of the four regions correct)	
(b)	$\frac{7}{10}$	M1	(ft diagram) for $\frac{a}{10}$ where $0 < a < 10$ and a is an integer or $\frac{7}{b}$ where $b > 7$ and b is an integer **or** $1 - \frac{3}{10}$ **or** 7 : 10	Repeated digits in the diagram should be counted as 2 elements
		A1	(ft diagram) for $\frac{7}{10}$ oe	Accept any equivalent fraction, or 0.7 or 70%

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
14 (a)	4^{-2}	B1	for 4^{-2}	Accept $n = -2$
(b)	5	M1	for $8^{\frac{5}{3}} = \left(\sqrt[3]{8}\right)^5$ or 2^5 or $\sqrt[3]{8^5}$ or $\sqrt[3]{32768}$ **or** $9^{\frac{3}{2}} = \left(\sqrt{9}\right)^3$ or 3^3 or $\sqrt{9^3}$ or $\sqrt{729}$	
		M1	for correctly evaluating $8^{\frac{5}{3}}$ **or** $9^{\frac{3}{2}}$, eg 32 or 27 seen	
		A1	cao	

13	Proof	M1 M1 C1	for giving a general term for a consecutive triangular number eg $\frac{(n+1)(n+2)}{2}$ or $\frac{(n-1)n}{2}$ for $\frac{n(n+1)}{2}+\frac{(n+1)(n+2)}{2}$ or $\frac{(n-1)n}{2}+\frac{n(n+1)}{2}$ oe for completing the proof to show a square number, eg $(n+1)^2$ or n^2	

2	(a)(i)	1	B1	cao	
	(ii)	$\frac{1}{25}$	B1	oe	
	(b)	2^6	M1	for a correct first step using a rule of indices, eg 2^{5+4} (= 2^9) or 2^{5-3} (= 2^2) or 2^{4-3} (= 2^1) **or** for 2×2×2×2×2×2 or 64	
			A1	for 2^6	Accept $n = 6$

Paper: 1MA1/1H

<table>
<tr>
<td>16</td>
<td>$\frac{1}{120}$</td>
<td>P1

P1

A1</td>
<td>for correctly identifying at least two of the number of options,
eg 2 options for 1st digit, 4 options for 2nd digit

for process to multiply the four numbers of options with at least three correct,
eg $2 \times 4 \times 3 \times 5$ (= 120)

or

for process to multiply 4 fractions with at least 3 correct $\frac{1}{2} \times \frac{1}{4} \times \frac{1}{3} \times \frac{1}{5}$ oe

for $\frac{1}{120}$ oe</td>
<td>May be seen as part of a calculation

<table>
<tr><th>Position of digit</th><th>Number of options</th></tr>
<tr><td>1st</td><td>2</td></tr>
<tr><td>2nd</td><td>4</td></tr>
<tr><td>3rd</td><td>3</td></tr>
<tr><td>4th</td><td>5</td></tr>
</table>
</td>
</tr>
</table>

Paper: 1MA1/3H

Question	Answer	Mark	Mark scheme	Additional guidance
13 (a)	No with reason	C1	for No with reason, **Acceptable examples** No, $4 \div 4 \neq 8 \div 6$ No, $4 \div 4 \neq 8 \div 6$ so they don't have the same constant No because $y = kx$ doesn't work No, it wont pass through (0, 0) **Not acceptable examples** Yes, ... No, they don't increase by the same number No, they don't have the same constant	
(b)(i)	49	M1	for stating a correct relationship, eg $w = k \times \sqrt{t}$ **or** $140 = k \times \sqrt{64}$ oe	Condone the use of 'α' instead of '=' for both M marks
		M1	for method to find constant of proportionality, eg $140 \div \sqrt{64}$ (= 17.5 oe)	Award of this mark implies the previous mark
		A1	cao	
(ii)	sketch	C1	for sketch	Sketch at end of mark scheme

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
11	125	P1	for setting up an equation with a constant term, eg $T=\frac{k}{w}$ or $w=K\sqrt[3]{d}$	Condone the use of 'α' instead of '=' for the first two P marks Equation can be implied by correct substitution
		P1	for a process to substitute values in one equation, eg $20=\frac{k}{6}$ or $k = 120$ or $1=K\sqrt[3]{8}$ or $K=\frac{1}{2}$ oe	
		P1	(dep P2) for combining the two equations ft their values of k and K, eg $T=\frac{"120"}{"\frac{1}{2}"\sqrt[3]{d}}$ oe **OR** for a correct process to find the value of w when $T = 48$, eg $w=\frac{"120"}{48}$ (= 2.5 oe)	$T=\frac{120}{w}$ $w=\frac{1}{2}\sqrt[3]{d}$
		P1	for substitution into their formula, eg $48=\frac{"120"}{"\frac{1}{2}"\sqrt[3]{d}}$ **OR** for substitution of found value of w in the equation for d, eg $"2.5"="\frac{1}{2}"\sqrt[3]{d}$	
		A1	cao	

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
11 (a)	2	M1	for a correct start, using one rule of indices, eg $3^n \times 3^{12} = 3^{14}$ or $3^n \times 3^{20} = 3^{22}$ or $3^{20} \div 3^8 = 3^{12}$ or $3^{14} \times 3^8 = 3^{22}$ or $\frac{3^{n+20}}{3^8} = 3^{14}$ or $3^{n-8} \times 3^{20} = 3^{14}$ **or** for forming an equation in n, eg $n + 20 - 8 = 14$ oe or (n =) $14 + 8 - 20$	
		A1	cao SCB1 for an answer of 3^2 if M0 scored	Allow an answer of $3^n = 3^2$ An answer of 9 or $3^n = 9$, $n \neq 2$ on its own is to be awarded 0 marks
(b)	$16m^8$	M1	for an intention to find the cube root and square, eg $\sqrt[3]{64m^{12}}^2$ or $\sqrt[3]{\left(64m^{12}\right)^2}$ or $\left(4m^4\right)^2$ or $\sqrt[3]{4096m^{24}}$ or for am^8 with $a \neq 16$ or $16m^b$ with $b \neq 8$	Do not condone missing brackets 16 or m^8 imply M1 Allow multiplication sign for M1
		A1	cao	Accept $a = 16$, $b = 8$

Paper: 1MA1/3H

edexcel

GCSE
Edexcel GCSE
Mathematics A 1387

July 2004

advancing learning, changing lives

Mark Scheme

Edexcel GCSE
Mathematics A 1387

London Qualifications is one of the leading examining and awarding bodies in the UK and throughout the world. It incorporates all the qualifications previously awarded under the Edexcel and BTEC brand. We provide a wide range of qualifications including general (academic), vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, our centres receive the support they need to help them deliver their education and training programmes to learners.

For further information please call Customer Services on 0870 240 9800, or visit our website at www.edexcel.org.uk

July 2004

Publications Code UG015616

All the material in this publication is copyright
© London Qualifications Ltd 2004

CONTENTS

Notes on Marking Principles....2
Paper 5514....4
Paper 5515....8
Paper 5516.... 11
Paper 5517.... 15
Paper 5518.... 19
Paper 5519.... 23

NOTES ON MARKING PRINCIPLES

NOTES ON MARKING PRINCIPLES

1 Three types of mark are available

M marks - awarded for correct working seen or implied.
A marks - conditional accuracy marks which are awarded for accurate working following the award of M marks.
B marks - unconditional accuracy marks (independent of M).

2 Abbreviations

cao - correct answer only.
ft - follow through.
$\surd$ - Denotes a "follow through" answer.

SC - special case.
isw - ignore subsequent working.
oe - or equivalent (and appropriate).
NB: a candidate cannot benefit from both isw and ft.

3 If no working shown, then

correct answers normally score full marks
incorrect (even though nearly correct) answers score no marks.

4 Marking instructions

Misread loses A marks (and sometimes B marks) on that part but ft can be allowed on subsequent parts. M marks can still be earned. If in doubt contact your team leader.
If there is a wrong answer in the answer space DO CHECK the working in the body of the script.

- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.
- If there is a wrong answer in the answer space DO CHECK the working in the body of the script.
- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.

5 Style of marking

Answer correct: tick and write part mark in margin NEXT TO BRACKETED MARK.
Answer incorrect: cross, but show M, A or B marks if any earned in body of script and transfer the total of these to the margin next to the bracket mark.
Total for each double page at bottom right page (except for back if used), FINAL TOTAL IN RELEVANT BOX ON FRONT COVER.
Nought in margin for fully incorrect question or page AND FOR NO ATTEMPT.
Where no attempt has been made a line should be put in the answer space and zero in the margin next to the bracketed mark
CHOICE OF METHOD – No marks unless one answer is in answer space – then mark that.
CROSSED OUT WORK - if not replaced this should be marked (if legible).
There must always be a mark next to bracketed mark in the margin/

6 Follow Through Marks

Follow throughs are guided by two principles:

(a) Follow throughs which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous, do not award.
(b) Follow throughs which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

7 Probability

Probability answers must be given as fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least two decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.

6 Linear Equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

8 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

PAPER 5501					
No		**Working**	**Answer**	**Mark**	**Notes**
1			K: 1020 L: 8.06	2	B1 for 1020 or 1,020 B1 for 8.06 Accept £8.06p and £8,06
2	(a)		28, 33	2	B1, B1 (B1 ft for "28" + 5 if both numbers >25)
	(b)			1	B1 for add 5, +5, for going up in 5's the difference is 5 oe
	(c)			1	B1 for they end in 3 and 8, it ends in 7; or refers to 383 and 388 are in sequence
3	(a)(i)		$\frac{1}{2}$	2	B2 for ½ accept half (B1 for an equivalent unsimplified fraction eg 4/8 or 50% or 0.5)
	(ii)		2 rectangles shaded	1	B1 for correct shading (any 2 rectangles)
	(b)(i)		40	4	B1 for 40 cao
	(ii)		150 000		B1 for 150 000 cao (accept 150,000 not 150.000)
	(iii)		6.55		B1 for 6.55 cao not $6.5^{½}$
	(iv)		$\frac{3}{8}$		B1 for $\frac{3}{8}$ oe accept 0.375
	(c)		Cross 3cm from *A*	1	B1 mark a cross 3 cm (±2 mm) from *A*
4	(a)(i)		metres	2	B1 for metres (m)
	(ii)		grams		B1 for grams (g)
	(b)		miles	1	B1 for miles
5	(a)		(3, 2)	1	B1 for (3, 2)
	(b)(i)		*Q* at (0, 3)	1	B1 for *Q* plotted correctly on *y*-axis at (0, 3) ±2mm
	(ii)		*R* at (−2, −3)	1	B1 for *R* plotted correctly at (−2, −3) ±2mm
6	(a)(i)		30	2	B1 for 30
	(ii)		25		B1 for 25
	(b)(i)		4 shapes	2	B1 for drawing 4 shapes
	(ii)		1½ shapes		B1 for drawing 1½ shapes

PAPER 5501					
	No	**Working**	**Answer**	**Mark**	**Notes**
7	(a)		30	1	B1 for 30
	(b)	$20 \times 3 + 5$	65	2	M1 for $20 \times 3 + 5$ oe A1 for 65
8	(a)		Draws perp.	1	B1 for correctly drawing perp must touch line or cut line *AB* ± 2mm
	(b)		Sketches a cylinder	1	B1 for sketching cylinder
9	(a)	$\begin{array}{r} 1974 \\ \underline{6580} \\ 8554 \end{array}$	855.4	3	M1 for complete method with relative place value correct, condone 1 error in multiplication A2 cao (A1 for digits 8554 seen or A1 for "855.4" dependent on 1 arithmetic error)
	(b)	990 ÷ 69 = 14.3…. or 14 rem 24	14	4	Method 1 - Everything excluding long division M2 for a valid method with no errors …need to see 966 or 1035 (M1 for a valid method, including estimation, that reaches an answer between 901 and 1099 inclusive. Ignore errors.) A2 for 14 with a fully correct method (A1 (dep on 1st M1) for an answer between 14 and 15 inclusive. Ignore errors. Cannot have used estimation.) Note: Estimation can only score a maximum of M1 unless accompanied by another method or 966 or 1035 see (award M2) Method 2 – long division method M2 for 990 ÷ 69 = integer answer between 12 and 19 inclusive and rem 30 seen (M1 for 990 ÷ 69 with 1 in tens column.) A2 for 14 with fully correct method (A1 (dep on 1st M1) for an answer between 14 and 15 inclusive.)

PAPER 5501					
	No	**Working**	**Answer**	**Mark**	**Notes**
10	(a)(i)		7	2	B1 for 7
	(ii)		−10		B1 for −10
	(b)(i)		6	2	B1 for 6 (accept −6)
	(ii)		8		B1 for 8 (accept −8)
	(c)		−7	1	B1 for −7 cao
11	(a)		B and D	2	B1, B1 (–1 each extra)
	(b)(i)		A	2	B1 for A
	(ii)		3		B1 for 3
12	(a)	800 − 144	656	2	M1 for at least 1 digit correct and in correct position needs to be 3 digit number A1 for 656
	(b)	144 is less than 200 so Trudy is wrong		2	B1 for 200 B1ft for 'correct' explanation based on cand's "200"
	(c)	$\frac{45\times800}{100}$	360	2	M1 for 45 × 800 ÷ 100 oe A1 for 360
	(d)	$\frac{176}{800}\times100$	22	2	M1 for $\frac{176}{800}\times100$ oe A1 for 22
13	(a)(i)		25	4	B1 for 25
	(ii)		28		B1 for 28
	(iii)		5 and 20		B1 for 5 and 20
	(iv)		26 and 33		B1 for 26 and 33
	(b)	$2^3 = 8$ or $2^3 = 2 \times 2 \times 2$ (which is not 6)		1	B1 for valid 'explanation'
14			10 m^2	2	B1 for 10 B1 indep for m^2.

PAPER 5501					
No		**Working**	**Answer**	**Mark**	**Notes**
15	(a)	$\frac{11}{12}-\frac{10}{12}$	$\frac{1}{12}$	2	M1 for correctly writing both fractions to a common denominator A1 for $\frac{1}{12}$ oe
	(b)	$\frac{70\times400}{200}$	140	2	B2 for 140 (accept 136) (B1 for sight of any two of 70, 400 or 200)
16	(a)		$2y$	1	B1 for $2y$ or $2\times y$, $y2$, $y\times2$
	(b)		$3p^2$.	1	B1 for $3p^2$ or $3\times p^2$ or $p^2\times3$ or $p^2 3$
	(c)		$x(x-3)$	2	B2 for $x(x-3)$ (B1 for x (x…….)) $\frac{25}{60}$ wrongly cancelled gets B1 ISW
17	(a)		2 23 9 34 15 2 9 26 17 25 18 60	3	B3 for all correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
	(b)		$\frac{25}{60}$	1	B1 for $\frac{25}{60}$ or $\frac{5}{12}$ oe $\frac{25}{60}$ wrongly cancelled gets B1 ISW

PAPER 5501					
	No	**Working**	**Answer**	**Mark**	**Notes**
18	(a)(i)		60	2	B1 for 60
	(ii)				B1 for all **angles equal** so equilateral triangle oe
	(b)(i)		130	3	B1 for 130
	(ii)				B1 for isosceles triangle oe or 2 angles equal accept $\angle Q = \angle R$ B1 for angles on a straight **line** add up to **180°**oe (180° could be in working)
	(c)		64	1	B1 for 64
19	(a)		1	1	B1 for 1
	(b)		3	1	B1 for 3
	(c)	$9\times1+3\times2+5\times3+3\times4$ (= 42) "42" ÷ 20	2.1	3	M1 for 9×1, 3×2, 5×3, 3×4 or for 42 seen M1 (dep) for "42" ÷ 20 A1 for 2.1 or $2\frac{1}{10}$ or $2\frac{2}{20}$
20	(a)		09 05	1	B1 for 09 05 oe
	(b)		7	1	B1 for 7
	(c)		10	1	B1 for 10
	(d)	7km in 20 mins	21	3	B1 for 20 (minutes) oe or $\times3$ seen M1 for $\frac{"7"}{"20"}$ A1 cao [SC: "7" × 3 seen gets B1 M1]



PAPER 5501				
No	**Working**	**Answer**	**Mark**	**Notes**
21	Vertices of new triangle at (2, −2) (2, 6) and (6, −2)	Correct triangle drawn	3	B3 cao (B2 for either correct triangle in wrong translated position or for 2 vertices correct) (B1 a right angled triangle with horizontal length 4 or vertical length 8 in the same orientation as the shaded triangle) SC B2 for s + 3 totally correct
22 (a) (b) (c)	 8560 ÷ (10 × 10)	8 Pentagon 85.6	1 1 2	B1 for 8 B1 for pentagon M1 for 8560 ÷ (10 × 10) oe A1 for 85.6
23	$2x + x + 100 + 47 = 360$ $2x + x = 360 - 100 - 47$ $x = 71$ Largest angle = $2x$ =	142	4	M1 for $2x + x + 100 + 47 = 360$ or 360 – 147 or 213 seen M1 dep for correctly separating x-terms and non x terms or "360 – 147" ÷ 3 A1 for $x = 71$provided M2 awarded A1 ft for 142

PAPER 5502					
	No	**Working**	**Answer**	**Mark**	**Notes**
1	(a)		Diagram	1	B1 cao
	(b)		22, 26	1	B1 cao
	(c)		46	1	B1 cao
2	(a)	£5.95 + £1.62 + 25p + 2 × 48p = £8.78	1.22	3	M1 for 3 of £5.95, £1.62, 25p, 48p seen with intention to add or 170 seen A1 for £8.78 A1 ft from " 8.78"
	(b)	160 ÷ 12 = 13.(333…)	14	2	M1 for 160 ÷ 12, or 13 or better or 12 × 13 or 12 × 14 A1 14 cao
	(c)	"14" × £12.20 = £170.8**0**	170.80	2	M1 for "14" × 12.20 A1 for 170.8**0** or 158.6**0** if 13 in (b) NB: 170.8 or 158.6 gets M1 A0
3	(a)(i)		1459	1	B1 cao
	(ii)		9541	1	B1 cao
	(b)		9 + 5 = 14	1	B1 cao
	(c)		0	1	B1 cao
4	(a)		1123	1	B1 cao
	(b)		1125	1	B1 cao
	(c)		43	1	B1 cao
	(d)	67 – "43"	24	2	M1 for 67 – "43" A1 ft from "43"
	(e)(i)		80	2	B1 cao
	(ii)		56 – 58		B1 answer in range 56 to 58 inc
	(f)		Completes graph	2	B2 for (Apr , 70) and (May , 60 < p < 65) plotted and joined (B1 for one point plotted)

PAPER 5502					
	No	Working	Answer	Mark	Notes
5	(a)(i)		4	1	B1 cao
	(ii)		10	1	B1 cao
	(b)		28	2	B2 for 28 (B1 for 29 or 14×2)
6	(a)		$\frac{3}{4}$	1	B1 accept 0.75, 75%
	(b)	$\frac{3}{4} \times 28 = 21$	15.54	3	M1 for $\frac{3}{4} \times 28$ or 21 seen M1 for "21" $\times$ 74 or 28×74 or 7×74 A1 cao
	(c)(i)		44 – 46	2	B1 44 – 46 inc
	(ii)		5.2 – 5.4		B1 5.2 – 5.4 inc
7	(a)		Correct shape	1	B1
	(b)		Correct shape	1	B1
8	(a)		Hexagon	1	B1
	(b)		Right	1	B1 cao
	(c)		Drawing	2	B2 for 7 additional hexagons with at least 2 points at which 3 hexagons meet (B1 for one point at which 3 hexagons meet)
9	(a)		$\frac{3}{4}$	2	M1 $\frac{80-20}{80}$ or $\frac{60}{80}$ or 0.75 (oe) A1 cao Do not accept equivalents
	(b)		15	1	B1 cao
	(c)		Iron and Cook set	1	B1 for both Accept mixer and fryer
	(d)		Microwave	1	B1 cao
	(e)		Bar chart	2	B1 one column (mixer): 70, 90 B1 second column (fryer): 70, 85 [SC: B1 columns correct but reversed]

PAPER 5502					
	No	Working	Answer	Mark	Notes
10	(a)		Acute	1	B1
	(b)		Reflex	1	B1
	(c)		reason	1	B1 reason eg 120 + 230 = 350 ≠ 360
11	(a)		$\frac{1}{2}$	1	B1 for $\frac{1}{2}$ oe
	(b)		reason	1	B1 for reason e.g could get 30 heads
12	(a)	7.5 × 4	30	2	M1 for 7.5 × 4 or $7\frac{1}{2} \times 4$ A1 cao
	(b)	√(42.25)	6.5	2	B2 for 6.5 (B1 for 42.25)
13	(a)		$6x$	1	B1 accept $6 \times x$, $x6$ oe
	(b)		$x - 4$	1	B1 cao
	(c)		$12(x - 4)$	1	B1 for $12(x - 4)$ or $12x - 48$ or $12 \times x - 48$ oe
	(d)		$18x - 48$	2	M1 for "$6x$" + "$12(x-4)$" or "$6x$" + "$12x - 48$" oe A1 ft $18x - 48$, $2(9x - 24)$, $3(6x - 16)$, $6(3x - 8)$
14	(a)		90	2	B2 for 90 ± 2 (B1 for 9 ± 0.2)
	(b)		317	1	B1 for 317 ± 4
15	(a)	Accurate drawing of triangle		2	B1 cao for 38° (±2°) B1 cao (indep) for BC drawn 7.3 cm (±2mm) and completing the triangle.
	(b)	Measure angle A	63°	1	B1 for 63° (±2°) or ft their diagram (±2°)

PAPER 5502					
	No	Working	Answer	Mark	Notes
16	(a)	0 × 1 = 0 1 × 1 = 1 2 × 4 = 8 3 × 8 = 24 4 × 8 = 32 5 × 7 = 35	100	3	M1 col 1 × col 2 (at least 3 shown) M1 (dep) sum of totals A1 cao SC: M1 M1 A0 for 101
	(b)	Either 35.50 ÷ 1.42 = 25; 26.99 – 25 = 1.99 Cheaper in the USA Or 26.99 × 1.42 = 38.33; 38.33 – 35.50 = 2.83 Cheaper in the USA		3	M1 35.50 ÷ 1.42 A1 25 seen B1 ft (dep on M1) £1.99 with conclusion (must have units) **OR** M1 26.99 × 1.42 A1 38.32 or 38.33… seen B1 ft (dep on M1) $2.83 or $2.82 with conclusion (must have units) must be to 2dp
17	(a)		-1, (1), (3), 5, 7, 9	2	B2 cao (B1 for 2 values)
	(b)		Graph	2	B1 ft for plotting points ±1/2 square B1 cao for line between $x = -2$ and $x = 3$
	(c)(i)		0.4	2	B1 for 0.4 or ft from single straight line with positive gradient
	(ii)		1.2		B1 for 1.2 or ft from single straight line with positive gradient

PAPER 5502					
	No	Working	Answer	Mark	Notes
18	(a)		Plots	1	B1 cao
	(b)		description	1	B1 dynamic relationship or "positive" (correlation)
	(c)		line of best fit	1	Line within overlay region, and to the extent of.
	(d)(i)	280 g →	reading	2	B1 ft from single straight line of positive gradient (±1/2 square)
	(ii)	120 pages →	reading		B1 ft from single straight line of positive gradient (±1/2 square)
19			200 150 225 150 10	3	B3 cao (B2 for three correct, or B1 for one correct)
20	(a)			2	B1 B1 for correct sketch – ignore additional internal lines B1 B0 for rectangle or right-angled triangle or correct shape without line NB 3-D sketch gets B0 B0
	(b)			2	B1 B1 for correct plan – ignore internal lines – accept a rotated plan, not reflected B1 B0 for a single rectangle drawn NB 3-D sketch gets B0 B0
21	(a)	163.25 – 35.50 = 127.75 127.75 ÷ 18.25 = 7	8	3	M1 163.25 – 35.50 (or sight of 127.75) M1 (dep) "127.75" ÷ 18.25 A1 cao SC: M2 for 7 days
	(b)	1.175 × £64 or $\frac{17.5}{100}$ × 64 or £11.20 seen	75.20	2	M1 1.175 × 64, 0.175 × 64 oe or 11.20 or 75.2 seen or 6.40, 3.20 and 1.60 seen A1 cao

PAPER 5502					
	No	Working	Answer	Mark	Notes
22	(a)		6	1	B1 cao
	(b)		$t^2 - 2t$	1	B1 oe
	(c)		$3(y - 4)$	1	B1 Accept $3 \times (y - 4)$ or $3(y - 4$
23	(a)		Reason	1	B1 Makes some mention of bias either directly or making reference to an insufficient or biased range of responses
	(b)		Reason	1	B1 (a) an insufficient range of responses (b) No mention of money (c) No time frame in the question (d) Misunderstanding of "A lot" and "Not much"



PAPER 5503				
No	**Working**	**Answer**	**Mark**	**Notes**
1		10.	1	B1 for 10
2 (a)(i)		$3e + 2f$	2	B1 for $3e + 2f$
(ii)		$3p^2$		B1 for $3p^2$
(b)	$5 \times -3 + 1$	-14	2	M1 for $5 \times -3 + 1$ A1 for -14
3	1974 $\frac{6580}{8554}$	855.4	3	M1 for complete correct method with relative place value correct, condone 1 error in multiplication. A2 cao (A1 for digits 8554 seen or A1 ft for "855.4" dependent on one arithmetic error only)
4 (a)		2 23 9 34 15 2 9 26 17 25 18 60	3	B3 all correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
(b)		$\frac{25}{60}$	1	B1 for $\frac{25}{60}$ or $\frac{5}{12}$ oe
5 (a)	$\frac{45}{100} \times 800$	360	2	M1 for $\frac{45}{100} \times 800$ oe A1 for 360
(b)	$\frac{176}{800} \times 100$	22	2	M1 for $\frac{176}{800} \times 100$ oe A1 for 22

PAPER 5503					
	No	Working	Answer	Mark	Notes
6	(a)	$\frac{11}{12}-\frac{10}{12}$	$\frac{1}{12}$	2	M1 for correctly writing both fractions to a common denominator A1 for $\frac{1}{12}$ oe
	(b)	$\frac{70\times400}{200}$	140	2	B2 for 140 (accept 136) (B1 for sight of any two of 70, 400 or 200)
7	(a)(i)		60	2	B1 for 60
	(ii)				B1 for all angles equal so equilateral triangle oe
	(b)(i)		130	3	B1 for 130
	(ii)				B1 for isosceles triangle oe B1 for angles on a straight line add up to 180°oe
	(c)		64	1	B1 for 64
8		$9\times1+3\times2+5\times3+3\times4$ (=42) "42" ÷ 20	2.1	3	M1 for 9×1, 3×2, 5×3, 3×4 or for 42 seen M1 (dep) "42" ÷20 A1 for 2.1 or $2\frac{1}{10}$ or $2\frac{2}{20}$
9	(a)		09 05	1	B1 for 09 05 oe
	(b)		7	1	B1 for 7
	(c)		10	1	B1 for 10
	(d)	7 km in 20 minutes	21	3	B1 for 20 (minutes) oe or $\times3$ seen M1 for $\frac{"7"}{"20"}$ A1 cao SC: "7" $\times3$ gets B1 M1

PAPER 5503					
No		**Working**	**Answer**	**Mark**	**Notes**
10		Vertices of new triangle at (2, −2) (2, 6) and (6, −2)	Correct triangle drawn	3	B3 cao (B2 for either correct triangle in wrong translated position or for 2 vertices correct) (B1 for a right-angled triangle with horizontal length 4 or vertical length 8 in the same orientation as shaded triangle) SC B2 for scale factor 3, totally correct
11		$2^2 + 3 =$	7 which is not even	2	B2 (B1 for correctly evaluating $n^2 + 3$ with a prime number value for n.)
12	(a)	360 ÷ 5	72	2	M1 for 360 ÷ 5 oe A1 for 72
	(b)	8560 ÷ (10 × 10)	85.6	2	M1 for 8560 ÷ (10 × 10) oe A1 for 85.6
	(c)(i)	Least length = 100.5	100.5	2	B1 for 100.5
	(ii)	Greatest length = 101.5	101.5		B1 for 101.5 or 101.499 or better
13		$2x + x + 100 + 47 = 360$ $2x + x = 360 - 100 - 47$ $x = 71$ Largest angle = $2x$ =	142	4	M1 for $2x + x + 100 + 47 = 360$ or 360 – 147 or 213 seen M1 (dep) for correctly separating x-terms and non x terms or $\frac{"360-147"}{3}$ A1 for $x = 71$ A1 ft (provided M2 awarded) for "142"
14	(a)(i)		2.21	2	B1 for 2.21
	(ii)		0.013		B1 for 0.013
	(b)	LCM = 3 × 13 × 17 = 3 × 221	663	2	M1 for 3 × 13 × 17 oe A1 for 663


PAPER 5503				
No	**Working**	**Answer**	**Mark**	**Notes**
15		Tick boxes 1,3 & 6	3	B1 for $\frac{\pi abc}{2d}$; B1 for $2a^2$; B1 for $2\left(c^2+d^2\right)$ (−B1 for each additional expression ticked (>3) to a minimum of 0)
16	200×0.2	40	2	M1 for 200×0.2 or $\frac{40}{200}$ seen A1 for 40
17 (a)	eg 2 | 108 2 | 54 3 | 27 3 | 9 3	$2^2 \times 3^3$	3	M2 for full systematic method of at least 4 divisions by prime numbers oe factor trees; condone 1 calculation error (M1 for 108 written as a correct product(s) with only one non-prime or equivalent division or a full process with 2 calculation errors.) A1 for $2^2 \times 3^3$ (accept $2 \times 2 \times 3 \times 3 \times 3$)
(b)		12	1	B1 for 12
18	Perpendicular from P to intersecting arcs (within tramlines); perpendicular at least 2cm long		2	M1 relevant pair of arcs crossing within tramlines A1 SC M1 AO for a full construction of a perpendicular to AB not passing through P.
19	15×10	150 cm^3.	3	M1 for 15×10 A1 for 150 B1 for cm^3.

PAPER 5503					
No		**Working**	**Answer**	**Mark**	**Notes**
20			$y = 2x + 6$	2	B2 for $y = 2x + 6$ (B1 for $y = 2x + k$, $k \neq 1$ or for $y = mx + 6$, m $\neq 0$ or for $2x + 6$)
21		$6\frac{2}{5} = \frac{32}{5}$ Area of triangle = $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ (=2) Length of a side of sq. = $\sqrt{18 \times "2"}$ (=6) Perimeter of square = 4 × 6	24	5	B1 for $6\frac{2}{5} = \frac{32}{5}$ oe or $3\frac{1}{5} = \frac{16}{5}$ or $\frac{30}{8} + \frac{2}{8}$ oe (or implied by area of triangle=2) M1 for $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ oe M1 for area of square = 18 × product of two lengths A1 for $\sqrt{18 \times "2"}$ A1 for 24
22	(a)		$x(x-3)$	2	B2 for $x(x-3)$ (B1 for x (…….))
	(b)		k^3.	1	B1 for k^3.
	(c)(i)	$4x + 20 + 3x - 21$	$7x - 1$	4	M1 for **three** of 4 terms $4x + 20 + 3x - 21$ (or better) A1 for $7x - 1$
	(ii)	$x^2 + 3yx + 2xy + 6y^2$	$x^2 + 5xy + 6y^2$		M1 for **three** of 4 terms $x^2 + 3yx + 2xy + 6y^2$ A1 for $x^2 + 5xy + 6y^2$
	(d)		$(p+q)(p+q+5)$	1	B1 for $(p+q)(p+q+5)$

PAPER 5503					
	No	Working	Answer	Mark	Notes
23	(a)		32	1	B1 for 32 (accept 31.5 to 33.5 inclusive)
	(b)			3	B1 for ends of whiskers at 9 and 57, (with a box) B1 for ends of box at 16 and 45 (± 0.5) B1 for median marked at 32 or a complete box and whisker diagram drawn with a median inside the box
	(c)	Median(B)>Median(G); on average boys take longer IQR(B)>IQR(G); times for boys have a greater spread		2	B1 eg for comparison of medians B1 eg for comparison of (interquartile) ranges
24	(a)	$6y = 15 - 5x$	$y = \frac{15-5x}{6}$	2	M1 for either $6y = 15 - 5x$ or -6 $y = 5x - 15$ or for $\frac{5x}{6} + y = \frac{15}{6}$ or a correct ft on sign error to $y=$ A1 for $y = \frac{15-5x}{6}$ oe
	(b)	$6k + 5(-21) = 15$	20	2	M1 for substitution of $x = -21$ (or $x = 21$) into given equation or candidate's answer to (a) A1 for $k = 20$
	(c)(i)		Region R indicated	3	B2 correct region shaded (accept unshaded if R clear) (B1 shaded or R region satisfies 3 of the 4 given inequalities with same boundaries)
	(ii)		(1,1)		B1 for (1,1)

PAPER 5503					
	No	Working	Answer	Mark	Notes
25	(a)		60	1	B1 for 60 B1 for *BDC* = 25
	(b)	eg Angle *BDC* = 25 *ADB* = 60 − 25	35	2	[Award the mark for equivalent approaches] B1 ft for (a) − 25
	(c)	Ben is correct; angle *DAB* = 65 + 25 = 90 and since angle in a semi-circle is 90°, *BD* must be a diameter		1	B1 for full valid justification

PAPER 5504					
	No	Working	Answer	Mark	Notes
1	(a)	Accurate drawing of triangle		2	B1 cao for 38º (±2º) B1 cao (indep) for BC drawn 7.3 cm (±2mm) and completing the triangle.
	(b)	Measure angle A	63º	1	B1 for 63º (±2º) or ft their diagram (±2º)
2	(a)	0 × 1 = 0 1 × 1 = 1 2 × 4 = 8 3 × 8 = 24 4 × 8 = 32 5 × 7 = 35	100	3	M1 col 1 × col 2 (at least 3 shown) could be implied by answers. M1 (dep) sum of totals A1 cao SC: M1 M1 A0 for 101
	(b)	Either $35.50 ÷ 1.42 = £25; £26.99 - £25 = £1.99 Cheaper in the USA Or £26.99 × 1.42= $38.33; $38.33 - $35.50 = $2.83 Cheaper in the USA		3	M1 35.50 ÷ 1.42 A1 25 seen B1 ft (dep on M1) £1.99 with conclusion; units needed OR: M1 26.99 × 1.42 A1 38.33 or 38.32…seen B1 ft (dep on M1) $2.83 or $2.82 with conclusion; units needed
3	(a)		-1, (1), (3), 5, 7, 9	2	B2 cao (B1 for 2 values)
	(b)		Graph	2	B1 ft for plotting points ±1/2 square B1 cao for line between $x = -2$ and $x = 3$
	(c)(i)		0.4	2	B1 0.4 or ft single straight line with positive gradient
	(ii)		1.2		B1 1.2 or ft single straight line with positive gradient

PAPER 5504					
	No	Working	Answer	Mark	Notes
4	(a)			2	B1 Rectangle drawn B1 Triangle attached Ignore additional internal lines; 3D sketch gets B0
	(b)			2	B1 single Rectangle drawn B1 Attached smaller rectangle drawn Accept rotated plan, Ignore internal lines down; Nets get B0
5	(a)	£163.25 - £35.50 = £127.75 £127.75 ÷ £18.25 = 7	8	3	M1 163.25 – £35.50 (or sight of £127.75) M1 (dep) "127.75" ÷ 18.25 A1 cao SC: M2 for 7 days
	(b)	1.175 × £64 or $\frac{17.5}{100}$ × 64 or £11.20 seen	75.20	2	M1 1.175 × 64.00 0.175 × 64 oe or 11.20 or 75.2 seen OR 6.40, 3.20, and 1.60 seen A1 cao
	(c)	54 × £2.38 =	128.52	2	M1 54 × £2.38 A1 cao
6		Tessellation	Drawing	2	B2 seven additional hexagons, with at least 2 points at which 3 hexagons meet (B1 one point at which 3 hexagons meet).
7			200 150 225 150 10	3	B3 cao (B2 for three correct, or B1 for one correct)
8	(a)(i)	Bearing measured	70	2	B1 70 ± 4
	(ii)	Bearing measured	317		B1 317 ± 4
	(b)	Locus (circle) drawn & shaded		2	B1 circle ±2 mm centre Manchester B1 shading with accurate or approximate circle within tolerance

PAPER 5504					
	No	Working	Answer	Mark	Notes
9	(a)		Plots	1	B1 cao
	(b)		Description	1	B1 dynamic relationship or "positive" (correlation)
	(c)		Line of best fit	1	Line within overlay region, and to the extent of.
	(d)(i)	280 g →	Reading	2	B1 ft from single straight line of positive gradient (±1/2 square) B1 ft from single straight line of positive gradient (±1/2 square)
	(ii)	120 pages →	Reading		
10	(a)	$\frac{1}{6}, \frac{3}{8}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}$		2	B2 for all correct (B1 for 1 error or all correct but wrong order, or use of a common denominator decimals).
	(b)	3/5, 65%, 2/3, 0.72, 3/4		2	B2 for all correct (B1 for 1 error or all correct but wrong order or conversions to decimals oe)
11	(a)		182, 178, 180, 184	2	M1 mean of any three consecutive months, eg (147 + 161 + 238) ÷ 3 oe A1 cao
	(b)	Sale price = 80% Fun Friday price = 70% of 80% = 56% (oe)		2	B1 B1 for a fully correct explanation involving a worked example (oe) B1 a partially complete explanation

PAPER 5504					
	No	Working	Answer	Mark	Notes
12	(a)		Reason	1	B1 makes a mention of bias, either directly, or making reference to an insufficient or biased range of responses.
	(b)(i)		Reason	1	B1 make some mention of any one of (a) an insufficient range of responses (b) no mention of money (c) no time frame in the question (d) misunderstanding of "a lot" or "not much"
	(ii)		Question	2	B2 B1 for each of the following upto B2 (a) an improved question eg time frame made clear (b) response boxes (imperfect) (c) response boxes no errors OR For suggesting a generally improved question (a) a question clearly in the context of changes to the canteen (b) at least 3 boxes showing a full range of responses
13	(a)	$20y - 18y = 16 - 9$ oe $2y = 7$	$3\frac{1}{2}$ oe	3	M1 $20y - 18y = 16 - 9$ oe M1 $2y = 7$ A1 cao
	(b)	$40 - x = 3(4 + x)$ $40 - x = 12 + 3x$ $40 - 12 = x + 3x$ $4x = 28$	7	3	M1 multiplying through by 3: $3 \times \frac{40 - x}{3} = 3 \times 4 + 3 \times x$ A1 $40 - 12 = x + 3x$ A1 cao
14	(a)		$12(x\text{-}4)$	1	B1 for $12(x - 4)$ or $12x - 48$ or $12 \times x - 48$ oe $x = 12(x - 4)$ gets B0
	(b)		$18x\text{-}48$	2	M1 $6x +$ "$12(x - 4)$" or $6x +$ "$12x - 48$" A1 ft a linear expression $18x - 48$, $2(9x - 24)$, $3(6x - 16)$, $6(3x - 8)$


PAPER 5504					
No		**Working**	**Answer**	**Mark**	**Notes**
15		x $x^3 - 2x$ 4.1 60.7(21) 4.2 65.6(88) 4.3 70.9(07) Accept answer to 1 dp 4.22 66.7(114) or more 4.23 67.2(269) 4.24 67.7(45) 4.25 68.2(656)	4.2	4	B2 for trial $4.1 \leq x \leq 4.3$ evaluated (B1 for trial $4 < x < 5$ evaluated) B1 for different trial $4.225 \leq x \leq 4.25$ evaluated B1 (dep on at least one previous B1) for 4.2 cao
16	(a)		1.0×10^{-9}	1	B1 accept 1×10^{-9} or just 10^{-9}
	(b)	$1 \div 5 \times 10^{-9}$	2×10^{8}	2	M1 for $1 \div$ ("5×10^{-9}") or digit 2 with zeros only seen Condone omission of bracket for M1. A1 cao
17	(a)	$\frac{6.27 \times 4.52}{4.81 + 9.63} = \frac{28.3404}{14.44} = 1.962631579$	1.9626…	2	B2 for 1.9626… (B1 for 28.34 or 14.44)
	(b)		1.96	1	B1 ft from (a) as rounded to 1dp or 2dp. Do not accept 2, 2.00, but accept 2.0
18			$d = 4n + 6$	2	B2 $d = 4n + 6$ oe (B1 $d = 4n + k$, k an integer or 0, $4n + 6$, $n = 4n + 6$)
19		Vol = $\pi \times 3.8^2 \times 2.5 = \pi \times 14.44 \times 2.5$ = 45.36… × 2.5 = 113.411 Mass = "113" × 1.5 = 170.1165	170	4	M1 for $\pi \times r^2 \times 2.5$ where r = is 3.8 or 7.6 A1 if $r = 3.8$ M1 for "113" × 1.5 A1 for 169.5 – 170.3 cao


PAPER 5504					
	No	Working	Answer	Mark	Notes
20	(a)	$10^2 + 6^2$ or 136 $\sqrt{(100+36)}$ or $\sqrt{136} = 11.66\ldots$	11.7	3	M1 $10^2 + 6^2$ or 136 seen M1 (dep) $\sqrt{100+36}$ or $\sqrt{136}$ A1 11.66 – 11.7
	(b)	$\cos x = \frac{8}{10}$ or 0.8 $x = \cos^{-1} 0.8 = 36.869^\circ$	36.9	3	M1 for $\cos = \frac{8}{10}$, cos = 0.8 (oe) M1 (dep) for $\cos^{-1}$ (oe) A1 for 36.86 – 36.9
21		$18x - 6y = 99$ $\quad 12x - 4y = 66$ $8x + 6y = 18$ $\quad 12x + 9y = 27$ $26x = 117$ $\quad -13y = 39$	$x = 4\frac{1}{2}$ $y = -3$	4	M1 correct process to eliminate either x or y (condone one error) A1 cao for non – eliminated one. M1 (dep on 1st M1) for correct substitution of their found value. A1 cao (need both)
22		$\frac{133}{72} = 1.8472, \frac{160}{82} = 1.9512$ OR $\frac{72}{133} = 0.54135, \frac{82}{160} = 0.5125$ OR $\frac{160}{133} = 1.203...., \frac{82}{72} = 1.1388....$ OR $\frac{133}{160} = 0.83125...., \frac{72}{82} = 0.878$	1.84..≠ 1.95.. 1.20..≠ 1.13..	3	M1 for $\frac{133}{72}$ (= 1.8472…) oe Accept 1.8, 1.85 M1 for $\frac{160}{82}$ (=1.9512…) oe consistent pairing Accept 2.0, 1.9 **OR** M1 for $\frac{160}{133}$ (= 1.203…) oe M1 for $\frac{82}{72}$ (=1.1388) oe A1 for enough decimal places to show that the ratios are not equal; since the scale factors are different the shapes cannot be similar. NB Do Not need conclusion

PAPER 5504					
	No	Working	Answer	Mark	Notes
23	(a)		0.4, 0.6 0.4, 0.6, 0.4	2	B1 for LHS: (0.6), 0.4 B1 for RHS: 0.6, 0.4, 0.6, 0.4
	(b)	(30 × 42) – (25 × 42.8) = 1260 – 1070 = 190 190 ÷ 5 =	38	3	M1 for (30 × 42) – (25 × 42.8) or 1260 – 1070 or 190 seen M1 (dep) for "190" ÷ 5 A1 cao
24	(a)	£12000 × 0.25 = £3000; £12000 – £3000 = £9000 £9000 × 0.25 = £2250; £9000 – £2250 = £6750 £6750 × 0.25 = £1687.50; £6750 – £1687.50 =	£5062.50	3	M1 for 12000 × 0.75 (=9000) oe or £3000 or £23437.50 seen M1 (dep) for at least two further depreciation calculations (complete steps) A1 cao **OR** M2 for $12000 \times (0.75)^3$ or 5062.50 seen (M1 for $12000 \times (0.75)^n$, $n = 2$ or 4)
	(b)	0.8 × 0.8 × 0.8 × 0.8 (oe)	0.4096	2	M1 0.8^4 (oe) A1 cao

PAPER 5505					
	No	Working	Answer	Mark	Notes
1	(a)(i) (ii) (b)	 LCM = 3 × 13 × 17 = 3 × 221	2.21 0.013 663	2 2	B1 for 2.21 B1 for 0.013 M1 for 3 × 13 × 17 oe A1 for 663
2			Tick boxes 1,3 & 6	3	B1 for $\frac{\pi abc}{2d}$; B1 for $2a^2$; B1 for $2(c^2 + d^2)$ (−B1 for each additional expression ticked (>3) to a minimum of 0)
3	(a) (b)	200 × 0.2 0.2 + 0.4	40 0.6	2 2	M1 for 200 × 0.2 or $\frac{40}{200}$ seen A1 for 40 M1 for 0.2 + 0.4 A1 for 0.6
4	(a) (b)	eg 2 | 108 2 | 54 3 | 27 3 | 9 3 12	$2^2 \times 3^3$ 12	3 1	M2 for full systematic method of at least 4 divisions by prime numbers oe factor trees; condone 1 calculation error (M1 for 108 written as a correct product with only one non-prime or equivalent division or a full process with 2 calculation errors.) A1 for $2^2 \times 3^3$ (accept 2 × 2 × 3 × 3 × 3) B1 for 12
5		Perpendicular from P to intersecting arcs (within tramlines); perpendicular at least 2 cm long		2	M1 relevant pair of arcs crossing within tramlines A1 SC M1A0 for full construction of a line perpendicular to AB not through P
6		15 × 10	150 cm^3.	3	M1 for 15 × 10 A1 for 150 B1 for cm^3

PAPER 5505					
	No	Working	Answer	Mark	Notes
7	(a)		k^3.	1	B1 for k^3.
	(b)(i)	$4x + 20 + 3x - 21$	$7x - 1$	4	M1 for **three** of 4 terms $4x + 20 + 3x - 21$ (or better) A1 for $7x - 1$
	(ii)	$x^2 + 3yx + 2xy + 6y^2$	$x^2 + 5xy + 6y^2$		M1 for **three** of 4 terms $x^2 + 3yx + 2xy + 6y^2$ A1 for $x^2 + 5xy + 6y^2$
	(c)		$(p+q)(p+q+5)$	1	B1 for $(p+q)(p+q+5)$
	(d)		m^8	1	B1 for m^8.
	(e)		$6r^3t^6$	2	B2 for $6r^3t^6$ (B1 for $.....r^3t^6$ or for $6....t^6$)
8	(i)	Least length = 100.5	100.5	2	B1 for 100.5
	(ii)	Greatest length = 101.5	101.5		B1 for 101.5; ACCEPT 101.499 or better
9		$6\frac{2}{5} = \frac{32}{5}$ Area of triangle = $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ (= 2) Length of a side of sq. = $\sqrt{18 \times "2"}$ (=6) Perimeter of square = 4 × 6	24	5	B1 for $6\frac{2}{5} = \frac{32}{5}$ oe or $3\frac{1}{5} = \frac{16}{5}$ oe or $\frac{30}{8} + \frac{2}{8}$ oe (or implied by area of triangle=2) M1 for $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ oe M1 for (area of square) = 18 × product of two lengths A1 = $\sqrt{18 \times "2"}$ A1 for 24

PAPER 5505					
	No	Working	Answer	Mark	Notes
10	(a)	$6y = 15 - 5x$	$y = \frac{15-5x}{6}$	2	M1 for either $6y = 15 - 5x$ or for $\frac{5x}{6} + y = \frac{15}{6}$ or $-6y = 5x - 15$ or a correct ft on sign error to $y =$ A1 for $y = \frac{15-5x}{6}$ oe
	(b)	$6k + 5(-21) = 15$	20	2	M1 for subst. of $x = -21$ (or $x = 21$) into given eq^n or answer to (a) A1 for $k = 20$
	(c)(i)		Region R indicated	3	B2 correct region shaded (accept unshaded if R clear) (B1 shaded (R) region satisfies 3 of the 4 given inequalities with same boundaries)
	(ii)		(1,1)		B1 for (1,1)
11	(a)		$y = 2x + 6$	2	B2 for $y = 2x + 6$ (B1 for $y = 2x + k$, $k \neq 1$ or for $y = mx + 6$, $m \neq 0$ or for $2x+6$)
	(b)	Grad of $AB = 2$; Grad of $BC = \frac{-1}{"2"}$	$y = -\frac{1}{2}x + 6$	2	M1 for Grad of $BC = \frac{-1}{"2"}, -\frac{1}{2}$ or grad of $BC = -\frac{1}{\text{grad of AB}}$ A1 ft for $y = -\frac{1}{2}x + 6$ oe ft on wrong coeff. of x in (a)
	(c)	Eg A rectangle is always a cyclic quadrilateral because the opposite angles of a rectangle always add up to 180° (90+90)		1	B1 for valid explanation. (eg lines from the pt of int. of diagonals of rect to all 4 vertices are equal(radii))
12		Vertices of new triangle at $(-1, -4)$ $(-4, 2)$ and $(2, 2)$	Correct triangle drawn	3	B3 cao (B2 for 3 correct vertices no triangle or triangle with 2 correct vertices (B1 for triangle with two of six co-ordinates correct from using P as centre or any isosceles triangle with base 6 and height 6)

PAPER 5505					
	No	Working	Answer	Mark	Notes
13	(a)		32	1	B1 for 32 (accept 31.5 to 33.5 inclusive)
	(b)			3	B1 for ends of whiskers at 9 and 57 (with a box) B1 for ends of box at 16 and 45/46 ($\pm$ 0.5) B1for median marked at "32"or complete box and whisker diagram drawn with a median inside the box
	(c)	Median(B)>Median(G); on average boys take longer IQR(B)>IQR(G); times for boys have a greater spread		2	B1 eg for comparison of medians (ft on diagrams) B1 eg for comparison of (interquartile) ranges (ft on diagram)
14		Reading top to bottom frequencies are 20; 18; 45; 52		2	B2 all correct (B1 for one frequency correct)
15	(a)	$1600 = p\,q^0$.; $400 = pq^2$ $q^2 = 0.25$	$p = 1600$ $q = 0.5$	3	M1 for either $400 = pq^2$ or $1600 = pq^0$ B1 for $p = 1600$ A1 for $q = 0.5$
	(b)	$V = pq^{-2}$	6400	2	M1 for recognition that t = –2 A1 for 6 400

PAPER 5505					
	No	Working	Answer	Mark	Notes
16	(a)	(I) $AB = AC$ (triangle ABC is isosceles) (II) PB = PC tangents (from a point to a circle are) equal (III) AP = AP (common side) so the 2 triangles are congruent ,SSS.		3	B3 for I, II, III with congruency reason (B2 for any two of I, II or III) (B1 for any one of the I, II or III)
	(b)	$BPC = 20°$ PBC (or PCB) = 90-1/2 "20" (= 80°) $BAC = PBC$ = "80"	50°	4	B4 for 50° (B3 for $BAC = 80°$) (B2 for $PBC = 80°$ or $PCB = 80°$) (B1 for $APC = 10°$ or $BPC = 20°$ or a middle angle = 90°) SC if clear numerical slip seen eg "PBC = 180 – 90 – 10 = 70" then goes on to get correct ft angle $ABC = 55$ deduct 1 from total so this cand would get B4–1 = B3
17	(a)	$\overrightarrow{OP} = \overrightarrow{OA} + \overrightarrow{AP}$ $= \overrightarrow{OA} + \frac{2}{3}(6\mathbf{c} - 6\mathbf{a})$ $= 6\mathbf{a} + 4\mathbf{c} - 4\mathbf{a}$	$2\mathbf{a} + 4\mathbf{c}$	3	M1 for $\overrightarrow{OP} = \overrightarrow{OA} + \overrightarrow{AP}$ or any correct vector journey involving $\overrightarrow{OP}$ M1 for $\overrightarrow{AP} = \frac{2}{3}(6\mathbf{c} - 6\mathbf{a})$ oe or $\overrightarrow{CP} = \frac{1}{3}(-6\mathbf{c} + 6\mathbf{a})$ oe or reverse vectors A1 for $2\mathbf{a} + 4\mathbf{c}$ oe (accept unsimplified)
	(b)	Eg $\overrightarrow{OM} = \overrightarrow{OC} + \overrightarrow{CM} = 6\mathbf{c} + 3\mathbf{a}$ $\overrightarrow{OM} = 1.5\overrightarrow{OP}$	$\overrightarrow{OM} = 1.5\overrightarrow{OP}$ so OPM is a straight line	2	B1 for $\overrightarrow{OM} = 6\mathbf{c} + \frac{1}{2}(6\mathbf{a})$ or $\overrightarrow{PM} = 2\mathbf{c} + \mathbf{a}$ unsimplified or reverse vectors B1 for a fully correct proof.

PAPER 5505					
	No	Working	Answer	Mark	Notes
18	(a)		4	1	B1 for 4 condone $\pm$ 4
	(b)		2	1	B1 for 2 condone ± 2
	(c)	$\sqrt{160} = 4\sqrt{10}$; $\left[\frac{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)-\sqrt{2}\times\sqrt{5}}{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)}\right]\times 100$ $\left[\frac{6\sqrt{10}-\sqrt{10}}{6\sqrt{10}}\right]\times 100$	$\frac{500}{6}$	4	B1 for **either** $\sqrt{160} = 4\sqrt{10}$ or $\sqrt{8} = 2\sqrt{2}$ **or** $\sqrt{20} = 2\sqrt{5}$ M1 for $\left[\frac{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)-\sqrt{2}\times\sqrt{5}}{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)}\right]$ oe $(\times 100)$ B1 for either $6\sqrt{10} - \sqrt{10}$ or $6\sqrt{10}$ A1 for $\frac{500}{6}$ (accept 83.3 if no obvious earlier error)

PAPER 5505					
	No	Working	Answer	Mark	Notes
19	(a)(i)		$(2x-7)(x-14)$	3	M1 x^2 term and constant term (± 98) obtained or $2x(x-14) - 7(x-14)$ or $x(2x-7) - 14(2x-7)$ A1 for $(2x-7)(x-14)$
	(ii)		$x = \frac{7}{2}$; $x = 14$		B1ft ft (i) provided of form $(2x\pm a)(x\pm b)$
	(b)(i)		$\frac{7}{n+7}$	3	B1 for $\frac{7}{n+7}$ oe
	(ii)	$\frac{7}{n+7} = \frac{2}{5} \Rightarrow 2(n+7) = 5 \times 7$ $2n = 21$	$n=10.5$ is not possible since n has to be an integer		M1 for $2(n+7) = 5\times 7$ or $n+7=5\times 3.5$ (can be implied) ft (b)(i) fractional in terms of n and < 1 A1 ft for $n=10.5$ not possible (since n not integer) oe
	(c)	$2\times\left(\frac{n}{n+7}\right)\times\left(\frac{7}{n+7}\right) = \frac{4}{9}$ $14n\times 9 = 4(n+7)^2$ $14n\times 9 = 4(n^2+14n+49)$ $4n^2+56n+196-126n = 0$	$2n^2 - 35n + 98 = 0$	5	M1 for $\left(\frac{n}{n+7}\right)\times\left(\frac{7}{n+7}\right)$ seen M1 for $2\times\left(\frac{n}{n+7}\right)\times\left(\frac{7}{n+7}\right)$ oe $(=\frac{4}{9})$ M1(dep on 1st M) elimination of fractions within an equation B1 3 terms correct in expansion of $(n+7)^2 = n^2 + 7n + 7n + 49$ A1 full valid completion to printed answer

PAPER 5505					
No		**Working**	**Answer**	**Mark**	**Notes**
	(d)	$\frac{7}{n+7}\times\frac{7}{n+7}=\frac{7}{21}\times\frac{7}{21}=$	$\frac{1}{9}$	2	M1 for $\frac{7}{n+7}\times\frac{7}{n+7}$ or better or ft [answer (b)(i)]^2 or $1-\frac{4}{9}-(\frac{n}{n+7})^2$ A1 for $\frac{1}{9}$ oe cao
20	(a)(i)		$y=1+\sin x$	2	B1 for $y=1+\sin x$
	(ii)		$y=2\sin x$		B1 for $y=2\sin x$ SC both (i) f(x) + 1, (ii) 2f(x) B1
	(b)	Stretch parallel to y-axis scale factor 3 Stretch parallel to x-axis scale factor $\frac{1}{2}$		3	M1 for 'stretch' A1 for Stretch parallel to y-axis scale factor 3 oe A1 for Stretch parallel to x-axis scale factor $\frac{1}{2}$ oe SC if M0 award BI for "sf 3 vertically" **and** "sf $\frac{1}{2}$ horizon."

PAPER 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
1		Question	2	B1 for each of the following upto B2 (a) an improved question eg time frame made clear (b) response boxes (imperfect) (c) response boxes no errors OR For suggesting a generally improved question (a) a question clearly in the context of changes to the canteen (b) at least 3 boxes showing a full range of responses
2	Locus (circle) drawn & shaded		2	B1 circle centre Manchester ±2mm B1 shading with accurate or approximate circle within tolerance
3 (a)		182, 178, 180, 184	2	M1 mean of any three consecutive months, eg (147 + 161 + 238) ÷ 3 oe A1 cao
(b)	Sale price = 80% Fun Friday price = 70% of 80% = 56% (oe)		2	B1 B1 for a fully correct explanation involving a worked example (oe) B1 a partially complete explanation
4	x $x^3 - 2x$ 4.1 60.7(21) 4.2 65.6(88) 4.3 70.9(07) 4.22 66.7(114) 4.23 67.2(269) 4.24 67.7(45) 4.25 68.2(656)	4.2	4	B2 for trial $4.1 \leq x \leq 4.3$ evaluated (B1 for trial $4 < x < 5$ evaluated) B1 for different trial $4.225 \leq x \leq 4.25$ evaluated B1 (dep on at least one previous B1) for 4.2 cao

PAPER 5506					
	No	Working	Answer	Mark	Notes
5	(a)		1.0×10^{-9}	1	B1 accept 1×10^{-9} or just 10^{-9}
	(b)	$1 \div (5 \times 10^{-9})$	2×10^{8}	2	M1 for $1 \div$ ("5×10^{-9}") or digit 2 with zeros only seen Condone omission of bracket for M1. A1 cao
6	(a)	$\frac{6.27 \times 4.52}{4.81 + 9.63} = \frac{28.3404}{14.44} = 1.962631579$	1.9626…	2	B2 for 1.9626… (B1 for 28.34… or 14.44)
	(b)		1.96	1	B1 ft from (a) as rounded to 1dp or 2dp. Do not accept 2, 2.00, but accept 2.0
7			$d = 4n + 6$	2	B2 $d = 4n + 6$ oe (B1 $d = 4n + k$, k an integer $\neq 6$, $4n + 6$, $n = 4n + 6$)
8		Vol = $\pi \times 3.8^2 \times 2.5 = \pi \times 14.44 \times 2.5$ $= 45.36\ldots \times 2.5 = 113.411$ Mass = "113" × 1.5 = 170.1165	170	4	M1 for $\pi \times r^2 \times 2.5$ where r = is 3.8 or 7.6 A1 if $r = 3.8$ M1 for "113" × 1.5 A1 for 169.5 – 170.3 cao

PAPER 5506					
	No	Working	Answer	Mark	Notes
9	(a)	$10^2 + 6^2$ or 136 $\sqrt{(100 + 36)}$ or $\sqrt{136} = 11.66\ldots$	11.7	3	M1 for $10^2 + 6^2$ or 136 seen M1 (dep) $\sqrt{100+36}$ or $\sqrt{136}$ A1 11.66 – 11.7
	(b)	Cos x = 8/10 or 0.8 $x = \cos^{-1} 0.8 = 36.869^o$	36.9	3	M1 for cos = $\frac{8}{10}$, cos = 0.8(oe) M1 (dep) for $\cos^{-1}$ (oe) A1 for 36.86 – 36.9 M1 Use of sine rule and x found M1 for $x = 90 - \sin^{-1}$("0.8") A1 for 36.86 – 36.9
10		$18x - 6y = 99$ $12x - 4y = 66$ $8x + 6y = 18$ $12x + 9y = 27$ $26x = 117$ $-13y = 39$	$x = 4\frac{1}{2}$ $y = -3$	4	M1 correct process to eliminate either x or y (condone one error) A1 cao for non – eliminated one. M1 (dep on 1st M1) for correct substitution of their found value. A1 cao (need both)

PAPER 5506					
	No	Working	Answer	Mark	Notes
11		$\frac{133}{72} = 1.8472\,,\ \frac{160}{82} = 1.9512$ OR $\frac{72}{133} = 0.54135, \frac{82}{160} = 0.5125$ OR $\frac{160}{133} = 1.203...., \frac{82}{72} = 1.1388....$ OR $\frac{133}{160} = 0.83125...., \frac{72}{82} = 0.878$	1.84..≠ 1.95.. 1.20..≠ 1.13..	3	M1 for $\frac{133}{72}$ (= 1.8472…) oe Accept 1.8, 1.85 M1 for $\frac{160}{82}$ (=1.9512…) oe consistent pairing Accept 2.0, 1.9 **OR** M1 for $\frac{160}{133}$ (= 1.203…) oe M1 for $\frac{82}{72}$ (=1.1388) oe A1 for enough decimal places to show that the ratios are not equal; since the scale factors are different the shapes cannot be similar. NB Do Not need conclusion
12	(a)	£12000 × 0.25 = £3000; £12000 – £3000 = £9000 £9000 × 0.25 = £2250; £9000 – £2250 = £6750 £6750 × 0.25 = £1687.50; £6750 – £1687.50 =	£5062.50	3	M1 for 12000 × 0.75 (=9000) oe or £3000 or £23437.50 seen M1 (dep) for at least two further depreciation calculations (complete steps) A1 cao **OR** M2 for 12000 × $(0.75)^3$ or 5062.50 seen (M1 for 12000 × $(0.75)^n$, $n = 2$ or 4)
	(b)	0.8 × 0.8 × 0.8 × 0.8 (oe)	0.4096	2	M1 0.8^4 (oe) A1 cao

PAPER 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
13	$r^2 = \frac{3\times10}{\pi h} = \frac{3\times10}{\pi 1.5} = \frac{3\times10}{4.712..} = 6.36, 6.37$ $r = \sqrt{6.366\ldots}$	2.52-2.54	3	M1 for correct rearrangement to give $r^2 = \frac{3V}{\pi h}$ or $30 \div 4.712..$ or $6.36 - 6.37$ *Allow 0.3, 0.33 for* $\frac{1}{3}$ M1 (dep) for $\sqrt{}$ A1 cao 2.52 – 2.54
14 (a)		0.4 0.6,0.4, 0.6,0.4	2	B1 for LHS: (0.6), 0.4 B1 for RHS: 0.6, 0.4, 0.6, 0.4
(b)	0.6×0.6	0.36	2	M1 $0.6 \times$ "0.6" [0 < "0.6" < 1] A1 cao
(c)	$(30 \times 42) - (25 \times 42.8) = (1260 - 1070 = 190$ $190 \div 5 =$	38	3	M1 for $(30 \times 42) - (25 \times 42.8)$ or $1260 - 1070$ or 190 seen M1(dep) for "190" ÷ 5 A1 cao 38

PAPER 5506					
	No	Working	Answer	Mark	Notes
15	(a)	$S = \frac{k}{f^2};\ 125 = \frac{k}{8^2};\ k = 8000$	$S = \frac{8000}{f^2}$	3	M1 for $S = \frac{k}{f^2}$ M1 for $125 = \frac{k}{8^2}$ A1 cao These marks can be awarded if the full formula appears in part (b), rather than in part (a). [SC: $S = \frac{125}{64} f^2$ M1 M0 A0, $S = 1.95(3125) f^2$ M1 M0 A0 $S = \frac{1000}{f}$ M1 M0 A0]
	(b)		500	1	B1 cao

PAPER 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
16	$0.5 \times 3.2 \times 8.4 \times \text{Sin } B = 10$ $\sin B = 0.74404\ldots$ 48.077 $AC^2 = 3.2^2 + 8.4^2 - 2 \times 3.2 \times 8.4 \times \cos B$ $AC^2 = 44.8815\ldots..$ $AC = 6.69\ (936\ldots)$ Perimeter = 18.3 Use the altitude AD, $\frac{h \times 8.4}{2} = 10 \Rightarrow h = (2.381)$ $BD = \sqrt{3.2^2 - h^2} = 2.139$ $DC = 6.261$ $AC = \sqrt{2.38^2 + '6.261'^2} = 6.69(936)$ Perimeter = 18.3	18.3	6	M1 for $0.5 \times 3.2 \times 8.4 \times \sin B$ (= 10) A1 for sin B = 0.74(404…) or B = 47.7 – 48.1 M1 for $3.2^2 + 8.4^2 - 2 \times 3.2 \times 8.4 \times \cos$ "48.077" M1(dep) for AC^2 = "44.8(815)"….. with correct order of evaluation A1 AC = 6.69 – 6.7 A1 18.29 – 18.3 **for** $\frac{h \times 8.4}{2} = 10 \Rightarrow h = (2.381)$ M1 for $BD^2 = 3.2^2 - "2.381"^2$ A1 BD = 2.1 – 2.2 M1 (dep) $AC^2 = "2.381"^2 + "6.261"^2$ A1 AC = 6.69 – 6.7 A1 18.29 – 18.3

PAPER 5506					
	No	Working	Answer	Mark	Notes
17	(a)	$(x + 8)^2 = x^2 + (x + 5)^2$ $x^2 + 16x + 64 = 2x^2 + 10x + 25$		4	B1 for angle $OTA = 90°$ (implied by use of Pythagoras with OA as hypotenuse) M1 $(x + 8)^2 = x^2 + (x + 5)^2$ oe M1 for correct squaring of x+8 or x+5 A1 for completion following correct working
	(b)	$(+6 + \sqrt{(6^2 - - 4\times39)})/2$ $= (6 \pm \sqrt{(36 + 156)})/2$ $= (6 \pm \sqrt{192})/2 = (6 + 13.856)/2$	9.93	3	M1 for substitution into quadratic formula, allow sign errors in b and c M1 for $x = (6 \pm \sqrt{192})/2$ [+ alone will do] A1 for 9.92 – 9.93
					M1 for $(x-3)^2-9-39$ M1 for $x = 3 \pm \sqrt{48}$ [+ alone will do it] A1 for 9.92 – 9.93 T&I= 9.93 gets M1,M0,A0
18		$EB = 60 \times \tan 30°$ $BD = \sqrt{(60^2 + 60^2)}$ $\tan BDE = 34.64 \div 84.85$	22.2°	4	M1 for $EB = 60\times\tan30$ M1 for $BD = \sqrt{(60^2 + 60^2)}$ M1 for $\tan BDE =$ "34.64" ÷ "84.85" A1 22.17 – 22.21
		$EB = 60 \times \tan 30°$ (= 34.64) $ED^2 = 60^2 + \left(\frac{60}{\cos 30}\right)^2$ $ED = \sqrt{8400} = (91.65)$ $\text{Angle} = \sin^{-1}\left(\frac{EB}{\sqrt{8400}}\right) = 22.2$			M1 for $EB = 60 \times \tan 30°$ oe M1 for fully correct method for ED M1 for $\sin BDE = \left(\frac{'34.84'}{'\sqrt{8400}'}\right)$ (oe) A1 22.17 – 22.21

PAPER 5506					
	No	Working	Answer	Mark	Notes
19	(a)	$x^2 + 6^2 = 25$ $x^2 = -11$		2	M1 substitutes y=6 to get x^2+36 = 25 A1 deduces x^2=25-36<0; impossible, (so line does not intersect curve)
	(b)	$y^2 = (2x - 2)^2$ $x^2 + (2x - 2)^2 = 25$ $x^2 + 4x^2 - 8x + 4 = 25$ $5x^2 - 8x + 4 - 25 = 0$ $5x^2 - 8x - 21 = 0$ $(5x + 7)(x - 3) = 0$ or $x = \frac{--8 \pm \sqrt{(-8)^2 - 4 \times 5 \times (-21)}}{10} = \frac{8 \pm \sqrt{484}}{10}$ So $x = 3$, or $x = -1.4$ Subs into either equation gives: $y = 4$, or $y = -4.8$	$x = 3$ $y = 4$ $x = -1.4$ $y = -4.8$	6	M1 $y^2=(2x-2)^2$ seen or implied A1 $x^2 + 4x^2 - 8x + 4 = 25$ (oe expanded form) M1 correct attempt to solve 3 term quadratic equation A1 $x = 3$, $x = -1.4$ M1 (dep on previous Ms) Sub one value of x into either equation gives: A1 $y = 4$, $y = -4.8$ NB: incorrectly matched pairs loses the final A1.

PAPER 5506					
	No	Working	Answer	Mark	Notes
20	(a)	400.005 ÷ 59.95 = 6.672310 6.672310 ÷ 1000 × 3600 = 24.02032 399.995 ÷ 60.05 = 6.661032 6.661032 ÷ 1000 × 3600 = 23.97972	24.020 23.980	5	B1 for 400.005 or 59.95 seen oe M1 for "400.005" ÷ "59.95" where "400.005" ∈ [400.005,400.5] and "59.95" ∈ [59.5,59.95] oe B1 for 399.995 or 60.05 seen oe M1 for "399.995" ÷ "60.05" where "399.995" ∈ [399.5,399.95] and "60.05" ∈ [60.05,60.5] oe A1 23.979-23.980 and 24.020-24.0204
	(b)	24.0 because to 1dp the answers are the same		1	B1cao for 24.0 with reason
	(c)	177 × 50/477 = 18.553 → 19 → 18 111 × 50/477 = 11.635 → 12 86 × 50/477 = 9.0147 → 9 82 × 50/477 = 8.595 → 9 21 × 50/477 = 2.201 → 2	18 12 9 9 2	3	M1 for $\frac{50}{"477"} \times 177$ or 111 or 86 or 82 or 21 A1 for all integers or better answers, at least 3 correct A1 cao
21	(a)	$40 - x = 3(4 + x)$ $40 - x = 12 + 3x$ $40 - 12 = x + 3x$ $4x = 28$	7	3	M1 multiplying through by 3: $3 \times \frac{40 - x}{3} = 3 \times 4 + 3 \times x$ A1 $40 - 12 = x + 3x$ A1 cao
	(b)	$\frac{2x(2x - 3)}{(2x - 3)(2x + 3)} = \frac{2x}{2x + 3}$	$\frac{2x}{2x + 3}$	3	B1 for $(2x-3)(2x+3)$ B1 for $2x(2x-3)$ or $(2x + 0)(2x + 3)$ B1 cao

PAPER 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
22	Distance from x axis is y. Distance from (0, 2) is $\sqrt{(x^2+(y-2)^2)}$ $y^2 = x^2 + (y-2)^2$ $y^2 = x^2 + y^2 - 4y + 4$ $0 = x^2 - 4y + 4$ $4y = x^2 + 4$ and finish		4	B1 for $(x-0)^2 + (y-2)^2$ or $\sqrt{((x-0)^2+(y-2)^2)}$ oe seen B1 for $y = \sqrt{(x-0)^2+(y-2)^2}$ or $y^2 = (x-0)^2 + (y-2)^2$ oe B1 $(y-2)^2 = y^2 - 4y + 4$ seen B1 for $4y = x^2 + 4$ and finish

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone: 01623 467467
Fax: 01623 450481
Email: publications@linneydirect.com

Order Code UG015616 July 2004

For more information on Edexcel and BTEC qualifications please contact
Customer Services on 0870 240 9800
or http://enquiries.edexcel.org.uk
or visit our website: www.edexcel.org.uk

London Qualifications Limited, trading as Edexcel. Registered in England and Wales No. 4496750
Registered Office: Stewart House, 32 Russell Square, London WC1B 5DN



edexcel

Edexcel GCSE

Mathematics A 1387

June 2007

advancing learning, changing lives

Mark Scheme

Edexcel GCSE

Mathematics A 1387

Table Of Contents

Notes on Marking Principles	3 - 4
5521 / 01 Mark Scheme	5 - 11
5521 / 02 Mark Scheme	12 – 17
5523 / 03 Mark Scheme	18 – 27
5523 / 04 Mark Scheme	28 – 35
5525 / 05 Mark Scheme	36 – 43
5525 / 06 Mark Scheme	44 - 52

NOTES ON MARKING PRINCIPLES

1 **Types of mark**
M marks: method marks
A marks: accuracy marks
B marks: unconditional accuracy marks (independent of M marks)

2 Abbreviations
cao – correct answer only
ft – follow through
isw – ignore subsequent working
SC: special case
oe – or equivalent (and appropriate)
dep – dependent
indep - independent

3 No working
If no working is shown then correct answers normally score full marks
If no working is shown then incorrect (even though nearly correct) answers score no marks.

4 With working
If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.
If there is no answer on the answer line then check the working for an obvious answer.

5 Follow through marks
Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

6 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

7 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

8 Linear equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.


9 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

Remember: if you are having difficulty making a decision on how you should mark a candidate response contact your Team Leader for advice, or send the item to review.


Paper 5521_01					
No		Working	Answer	Mark	Notes
1		Cat |||| /||| 8 Dog |||| | 6 Fish || 2 Hamster |||| 4	8, 6, 2, 4	3	M1 for attempt to tally or one frequency correct in either column A1 for 1 frequency correct or all tallies correct in correct column A1 for all frequencies correct (accept if /20)
2			27, 35, 42, 67, 118	1	B1 cao
3	(a)		Diameter drawn	1	B1 for a diameter
	(b)		Right angle marked	1	B1 *R* marked correctly
	(c)		Rectangle drawn	1	B1 for a rectangle
4	(a)		40	1	B1 cao
	(b)		50	1	B1 cao
	(c)		3 full loaves 4 full loaves + 1 half loaf	2	B1 for 3 full loaves B1 for 4 full loaves + 1 half loaf
5	(a)		7252	1	B1 cao
	(b)		Three thousand and eighty six	1	B1 accept 3 thousand and eighty six (condone 0 hundred)
	(c)		4600	1	B1 accept 4600
	(d)		200	1	B1 for 200 or 2 hundred or 100 or hundred
6	(i)		Cube	2	B1 for 'cube' (accept 'cuboid') ignore spelling
	(ii)		Cylinder		B1 for 'cylinder' ignore spelling
7		5×100	500	2	B2 for 490 or 500 or 510 (B1 for either 5 or 5.0 or 100 seen)

Paper 5521_01					
No		Working	Answer	Mark	Notes
8	(a)		8.4 cm or 84 mm	2	B2 allow ±2mm (B1 for 8.4 or 84, B1 for appropriate unit)
	(b)		37°	1	B1 allow ±2°
9	(a)		Carbon black	1	B1 accept 'black carbon' accept 26%
	(b)		0.1(0)	1	B1 cao
	(c)		0.04	1	B1 cao
	(d)	$\frac{26}{100}$	$\frac{13}{50}$	2	M1 for $\frac{26}{100}$ A1 cao
10	(a)		97	1	B1 cao
	(b)		London Reading	1	B1 cao
	(c)	41 + 57 + 58	156	3	M1 for two of 41, 57, 58 M1(dep) for '41' + '57' + '58' A1 cao
11	(a)		3	1	B1 cao allow ± 0.2
	(b)		–5	1	B1 cao allow ±0.2

Paper 5521_01					
No		**Working**	**Answer**	**Mark**	**Notes**
12	(a)		Centimetres (cm) miles litres (*l*)	3	B3 (B1 for each correct answer) accept abbreviations
	(b)		300	1	B1 cao
	(c)	'1500>1400' or '1.5>1.4'	Reason	1	B1 for No and '1500>1400' or '1.5>1.4'
13	(a)	Height of bars 12, 8, 6 lines drawn between points	Bars drawn	2	B2 for 3 bars correctly drawn (B1 for 2 bars correct)
	(b)		July and August	1	B1 oe
	(c)	24 – 4	20	1	B1 (Accept '4 to 24' oe)
	(d)	The temperatures are rising	Temp's rising oe	1	B1 for reason
14	(a) (i)		$(-4, 3)$	2	B1 cao
	(ii)		$(2, -1)$		B1 cao
	(b) (i)	*D* marked at $(-4, -1)$	Point marked on grid	2	B1 for point marked at $(-4, -1)$ cao
	(ii)		$(-4, -1)$		B1 ft
15	(a)			1	B1 cao
	(b)			1	B1 cao

Paper 5521_01				
No	**Working**	**Answer**	**Mark**	**Notes**
16 (a)		(Pat +) reason	1	B1 correct comment (Pat may be implied)
(b)	21 ÷ 3	7	1	B1 cao
17 (i)		S marked at 1	3	B1 for S within ½ cm of 1
(ii)		P marked at 0		B1 for P marked at 0 cao
(iii)		Q marked at "1/3"		B1 for Q marked at 1/3 ± 1 cm use overlay
18	6 + 6 + 3 or $2\frac{1}{2}\times 6$	15	3	M1 for realizing 6 glasses in one bottle M1 for realizing 3 glasses in ½ a bottle A1 cao (M2 for attempt to find $2\frac{1}{2}\times 6$) oe
19	15 and 16 parts shaded Alternative 1 $\frac{3}{4}=0.75,\ \frac{4}{5}=0.8$ Alternative 2 $\frac{3}{4}=\frac{15}{20},\ \frac{4}{5}=\frac{16}{20}$	$\frac{4}{5}$ + reason	3	M1 for correctly shading 15 parts for 3/4 M1 for correctly shading 16 parts for 4/5 A1 (dependent on M2) for selection of 4/5 Alternative 1 M1 for $\frac{3}{4}=0.75$ M1 for $\frac{4}{5}=0.8$ A1 (dep on M2) for selection of 0.8 or $\frac{4}{5}$ or $\frac{16}{20}$ Alternative 2 M1 for $\frac{3}{4}=\frac{15}{20}$ M1 for $\frac{4}{5}=\frac{16}{20}$ A1(dep on M2) for selection of $\frac{4}{5}$ or $\frac{16}{20}$

Paper 5521_01				
No	**Working**	**Answer**	**Mark**	**Notes**
20 (a)		−5 (−3) (−1) 1 3 5	2	B2 cao (B1 for any 2 or 3 correct)
(b)	Points plotted	Line	2	B2 for line from (−1, −5) to (4, 5) (B1 ft for plotting at least 5 "points")
21	Shapes shaded on grid	6 tessellating shapes	2	B2 for fully correct with 5 or more additional shapes, no gaps (B1 for 4 shapes tessellating with at least one shape inverted, with or without the given shape ignore extras)
22 (a)		$\frac{7}{20}$	1	B1 for $\frac{7}{20}$ oe
(b)	7 + 2 (or 20 – 11) are not lime flavour	$\frac{9}{20}$	1	B1 for $\frac{9}{20}$
(c)		0	1	B1 for 0, zero or nought ($\frac{0}{20}$ gets B0)
23 (a)		$4a$	1	B1 accept $4\times a$, $a \times 4$, $a4$
(b)		$12b$	1	B1 accept $12\times b$, $b \times 12$, $b12$
(c)		$2a + 6b$	2	B2 cao (B1 for $2a$ or $6b$ seen)
(d)		$x(x-6)$	2	B2 cao (B1 for $x(ax+b)$ where a, b are numbers not equal to 0 or $x - 6$ seen on its own, or as part of an expression)
24 (a)		40	1	B1 cao
(b)		45	1	B1 for 42 – 48 accept ¾ hour
(c)	40×2 or $\frac{40}{30}\times 60$ or $40\div\frac{1}{2}$	80	2	M1 for 40×2 or $\frac{40}{30}$ or $40\div\frac{1}{2}$ A1 cao NB $\frac{40}{45}\times 60$ gets M0 A0

Paper 5521_01				
No	**Working**	**Answer**	**Mark**	**Notes**
25 (a)		4560	1	B1 cao
(b)		45.6	1	B1 cao
(c)		2.4	1	B1 cao
26	$-7 - 3 = -10$ $2 \times -10 = -20$ $-20 \div 4$	-5	3	M1 for substitution of 2 and -7 into $p(q - 3)$ or sight of -20 or $-14 - 6$ M1dep for '-20' $\div 4$ A1 cao B1 SC for sight of -10 if M0 awarded
27 (a)		Reflection in y-axis	1	B1 cao
(b)		Rotation by half turn about (0,0)	2	B2 cao (B1 for half turn not about (0,0).)
(c)		Enlargement Scale factor 3 Centre (0,0)	3	B1 for 'enlargement' B1 for "scale factor 3" or 3 seen B1 for 'centre (0,0)
28 (a)		Reason	1	B1 for 'The frequencies are nearly equal' oe
(b)	$1\times26+2\times26+3\times23+4\times25 = 247$ $247/100$	2.47	3	M1 for fx (attempting at least 2 relevant products) M1 for $\sum fx \div 100$ A1 2.47 cao

Paper 5521_01				
No	**Working**	**Answer**	**Mark**	**Notes**
29	5 × 5 × 6	150	4	M1 for attempt at 1 division (e.g. 40 ÷ 8), may be implied by marks or number on one edge of diagram or by 5 or 6 seen M1 for attempt at 3 divisions (40÷8, 40÷8, 60÷10), may be implied by marks or numbers on diagram or by 5,5 and 6 seen. M1 (dep on 1st M1) for "5" × "5" × "6" A1 cao Alternatively M1 for 40 × 40 × 60 **or** 8 × 8 × 10 **or** 96000 or 640 seen M1 for 40 × 40 × 60 **and** 8 × 8 × 10 or 96000 **and** 640 seen M1 (dep on 1st M1) for "(40 × 40 × 60)" ÷ "(8 × 8 × 10)" A1 cao SC:B1 for dividing area of one carton face by area of corresponding box face if M0

Paper 5521_02					
No		Working	Answer	Mark	Notes
1	(a)	Draw diagram.	Diagram	1	B1 cao
	(b)		13,16	1	B1 cao
	(c)		31	1	B1 cao
2	(a)		8	1	B1 cao
	(b)		14	2	B2 for 14 (B1 for 13 or 15)
	(c)		16	2	B2 for 16 (B1 for 15, 17 or 8)
3	(a)		4,7 drawn	2	B2 for car height 4 and bus height 7, (B1 for one correct)
	(b)		6	1	B1 cao
	(c)		Walk	1	B1
	(d)		27	1	B1 cao
4	(a)		40	1	B1 for 40–41 inclusive
	(b)		12	1	B1 for 11.5 – 12.5 inclusive
5	(a)		Row complete	2	B2 for 1+..+11; 36 (B1 for one of the 2 cells complete)
	(b)		Square	1	B1 "square"
6		One line of symmetry		1	B1 within 2mm of centre of base / 2mm of vertex

Paper 5521_02				
No	**Working**	**Answer**	**Mark**	**Notes**
7 (a) (i)		09 06	2	B1 (accept 9 06 oe)
(ii)		39		B1 cao
(b)		06 55	1	B1 (accept 6 55 oe)
(c)		2h 6min	1	B1 cao
(d)		15 min	2	M1 for 0906 – 0645 – "(c)" or 0906 – 0645 − 2hr 6min or 2hr 21min − "(c)" or 2hr 21min − 2hr 6 min or 141 − 126 or 20 − 5 A1 cao SC: B1 for 55 or 75 or 93 seen
8 (i)		8,10,12,20 or 30	3	B1 at least one of 8, 10, 12, 20, 30 (no extras)
(ii)		8,12 or 20		B1 at least one of 8,12, 20 (no extras)
(iii)		3 or 5		B1 3 or 5 or both (no extras)
9 (a)		C or G	1	B1 at least one of C or G (no extras)
(b)		A and F	1	B1 cao
(c)		2	1	B1 (accept –2)
10 (a)		$\frac{7}{10}$	1	B1 7/10 oe
(b)(i)		4 squares	2	B1 4 squares shaded
(ii)		80%		B1 80% or ft from unshaded part (no ft from 0% or 100%)
(c)(i)		2.5	2	B1 2.4–2.6 inclusive
(ii)		1.7		B1 1.6–1.8 inclusive
11 (a)		2	1	B1 for 2 or –2
(b)		14	1	B1 for 14 or –14

Paper 5521_02					
No		**Working**	**Answer**	**Mark**	**Notes**
12		2 × 8.50 = 17.00 3 × 4.50 = 13.50 Total = 30.50 50.00 – 30.50	19.5(0)(*p*)	3	M1 for adding 5 correct values or 2 × 8.50 + 3 × 4.50 (ignore units) or 30.5(0) or 3050 seen M1 dep for 50 − "30.50" (ignore units) (OR M1for adding at least 1 adult ticket and at least 1 child ticket **and** subtracting from 50) A1 cao SC: B1 for 24 or 37 or 2400 or 3700 seen
13	(a)		Hexagon	1	B1
	(b)(i)		120	2	B1 cao
	(ii)		Str line		B1 reference to a (straight) line and 180°
	(c)		Obtuse	1	B1 Accept "interior"
14	(a)	2, 2, 3, 3, 3, 4, 4, 4, 5, 6	3.5	2	M1 ordering the numbers (condone 1 error or omission) A1 cao
	(b)	36 ÷ 10	3.6	2	M1 sum of numbers ÷ 10 A1 cao SC B1 for 3r 6
	(c)	6 – 2	4	1	B1 cao
15	(a)		Paul	1	B1 cao
	(b)	36 ÷ 2 oe	18	1	B1 cao
	(c)	60/360 =	1/6	2	M1 60/360 oe A1 cao

Paper 5521_02					
No		**Working**	**Answer**	**Mark**	**Notes**
16		4.7 ÷ 5.9 = 0.796610169	0.7966..	2	B2 for 0.7966 or better (B1 for 0.8, 0.80, 0.79, 0.796, 0.797 or digits 59 seen
17		$6x - 7 + 7 = 38 + 7$ $6x = 45$	7.5	2	M1 $6x = 45$ or +7 both sides A1 7.5 oe; accept 45/6 oe
18		**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	3	B3 all six entries correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
19	(a)	900 × 1.70 =	1530	2	M1 900 × 1.7(0) or digits 153(0) seen A1 cao
	(b)	160 ÷ 1.70 =	£94.12 or £94.11	2	M1 160 ÷ 1.7(0) or digits 941(....) seen A1 cao
20	(a)(i)	180 – 54 (=126) "126" ÷ 2	63	3	M1 for (180–54) ÷ 2 A1 cao
	(ii)		Reason		B1 (indep) angles in triangle add to 180 OR equal angles in isosceles triangle OR equal angles and 2 sides the same (B0 if any incorrect reasoning given eg parallel, equilateral triangle)
	(b)	180 – "x"	117	1	B1 117 or ft 180 – "x" if $x < 90$
21	(a)	3 × 35 + 50	155	2	M1 for 3 × 35 + 50 or digits 155 seen A1 cao
	(b)	260 – 50 = 210 210 ÷ 35 =	6	3	M1 for 260–50 or 210 seen. M1 for "260-50" ÷ 35 or 210÷35 A1 cao SC B1 for starting at a number between 100 and 170 and adding at least two 35's and showing a total between 230 and 290 OR For adding at least three 35's, perhaps with other numbers, and showing a total between 180 and 240 (or between 230 and 290 if 50 is included in the sum)

Paper 5521_02				
No	**Working**	**Answer**	**Mark**	**Notes**
(c)		P=35h + 50	3	B3 for P=35h+50 or P=35×h+50 oe (B2 for correct RHS or P=h + 50 ×35 or P=35h+k where k is numerical oe) (B1 for P = some other linear expression in h, OR h + 50 ×35 OR 35h seen) NB: P=h scores no marks; ignore £ signs. SC B2 for $h = \frac{P-50}{35}$
22 (a)		Elevation	2	B2 for 4 vertical squares. Accept 4 by 1 rectangle. (B1 for 4 vertical squares with one square added or one parallelogram added at the top, or 3 vertical squares, or 4 horizontal squares)
(b)		Plan	2	B2 for 2 adjacent squares, vertical or horizontal. Accept 2 by 1 rectangle. (B1 for 3 adjacent horizontal or vertical squares or a rectangle with sides in the ratio 2:1)
23 (i) (ii) (iii)		5 9 6	3	B1 cao B1 cao B1 cao
24	$45.00 + 45.00 \times \frac{15}{100} =$ $45.00 + 6.75 =$	51.75	3	M2 for $45.00 + 45.00 \times \frac{15}{100}$ oe or 45.00×1.15 oe OR $45.00 + 6.75$ OR complete method or 5175 seen. (M1 for $45.00 \times \frac{15}{100}$ oe OR 6.75 seen OR 675 seen OR correct method for calculating 15% of 45) A1 cao SC Award B2 for an answer of 38.25

Paper 5521_02				
No	**Working**	**Answer**	**Mark**	**Notes**
25 (a)		Points	1	B1 all three points ±1 full square
(b)		Negative	1	B1 Negative (ignore additional descriptors unless contradictory)
(c)		lobf	1	B1 A single straight line drawn to cross between (5,30), (5,40) and (40,0), (40,15); accept freehand if considered to be straight.
(d)(i)		18–25	2	B1 18g–25g inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient.
(ii)		30–40		B1 30–40 min inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient
26		300 3 75 150	3	B3 for 4 correct answers (B2 for 2 or 3 correct answers) (B1 for 1 correct answer)
27	π × 0.65	2.04–2.05	2	M1 for π × 0.65 or 3.14 x 0.65 or 3.142 x 0.65 A1 for 2.04–2.05 SC Award B1 for 2.0 seen (not 2)
28	5 miles = 8 km 70mph÷5×8 = 112 km/h OR 120km/h ÷ 8 × 5 = 75 mph Faster than 70 mph	70mph (Great Britain) (112 km)	3	M1 5 miles = 8 km; OR 70 mph is about 100 km/h OR 1km=0.6(25) miles OR 1mile=1.6km oe M1 70 ÷ 5 × 8 (=112) or 120 ÷ 8 × 5 (=75) A1 (dep on at least M1) GB or 70 mph Refer to both answer line and working. NB GB or 70 mph without working scores 0 marks

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
1		Cuboid drawn	2	B2 for correct isometric drawing in any orientation (ignore points 'behind', mark 7 vertices only); accept lines drawn near to dots as long as there is no ambiguity. (B1 for one of the three faces drawn correctly **or** for an isometric drawing of any cuboid)
2	Different makes of car Tally Frequency	Make of car Tally Frequency	3	B1 for make of car or list of at least 3 different makes B1 for tally or tally marks B1 for frequency or totals
3		6 tessellating shapes	2	B2 for fully correct with 5 or more additional shapes, no gaps (B1 for 4 or more shapes tessellating, with at least one shape inverted, with or without the given shape, ignore extras)
4	24.90 ÷ 3 or 8.30 24.90 – '8.30' or 2 × 8.30	16.6(0)	3	M1 for 24.90 ÷ 3 or 8.30 M1 (dep) for 24.90 – “8.30” or 2 × “8.30” A1 for 16.60 or 16.6

Paper 5523_03



No	Working	Answer	Mark	Notes
5	315 × 24 24 × 315 1260 6300 7560 7200 7560 **3 1 5** 0 6 | 0 2 | 1 0 — **2** 1 2 | 0 4 | 2 0 — **4** 7 5 6 0 **300 10 5** 6000 200 100 — **20** 1200 40 20 — **4** 6000+200+100+1200+40+20 = 7560 **3 0.1 0.05** 60 2 1 — **20** 12 0.4 0.2 — **4** 60 + 2 + 1 + 12 + 0.4 + 0.2 = 75.6	75.6(0)	3	M1 for a complete method with relative place value correct. Condone 1 multiplication error, addition not necessary. **OR** M1 for a complete grid with not more than 1 multiplication error, addition not necessary. **OR** M1 for sight of a complete partitioning method, condone 1 multiplication error, final addition not necessary. A1 for 7560 or digits 756(0) A1 (dep on M1, but not previous A1) for correct placement of decimal point.

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
6	15 and 16 parts shaded Alternative 1 $\frac{3}{4} = 0.75$ or 75%, $\frac{4}{5} = 0.8$ or 80% Alternative 2 $\frac{3}{4} = \frac{15}{20}$, $\frac{4}{5} = \frac{16}{20}$	$\frac{4}{5}$ + reason	3	M1 for shading 15 parts for $\frac{3}{4}$ M1 for shading 16 parts for $\frac{4}{5}$ A1 (dep on M2) for selection of $\frac{4}{5}$ with correct shading Alternative 1 M1 for $\frac{3}{4} = 0.75$ or 75% M1 for $\frac{4}{5} = 0.8$ or 80% A1 (dep on M2) for selection of 0.8 or 80% or $\frac{4}{5}$ with correct decimals or percentages Alternative 2 M1 for $\frac{3}{4} = \frac{15}{20}$ oe M1 for $\frac{4}{5} = \frac{16}{20}$ oe A1 (dep on M2) for selection of $\frac{4}{5}$ or $\frac{16}{20}$ with equivalent fractions

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
7		$5 \times 5 \times 6$	150	4	M1 for attempt at 1 division (e.g. $40 \div 8$), may be implied by marks or number on one edge of diagram or by 5 or 6 seen M1 for attempt at 3 divisions ($40 \div 8$, $40 \div 8$, $60 \div 10$), may be implied by marks or numbers on diagram or by 5,5 and 6 seen. M1 (dep on 1st M1) for "5" × "5" × "6" A1 cao Alternatively M1 for $40 \times 40 \times 60$ **or** $8 \times 8 \times 10$ **or** 96000 or 640 seen M1 for $40 \times 40 \times 60$ **and** $8 \times 8 \times 10$ or 96000 **and** 640 seen M1 (dep on 1st M1) for "$(40 \times 40 \times 60)$" ÷ "$(8 \times 8 \times 10)$" A1 cao SC:B1 for dividing area of one carton face by area of corresponding box face if M0
8	(a)		$\frac{7}{20}$	1	B1 for $\frac{7}{20}$ oe
	(b)	7 + 2 (or 20 – 11) are not lime flavour	$\frac{9}{20}$	1	B1 for $\frac{9}{20}$ oe
	(c)		0	1	B1 for 0, zero or nought ($\frac{0}{20}$ gets B0)
9	(a)		$80x$	1	B1 for $80x$ (accept $80 \times x$, $x80$, $x \times 80$) seen
	(b)		$95y$	1	B1 for $95y$ (accept $95 \times y$, $y95$, $y \times 95$) seen
	(c)		$80x + 95y$	2	M1ft for adding "$80x$" and "$95y$" (algebraic expressions only) A1 for $80x + 95y$

Paper 5523_03

No	Working	Answer	Mark	Notes
10 (a)		40	1	B1 cao
(b)		45	1	B1 for 42 to 48 (accept 3/4 hour)
(c)	40×2 or $\frac{40}{30}\times60$ or $40\div\frac{1}{2}$	80	2	M1 for 40×2 or $\frac{40}{30}$ or $40\div\frac{1}{2}$ A1 cao NB $\frac{40}{45}\times 60$ gets M0 A0
11 (a)	$3\times3-4\times2$ or $9-8$	1	2	M1 for substitution of 3 and 2 into expression or 9 and 8 seen A1 cao
(b)	$-7-3=-10$ $2\times-10=-20$ $-20\div4$	-5	3	M1 for substitution of 2 and -7 into $p(q-3)$ or sight of -20 or $-14-6$ M1 (dep) for "-20" $\div 4$ A1 cao SC: B1 for -10 seen if M0
12 (a)	6 | 8 9 7 7 | 8 5 9 6 3 8 | 1 3 1 7 1 9 | 0 1	6 | 7 8 9 7 | 3 5 6 8 9 8 | 1 1 1 3 7 9 | 0 1	3	M1 for unordered diagram (condone one error) A1 cao B1 for key (eg 6 | 7 = 67)
(b)(i)		Explanation	2	B1 for '(order numbers and) select middle value' oe
(ii)		79		B1 cao

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
13	(a)		Reflection in y-axis	1	B1 for triangle with vertices at (-1, 1) (-3, 1) and (-1,4)
	(b)		Rotation by half turn about (0, 0)	2	B2 for triangle with vertices (-1, -1) (-3, -1) and (-1, -4) (B1 for half turn not about (0,0))
	(c)		Enlargement Scale factor 3 Centre (0, 0)	3	B1 for 'enlargement' B1 for 'scale factor 3' or 3 seen B1 for 'centre (0,0)' B0 for any combination of transformations
14	(a)		4560	1	B1 cao
	(b)		45.6	1	B1 cao
	(c)		2.4	1	B1 cao
15	(a)	$4a-2a+5b+b$	$2a+6b$	2	B2 cao (B1 for $2a$ or $6b$ seen)
	(b)		$x(x-6)$	2	B2 cao (B1 for $x(ax+b)$ where a, b are numbers not equal to zero or $x-6$ seen on its own, or part of an expression)
	(c)		$3x-2x^3$	2	B2 cao (B1 for $3x$ or $2x^3$)
	(d)		$4x(3y+x)$	2	B2 cao (B1 for $2(6xy+2x^2)$ or $4(3xy+x^2)$ or $x(12y+4x)$ or $2x(6y+2x)$ or $4x(\quad)$)

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
16	(a)	$1 - (0.2 + 0.3 + 0.1)$	0.4	2	M1 for $1 - (0.2 + 0.3 + 0.1)$ A1 for 0.4 oe, accept $\frac{0.4}{1}$
	(b)	0.2×200	40	2	M1 for 0.2×200 A1 cao NB $\frac{40}{200}$ is M1 A0, 40 out of 200 is M1 A1
17	(a) (i)	$180 - 2 \times 25$	130	3	M1 for $180 - 2 \times 25$ A1 cao
	(ii)		Reason		B1 for mentioning isosceles and equal (or base) angles or equal sides and equal (or base) angles
	(b)	$180 - 95$	85	1	B1 cao
18	(a) (i)		7^5	3	B1 cao
	(ii)		7^4		B2 cao (B1 for sight of 7^5 or 7^{2+3} or 7×7^3 or $7^1 \times 7^3$ or $7^2 \times 7^2$ or 7^{2+3-1})
	(b)		$\frac{1}{2}$	1	B1 for $\frac{1}{2}$ or 0.5 or 2^{-1}
19	(a)		3×10^7	1	B1 cao
	(b)		0.002	1	B1 cao

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
20		Box plot	2	3 aspects: 1st aspect – vertical line for median 2nd aspect – box using correct quartiles 3rd aspect – whiskers (could be single line) drawn with correct end points B2 for fully correct box plot (B1 for 1 aspect)
21 (a)	e.g. 2 | 126 3 | 63 3 | 21 7	$2\times3\times3\times7$	2	M1 for a systematic method of at least 2 correct divisions by a prime number oe factor trees; can be implied by digits 2, 3, 3, 7 on answer line. A1 for $2\times3^2\times7$ or $2\times3\times3\times7$
(b)	$2\times3\times7$	42	2	B2 cao (B1 for 6, 14, 21 or $2\times3\times7$)
22	$\frac{8}{3}\times\frac{5}{4}=\frac{8\times5}{3\times4}=\frac{40}{12}$	$3\frac{1}{3}$	3	B1 for $\frac{8}{3}$ oe improper fraction or $\frac{5}{4}$ oe improper fraction M1 (dep on B1) for multiplying numerator and denominator of "$\frac{8}{3}$" and "$\frac{5}{4}$" A1 for $3\frac{1}{3}$ oe mixed number or $\frac{10}{3}$ OR B1 for 1.25 and 2.67 or 2.66(…) M1 (dep on B1) for correct method of multiplication A1 for $3.\dot{3}$

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
23			2	M1 for a relevant pair of intersecting arcs A1 for line drawn within guidelines, at least 3cm in length, accept broken line [SC: B1 for line drawn within guidelines if M0]
24 (a)		$-1, 0, 1, 2, 3$	2	B2 cao (-1 each error or omission)
(b)(i)		$x \geq \frac{7}{2}$	3	M1 for $2x \geq 7$, condone use of = sign or wrong equality A1 for $x \geq \frac{7}{2}$ oe as final answer
(ii)		4		SC:B1 for 3.5 or $\frac{7}{2}$ seen if M0 B1 ft from $x \geq$ "$\frac{7}{2}$"
25	$4x + 2y = 8$ $4x - 10y = 20$ $12y = -12$ $y = -1$ $4x + 2(-1) = 8$ $x = 2.5$	$x = 2.5$ $y = -1$	3	M1 for correct process to eliminate either x or y (condone one arithmetical error) M1 (dep) for substituting found value into either equation A1 for $x = 2.5$, $y = -1$ [SC: B1 for $x = 2.5$ or $y = -1$ if M0]

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
26		Interior angle of hexagon = 180 – (360 ÷ 6) = 120 360 – (90 + 120)	150	4	Alternative 1 M1 for 360 ÷ 6 A1 for 60 M1 (dep on M1) for "60" + 90 A1 cao Alternative 2 M1 for 360 ÷ 6 A1 for 60 M1 (dep on M1) for 360 – (2 × "60" + 90) A1 cao Alternative 3 M1 for (6 – 2) × 180 ÷ 6 A1 for 120 M1 (dep on M1) for 360 – (90 + "120") A1 cao
27	(a)		(16), 50, 82, 96, 100	1	B1 cao
	(b)	Cumulative freq. diag. curve/ segments	Cum. freq graph	2	B1 for 4 or 5 points plotted correctly ± 1 full (2mm) square depending on sensible table (condone 1 addition error) B1 (dep) for points joined by curve or line segments provided no gradient is negative - ignore any part of graph outside range of their points. (SC:B1 if 4 or 5 points plotted not at end but consistent within each interval and joined)
	(c)	100 – 42	58	2	M1 (ft dep on graph being cf) for reading from graph at 18 or 19, can be implied by answer in range 40 to 46 A1 for answer in range 56 to 60 or ft for 100 – '42' ±1 full (2mm) square

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
1	(a)	900 × 1.70 =	1530	2	M1 900 × 1.7(0) or digits 153(0) seen A1 cao
	(b)	160 ÷ 1.70 =	£94.12 or £94.11	2	M1 160 ÷ 1.7(0) or digits 941(....) seen A1 cao
2	(a)	4.7 ÷ 5.9 = 0.796610169	0.7966..	2	B2 for 0.7966 or better (B1 for 0.8, 0.80, 0.79, 0.796, 0.797 or digits 59 seen)
	(b)	0.82, 0.8, 0.85, 0.66, 0.875 0.66, 0.8, 0.82, 0.85, 0.875 2/3, 4/5, 0.82, 85%, 7/8	2/3, 4/5, 0.82, 85%, 7/8	2	B2 correct order (oe decimals in order) (B1 correct order reversed, or one error in ordered listing) with or without decimal equivalents. NB Accept 0.67 or 0.66
3		**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	3	B3 all 6 correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
4	(a)	–3 (–1) (1) 3 5 7	**–3**, 3, **5**, **7**	2	B2 all correct (B1 2,3 correct)
	(b)	(-2,-3),(-1,-1),(0,1),(1,3),(2,5),(3,7)	line	2	B2 cao for line from $x = -2$ to $x = 3$ (B1 plotting at least 5 points correctly or single line passing through (0, 1) or single line of gradient 2) The six possible points are: (-2,-3), (-1,-1), (0,1), (1,3), (2,5), (3,7)
	(c)(i)		–2	2	B1 $y = -2$ or ft from line segment
	(ii)		2.5		B1 $x = 2.5$ or ft from line segment

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
5	(a) (i)	180 – 54 (=126) "126" ÷ 2	63	3	M1 for (180–54) ÷ 2 A1 cao
	(ii)		Reason		B1 (indep) angles in triangle add to 180 OR equal angles in isosceles triangle OR equal angles and 2 sides the same (B0 if any incorrect reasoning given eg parallel, equilateral triangle)
	(b)	180 – "x"	117	1	B1 117 or ft 180 – "x" if $x < 90$
6		15/120×360 = 45 Plain 40/120×360 = 120 Cheese & Onion 55/120×360 = 165 Salt & Vinegar 10/120 × 360 = 30 Beef	45°, 120°, 165°, 30°	4	M1 evidence of method for at least one angle (could be implied by one correct angle of four on pie chart or in the table) A2 All four angles drawn ±2° tolerance, any order (A1 at least 2 angles correctly drawn ±2°, or all 4 angles in the table) B1 (dep on at least 1 angle drawn correctly, and exactly 4 sectors) for labels (flavour or frequency; initials will do) NB: Ignore the table if the pie chart provides the marks.
7	(a)	260 – 50 = 210 210 ÷ 35 =	6	3	M1 for 260–50 or 210 seen. M1 for "260-50" ÷ 35 or 210÷35 A1 cao
	(b)		P=35h + 50	3	B3 for P=35h+50 or P=35×h+50 oe (B2 for correct RHS or P=h + 50 ×35 or P=35h+k where k is numerical oe) (B1 for P = some other linear expression in h, OR h + 50 ×35 OR 35h seen) NB: P=h scores no marks; ignore £ signs. SC B2 for $h = \frac{P-50}{35}$
8	(a)		Elevation	2	B2 for 4 vertical squares. Accept 4 by 1 rectangle. (B1 for 4 vertical squares with one square added or one parallelogram added at the top, or 3 vertical squares, or 4 horizontal squares)

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
(b)		Plan	2	B2 for 2 adjacent squares, vertical or horizontal. Accept 2 by 1 rectangle. (B1 for 3 adjacent horizontal or vertical squares or a rectangle with sides in the ratio 2:1)
9 (a)		Points	1	B1 all three points ±1 full square.
(b)		Negative	1	B1 Negative. Ignore other descriptors unless contradictory.
(c)		lobf	1	B1 A single straight line drawn to cross between (5,30), (5,40) and (40,0), (40,15); accept freehand if considered to be straight.
(d)(i) (ii)		18–25 30–40	2	B1 18g–25g inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient. B1 30–40 min inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient.
10		300, 3, 75, 150	3	B3 for 4 correct (B2 for 2 or 3 correct) (B1 for 1 correct)
11 (a)	$6x - 7 + 7 = 38 + 7$ $6x = 45$	7.5	2	M1 $6x = 45$ or +7 both sides A1 7.5 oe; accept 45/6
(b)	$5y - 2 = 10$ or $20y - 8 = 40$ $5y = 12$ $20y = 48$	$2\frac{2}{5}$	3	M1 $20y–8(=40)$ or $\frac{4(5y-2)}{4} = \frac{40}{4}$ or $5y–2=10$ M1 (indep) for correct rearrangement into the form $ay=b+c$ or better (eg $20y=40+8$ or $5y=10+2$, using own terms) A1 for $2\frac{2}{5}$, 2.4 oe
12	5 miles = 8 km 70mph÷5×8 = 122 km/h OR 120km/h ÷ 8 × 8 = 75 mph Faster than 70 mph	70mph (Great Britain) (112 km)	3	M1 5 miles = 8 km; OR 70 mph is about 100 km/h OR 1km=0.6(25) miles OR 1mile=1.6km oe M1 70 ÷ 5 × 8 (=112) or 120 ÷ 8 × 5 (=75) oe A1 (dep on at least M1) GB or 70 mph Refer to both answer line and working. NB GB or 70 mph without working scores 0 marks.

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
13 (a)	$\pi \times 0.65$	2.04–2.05	2	M1 for $\pi \times 0.65$ or 3.14×0.65 or 3.142×0.65 oe A1 2.04–2.05 SC Award B1 for 2.0 seen (not 2)
(b)	1000 ÷ "(a)"	487–491	2	M1 for 1000 (or 100) ÷ "(a)" A1 for 487–491
14	$45.00 + 45.00 \times \frac{15}{100} =$ $45.00 + 6.75 =$	51.75	3	M2 for $45.00 + 45.00 \times \frac{15}{100}$ oe or 45.00×1.15 oe OR $45.00 + 6.75$ OR complete method or 5175 seen. (M1 for $45.00 \times \frac{15}{100}$ oe OR 6.75 seen OR 675 seen OR correct method for calculating 15% of 45) A1 cao SC Award B2 for an answer of 38.25
15		273	2	M1 for 728 ÷ 8 or 728 ÷ "3+5" or 91 A1 cao SC B1 for 455, or for 273:455
16	3 24 3.7 46.9(53) 4 60 3.8 51.0(72) 3.1 26.6(91) 3.9 55.4(19) 3.2 29.5(68) 3.21 29.8(66…) 3.3 32.6(37) 3.22 30.1(66…) 3.4 35.9(04) 3.23 30.4(68..) 3.5 39.3(75) 3.24 30.7(72...) 3.6 43.0(56) 3.25 31.0(78...) or 31	3.2	4	B2 for trial between 3.2 and 3.3 inclusive (B1 for trial between 3 and 4 inclusive) B1 for different trial between 3.21 and 3.25 inclusive B1 (dep on at least one previous B1) cao for 3.2 as final answer. NB: embedded answers: –B1; award Bs for evaluations rounded or truncated to at least 1 dp or for 31

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
17	$45^2 + 34^2 =$ $2025 + 1156 = 3181$ $\sqrt{3181} = 56.4$	56	4	M1 for $45^2 + 34^2$ M1 (dep) for $\sqrt{(2025 + 1156)}$ A1 for 56.4 … B1 for rounding their diagonal to the nearest integer (dep on evidence from decimal) NB 56 as the final answer gets full marks. NB Scale drawings result in 0 marks.
18	$2000 \times (1.055)^3$ Interest = 2348.48 – 2000 =	348.48	3	M1 for 5.5/100 × 2000 (oe) or 330 or 2330 or 110 or 2110 M1 (dep) for 5.5/100 × (2000 + "110" + "116.05") or 122.4… seen A1 cao (accept only 348.48 or 348.49) **OR** M2 for $2000 \times (1.055)^3$ or 2348.48(…) or 2348.49 seen (M1 for $2000 \times (1.055)^n$, $n \neq 3$) A1 for 348.48 or 348.49 [SC: B2 for 2348.48 - 2348.49]
19		Line	2	B2 line fully within tramlines, crossing AB and CD (B1 a straight line which crosses AB within the tramline, and also crosses CD) NB: Accept dotted or dashed lines, but not curves; accept freehand if considered to be straight. SC B1 for a perpendicular bisector of AB that is at least half way from AB to CD within the tramlines
20 (a)		$90<t\leq100$	1	B1 for $90<t\leq100$; accept 90-100.
(b)	65×12=780 75×22=1650 85×23=1955 95×24=2280 105×19=1995 8660/100=	86.6	4	M1 for use of fx with x consistent within intervals (including end points). Allow one slip. M1 (dep) for use of midpoints M1 (dep on 1st M1) for use of $\sum fx/100$ or $\sum fx/\sum f$ A1 86.6

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
21	(a)	$8 \times \frac{10}{4}$	20	2	M1 $\frac{10}{4}$ or $\frac{4}{10}$ or 0.4 or 2.5 oe seen A1 cao NB ratios get M0 unless of the form 1:*n* OR M1 $\frac{8}{4}, \frac{4}{8}$ oe seen A1 cao
	(b)	$15 \times \frac{4}{10}$	6	2	M1 $15 \times \frac{4}{10}$ oe A1 cao
22	(a)	$x^2 - 4x + 3x - 12 = x^2 - x - 12$	$x^2 - x - 12$	2	M1 for exactly 4 terms correct ignoring signs (eg x^2, 4*x*, 3*x*, 12) or 3 correct terms out of 4 terms with correct signs (eg 3 out of 4 of x^2, –4*x*, +3*x*,–12) A1 cao
	(b)	$(x+2)(x+5)$	$(x+2)(x+5)$	2	B2 cao (B1 for exactly one of $(x+2)$, $(x+5)$)
	(c)	$6=15+4q-20$ $6-15=4(q-5)$ $p-3t=4q-4t$ $6-3\times5=4(q-5)$	2 ¾	3	M1 for correct substitution of *p* and *t*. M1 for correct expansion of $4(q-t)$ oe (eg $4q-20$, $4q-4t$) A1 11/4 or 2 ¾ or 2.75 OR M1 for correct substitution of *p* and *t*. M1 for $\frac{p-3t}{4} = q-t$ oe A1 11/4 or 2 ¾ or 2.75

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
23	(a)	$9.9\times10^8 - 6.0\times10^7$	9.3×10^8	2	M1 for $99\times10^7 - 6\times10^7$ or $9.9\times10^8 - 0.6\times10^8$ or conversion of either to an ordinary number, or 930000000 or 93×10^7 or 9.3×10^n where n is any positive integer A1 cao
	(b)	$\frac{6.0-4.5}{4.5}\times100 = \frac{1.5}{4.5}\times100 =$ or $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}\times100$	33.3%	3	M2 $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}\times100$ oe (M1 for $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}$ or $\frac{6.0\times10^7 - 4.5\times10^7}{6.0\times10^7}\times100$ oe A1 cao OR M2 $\frac{6.0\times10^7}{4.5\times10^7}\times100$ -100 = (M1 $\frac{6.0\times10^7}{4.5\times10^7}\times100$ or 133.33(%)) A1 cao NB Accept any of the above expressions without any reference to 10^7.
24		LCM (40,24) = 120 Bread buns 120÷40 Burgers 120÷24 OR Bread buns: 40 is $2\times2\times2$ (×5) Burgers: 24 is $2\times2\times2$ (×3)	Bread buns 3 Burgers 5	3	M1 attempt to find LCM by eg lists of multiples, or summing of 40 and 24. A1 identify 120 (as LCM) A1 cao (both) OR M1 expansion of either number into its prime factors in a factor tree or 8×5 or 8×3 A1 both expansions correct A1 cao (both) SC B2 if answers given the wrong way around

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
25	8 × 41 = 328 2 × 29 = 58 328 – 58 = 270 270 ÷ 6 = 45	45	3	M1 for either 8 × 41 (=328) or 2 × 29 (=58) M1 (dep) “328” – “58” (=270) A1 cao NB 328 and/or 58 on the answer line gets M1 (implied); 270 on the answer line gets M2 (implied)

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
1 (a)	$1 - (0.2 + 0.3 + 0.1)$	0.4	2	M1 for $1 - (0.2 + 0.3 + 0.1)$ A1 for 0.4 oe, accept $\frac{0.4}{1}$
(b)	0.2×200	40	2	M1 for 0.2×200 A1 cao NB $\frac{40}{200}$ is M1 A0, 40 out of 200 is M1 A1
2	$650 - 430 = 220$ 1 choc ice costs 110p $650 - 5 \times 110 = 100$p	50	3	M1 for $650 - 430$ or 220 or 110 oe seen M1 for $650 - 5 \times \frac{'220'}{2}$ or $430 - 3 \times \frac{'220'}{2}$ oe A1 for 50p or £0.50 or £0.5 *Alternative scheme* $2x + 5y = 650$ $2x + 3y = 430$ oe M1 for subtracting two simultaneous equations to eliminate x (lollies)(2 or 3 terms correct) M1 for $650 - 5 \times 'y'$ or $430 - 3 \times 'y'$ oe A1 for 50p or £0.50 or £0.5 *Alternative scheme* M1 for $3\times (2x + 5y = 650)$ evaluated and $5 \times (2x + 3y = 430)$ evaluated oe (5 or 6 terms correct) M1 for subtraction of equations to eliminate y (choc ices)(2 or 3 terms correct ft) A1 for 50p or £0.50 or £0.5

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
3		question + response boxes oe	2	1st aspect: One question (eg 'how long does it take you to travel to school?' or 'What time did you leave home to get to school?'); ignore other questions. 2nd aspect: Response list (at least two), not overlapping. 3rd aspect: Some mention of units (eg minutes) in either question or responses B2 for all three aspects, or B1 for just one aspects.
4	2 [(3×1) + (4×1)] + (3×6) + (1×6) + (2×6) + (4×6) + (1×6) + (5×6)	110 cm^2	4	M1 for attempt to find the area of one face M1 for at least 6 faces with intention to add A1 cao B1 (indep) for cm^2 (with or without numerical answer)
5	$\left(\frac{-6+4}{2}\right), \left(\frac{5+3}{2}\right)$	(–1, 4)	2	B2 cao [B1 for (–1,*a*) or (b,4) or (4,-1)]
6		Box plot	2	3 aspects: 1st aspect – vertical line for median 2nd aspect – box using correct quartiles 3rd aspect – whiskers (could be single line) drawn with correct end points B2 for fully correct box plot (B1 for 1 aspect)
7			2	M1 for a relevant pair of intersecting arcs A1 for line drawn within guidelines, at least 3cm in length, accept broken line [SC: B1 for line drawn within guidelines if M0]

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
8	(a)	e.g. 2 | 126 3 | 63 3 | 21 7	$2\times3\times3\times7$	2	M1 for a systematic method of at least 2 correct divisions by a prime number oe factor trees; can be implied by digits 2, 3, 3, 7 on answer line. A1 for $2\times3^2\times7$ or $2\times3\times3\times7$
	(b)	$2\times3\times7$	42	2	B2 cao (B1 for 6, 14, 21 or $2\times3\times7$)
9	(a)		$-1,0,1,2,3$	2	B2 cao (-1 each error or omission)
	(b)(i)		$x\geq\frac{7}{2}$	3	M1 for $2x\geq7$, condone use of = sign or wrong equality A1 for $x\geq\frac{7}{2}$ oe as final answer
	(ii)		4		SC:B1 for 3.5 or $\frac{7}{2}$ seen if M0 B1 ft from $x\geq$ "$\frac{7}{2}$" or $x>$"$\frac{7}{2}$"
10	(a)(i)		7^5	3	B1 cao
	(ii)		7^4		B2 cao (B1 for sight of 7^5 or 7^{2+3} or 7×7^3 or $7^1\times7^3$ or $7^2\times7^2$ or 7^{2+3-1})
	(b)		$\frac{1}{2}$	1	B1 for $\frac{1}{2}$ or 0.5 or 2^{-1}
11	(a)	$5n=m+21$	$n=\frac{m+21}{5}$	2	M1 for $5n=m+21$ or for attempt to divide three terms by 5 A1 $n=\frac{m+21}{5}$ oe

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
(b)	$4p-8q=3p+2$ $p-8q=2$ $p=8q+2$	$p=8q+2$	3	M1 for $4p-aq$ or $\frac{3}{4}p+b$ where a is an integer and b is a number M1 (dep) for taking one term correctly to LHS or RHS of expression A1 $p=8q+2$ oe
12		$y=\frac{1}{2}x+3$	2	B2 for $y = ½x + 3$ oe (B1 for $y = ½x + c$, $c \neq 7$ or $y = mx + 3$ oe or $½x + 3$ or $M = ½x + 3$)
13	$\frac{8}{3}\times\frac{5}{4}=\frac{8\times5}{3\times4}=\frac{40}{12}$	$3\frac{1}{3}$	3	B1 for $\frac{8}{3}$ oe or $\frac{5}{4}$ oe M1 (dep on B1) for multiplying numerator and denominator of "$\frac{8}{3}$" and "$\frac{5}{4}$" A1 for $3\frac{1}{3}$ oe mixed number or $\frac{10}{3}$ *OR* B1 for 1.25 and 2.67 or 2.66(…) M1(dep on B1) for correct method of multiplication A1 for $3.\dot{3}$
14	$4x+2y=8$ $4x-10y=20$ $12y=-12$ $y=-1$ $4x+2(-1)=8$ $x=2.5$	$x=2.5$ $y=-1$	3	M1 for correct process to eliminate either x or y (condone one arithmetical error) M1 (dep) for substituting found value into either equation A1 for $x = 2.5$, $y = -1$ [SC: B1 for $x = 2.5$ or $y = -1$ if M0]

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
15		Interior angle of hexagon = 180 – (360 ÷ 6) = 120 360 – (90 + 120)	150	4	Alternative 1 M1 for 360 ÷ 6 A1 for 60 M1 (dep on M1) for "60" + 90 A1 cao Alternative 2 M1 for 360 ÷ 6 A1 for 60 M1(dep on M1) for 360 – (2 × "60" + 90) A1 cao Alternative 3 M1 for (6 – 2) × 180 ÷ 6 A1 for 120 M1(dep on M1) for 360 – (90 + "120") A1 cao
16	(a)		(16), 50, 82, 96, 100	1	B1 cao
	(b)	Cumulative freq. diag. curve/ segments	Cum. freq graph	2	B1 ft for 4 or 5 points plotted correctly ± 1 full (2mm) square depending on sensible table (condone 1 addition error) B1 (dep) for points joined by curve or line segments provided no gradient is negative - ignore any part of graph outside range of their points. (SC:B1 if 4 or 5 points plotted not at end but consistent within each interval and joined)

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
(c)	$100 - 42$	58	2	M1 (ft dep on graph being cf) for reading from graph at 18 or 19, can be implied by answer in range 40 to 46 A1 for answer in range 56 to 60 or ft for 100 –'42' ±1 full (2mm) square
17 (a)		0.6 and 0.7,0.3,0.7	2	B1 for 0.6 on LH branch B1 for 0.7, 0.3 and 0.7 on RH branches
(b)	0.4×0.3	0.12	2	M1 for 0.4×0.3 A1 0.12 oe
(c)	$0.4\times0.7+0.6\times0.3$	0.46	3	M1 for '0.4×0.7' or '0.6×0.3' M1 for addition of two products from correct branches A1 0.46 oe Alternative M2 for an attempt to evaluate 1-(0.3×0.4 +'0.6×0.7') A1 cao
18	$x = 0.4545...$ so $100x = 45.4545...$ $99x = 45$ $x = \frac{45}{99} = \frac{15}{33}$	proof	3	M1 for $100x = 45.45\ldots$ or $10000x = 4545.45\ldots$ M1 (dep) for subtraction of both sides A1 for $\frac{15}{33}$ from correct proof
19	$\left(\sqrt{3}\right)^2 + \sqrt{3}\sqrt{2} - \sqrt{3}\sqrt{2} - \left(\sqrt{2}\right)^2$ $= 3 - 2$	1	2	B2 cao (B1 for $\sqrt{3}\sqrt{3} + \sqrt{3}\sqrt{2} - \sqrt{3}\sqrt{2} - \sqrt{2}\sqrt{2}$ oe, $\sqrt{3}\sqrt{3} - \sqrt{2}\sqrt{2}$ oe, or for 2,3, $\sqrt{4}$ $\sqrt{6}$ $\sqrt{9}$ seen)
20 (a)	$\frac{96}{24}$ or 4 $\sqrt{4}$ or 2	8	3	M1 for $\frac{96}{24}$ or $\frac{24}{96}$ or 4 or $\frac{1}{4}$ oe M1 for $\sqrt{\frac{96}{24}}$ or $\sqrt{\frac{24}{96}}$ or $\sqrt{'4'}$ or $\frac{1}{\sqrt{'4'}}$ or 2 or $\frac{1}{2}$ oe A1 cao

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
	(b)	12×2^3	96	2	M1 for '2'^{3} or 8 A1 cao
21	(a)	$x\times3-x\times2x^2$	$3x-2x^3$	2	B2 cao (B1 for a two term expression with either $3x$ or $2x^3$)
	(b)		$4x(3y+x)$	2	M1 for taking out a factor of x, $2x$, 2, 4 or $4x$ A1 cao
	(c)	$\frac{5a}{b^2}$	$\frac{5a}{b^2}$	2	B2 for $\frac{5a}{b^2}$ or $5ab^{-2}$ (accept $\frac{5a}{1b^2}$) (B1 for either dealing with the numbers or dealing with the powers of a)
	(d)	$\frac{\cancel{x-3}}{\cancel{(x-3)}(x+3)}$	$\frac{1}{x+3}$	2	M1 for $(x-3)(x+3)$ A1 cao
22	(a)		$\mathbf{a}+\mathbf{b}$	1	B1 $\mathbf{a}+\mathbf{b}$ oe
	(b)	$ED=\mathbf{a}$ $DX=-2\mathbf{b}+2\ AC=2\mathbf{a}$ (So, $DX=2ED$)	$2\mathbf{a}$	3	M1 for $(DX=)\ DA+AX$ A1 for $(DX=)\ -2\boldsymbol{b}+2$ ("$\boldsymbol{a}+\boldsymbol{b}$") A1 $2\boldsymbol{a}$ from fully correct proof
23		$\pi(2x)^2h=\frac{4}{3}\pi(3x)^3$ $h=\frac{\frac{4}{3}\pi(3x)^3}{\pi(2x)^2}$	$9x$	3	M1 for $\pi(2x)^2h=\frac{4}{3}\pi(3x)^3$ (condone absence of brackets) M1 (dep) for valid algebra that gets to $h=ax$ (condone one error in powers of numerical constants) A1 cao

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
24	(i)	$n^2+(n+1)^2=2(n^2+n)+1$	$2n^2+2n+1$	4	M1 for at least 3 terms correct from $n^2+n+n+1$ A1 for $2n^2+2n+1$ oe
	(ii)		$2(n^2+n)$ is always even so $2(n^2+n)+1$ is always odd		M1 for recognizing $2n^2$ is always even A1ft complete proof for their quadratic *Alternative method* M1 for recognizing that if n^2 is odd then $(n+1)^2$ is even or vice versa A1 for complete proof
25			42 318	2	B1 for answer in range 36 – 48 B1 for answer in range 312 – 324
26	(a)	$(x-3)^2-3^2+15$	$p=3, q=6$	2	B2 for $p=3$ and $q=6$ (B1 for $p=3$ OR $q=6$) SC: award B2 for $(x-3)^2+6$ if p and q are not identified
	(b)		Sketch	2	B1 for U shaped curve B1 ft for TP in first quadrant (ft if TP not in first quadrant)
27	(a)	Graph translated 2 units upwards through points (–4, 2), (–2, 4), (0,2) and (3,5)	Sketch	2	M1 for a vertical translation A1 curve through points (–4, 2), (–2, 4), (0,2) and (3,5) ± ½ square
	(b)	Graph reflected in x-axis through points (–4,0), (–2, –2), (0, 0) and (3, -3)	Sketch	2	M1 for reflection in x-axis or y-axis A1 curve through points (–4,0), (–2, –2), (0, 0) and (3, -3) ± ½ square

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
1		273	2	M1 for 728 ÷ 8 or 728 ÷ "3+5" or 91 A1 cao SC B1 for 455 or 273:455
2		$3n - 1$	2	B2 for $3n - 1$ oe (B1 for $3n + k$ where $k \neq -1$ but k could be 0)
3	3 24 3.7 46.9(53) 4 60 3.8 51.0(72) 3.1 26.6(91) 3.9 55.4(19) 3.2 29.5(68) 3.21 29.8(66…) 3.3 32.6(37) 3.22 30.1(66…) 3.4 35.9(04) 3.23 30.4(68..) 3.5 39.3(75) 3.24 30.7(72...) 3.6 43.0(56) 3.25 31.0(78...) or 31	3.2	4	B2 for trial between 3.2 and 3.3 inclusive (B1 for trial between 3 and 4 inclusive) B1 for different trial between 3.21 and 3.25 inclusive B1 (dep on at least one previous B1) cao for 3.2 NB: embedded answers: –B1; award Bs for evaluations rounded or truncated to at least 1 dp or for 31
4	2 is the only even prime number and the product of 2 odd numbers is odd	Yes	2	B2 for 'yes' and '2 is the only even prime number and the product of two odd numbers is odd' oe (B1 for 'yes' and either '2 is the only even prime number' oe **or** 'the product of two odd numbers is odd' oe)
5	$45^2 + 34^2 =$ $2025 + 1156 = 3181$ $\sqrt{3181} = 56.4$	56	4	M1 for $45^2 + 34^2$ M1 for $\sqrt{(2025 + 1156)}$ A1 for 56.4 … B1 for rounding their diagonal to the nearest integer (dep on evidence from decimal) NB 56 with no incorrect working as the final answer gets full marks. NB Scale drawings result in 0 marks.

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
6	2000 × $(1.055)^3$ Interest = 2348.48 – 2000 =	£348.48	3	M1 for 5.5/100 × 2000 (oe) or 330 or 16.5/100×2000 or 2330 or 110 or 2110 M1 (dep) for 5.5/100 × (2000 + "110" + "116.05") or 122.4… seen A1 cao (accept only 348.48 or 348.49) **OR** M2 for 2000 × $(1.055)^3$ or 2348.48(…) or 2348.49 seen (M1 for 2000 × $(1.055)^n$, $n \neq 3$) A1 for 348.48 or 348.49 [SC: B2 for 2348.48 - 2348.49]
7 (a)		Line	2	B2 line fully within tramlines, crossing AB and CD (B1 a straight line which crosses AB within the tramline, and also crosses CD) NB: Accept dotted or dashed lines, but not curves; accept freehand if considered to be straight. SC B1 for the perpendicular bisector of *AB* reaching halfway or more from *AB*
(b)		Region		B2 correct arc ±2mm and shaded within. Allow dotty or continous arc. (B1 inaccurate arc and shaded or accurate arc unshaded)
8	65×12=780 75×22=1650 85×23=1955 95×24=2280 105×19=1995 8660/100=	86.6	4	M1 for use of fx with x consistent within intervals (including end points). Allow one slip even if outside interval M1 (dep) for use of midpoints M1 (dep on 1st M1) for use of $\sum fx/100$ or $\sum fx/\sum f$ A1 86.6

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
9		Distance ÷ time: 1400 ÷ 2 h 20 min 20 mins is $\frac{1}{3}$ hour 1400 × 3 ÷ 7 =	600 kph	3	B1 20 mins as $\frac{1}{3}$ hour or as 0.33……hour M1 for distance ÷ time eg 1400 ÷ "2h 20 min" A1 cao OR B1 2 hour 20min = 140 (min) M1 Speed = $\frac{1400}{140}$ = (10 km per minute) A1 cao
10	(a)	$(x+18)+2x+(2x+7)=180$	Equation	2	B2 for $(x+18)+2x+(2x+7)=180$ oe (B1 for $(x+18)+2x+(2x+7)$)
	(b)	$5x+25 = 180$ $5x = 155$	31	2	M1 for simplifying to at least $5x+25=180$ or 360 (may be earned in (a)) A1 for $x=31$
11	(a)	(–14) –4 (0) 4 14	–4, 4, 14	2	B2 for all 3 values correct (B1 for 1 or 2 values correct)
	(b)		curve	2	B1 for all 5 points plotted correctly ± ½ square (ft from table if at least B1 awarded in (a)) B1 (indep) ft for any smooth curve through their points

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
12	(a)	$8 \times \frac{10}{4}$	20	2	M1 $\frac{10}{4}$ or $\frac{4}{10}$ or 0.4 or 2.5 oe seen A1 cao NB ratios get M0 unless of the form 1:*n* OR M1 $\frac{8}{4}, \frac{4}{8}$ oe seen A1 cao
	(b)	$15 \times \frac{4}{10}$	6	2	M1 $15 \times \frac{4}{10}$ oe A1 cao
13	(a)	$9.9\times10^8 - 6.0\times10^7$	$\pm 9.3\times10^8$	2	M1 for $99\times10^7 - 6.0\times10^7$ or $9.9\times10^8 - 0.60\times10^8$ or correct conversion of either to an ordinary number or 930000000 or 9.3×10^n where *n* is any positive integer A1 cao
	(b)	$\frac{6.0-4.5}{4.5} \times 100 = \frac{1.5}{4.5} \times 100 =$ or $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7} \times 100$	33.3%	3	M2 $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7} \times 100$ oe (M1 for $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}$ or $\frac{6.0\times10^7 - 4.5\times10^7}{6.0\times10^7} \times 100$ oe A1 33.3 – 33.4 OR M2 $\frac{6.0\times10^7}{4.5\times10^7} \times 100$ -100 or 33.3% (M1 $\frac{6.0\times10^7}{4.5\times10^7} \times 100$ or 133.3%) A1 33.3-33.4 NB Accept any of the above expressions without any reference to 10^7.

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
14	LCM (40,24) = 120 Bread buns 120÷40 Burgers 120÷24 OR Bread buns: 40 is $2 \times 2 \times 2$ (×5) Burgers: 24 is $2 \times 2 \times 2$ (×3)	Bread Buns 3 Burgers 5	3	M1 attempt to find LCM by eg lists of multiples, or summing of 40s and summing of 24s, with at least 3 numbers in each list A1 identify 120 as LCM A1 cao (both) **OR** M1 expansion of either number into its prime factors in a factor tree or 8×5 or 8×3 A1 both expansions correct A1 cao (both) SC B2 if answers given the wrong way around
15	$\sin 32 = \frac{AB}{12}$ $AB = 12 \times \sin 32$ $AB = 6.35903\ldots$	6.36	3	M1 $\sin 32 = \frac{AB}{12}$ (accept $\text{Sin}\ \frac{AB}{12}$) M1 $12 \times \sin 32$ or $12 \times 0.5299..$ A1 accept 6.359 – 6.360 SC Gradians 5.78(1…) Radians 6.62 Get M1M1A0 **OR** Use of Sine Rule $\frac{\sin 32}{AB} = \frac{\sin 90}{12}$ or $\frac{AB}{\sin 32} = \frac{12}{\sin 90}$ M1 $AB = \frac{12 \times \sin 32}{\sin 90}$ M1 AB = 6.359 – 6.36 A1 SC Gradians 5.85(…) Radians 7. 40(…) M1M1A0

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
16		$8 \times 41 = 328$ $2 \times 29 = 58$ $328 - 58 = 270$ $270 \div 6 = 45$	45	3	M1 for either 8×41 (=328) or 2×29 (=58) M1 (dep) "328" – "58" (=270) A1 cao NB 328 and /or 58 on the answer line gets M1 (implied); 270 on the answer line gets M2 (implied)
17	(a)	$x^2 - 4x + 3x - 12 = x^2 - x - 12$	$x^2 - x - 12$	2	M1 for exactly 4 terms correct ignoring signs (x^2, $4x$, $3x$, 12) or 3 out of 4 terms with correct signs (x^2, $-4x$, $+3x$, -12) A1 cao
	(b)	$6x^2 - 8x + 15x - 20 = 6x^2 + 7x - 20$	$6x^2 + 7x - 20$	2	M1 for exactly 4 terms correct ignoring signs ($6x^2$, $8x$, $15x$, 20) or 3 out of 4 terms with correct signs ($6x^2$, $-8x$, $+15x$, -20) A1 cao
	(c)	$(x+2)(x+5)$	$(x+2)(x+5)$	2	B2 cao (B1 for exactly one of $(x + 2)$, $(x + 5)$)
	(d)		$12p^8q^3$	2	B2 cao (B1 for any 2 out of 3 terms correct in a product or 3 terms correct in a sum or part product)
	(e)	$6 = 15 + 4(q - 5)$ $6 = 15 + 4q - 20$ $11 = 4q$	$2\frac{3}{4}$	3	M1 for correct substitution of p and t. M1 for correct expansion of $4(q-t)$ oe (eg $4q-20$, $4q-4t$) A1 11/4 or 2 ¾ or 2.75 OR M1 for correct substitution of p and t. M1 for $\frac{p-3t}{4} = q - t$ oe A1 11/4 or 2 ¾ or 2.75
18	(a)	$T=kx$; $150=6k$; $k=25$	$T=25x$	3	M1 for $T=kx$, k algebraic M1 subs $T=150$ and $x=6$ into $T = kx^n$ $(n \neq 0)$ A1 for $T=25x$ oe SC B1 $T \propto 25x$ oe

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
	(b)	$T = 25 \times 15 =$	375	1	B1 ft on k , $k \neq 1$
	(c)	$600 = 25x$; $x = 600 \div 25 =$	24	1	B1 ft on k , $k \neq 1$
19		$\frac{120}{570} \times 50 = 10.5263$ $\frac{250}{570} \times 50 = 21.9298$ $\frac{200}{570} \times 50 = 17.5438$	10 22 18	3	M1 method shown eg $\frac{120}{120+250+200} \times 50$ or one of 10.5(263) 21.9(298), 17.5(438) **OR** 570 ÷ 50 = 11.4 and one of 120 ÷ 11.4 =10.526…) , 270 ÷ 11.4 (=21.9298…) , 200 ÷ 11.4 (=17.5438…) A1 for 10.5(263), 21.9(298), 17.5(438) or 11, 22, 18 B1 correction to add to 50: 10, 22, 18 or 11, 22, 17
20		$x(x+1) + 4(2x–3) = (2x–3)(x+1)$ $x^2 + x + 8x – 12 = 2x^2 + 2x – 3x – 3$ $x^2 – 10x + 9 = 0$ $(x – 9)(x – 1) = 0$	$x = 1, 9$	5	M1 for multiplying through by common denominator $(2x–3)(x+1)$ M1 (dep)for either $x^2 + x + 8x – 12$ or $2x^2 + 2x – 3x – 3$ oe A1 for correct quadratic (= 0) M1 for a correct method to solve 3 term quadratic A1 cao for both solutions
21		$AB^2 = 8^2 + 9^2 – 2\times8\times9\times\cos40$ $AB^2 = 64 + 81 – 144\times\cos40$ $AB^2 = 145 – 144 \times 0.766$ $AB^2 = 145 – 110.31… = 34.6896$ $AB = \sqrt{34.6796} = 5.8897877$	5.89	3	M1 Subs in Cos Rule: $8^2 + 9^2 – 2\times8\times9\times\cos40$ M1 correct order of evaluation of $8^2 + 9^2 – 2\times8\times9\times\cos40$ A1 cao 5.88 – 5.89 SC: Award B2 for one of $AB^2 = 241.03…$ or $AB = 15.525…$ (radians) $AB^2 = 28.50…$ or $AB = 5.33…$ (gradians)
22	(a)		24 30	2	B1 24 cao B1 30 cao
	(b)	Column $60 \leq w < 75$ at 10 Column $75 \leq w < 95$ at 4	10 4	2	B1 10 and correct width (tol $\pm\frac{1}{2}$ small square) B1 4 and correct width (tol $\pm\frac{1}{2}$ small square)

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
23 (a)		100.5	1	B1 cao
(b)		10.515	1	B1 cao
(c)	$\frac{100.5}{10.515}$	9.5577746	2	M1 for greatest distance divided by least time Where 100 < greatest distance ≤ 100.5, 10.51 ≤ least time < 10.52 A1 for 9.555 – 9.56
(d)	$\frac{99.5}{10.525}$	9.45368..	2	M1 for least distance divided by greatest time Where 99.5 ≤ least distance < 100, 10.52 < greatest time ≤10.53 A1 for 9.45 – 9.455
24	$\frac{4}{7}\times\frac{4}{7}\times\frac{3}{7} + \frac{3}{7}\times\frac{3}{7}\times\frac{4}{7} =$ $\frac{48+36}{343} = \frac{84}{343}$ But there are three ways this can be achieved: BBG, BGB, GBB So the probability is $\frac{84}{343}\times 3$ OR $1-\frac{64}{343}-\frac{27}{343}$ NB: 84/343 = 0.244897; 252/343 = 0.73469 $\frac{4}{7} = 0.57(142...)$, $\frac{3}{7} = 0.42(857...)$	$\frac{252}{343}$	3	M1 for $\frac{4}{7}\times\frac{4}{7}\times\frac{3}{7}$ or $\frac{3}{7}\times\frac{3}{7}\times\frac{4}{7}$ oe or $\left(\frac{4}{7}\right)^3$ oe or $\left(\frac{3}{7}\right)^3$ oe or $\frac{91}{343}$ or 0.10(49…) or 0.13(99….) M1 (indep) for identification of all 6 outcomes (M2 for $1-\left[\left(\frac{4}{7}\right)^3+\left(\frac{3}{7}\right)^3\right]$) oe A1 $\frac{252}{343}$, $\frac{36}{49}$, 0.73(469…) oe

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
25	(a)	$\frac{1}{3} \times \pi \times 5^2 \times 8 = \pi \times 25 \times 8 \div 3 =$ 209.4395	209–210	2	M1 for $\frac{1}{3} \times \pi \times 5^2 \times 8$ A1 for 209–210
	(b)	Base radius = $\frac{216}{360} \times 15 = 9$ Height = $\sqrt{(15^2 - 9^2)} =$	12	4	M1 for 216 ÷ 360 A1 for 9 M1 for $\sqrt{(15^2 - \text{“9”}^2)}$, where “9” < 15 A1 cao

PMT

Mark Scheme (Results)

Summer 2016

Pearson Edexcel GCSE
In Mathematics B (2MB01)
Higher (Calculator) Unit 3

PMT

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2016
Publications Code 5MB3H_01_1606_MS
All the material in this publication is copyright
© Pearson Education Ltd 2016

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Note that in some cases a correct answer alone will not score marks unless supported by working; these situations are made clear in the mark scheme. Examiners should be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will award marks for the quality of written communication (QWC).
The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labelling conventions.

ii) *select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 **With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Partial answers shown (usually indicated in the ms by brackets) can be awarded the method mark associated with it (implied).
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks; transcription errors may also gain some credit. Send any such responses to review for the Team Leader to consider.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 **Follow through marks**

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 **Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.

10 **Probability**

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded (embedded answers).

12 Parts of questions
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers
Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

Guidance on the use of codes within this mark scheme
M1 – method mark A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

PMT

PAPER: 5MB3H/01					
Question		**Working**	**Answer**	**Mark**	**Notes**
1*			Decision (supported)	3	M1 for correct first step e.g. 15 ÷ 3 M1 for complete method C1 for correct decision with accurate figures
2	(a)		Correct diagram	2	B2 cao (B1 for correct orientation)
	(b)		Enlargement, sf 4, centre (0, 1)	3	B1 for enlargement B1 for scale factor 4 B1 for centre (0, 1)
3			$w = \frac{d+5}{2}$	2	M1 for attempt to add 5 to both sides or divide both sides by 2 as a first step, eg $d + 5 = 2w$ oe or $\frac{d}{2} = w - \frac{5}{2}$ oe A1 for $w = \frac{d+5}{2}$ oe

PMT

PAPER: 5MB3H/01					
Question		Working	Answer	Mark	Notes
4			22.5	4	M1 for $4(x - 8) = 2x + 13$ M1 for expansion of bracket or division of all terms by 4, eg $4x - 32 = 2x + 13$, or $x - 8 = \frac{2x}{4} + \frac{13}{4}$ M1 for isolating x and number terms eg $2x = 45$, $\frac{x}{2} = \frac{45}{4}$ A1 for $\frac{45}{2}$ or 22.5 OR M1 for $(180 - 64) \div 2$ (= 58) M1 for $4(x - 8)$ = "58" or $2x + 13$ = "58" or "58" – 13 (= 45) M1 for isolating x and number terms eg $4x = 90$, $2x = 45$ or "45" ÷ 2 A1 for $\frac{45}{2}$ or 22.5 OR M1 for $64 + 4(x - 8) + 2x + 13$ M1 for $64 + 4(x - 8) + 2x + 13 = 180$ M1 for isolating x and number terms eg $6x = 135$ A1 for $\frac{45}{2}$ or 22.5
5			Correct region shaded	3	M1 for a circle centre M or N (accept arc of sufficient length to define the region) M1 for circle centre M radius 5 cm **and** circle centre N radius 3 cm (accept arc of sufficient length to define the region) A1 for correct region shaded

PAPER: 5MB3H/01					
Question		Working	Answer	Mark	Notes
*6			Conclusion (supported)	4	M1 for $\pi \times 120^2$ (= 45 216 – 45 249) M1 for "$\pi \times 120^2$" ÷ 1800 A1 for 25 – 25.2 C1 ft (dep on M2) for appropriate conclusion from their figures OR M1 for $\pi \times 120^2$ (= 45 216 – 45 249) M1 for 20 × 1800 A1 for 36 000 and 45 216 – 45 249 C1 ft (dep on M2) for appropriate conclusion from their figures OR M1 for $\pi \times 120^2$ (= 45 216 – 45 249) M1 for "$\pi \times 120^2$" ÷ 20 A1 for 2260 – 2263 C1 ft (dep on M2) for appropriate conclusion from their figures OR M1 for 1800 × 20 M1 for 36 000 ÷ π (=11 457 – 11 465) A1 for 107(.0…) C1 ft (dep on M2) for appropriate conclusion from their figures
7			161.50	5	M2 for a correct method to decrease 6720 by 20%, eg 6720 × 0.8 (= 5376) or 6720 × 0.2 (= 1344 and 6720 – 1344 (= 5376)) (M1 for a correct method to find 20% of 6720 eg 6720 × 0.2 or $\frac{20}{100} \times 6720$ (= 1344)) M1 for subtracting 1500 (= 3876) after a percentage calculation M1 "3876" ÷ 24 after the subtraction of 1500 A1 for 161.5(0)

PAPER: 5MB3H/01					
Question		**Working**	**Answer**	**Mark**	**Notes**
8		3 42 3.1 45.(291) 3.2 48.(768) 3.3 52.(437) 3.4 56.(304) 3.5 60.(375) 3.6 64.(656) 3.7 69.(153) 3.8 73.(872) 3.9 78.(819) 4 84 3.71 69.6(14811) 3.72 70.(078848) 3.73 70.5(45117) 3.74 71.(013624) 3.75 71.4(84375)	3.7	4	B2 for a trial between 3 and 4 exclusive (B1 for a trial at 3 or 4) B1 for a different trial of $3.7 < x \leq 3.75$ B1 (dep on at least one previous B1) for 3.7 NB. Trials should be evaluated to at least 2 sf truncated or rounded for values of x correct to 1 decimal place. Trials should be evaluated to at least 1 dp truncated or rounded for values of x correct to 2dp. **NB** No working scores 0 marks even if answer is correct
9			618	4	M1 for a method to find an area which is part of the cross section M1 (dep) for a complete method to find the total area of the cross section M1 (dep M1) for their cross sectional area × 12 A1 cao **OR** M1 for a method to find the volume of cuboid, 8 × 8 × 12 (= 768) M1 for a method to find the volume of the triangular prism, ½ ×5 × 5 × 12(= 150) M1 (dep M2) for a complete method for the volume of the prism A1 cao

PMT

5MB3H_01					
Question		Working	Answer	Mark	Notes
10	(a)		15.2	3	M1 for $12.7^2 + 8.3^2$ or 161.29 + 68.89 or 230.18 M1 for √"$(12.7^2 + 8.3^2)$" or √"161.29 + 68.89" or $\sqrt{230.18}$ A1 for 15.2 or 15.17(168…)
	(b)		16.1	3	M1 for $\sin 40° = \frac{DE}{25}$ oe or $\frac{DE}{\sin 40} = \frac{25}{\sin 90}$ oe M1 for 25 sin 40° A1 for 16.1 or 16.06(969…)
11	(a)		4.081	2	M1 for 8.875 or $\frac{71}{8}$ or 2.979…or 4.08 A1 for 4.081 or 4.0809(5…)…
	(b)		4×10^{-3}	2	M1 for 4×10^n, $n \neq -3$ or $a \times 10^{-3}$, $a \neq 4$ or 0.004 or 0.4×10^{-2} or 1/250 A1 cao
12			$x < 2$	2	M1 for an attempt to isolate x and number terms or multiply all terms by 2, or for $\frac{3}{2}x < 3$ or $-\frac{3}{2}x < -3$ or $x = 2$ A1 cao
13	(a)		0, 6, 4, –6	2	B2 for all values correct (B1 for any one value correct)
	(b)		Graph drawn	2	M1 ft (provided B1 in (a) for at least 6 points plotted correctly from their table A1 cao for correct curve drawn from (–4, –6) to (3, –6)
	(c)		1.5 to 1.6 and –2.5 to –2.6	2	M1 for correct use $y = 2$ (may be implied by one correct estimate) or correct use of formula. A1 for 1.5 to 1.6 and –2.5 to –2.6

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
14*		Sher Yas **Old 19300 12560** New 20265 13502 **Inc 965 942**	Correct comparison (supported)	3	M1 for a correct method to calculate the population at the beginning of the 10 year period for at least one place. eg $\frac{20265}{1.05}$ (= 19 300) oe or $\frac{13502}{1.075}$ (= 12 560) oe A1 for 965 (Sherbury) **and** 942(Yaston) C1 (dep on M1) ft for statement comparing increases leading to conclusion based on two comparable amounts eg increase in Sherbury is greater than increase in Yaston
15		$20x + 10y = 35$ $6x - 10y = -48$ $x = -\frac{1}{2}$ $y = \frac{9}{2}$ OR $12x + 6y = 21$ $12x - 20y = -96$ $26y = 117$ $y = \frac{9}{2}$ $x = -\frac{1}{2}$	$-\frac{1}{2}, \frac{9}{2}$	4	M1 for a correct process to eliminate either variable (condone one arithmetic error) A1 cao for either x or y M1 (dep on M1) for correct substitution of found value into one of the equations or appropriate method after starting again (condone one arithmetic error) A1 cao OR M1 for full method to rearrange and substitute to eliminate either variable (condone one arithmetic error) A1 cao for either x or y M1 (dep on M1) for correct substitution of found value into one of the equations or appropriate method after starting again (condone one arithmetic error) A1 cao **NB** Trial and improvement methods score 0 marks unless both x and y are correct
16			226	2	M1 for angle $ABC = 180 - 67$ (= 113) A1 cao OR M1 for obtuse angle $AOC = 2 \times 67$ (= 134) A1 cao

PMT

5MB3H_01					
Question		Working	Answer	Mark	Notes
17			– 2.82, 19.82	3	M1 for $\frac{--17 \pm \sqrt{(-17)^2 - 4\times1\times-56}}{2\times1}$ (condone one sign error) or $\left(x - \frac{17}{2}\right)^2$ oe M1 for $\frac{17 \pm \sqrt{513}}{2}$ or for method leading to $\frac{17}{2} \pm \sqrt{\frac{513}{4}}$ oe A1 for answers in the range 19.82 to 19.825; – 2.82 to –2.825
18			21.2	4	M1 for correct use of sine rule to find *AB* or *BC*, eg $\frac{6}{\sin 25} = \frac{AB}{\sin 120}$ or $\frac{6}{\sin 25} = \frac{BC}{\sin(180-120-25)}$ M1 for (*AB* =) $\frac{6}{\sin 25} \times \sin 120$ (= 12.2(951…)) or (*BC* =) $\frac{6}{\sin 25} \times \sin 35$ (= 8.1(431…) M1(dep M2) for method to find area, eg 0.5 × 6 × "*AB*" sin35 or 0.5 × 6 × "*BC*" sin 120 A1 for answer in the range 21.1 – 21.2 OR (with perpendicular from *A* meeting *BC* extended at a point *X*) M1 for method to find *AX* M1 for method to find *BX* M1(dep M2) for 0.5 × "*AX*" × "*BC*" A1 for answer in the range 21.1 – 21.2
*19			0.38 because the LB and UB agree to that number of figures	5	B1 for 10.75 or 10.85 or 10.84999… B1 for 75.055 or 75.065 or 75.064999… M1 for $\sqrt{\frac{10.85}{75.055}}$ as UB **OR** $\sqrt{\frac{10.75}{75.065}}$ as LB A1 (dep on all previous marks) for both 0.3802… and 0.3784 must clearly come from working with correct values C1 for 0.38 from 0.3784… and 0.3802 … **and** "both LB and UB round to 0.38" **NB** An answer of 038(932….) without working scores no marks

5MB3H_01					
Question		Working	**Answer**	**Mark**	**Notes**
20			7.35	3	M1 for $y \propto \frac{1}{\sqrt{x}}$ or $y = \frac{k}{\sqrt{x}}$ or $9 = \frac{k}{\sqrt{4}}$ M1 for $k = 9\sqrt{4}$ (=18) A1 for 7.35 or 7.348…
*21			$\frac{2}{3}$ (7**a** + 3**b**) and correct conclusion	4	M1 for correct vector for $\overrightarrow{OY}$, or $\overrightarrow{AP}$ eg $(\overrightarrow{OY}) = \overrightarrow{OA} + \overrightarrow{AY}$, or $\overrightarrow{OY} = \overrightarrow{OA} + \frac{2}{3}\overrightarrow{AP}$ may include terms in **a** and **b,** eg. 6**a** + $\overrightarrow{AY}$, 4**b** + 4**a** – **b** + $\overrightarrow{PY}$ or $\overrightarrow{AP} = \overrightarrow{AO} + \overrightarrow{OB} + \overrightarrow{BP}$ or –6**a** + 4**b** + 4**a** – **b** or – 2**a** + 3**b** M1 for $(\overrightarrow{AY}) = \frac{2}{3}\overrightarrow{AP}$ or ⅔(–6**a** + 4**b** + 4**a** – **b**) or ⅔(–2**a** + 3**b**) or $(\overrightarrow{PY} = \frac{1}{3}\overrightarrow{PA}$ or ⅓(– 4**a** + **b** – 4**b** + 6**a**) or ⅓(2**a** – 3**b**) M1 for correct expression for $\overrightarrow{OY}$ in terms of **a** and **b,** eg $\frac{14}{3}$ **a** + 2**b** C1 for $\frac{2}{3}$ (7**a** + 3**b**) and "$\overrightarrow{OY}$ is parallel to the vector 7**a** +3**b**" **oe**

Modifications to the mark scheme for Modified Large Print (MLP) papers.

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q2	(a)	Grid enlarged. Shading changed to dotty shading. Labels added to shapes. Image labelled shape A. Rotation given and labelled shape B. Question changed to 'Describe fully the single transformation which maps shape A to shape B. One row top and bottom, one column left and right has been removed. Three answer lines given.	2 cao (B1 for correct orientation in third quadrant or correct quarter turn)
Q2	(b)	Grid enlarged. Shading changed to dotty shading. Labels added next to the triangles 'triangle P' and 'triangle Q'. One column removed from the left.	B1 for enlargement B1 for scale factor 4 B1 for centre (0, 1)

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q4		Diagram enlarged. MLP only – x changed to y. Braille only – measurements given in the text.	M1 for $4(x - 8) = 2x + 13$ M1 for expansion of bracket or division of all terms by 4, eg $4x - 32 = 2x + 13$, $x - 8 = 2x/4 + 13/4$ M1 for isolating x and number terms eg $2x = 45$, $x/2 = 45/4$ A1 for 45/2 or 22.5 OR M1 for $(180 - 64) \div 2$ (= 58) M1 for $4(x - 8) = 58$ or $2x + 13 = 58$ M1 for isolating x and number terms eg $4x = 90$, $2x = 45$ A1 for 45/2 or 22.5 OR M1 for $64 + 4(x - 8) + 2x + 13$ M1 for $64 + 4(x - 8) + 2x + 13 = 180$ M1 for isolating x and number terms eg $6x = 135$ A1 for 45/2 or 22.5
Q5		Diagram kept the same size as original. Scale moved above the diagram and also put in the question paper.	B1 for circle centre M radius 5 cm B1 for circle centre N radius 3 cm B1 for correct region shaded

PMT

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q6		Diagram enlarged. Cross changed to a filled in circle. Arrow heads removed.	M1 for $\pi \times 120^2$ (= 45238….) M1 for "$\pi \times 120^2$" ÷ 1800 A1 for 25.0 – 25.2 C1 for "surface is large enough as 25 > 20" oe OR M1 for $\pi \times 120^2$ (= 45238….) M1 for 20 × 1800 A1 for 36000 C1 for "surface is large enough as 36000 < 45238..." oe OR M1 for $\pi \times r^2 = 1800 \times 20$ M1 for 36000 ÷ π (=11459..) A1 for 107.0… C1 for "surface is large enough as 107…< 120" oe
Q9		Model provided for all candidates. Diagram enlarged and provided for MLP.	M1 for a method to find an area which is part of the cross section M1 (dep) for a complete method to find the total area of the cross section M1 (dep M2) for a complete method for the volume of the prism A1 cao OR M1 for a method to find the volume of cuboid, 8 × 8 × 12 M1 for a method to find the volume of the triangular prism M1 (depM2) for a complete method for the volume of the prism A1 cao

PMT

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q10	(a)	Diagram enlarged. Braille only: measurements given in the text.	M1 for $12.7^2 + 8.3^2$ or $161.29 + 68.89$ or 230.18 M1 for $\sqrt{161.29 + 68.89}$ or $\sqrt{230.18}$ A1 for 15.2 or 15.17(168…)
Q10	(b)	Diagram enlarged. Braille only – measurements given in the text.	M1 for $\sin 40° = \frac{DE}{25}$ oe M1 for $25 \sin 40°$ A1 for 16.1 or 16.06(969…)
Q13	(a)	Wording added 'There are four spaces to fill.'	B2 for all values correct (B1 for any one value correct)
Q13	(b)	Grid enlarged.	M1 ft for at least 6 points plotted correctly from their table A1 cao for correct curve drawn from (–4, –6) to (3, –6)
Q16		Diagram enlarged. Dot added at centre.	M1 for angle ABC = 180 – 67 (= 113) A1 cao OR M1 for obtuse angle AOC = 2 × 67 (= 134) A1 cao

PMT

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q18		Diagram enlarged.	M1 for correct use of sine rule to find *AB* or *BC*, eg $\frac{6}{\sin 25} = \frac{AB}{\sin 120}$ M1 for (*AB* =) $\frac{6}{\sin 25} \times \sin 120$ or (*BC* =) $\frac{6}{\sin 25} \times \sin 35$ M1(dep M2) for method to find area, eg 0.5×6×*AB*sin35 or 0.5×6×*BC* sin 120 A1 for answer in the range 21.1 – 21.2 OR (with perpendicular from *A* meeting *BC* extended at a point *X*) M1 for method to find *AX* M1 for method to find *BX* M1(dep M2) for 0.5 × "*AX*" × "*BC*" A1 for answer in the range 21.1 – 21.2
Q21		Diagram enlarged.	M1 for correct vector for $\overrightarrow{OY}$, eg $(\overrightarrow{OY}) = \overrightarrow{OA} + \overrightarrow{AY}$, may include terms in **a** and **b,** eg. 6**a** + $\overrightarrow{AY}$, 4**b** + 4**a** – **b** + $\overrightarrow{PY}$ M1 for $(\overrightarrow{AY} =) \frac{2}{3}\vec{A}P$ or $(\overrightarrow{PY} =) \frac{1}{3}\overrightarrow{PA}$ M1 for correct expression for $\overrightarrow{OY}$ in terms of **a** and **b,** eg $\frac{14}{3}$**a** + 2**b** C1 for $\frac{2}{3}$(7**a** + 3**b)** and "$\overrightarrow{OY}$ is parallel to the vector 7**a** +3**b" oe**

PMT

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London WC2R 0RL

Question 19

The correct answer to this question was seen only very occasionally. Some students were able to gain credit for $9^{-\frac{1}{2}} = \frac{1}{3}$ or equivalent. Only a very few were able to write $27^{\frac{1}{4}}$ in any useful format.

Question 11

In part (a), many students were able to gain at least one of the two marks. Most commonly one mark was awarded for working out $\sqrt[4]{81}$ as 3 with fewer students able to work out $\sqrt[4]{10^8}$ as 10^2. 3×10^8 and 3×10^4 were common incorrect answers. Those students who started by writing 81×10^8 as 8100000000 rarely went on to get the correct answer but they often gained one mark when their attempt at finding the 4th root resulted in an answer of 3 followed by an incorrect number of zeros.

In part (b), many students were able to gain at least one of the two marks. Most commonly one mark was awarded for evaluating $64^{\frac{1}{2}}$ as 8 with fewer students able to interpret a negative index as a reciprocal. Negative values given as answers were very common. A number of students merely found $\frac{1}{2}$ of 64 instead of using indices properly.

Part (c) was poorly answered. Many of the students who gained the first mark for writing 9^{n-1} as $(3^2)^{n-1}$ or as 3^{2n-2} were not able to complete the method. Attempts at subtracting $2n - 2$ from n often resulted in sign errors. Some of the students with some idea wrote 9^{n-1} as 3^{2n-1}. Many students had no idea how to start.

Question 17

Many students were able to set up an equation with a constant term and they often wrote down both $y = k\sqrt{t}$ and $t = k/x^3$. Most then went on to gain the second mark for substituting values in at least one equation. The values found for the constants were usually correct although $8 = k/8$ did sometimes lead to $k = 1$. Having found the values of the constants many students did not use them to write down the two equations $y = 5\sqrt{t}$ and $t = \frac{64}{x^3}$. Centres should encourage students to clearly show the equations as doing so would have gained the third mark and it might also have helped them to find a formula for y in terms of x. Many simply stopped after finding the values of the constants and went no further. Those who did continue and gave $y = 5\sqrt{\frac{64}{x^3}}$ as the final answer were not awarded the accuracy mark because the question required the answer to be given in its simplest form. Simplifying $\sqrt{\frac{64}{x^3}}$ to $\frac{8}{x^2}$ was quite a common mistake. Students should ensure that they read the question carefully as a significant proportion failed to recognise "root", "inversely" and "cube". Notation when attempting to find the constants was often poor and some students omitted the constant from their initial equation.

Question 9

Just over a quarter of all students completed part (a) of this question successfully to gain all 3 marks. A commonly seen final answer was 10^{120}. This could only score 1 mark. However, about two thirds of all students got as far as simplifying the expression to $\sqrt{10^{120}}$. Some of these students left this as their answer while other students went on to evaluate it incorrectly.

Answers to part (b) were often marred by a lack of clarity. Answers such as "Liam should have multiplied the numbers inside and outside the brackets" were commonplace but could not usually be rewarded with the mark available. Another common incorrect response was for a student to state that the 12 should be squared. More successful responses usually referred to the answer Liam should get, that is 12^{100} or that Liam should have multiplied 50 and 2, not worked out 50^2.

Question 3

Many students made a good start by working out $2^{-5} \times 2^{10}$ as 2^3. Others gained the first mark for squaring 2^{-5} or 2^8. Not all these students went on to complete the simplification correctly. A common mistake was $(2^3)^2 = 4^6$. A significant number of students could not show a correct first step and gained no marks. In some responses 2^{-5} was identified as –32.

Question 5

A good proportion of students scored full marks in part (a) for correctly placing the ten numbers in the Venn diagram. The most common error was not recognising that 1 is a square number and writing 1 in the region $A \cap B'$ rather than in the region $A \cap B$. Students who

wrote the correct numbers in two or three of the four regions gained 2 marks. It should be emphasised to students that each number in the universal set should appear just once in a Venn diagram and it might be advisable for students to cross off numbers as they are being added to the Venn diagram.

In part (b) many students scored one mark for the correct denominator of 10 or for a denominator that followed through from their Venn diagram. A correct numerator was seen less frequently. It was evident that many students were unable to identify the region B' and the most common mistake was to give the probability that the number is in set B.

Question 14

Part (a) was answered quite well with many students able to write $\frac{1}{16}$ as 4^{-2}. Common incorrect answers included $4^{-4}, 4^{-\frac{1}{2}}$ and 4^{2}.

In part (b), students who interpreted $8^{\frac{5}{3}}$ as $\left(\sqrt[3]{8}\right)^{5}$ or $9^{\frac{3}{2}}$ as $\left(\sqrt{9}\right)^{3}$ gained the first mark and usually went on to gain at least two of the three marks for evaluating 2^{5} as 32 or 3^{3} as 27. Most went on to evaluate both correctly and gave a final answer of 5. When only one term was evaluated correctly it was more often $9^{\frac{3}{2}}$ as 27. Mistakes were more common in the evaluation of $8^{\frac{5}{3}}$ and $2^5 = 64$ was frequently seen. Some students found $\sqrt[3]{8} = 2$ or $\sqrt{9} = 3$ (or both) but then incorrectly multiplied by 5 and 3 respectively instead of raising to a power. Those students who interpreted $8^{\frac{5}{3}}$ as $\sqrt[3]{8^5}$ or $9^{\frac{3}{2}}$ as $\sqrt{9^3}$ gained the first mark but were rarely able to evaluate either $\sqrt[3]{8^5}$ or $\sqrt{9^3}$ correctly.

Question 13

A minority of students scored full marks for their responses to this question and there were more students than average that did not attempt the question. Weaker responses include those where a student had substituted a value or values for n, but of course, this could not serve as a proof and so could not be given any marks. Students who did approach the question more generally and attempted to write down an expression for the next triangular number term often wrote down incorrect expressions such as $\frac{n(n+2)}{2}$, $\frac{n+1(n+2)}{2}$ or $\frac{n(n+1)}{2} + 1$. Students writing down the expression $\frac{n+1(n+2)}{2}$ sometimes recovered their error in omitting brackets and went on to give a correct proof. Not all students who obtained the expression $n^2 + 2n + 1$ could factorise it to prove it was a square number.

Question 16

Where lower attaining students attempted this question their answers were often restricted to listing the possible numbers for each digit of the passcode without writing down the number of options for each digit. These students could not be awarded any credit. Having said this, the question discriminated well between higher attaining students with each of the possible marks often being awarded. Only a small proportion of students scored full marks for a correct probability. A significant number of students used fractions with denominator 9 in their calculations. They did not appear to appreciate the importance of "Zia tells her friend Amber that..." The most common error leading to the loss of the final mark was for students to include 1 in their list of prime numbers which led to an answer of $\frac{1}{150}$

Question 13

Much like question 4, in part (a) learners often struggled to articulate themselves clearly enough to gain this mark. Learners had two real avenues for a successful response. The first, and most common, was to work with $y = kx$ and show the value of k was not constant. To do this they needed to complete at least 2 calculations, and this was often not done, meaning the mark was not awarded. A second approach was to realise that the graph of $y = kx$ should pass through the origin, and that is not the case here. Fewer learners seemed to use this approach.
Part (b)(i) was a much more familiar question, and learners experienced more success. Many gaining at least one for setting up a correct equation, and then a good number using this to find the value of k, many of these then went on to gain the third mark too. The most common error seen was for learners to either just work with t, and not $\sqrt{t}$, or when rearranging they worked out $140 \times \sqrt{64}$ instead of $140 \div \sqrt{64}$.
In part (b)(ii) the most common wrong graph was a direct proportion graph for a linear relationship, but various other incorrect sketches of familiar shape were seen.

Question 11

Many responses set up an equation with a constant term and they often wrote down both $T = \frac{k}{w}$ and $w = k\sqrt[3]{d}$. Terminology was sometimes an issue with responses using d^3 or $\sqrt{d}$ instead of $\sqrt[3]{d}$ or incorrectly interpreting inverse proportion as direct proportion and writing $T = kw$. Most then went on to gain the second mark for substituting the given values in at least one equation and the values found for the constants were usually correct. Having found the values of the constants there were two routes available. Some responses combined $T = \frac{120}{w}$ and $w = \frac{1}{2}\sqrt[3]{d}$ to find a formula for T in terms of d and this gained the third mark. Substituting $T = 48$ in the formula earned the fourth mark and gave them an equation to solve to find the value of d. Having found the values of the constants some responses earned the third mark for a correct process to find the value of w when $T = 48$ and the fourth mark was then awarded for substituting the found value of w in the equation $w = \frac{1}{2}\sqrt[3]{d}$. Some responses got as far as $48 = \frac{120}{w}$ but did not rearrange the equation to find or isolate w. Whichever of the two approaches was used, errors were often made when solving the final equation. It was not uncommon for 2.5 to be multiplied by 0.5 instead of being divided by 0.5 or for responses to cube root the answer rather than cube it to find the value of w.

Question 11

In part (a), many responses demonstrated a secure understanding of index laws, with common approaches including applying the rule for multiplying powers with the same base (e.g. $3^n \times 3^{12} = 3^{14}$ or $3^n \times 3^{20} = 3^{22}$) or forming a linear equation such as $n + 20 - 8 = 14$. Clear and methodical working was often seen in more successful responses. Some responses included errors such as multiplying the bases while adding the powers (e.g. simplifying $3^{14} \times 3^8$ to 9^{22}) or incorrectly multiplying the indices (e.g. $14 \times 8 \div 20$), which led to inaccurate answers such as 5.6.

In part (b), most responses showed understanding of fractional indices and recognised the need to apply both cube root and square operations. Stronger responses often separated the numerical and algebraic components, for example: $(\sqrt{64})^2 = 16$ and $(m^{12})^{\frac{2}{3}} = m^8$ leading to the final expression $16m^8$. Less successful responses tended to complete only one of the two operations or applied the correct process to only one part of the expression (either 64 or m^{12}), leaving the other unchanged. Partial simplifications such as 16 or m^8 were common and credited where appropriate.

Centres may wish to encourage step-by-step presentation of working, particularly when applying fractional powers. A secure understanding of how powers apply to both numbers and algebraic terms can help improve accuracy and clarity in future responses.