Chhetri Academy

chhetriacademy.com

GCSE MATHS

Number

Edexcel 1MA1 | Subtopic collection Topic 06

Surds

Higher tier

Answers and examiner comments

49	2003-2025	Higher
question groups	available papers	tier

Using this booklet

1. Answer numbers match the companion question booklet.
2. Follow the original question number on marking tables.
3. Examiner comments are included where available.

Name _____ **Date** _____

Some entries use full source marking material in an appendix. Any unavailable mark schemes are identified beside the relevant answer.

Original question 17

No matching mark scheme was available in the supplied archive for this question.

Original question 18

The full source mark scheme begins on page 80. Use the original paper code, tier and question number shown above.

Original question 21

No matching mark scheme was available in the supplied archive for this question.

Original question 19

The full source mark scheme begins on page 131. Use the original paper code, tier and question number shown above.

Original question 14

The full source mark scheme begins on page 183. Use the original paper code, tier and question number shown above.

Question		Working	Answer	Mark	Notes
21		$6^2-(2\sqrt{3})^2=36-12=24$ Area $=\frac{1}{2}\times 2\sqrt{3}\times\sqrt{24}=\sqrt{72}$ $=\sqrt{36\times 2}=6\sqrt{2}$	proof	5	M1 $6^2-(2\sqrt{3})^2$ or $\sqrt{48}$ seen or $(2\sqrt{3})^2+x^2=6^2$ oe A1 $\sqrt{24}$ oe M1(dep on M1) $\frac{1}{2}\times 2\sqrt{3}\times$ '$\sqrt{24}$' A1 $\sqrt{72}$ oe A1 $6\sqrt{2}$ or $(k)=6$

13		$\frac{3}{\sqrt{7}} \times \frac{\sqrt{7}}{\sqrt{7}}$	$\frac{3\sqrt{7}}{7}$	2	M1 for $\times \frac{\sqrt{7}}{\sqrt{7}}$ A1 cao

Question		Working	Answer	Mark	Notes
16	(a)	$\frac{15}{\sqrt{5}} \times \frac{\sqrt{5}}{\sqrt{5}}$	$3\sqrt{5}$	2	M1 for $\frac{15}{\sqrt{5}} \times \frac{\sqrt{5}}{\sqrt{5}}$ A1 for $\frac{15\sqrt{5}}{5}$ or better
	(b)	$(1 + \sqrt{3})(1 + \sqrt{3})$ $= 1 + \sqrt{3} + \sqrt{3} + 3$ $= 4 + 2\sqrt{3}$	4, 2	2	M1 for $1\times1 + 1\times\sqrt{3} + 1\times\sqrt{3} + \sqrt{3}\times\sqrt{3}$ oe A1 cao

Original question 14

The full source mark scheme begins on page 196. Use the original paper code, tier and question number shown above.

Question		Working	Answer	Mark	Notes
26	(a)		$\frac{5\sqrt{2}}{2}$	2	M1 for $\frac{5}{\sqrt{2}}\times\frac{\sqrt{2}}{\sqrt{2}}$ oe A1 for $\frac{5\sqrt{2}}{2}$ oe
	(b)		$8\sqrt{3}$	2	M1 for $2\times2+2\sqrt{3}+2\sqrt{3}+\sqrt{3}\times\sqrt{3}$ or $\left(4+4\sqrt{3}+3\right)-\left(4-4\sqrt{3}+3\right)$ or $2\times2-2\sqrt{3}-2\sqrt{3}+\sqrt{3}\times\sqrt{3}$ at least three terms in either correct; could be in a grid. A1 cao **OR** Difference of two squares M1 for $\left(\left(2+\sqrt{3}\right)-\left(2-\sqrt{3}\right)\right)\left(\left(2+\sqrt{3}\right)+\left(2-\sqrt{3}\right)\right)$ A1 cao

17		$6 \times 6 + 6 \times \sqrt{5} - 6 \times \sqrt{5} - \sqrt{5} \times \sqrt{5}$ $\frac{31}{\sqrt{31}} \times \frac{\sqrt{31}}{\sqrt{31}}$	$\sqrt{31}$	3	M1 for $6 \times 6 + 6 \times \sqrt{5} - 6 \times \sqrt{5} - \sqrt{5} \times \sqrt{5}$ or $6^2 - \left(\sqrt{5}\right)^2$ (for 3 out of not more than 4 terms including signs or 4 terms correct ignoring signs) M1 $\frac{"31"}{\sqrt{31}} \times \frac{\sqrt{31}}{\sqrt{31}}$ or for $\frac{\text{[expression in surd form]}}{\sqrt{31}} \times \frac{\sqrt{31}}{\sqrt{31}}$ A1 cao

25			$1 + \sqrt{5}$	5	M1 for $\frac{1}{2} \times x \times x \times \sin 30°$ oe M1 for $\frac{1}{2}(x-2)(x+1)$ oe **or** $\frac{1}{2} \times (x-2) \times (x+1) \times \sin 90$ M1 (dep on at least one previous M1) for formation of equation from equating areas with x as the only variable A1 for $x^2 - 2x - 4 = 0$ oe in the form $ax^2 + bx + c = 0$ or $ax^2 + bx = c$ A1 cao

17	(a)		$\frac{1}{10}$	1	B1 for 0.1 or $\frac{1}{10}$ oe
	(b)		9	2	M1 for $(\sqrt[3]{27})^2$ or $\sqrt[3]{27^2}$ oe or $\sqrt[3]{27} = 3$ A1 cao
	(c)	$\sqrt{75} = \sqrt{25}\sqrt{3}$	$5\sqrt{3}$	2	M1 for $\sqrt{25 \times 3}$ or $\sqrt{25}\sqrt{3}$ oe A1 cao

13			7.5	3	B1 for length given as $\frac{\sqrt{120}}{4}$ oe M1 for squaring $\frac{\sqrt{120}}{4}$ or $\frac{120}{4\times4}$ oe A1 for $\frac{120}{16}$ oe or $7\frac{1}{2}$ or 7.5 oe SC B1 for $\sqrt{30} \times \sqrt{30}$

PAPER: 5MB2H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
*14			113	5	B1 for stating angle *TAO* = 90 M1 for stating angle *OBA* or angle *OAB* = 90 – 58 (=32) M1 for stating angle *ABT* = 180 − 58 – 41 (=81) or angle *AOB* = 180 – 64 (=116) A1 for 113 clearly identified as the answer C1 (dep on M1) for correct statements for method used: angle between tangent and radius = 90° AND at least one of base angles of an isosceles triangle are equal sum of angles in a triangle is 180 sum of angles in a quadrilateral is 360 NB angles may be seen in diagram

25	(a)		$4\sqrt{3}$	2	M1 for $\frac{12}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}}$ A1 for $\frac{12\sqrt{3}}{3}$ oe with a rational denominator
	(b)		18	2	M1 for $\sqrt{2}\times\sqrt{2} + \sqrt{2}\times\sqrt{8} + \sqrt{8}\times\sqrt{2} + \sqrt{8}\times\sqrt{8}$ oe A1 cao OR M1 for $(\sqrt{2} + 2\sqrt{2})^2$ A1cao

PAPER: 1MA0_1H

16			$30 - 10\sqrt{5}$	2	M1 for 4 terms correct with or without signs or 3 out of exactly 4 terms correct (the terms may be in an expression or table) or $25 - 10\sqrt{5} + 5$ A1 cao

21		$3-\sqrt{2}+3\sqrt{2}-\sqrt{2}\sqrt{2}$	$1+2\sqrt{2}$	2	M1 for 4 terms correct ignoring signs or 3 out of no more than 4 terms correct A1 cao

PAPER: 1MA0_1H

10	(a)		1	1	B1
	(b)		$\frac{1}{16}$	1	B1
	(c)		$2\sqrt{7}$	2	M1 for multiplying numerator and denominator by $\sqrt{7}$ A1 for $2\sqrt{7}$

PAPER: 5MB2H_01

22	(a)		$\frac{1}{8}$	1	B1 for $\frac{1}{8}$ oe
	(b)		1.5	1	B1 for 1.5 oe
	(c)		3	2	M1 for √12×√12 – √12×√3 – √3×√12 + √3×√3 or √144 – √36 – √36 + √9 oe. with no more than one sign error A1 cao OR M1 for writing (√12 – √3) as (2√3 – √3) (= √3) A1cao

PAPER: 1MA0_1H

15	(a)		$(y-7)(y+2)$	2	B2 cao (B1 for $(y \pm 7)(y \pm 2)$)
	(b)		$\sqrt{5}+29$	2	M1 expand brackets, with at least 3 correct terms including signs or 4 correct terms ignoring signs eg $2\sqrt{5}\times 3\sqrt{5}-2\sqrt{5}+3\sqrt{5}-1\times 1$ A1 for $\sqrt{5}+29$ or $29+\sqrt{5}$
	(c)		$\sqrt{3}$	2	M1 for $\frac{6}{\sqrt{12}}\times\frac{\sqrt{12}}{\sqrt{12}}$ oe or $\sqrt{12}=2\sqrt{3}$ A1 cao

5MB2H/01 June 2015

1MA0 1H November 2015

Question		Working	Answer	Mark	Notes
18			$2\sqrt{5}$	2	M1 for multiplication of denominator and numerator by $\sqrt{5}$ A1 cao

Original question 16

The full source mark scheme begins on page 215. Use the original paper code, tier and question number shown above.

Question	Working	Answer	Mark	Notes
25		$\frac{1}{4}-\frac{\sqrt{6}}{12}$	3	M1 for $\frac{1}{2}\times\frac{\sqrt{2}}{2}\times\frac{\sqrt{2}}{2}$ or $\frac{1}{2}\times\frac{\sqrt{2}}{2}\times\frac{\sqrt{3}}{3}$ M1 for $\frac{1}{2}\times\frac{\sqrt{2}}{2}\times\frac{\sqrt{2}}{2}-\frac{1}{2}\times\frac{\sqrt{2}}{2}\times\frac{\sqrt{3}}{3}$ A1 for $\frac{1}{4}-\frac{\sqrt{6}}{12}$ oe OR M1 for (BC =) $\frac{\sqrt{2}}{2}-\frac{\sqrt{3}}{3}$ M1 for $\frac{1}{2}\times\left\{\frac{\sqrt{2}}{2}-\frac{\sqrt{3}}{3}\right\}\times\frac{\sqrt{2}}{2}$ A1 for $\frac{1}{4}-\frac{\sqrt{6}}{12}$ oe

Modifications to the mark scheme for Modified Large Print (MLP) papers.

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 1MA0_1H			
Question		**Modification**	**Notes**
1		Model provided for all candidates. Diagram enlarged and also provided for MLP.	Standard mark scheme
2	(a)	Reflection drawn on the diagram. Shading changed to dotty shading. Wording changed 'It shows shape P and shape Q given on a grid. Describe fully the single transformation that maps shape P onto shape Q.' Grid enlarged. y axis is cut at -4. 3 answer lines given.	Marked scheme changed to: B1 Reflection B1 in the line $x = -1$
	(b)	Rotation drawn on the diagram. Wording changed 'It shows shape P and shape S given on a grid.' Describe fully the single transformation that maps shape P onto shape S.' 3 answer lines given.	Mark scheme changed to: B1 Rotation or rotate B1 90° anticlockwise B1 about (0, 1)
7		Diagram enlarged. Braille only: Plan of floor labelled ABCDEFGH as 1F Q24.	Standard mark scheme
10		Diagram enlarged. Crosses changed to filled in circles. Arrows heads changed to open headed arrows.	Standard mark scheme
12		Diagram enlarged. Shading changed to dotty shading. Braille only: Diagram turned into horizontal position (circle inside square).	Standard mark scheme

<table>
<tr><th colspan="4">PAPER: 1MA0_1H</th></tr>
<tr><th colspan="2">Question</th><th>Modification</th><th>Notes</th></tr>
<tr><td>14</td><td>(a)

(b)</td><td>Diagram lines made longer and thicker
Braille only: Angle A 75° exactly for bisection.
Diagram kept the same size. Point P has been moved to the left.</td><td>Standard mark scheme</td></tr>
<tr><td>15</td><td>(a)

(b)</td><td>Grid enlarged. Small squares removed. Lines of the box plot have been moved.
‘16 year old workers’ lines are at ‘75,125,175,200 and 300’.
‘18 year old workers’ lines are at ‘125, 200, 225, 275 and 375’.

‘£130’ changed to ‘£125’</td><td>Standard mark scheme with these following values

<table><tr><td></td><td>Age 16</td><td></td><td>Age 18</td></tr><tr><td>HV</td><td>300</td><td><</td><td>375</td></tr><tr><td>LV</td><td>75</td><td><</td><td>125</td></tr><tr><td>Median</td><td>175</td><td><</td><td>225</td></tr><tr><td>Range</td><td>225</td><td><</td><td>250</td></tr><tr><td>IQR</td><td>75</td><td>=</td><td>75</td></tr></table></td></tr>
<tr><td>14</td><td></td><td>Grid enlarged. Small squares removed. [Leeway needed for answering the question.]</td><td>Standard mark scheme</td></tr>
<tr><td>17</td><td>(a)
(b)</td><td>Wording added above the table ‘There are four spaces to fill.’
Grid enlarged. Right axis labelled.</td><td>Standard mark scheme</td></tr>
<tr><td>18</td><td></td><td>Diagram enlarged.</td><td>Standard mark scheme</td></tr>
</table>

PAPER: 1MA0_1H			
Question		**Modification**	**Notes**
19		Grid enlarged. [Leeway need for answering the question.]	Standard mark scheme
20		Diagram enlarged. Cross changed to filled in circle.	Standard mark scheme
21		Diagram enlarged. Wording added 'There are six spaces to fill.'	Standard mark scheme
23		Diagram enlarged	Standard mark scheme

Question	Working	Answer	Mark	Notes
26		8	4	M1 for $(2\sqrt{10})^2 - 2^2$ (= 36) A1 for (CD =) 6 M1 (dep on M1) for '6' × 4 – $\frac{1}{2}$ × '6' × 2 – $\frac{1}{2}$ × 2 × 2 – $\frac{1}{2}$ × ('6' – 2) × 4 C1 for area of 8 from fully correct working

Question	Working	Answer	Mark	Notes	Type
16	$\frac{(\sqrt{5}+\sqrt{5}+6)}{2} \times (\sqrt{5}-2)$ $(\sqrt{5}+3)(\sqrt{5}-2)$ $5+3\sqrt{5}-2\sqrt{5}-6$ $\sqrt{5}(\sqrt{5}-2)+\frac{6(\sqrt{5}-2)}{2}$ $5-2\sqrt{5}+3\sqrt{5}-6$	$\sqrt{5}-1$	3	M1 for $\frac{(\sqrt{5}+\sqrt{5}+6)}{2} \times (\sqrt{5}-2)$ M1 for expansion $5+3\sqrt{5}-2\sqrt{5}-6$ with 3 terms out of 4 correct including signs or all 4 terms correct ignoring signs A1 cao OR M1 for $\sqrt{5}(\sqrt{5}-2)+\frac{6(\sqrt{5}-2)}{2}$ M1 for expansion $5-2\sqrt{5}+3\sqrt{5}-6$ with 3 terms out of 4 correct including signs or all 4 terms correct ignoring signs A1 cao	E

PAPER: 5MB3H_01

Question	Working	Answer	Mark	Notes
19 (a)		5 and −7	3	M1 for $(x \pm 5)(x \pm 7)$ M1 for $(x - 5)(x + 7)$ A1 cao OR M1 for correct substitution into formula M1 for reduction to $\frac{-2 \pm \sqrt{144}}{2}$ A1 cao
(b)		$\pm\sqrt{3}$	4	M1 for multiplying through by a common denominator eg by $(x + 1)(2x + 3)$ M1 (dep) for reduction eg to $4x + 6 + x^2 + x$ or $2x^2 + 3x + 2x + 3$ M1 for $x^2 - 3$ (= 0) A1 cao for both answers; accept decimals to at least 2 dp

21		$16\sqrt{2}$	4	M1 for method to expand $(\sqrt{8}+2)^2$ with at least 3 correct terms out of 4 terms M1 for method to expand $(\sqrt{8}-2)^2$ with at least 3 correct terms out of 4 terms M1 (dep on M2) for a method to subtract the two expressions and use of $\sqrt{8}=2\sqrt{2}$ A1 cao OR M1 for factorising $a^2-b^2=(a+b)(a-b)$ M1 for substituting for a and b with simplification (at least 1 of the two terms correct) M1 (dep on M2) for multiplying the 2 terms together and use of $\sqrt{8}=2\sqrt{2}$ A1 cao

Paper: 1MA0/1H				
Question	**Working**	**Answer**	**Mark**	**Notes**
*22		155°	5	M1 for a method to find angle *AOD* e.g 90 − 40 (= 50) M1 for a complete method to find angle *BCD* eg 360 − '50' (= 310) and '310' ÷ 2 (= 155) A1 for 155 C2 for complete reasons for their method Angle between tangent and radius = 90 Angle at the centre is twice the angle at the circumference oe Angle sum of a triangle = 180 Sum of angles round a point = 360 (C1 for at least two reasons, one of which must be a circle theorem) OR M1 for a method to find angle *AOD* eg 90 − 40 (= 50) M1 for a complete method to find angle *BCD* eg 50 ÷ 2 (= 25) and 180 − '25' (= 155) A1 for 155 C2 for complete reasons for their method Angle between tangent and radius = 90 Angle at the centre is twice the angle at the circumference oe Opposite angles of a cyclic quadrilateral add up to 180 Angle sum of a triangle = 180 (C1 for at least two reasons, one of which must be a circle theorem)

Paper: 1MA0/1H

Question	Working	Answer	Mark	Notes
21	$\frac{6-\sqrt{8}}{\sqrt{2}-1} \times \frac{\sqrt{2}+1}{\sqrt{2}+1}$ $=\frac{6\sqrt{2}+6-\sqrt{8}\sqrt{2}-\sqrt{8}}{2-1}$ $=6\sqrt{2}+6-4-2\sqrt{2}$	$2 + 4\sqrt{2}$	M1 M1 A1	for correct first step eg multiplies numerator and denominator by $\sqrt{2}$ +1 condone missing brackets (dep) for expansion of numerator with 4 terms correct with or without signs **or** 3 out of exactly 4 terms correct for $2 + 4\sqrt{2}$ oe **or** for stating $a = 2$ and $b = 4$

13	5	M1 M1 A1	for $\sqrt{40}$ **or** $\sqrt{90}$ **OR** $2\sqrt{2}$ **or** $3\sqrt{2}$ for $2\sqrt{10}$ **or** $3\sqrt{10}$ **or** $\sqrt{4}\times\sqrt{10}$ **or** $\sqrt{9}\times\sqrt{10}$ **or** $\sqrt{4\times10}$ **or** $\sqrt{9\times10}$ **OR** $2\sqrt{2}+3\sqrt{2}$ cao	Answer of $5\sqrt{10}$ from correct working gets M2 A0

Paper: 1MA1/1H

20	(a)	explanation	C1	for a correct explanation, eg $\sqrt{3} \times -\sqrt{3} = -3$, not 3	
	(b)	explanation	C1	for correct explanation, eg $\sqrt{12} = 2\sqrt{3}$, not $3\sqrt{2}$	

Paper: 1MA1/2H

Question	Answer	Mark	Mark scheme	Additional guidance
20	fully correct working leading to $16\left(1+\sqrt{2}\right)$	C1	for expanding the numerator, eg $18+2\sqrt{2}\sqrt{18}+2$ or $\sqrt{324}+\sqrt{36}+\sqrt{36}+\sqrt{4}$ $(=32)$ **or** for simplifying $\sqrt{18}$, eg. $\sqrt{18}=3\sqrt{2}$ or $\sqrt{18}+\sqrt{2}=4\sqrt{2}$	Expanded terms need not be simplified
		C1	(indep) for method to rationalise the denominator, eg. $\frac{\text{"numerator"}}{\sqrt{8}-2}\times\frac{\sqrt{8}+2}{\sqrt{8}+2}$	
		C1	for fully correct working leading to $16\left(1+\sqrt{2}\right)$	Accept $a = 16$, $b = 1$

Question	Answer	Mark	Mark scheme	Additional guidance
14	$\frac{1}{2}$	M1	for $\frac{1}{\sqrt{3}} \times \frac{\sqrt{3}}{2}$ **or** $\frac{\sqrt{3}}{3} \times \frac{\sqrt{3}}{2}$ **or** $(\frac{1}{2} \div \frac{\sqrt{3}}{2}) \times \frac{\sqrt{3}}{2}$ **OR** $\tan 30 = \frac{1}{\sqrt{3}}$ oe **or** $\sin 60 = \frac{\sqrt{3}}{2}$	
		A1	for $\frac{1}{2}$ or 0.5	

Question	Answer	Mark	Mark scheme	Additional guidance
18 (a)	$3\sqrt{3}$	M1	for working unambiguously with $\sqrt{12}$, eg $\sqrt{4\times3}$ **or** $\sqrt{4}\times\sqrt{3}$ **or** $2\sqrt{3}$	
		A1	cao	
(b)	$\frac{\sqrt{3}}{81}$	M1	for simplifying the power eg $(\sqrt{3})^7 = 27\sqrt{3}$	
		M1	for method to rationalise the denominator eg multiplying by $\frac{\sqrt{3}}{\sqrt{3}}$	May be seen as the first step
		A1	for $\frac{\sqrt{3}}{81}$ **or** equivalent fraction in form $\frac{\sqrt{b}}{c}$, eg $\frac{\sqrt{2187}}{2187}$	

16	(a)	$2\sqrt{11}$	M1	for method to multiply numerator and denominator by $\sqrt{11}$ or a multiple of $\sqrt{11}$, eg $\frac{22}{\sqrt{11}} \times \frac{\sqrt{11}}{\sqrt{11}}$	
			A1	for $2\sqrt{11}$	
	(b)	$\frac{6+\sqrt{3}}{11}$	M1	for method to multiply numerator and denominator by $2\sqrt{3}+1$ or a multiple of $2\sqrt{3}+1$, eg $\frac{\sqrt{3}}{2\sqrt{3}-1} \times \frac{2\sqrt{3}+1}{2\sqrt{3}+1}$	
			M1	(dep) for $\sqrt{3} \times 2\sqrt{3} = 6$ or $2\sqrt{3} \times 2\sqrt{3} = 12$	
			A1	for $\frac{6+\sqrt{3}}{11}$ (accept $a = 6$ and $b = 11$)	

Paper: 1MA1/1H

20	$1+\frac{\sqrt{5}}{5}$	P1	for writing $\sqrt{180}$ as $6\sqrt{5}$	This process mark can be awarded whenever this is seen, which might be later in the process.
		P1	for process to rationalising the denominator eg $\frac{\sqrt{180}-2\sqrt{5}}{5\sqrt{5}-5}\times\frac{5\sqrt{5}+5}{5\sqrt{5}+5}$ **or** $\frac{4\sqrt{5}}{5\sqrt{5}-5}\times\frac{5\sqrt{5}+5}{5\sqrt{5}+5}$ oe	
		P1	(dep on previous P1) for expanding terms eg $\frac{5\sqrt{5}\sqrt{180}+5\sqrt{180}-50-10\sqrt{5}}{125-25}$ **or** $\frac{100+20\sqrt{5}}{100}$ oe	
		A1	for $1+\frac{\sqrt{5}}{5}$	Accept written as $a = 1$, $b = 5$

Paper: 1MA1/1H

Paper: 1MA1/3H

Question	Answer	Mark	Mark scheme	Additional guidance
19	Proof	P1	for start to process to find area of *ABCDEF*, eg area of equilateral triangle $= \frac{1}{2} \times x \times x \times \sin 60$ $\left(= \frac{\sqrt{3}}{4}x^2\right)$ **OR** for start to process to find area of *FGHIJK*, eg area of equilateral triangle $= \frac{1}{2} \times px \times px \times \sin 60$ $\left(= \frac{\sqrt{3}}{4}p^2x^2\right)$	Any correct process to find the area of part of the hexagon is acceptable for this mark, eg $\frac{1}{2} \times x \times x \times \sin 120$ or $\frac{1}{2} \times (x + 2x) \times \frac{\sqrt{3}}{2}x$ Allow sin 60 left in expressions for the first 3 marks.
		P1	for complete process of finding area of *ABCDEF*, eg $6 \times \frac{1}{2} \times x \times x \times \sin 60$ or $6 \times \frac{1}{2} \times x \times x \times \frac{\sqrt{3}}{2}$ $\left(= \frac{3\sqrt{3}}{2}x^2\right)$ oe **OR** for complete process of finding area of *FGHIJK*, eg $6 \times \frac{1}{2} \times px \times px \times \frac{\sqrt{3}}{2}$ $\left(= \frac{3\sqrt{3}}{2}p^2x^2\right)$ oe	
		P1	for process of finding area of *ABCDEF* eg $\frac{3\sqrt{3}}{2}x^2$ oe **AND** for process of finding area of *FGHIJK*, eg $p^2 \times \frac{3\sqrt{3}}{2}x^2$ oe	
		C1	correct algebra leading to given result, $\frac{3\sqrt{3}}{2}(p^2 - 1)x^2$	Accept $\frac{3\sqrt{3}}{2}x^2(p^2 - 1)$ as final result.

Paper: 1MA1/3H

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
19	Result shown	M1	(indep) for writing $\sqrt{12}$ as $2\sqrt{3}$	This mark can be awarded whenever this is seen, which might be later in the process.
		M1	for method to rationalise the denominator eg $\frac{8+\sqrt{12}}{5+\sqrt{3}}\times\frac{5-\sqrt{3}}{5-\sqrt{3}}$ **or** $\frac{8+2\sqrt{3}}{5+\sqrt{3}}\times\frac{5-\sqrt{3}}{5-\sqrt{3}}$ oe	
		M1	(dep on previous M1) for expanding terms, condone one error in numerator or denominator eg $\frac{40-8\sqrt{3}+5\sqrt{12}-\sqrt{12}\sqrt{3}}{25-5\sqrt{3}+5\sqrt{3}-\sqrt{3}\sqrt{3}}$ **or** $\frac{40-8\sqrt{3}+10\sqrt{3}-2\sqrt{3}\sqrt{3}}{25-5\sqrt{3}+5\sqrt{3}-\sqrt{3}\sqrt{3}}$ **or** $\frac{34+2\sqrt{3}}{22}$ oe	
		A1	for a complete chain of reasoning leading to $\frac{17+\sqrt{3}}{11}$	

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
19	$\frac{5\pm\sqrt{15}}{2}$	M1	for using a common denominator eg $\frac{x-1}{(2x-1)(x-1)}+\frac{3(2x-1)}{(2x-1)(x-1)}$ (= 1) **or** $(x-1)+3(2x-1)=(2x-1)(x-1)$	
		M1	for expanding and rearranging to get $2x^2-10x+5$ (= 0)	Note we don't need to see "= 0"; just the LHS is sufficient Accept other forms of the 3 term quadratic, eg $2x^2-10x=-5$
		M1	(dep M1) ft for a method to solve their 3 term quadratic equation eg $\frac{10\pm\sqrt{(-10)^2-4\times2\times5}}{2\times2}$ or $\frac{10\pm\sqrt{60}}{4}$ **or** $2\left[\left(x-\frac{5}{2}\right)^2-\left(\frac{5}{2}\right)^2\right]+5=0$ oe	Correct use of formula or completing the square
		A1	cao	

Paper: 1MA1/1H

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
19	$-\frac{1}{2}\pm\frac{1}{2}\sqrt{2}$	P1	for using a common denominator, eg $\frac{x+1}{x(x+1)}-\frac{x}{x(x+1)}$ (= 4) or $\frac{x+1-x}{x(x+1)}$ (= 4) **or** $x+1-x=4x(x+1)$	
		P1	for expanding and rearranging to get $4x^2+4x-1$ (= 0)	Note we don't need to see "= 0"; just the LHS is sufficient Accept other forms of the 3 term quadratic, eg $4x^2+4x=1$
		P1	(dep P1) ft for a method to solve their 3 term quadratic equation, eg $\frac{-4\pm\sqrt{4^2-4\times4\times-1}}{2\times4}$ or $4\left[\left(x+\frac{1}{2}\right)^2-\left(\frac{1}{2}\right)^2\right]-1=0$ oe	Correct use of formula or completing the square
		A1	for values of x, eg $\frac{-4\pm\sqrt{32}}{8}$ or $\pm\sqrt{\frac{1}{2}}-\frac{1}{2}$ oe	
		A1	for $-\frac{1}{2}\pm\frac{1}{2}\sqrt{2}$ oe in the form $a\pm b\sqrt{2}$ where are a and b are fractions	Accept $a=-\frac{1}{2}$, $b=\frac{1}{2}$ or $a=-\frac{1}{2}$, $b=-\frac{1}{2}$

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
16	$10\sqrt{2}$	P1	for process to find total surface area of sphere, eg $75\pi \div 3 \times 8\ (= 200\pi)$ or $75 \div 3 \times 8\ (= 200)$ **or** for setting up an equation, eg $\frac{3}{8} \times 4\pi r^2 = 75\pi$ or $\frac{3}{8} \times 4r^2 = 75$	Could work without π for the three P marks
		P1	for process to find r^2, eg $(r^2 =) \frac{"200\pi"}{4\pi}$ oe or $(r^2 =) \frac{75\pi}{4\pi} \times \frac{8}{3}$ oe or $r^2 =$ "50"	
		P1	for process to find radius, eg $(r =) \sqrt{\frac{"200\pi"}{4\pi}}$ oe or $\sqrt{"50"}$ or $5\sqrt{2}$ **or** for $(2r =) = \sqrt{"200"}$	
		A1	cao	

Question	Answer	Mark	Mark scheme		Additional guidance
23	$\frac{10\sqrt{3}+27}{39}$	M1	for method to rationalise one of the fractions, eg $\frac{3\sqrt{3}}{4-\sqrt{3}}\times\frac{4+\sqrt{3}}{4+\sqrt{3}}$ $(=\frac{12\sqrt{3}+9}{16-3})$ **or** $\frac{2}{\sqrt{3}}\times\frac{\sqrt{3}}{\sqrt{3}}$ **(=** $\frac{2\sqrt{3}}{3}$**)** oe	for method to write fractions with a common denominator, eg $\frac{3\sqrt{3}\times\sqrt{3}}{\sqrt{3}(4-\sqrt{3})}-\frac{2\times(4-\sqrt{3})}{\sqrt{3}(4-\sqrt{3})}$ oe	
		M1	for method to rationalise both of the fractions eg $\frac{3\sqrt{3}}{4-\sqrt{3}}\times\frac{4+\sqrt{3}}{4+\sqrt{3}}$ $(=\frac{12\sqrt{3}+9}{16-3})$ **and** $\frac{2}{\sqrt{3}}\times\frac{\sqrt{3}}{\sqrt{3}}$ **(=** $\frac{2\sqrt{3}}{3}$**)** oe	for writing as a **correct** single fraction without brackets, eg $\frac{9-8+2\sqrt{3}}{4\sqrt{3}-3}$ oe	
		M1	for method to write **correct** fractions with a common denominator eg $\frac{3(12\sqrt{3}+9)}{13\times3}-\frac{13\times2\sqrt{3}}{13\times3}$ or $\frac{36\sqrt{3}+27}{39}-\frac{26\sqrt{3}}{39}$ oe	for method to rationalise a fraction of the form $\frac{a+b\sqrt{3}}{c\sqrt{3}-d}$ where a, b, c and d are integers, eg $\frac{1+2\sqrt{3}}{4\sqrt{3}-3}\times\frac{4\sqrt{3}+3}{4\sqrt{3}+3}$ or $\frac{4\sqrt{3}+3+24+6\sqrt{3}}{48+12\sqrt{3}-12\sqrt{3}-9}$ oe	
		A1	for $\frac{10\sqrt{3}+27}{39}$ oe		

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
24	$4\left(2+\sqrt{2}\right)a^2$	P1	for process to find area required and identifying 135° or 45° or 90° angle(s), eg splitting shape into square and 4 triangles and an angle relevant to the triangle clearly identified.	90° must be in a triangle to gain credit. May be seen on diagram.
		P1	for process to find the area of a relevant triangle using 45°, eg $\frac{1}{2}\times a\times\left(2\times\frac{a}{\sqrt{2}}\right)\times\frac{1}{\sqrt{2}}\left(=\frac{a^2}{2}\right)$ or using 90°, eg $\frac{1}{2}\times a\times a\left(=\frac{a^2}{2}\right)$ or process to find the area of a square made from 2 small triangles, eg $a\times a\left(=a^2\right)$	Accept $0.49...a^2$ May be seen as the area of 2 squares (from 4 small triangles)
		P1	for process to find the length of the square, eg $a+a+\sqrt{a^2+a^2}\left(=2a+a\sqrt{2}\right)$	Accept $3.41a$
		P1	for process to find the total area, eg $\left("\,2a+a\sqrt{2}\,"\right)^2+4\times"\,\frac{a^2}{2}\,"$	Accept $(11.655+4\times0.49)\,a^2$
		A1	(dep on P3) for $4\left(2+\sqrt{2}\right)a^2$	Answer only award no marks. If working in decimals accept $\frac{13.656}{2+\sqrt{2}}$ leading to 4 Accept $p=4$ if supported by correct working

Modifications to the mark scheme for Modified Large Print (MLP) papers: 1MA1 2H

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme. Notes apply to both MLP papers and Braille papers unless otherwise stated.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

		Modification	Mark scheme notes
4	(b)	Wording added 'Look at the diagram for Question 4(b) in the Diagram Booklet. It shows a number line'. Wording removed 'below'. Diagram enlarged. Open headed arrow. Axis label moved to the right of the horizontal axis.	Standard mark scheme
4	(c)	Letter 'g' changed to 't'.	Standard mark scheme but note letter change.
5		Wording added 'Look at the diagram for Question 5 in the Diagram Booklet. It shows a triangle ABC and a rectangle PQRS.' Wording removed 'Here is a triangle and a rectangle.' Triangle labelled ABC. Rectangle labelled PQRS. Diagram enlarged. Diagrams stacked vertically. Right angle made more obvious. Shape labels moved to the left and to above the rectangle. Wording added 'In triangle ABC: AB is marked $6x$; BC is marked 8; Angle ABC is a right angle. In rectangle PQRS: PQ is marked 5; PS is marked $4x - 1$'	Standard mark scheme
7		Letter 'd' changed to 'n'.	Standard mark scheme but note letter change.
9		Wording added 'Look at the diagram for Question 9 in the Diagram Booklet. It shows a grid.' Wording added 'below' Table enlarged and left aligned. Letter 'a' changed to 'A'. Values changed in the table: 48 changed to 50; 64 changed to 65 Wording removed 'opposite' and replaced with 'in the Diagram Booklet.' Grid enlarged. No small squares. Black grid lines added. Open headed arrows. Right axis labelled. Axes labels moved to the top of the vertical axis and to the left of the horizontal axis	(a) Standard mark scheme using the amended numbers n in the table. (b) B1 for answer in the range 32.5 to 38.5 or ft their cf graph with a tolerance of ±2.5 (5 mm)
10		Wording added 'Look at the table for Question 10 in the Diagram Booklet.' Table enlarged and turned vertical.	Standard mark scheme
12		Wording added 'Look at the diagram for Question 12 in the Diagram Booklet. It shows the'. Wording removed 'The'; Wording removed 'lie'; Wording added 'the circumference of'. Diagram enlarged. Cross changed to a solid dot. Wording 'BC' removed and replaced with 'AD'. Labels moved to the left of the rectangle and inside the rectangle.	Standard mark scheme

		Modification	Mark scheme notes
13		Wording added 'Look at the diagram for Question 13 in the Diagram Booklet. It shows triangle ABC.'; Wording removed 'ABC is a triangle.' Diagram enlarged and rotated so AC is horizontal. Angle moved outside angle arc and angle arc made smaller. Wording added 'AB = 15 cm; AC = 18 cm; Angle ABC = 70°'	Standard mark scheme
16		Wording added 'Look at the diagram for Question 16 in the Diagram Booklet. It shows an incomplete histogram.' Diagram enlarged. Small squares removed. Bar heights changed: 0-20 is now 3 cm high, 20-30 is now 7 cm high, 30-40 is now 6 cm high. Open headed arrows. Shading changed. Right axis labelled. Axes labels moved to the top of the vertical axis and to the left of the horizontal axis. Value 28 changed to 30 and value 24 changed to 20. (a) Wording added 'in the Diagram Booklet'.	(a) M1 for using area to represent frequency eg 12 (cm^2) represents 30 (students) or showing a correct scale on fd axis A1 Bar of height 2 cm drawn for interval 40 – 60 hours (b) M1 for method to find total area under the histogram, eg $(4 \times 3) + (2 \times 7) + (2 \times 6) + (4 \times 2)$ (= 46) oe **or** for method to find total frequency, eg 30 + 35 + 30 + 20 oe A1 for 115 cao
18		Letter '*a*' changed to '*p*'; Letter '*b*' changed to '*q*'; Letter '*c*' changed to '*r*'.	Standard mark scheme but note change of letters.
19		Wording added 'Look at the diagram for Question 19 in the Diagram Booklet. It shows a right-angled triangle ABC.' Wording removed 'ABC is a right-angled triangle.' Diagram enlarged. Right angle made more obvious. Angle label moved outside angle arc and angle arc made smaller. Wording added 'Angle ABC is a right angle. Angle ACB is marked *x*.'	Standard mark scheme

		Modification	Mark scheme notes
20		Wording added 'Look at the diagram for Question 20 in the Diagram Booklet. It shows a triangle ORT.' Wording removed 'ORT is a triangle.' Diagram enlarged. Open headed arrows. Arrow heads enlarged. Line at M changed to a solid circle. Dashed line added for MT.	Standard mark scheme
21	(a)	Wording removed 'Here is the graph of $y = f(x)$'. Diagram removed. Wording added 'Look at the diagram for Question 21(a) in the Diagram Booklet. It shows the graph of $y = f(x)$' Wording 'grid below' removed and replaced with 'diagram'. Axes labels moved to the top of the vertical axis and to the right of the horizontal axis. Grid enlarged. $y = f(x)$ graph line added.	Standard mark scheme
21	(b)	Wording added 'Look at the diagram for Question 21(b) in the Diagram Booklet. It shows the graph of $y = f(x)$' Wording 'grid below' removed and replaced with 'diagram'. Axes labels moved to the top of the vertical axis and to the right of the horizontal axis. Grid enlarged. $y = f(x)$ graph line added.	Standard mark scheme
23		Wording added 'Look at the diagram for Question 23 in the Diagram Booklet. It shows'. Wording removed 'Here are'. Diagram enlarged. Arrow heads enlarged.	Standard mark scheme
24		Wording added 'Look at the diagram for Question 24 in the Diagram Booklet. It'. Wording removed 'The diagram'. Diagram enlarged. Shading changed. Letter 'a' changed to 't'.	Standard mark scheme but note change of letters.

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
20	120	P1	for using the cosine rule for any angle, eg $(5\sqrt{7})^2 = 10^2 + 5^2 - 2\times10\times5\times\cos B$	May be any angle; check diagram for working.
		P1	for a complete process to find $\cos B$ eg $(\cos B =)\dfrac{10^2 + 5^2 - (5\sqrt{7})^2}{2\times10\times5}$ oe	
		P1	for $\cos^{-1}(\frac{1}{2}) = 60$ **or** $\cos B = -0.5$	
		A1	cao	A correct answer with no supportive working gets 0 marks

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
22	$3+\sqrt{2}$	P1	for start of process to find the common ratio, eg writes $\frac{13+9\sqrt{2}}{3+2\sqrt{2}}$	
		P1	for process to rationalise the denominator, eg $\frac{13+9\sqrt{2}}{3+2\sqrt{2}}\times\frac{3-2\sqrt{2}}{3-2\sqrt{2}}$	Award P1 for process to rationalise the denominator of $\frac{3+2\sqrt{2}}{13+9\sqrt{2}}$
		P1	(dep P2) for expanding terms, condone one error in expansion of numerator or denominator, eg $\frac{39+27\sqrt{2}-18\sqrt{2}\sqrt{2}-26\sqrt{2}}{9+6\sqrt{2}-6\sqrt{2}-4\sqrt{2}\sqrt{2}}$ **or** $\frac{39+27\sqrt{2}-26\sqrt{2}-36}{9-8}$ oe	
		A1	cao	A correct answer with no supportive working gets 0 marks.

Modifications to the mark scheme for Modified Large Print (MLP) papers: 1MA1 1H

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme. Notes apply to both MLP papers and Braille papers unless otherwise stated.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 1MA1_1H			
Question		**Modification**	**Mark scheme notes**
5		Wording added 'Look at the diagram for Question 5 in the Diagram Booklet. It shows'. Wording removed 'The'. Wording removed 'lies'. Diagram enlarged. Cross changed to a dot.	Standard mark scheme
6		Wording added 'Look at the diagram for Question 6 in the Diagram Booklet. It'. Wording removed 'The diagram'. Diagram enlarged. Diagram rotated such that ABC is horizontal. Angles moved outside of the angle arcs and angle arcs made smaller. Wording added 'Angle DAB = x° Angle DBA = y° Angle DBC = w°'	Standard mark scheme
7		Letter 'x' changed to 'y'.	Standard mark scheme but note change of letter
9		Wording added 'Look at the diagram for Question 9 in the Diagram Booklet. It shows a grid.' Wording added 'on the grid in the Diagram Booklet'. Diagram enlarged. Axis labels moved to the top of the vertical axis and to the left of the horizontal axis. x and y labels moved to the top of the vertical axis and to the right of the horizontal axis. Small squares removed. Right axis labelled. Open headed arrows.	Standard mark scheme
11		Wording added 'Look at the diagram for Question 11 in the Diagram Booklet. It shows an incomplete box plot diagram.' Wording added 'below'. Wording added 'in the Diagram Booklet'. Table enlarged and left aligned. Diagram enlarged. Label moved to the left. Lower quartile changed from 780 to 700. For Braille i & ii added to table and answer. In (a)(ii) wording added 'There are two spaces to fill.'	Standard mark scheme but note change of 780 to 700 for the LQ.
15		Wording added 'Look at the diagram for Question 15 in the Diagram Booklet.' Wording 'Here is' removed and replaced with 'The diagram shows'. Diagram enlarged. Axis labels moved to the top of the vertical axis and to the left of the horizontal axis. d and t labels moved to the top of the vertical axis and to the right of the horizontal axis. Small squares removed. Label right axis. Open headed arrows	Standard mark scheme

PAPER: 1MA1_1H			
Question		**Modification**	**Mark scheme notes**
19		Wording added 'Look at the diagram for Question 19 in the Diagram Booklet. It'. Wording removed 'Here is'. Diagram enlarged. Axis labels moved to the top of the vertical axis and to the right of the horizontal axis. Open headed arrows. Value '0.3' changed to '0.75'	(a) M1 line drawn at 0.75 to sine curve **or** one correct value A1 for answer in the range 45 to 59 and 121 to 135 (b) Standard mark scheme
20		Wording added 'Look at the diagram for Question 20 in the Diagram Booklet. It shows'. Wording removed 'Here is'. Wording added 'BC = 5 cm BA = 10 cm AC = 5 $\sqrt{7}$ ' Diagram enlarged.	Standard mark scheme
21	(a)	Wording added 'Look at the diagram for Question 21 in the Diagram Booklet. It shows a grid.' Value '169' changed to '36', to reduce the size of the circle the candidate draws, and such that the grid fits in the Diagram Booklet. Diagram enlarged. Diagram cut at y=9, y=-9, x=9, and x=-9. Small squares removed. Axis labels moved to the top of the vertical axis and to the right of the horizontal axis.	Standard mark scheme but note the radius is now 6
21	(b)	Value '169' changed to '36'.	Standard mark scheme but note the solutions are now as follows: $x = 3.4$, $y = 5.0$ $x = -3.4$, $y = -5.0$

17	(a)	$\frac{\sqrt{7}}{7}$	B1	for $\frac{\sqrt{7}}{7}$ **or** $\frac{k\sqrt{7}}{7k}$ **or** $\frac{\sqrt{7k^2}}{7k}$ where k is an integer not equal to 0	
	(b)	$3\sqrt{5}$	M1	for writing $\sqrt{80}$ as $\sqrt{16\times5}$ **or** $\sqrt{16}\times\sqrt{5}$ **or** $4\sqrt{5}$	
			A1	for $3\sqrt{5}$ or $\sqrt{45}$	

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
15	$\frac{\sqrt{a}-1}{a-1}$	M1	for a correct method to rationalise the denominator, eg, $\frac{1}{\sqrt{a}+1}\times\frac{\sqrt{a}-1}{\sqrt{a}-1}$ or $\frac{1}{\sqrt{a}+1}\times\frac{1-\sqrt{a}}{1-\sqrt{a}}$	Condone use of a prime number in place of a for the M1 Do not ISW
		A1	for $\frac{\sqrt{a}-1}{a-1}$ or $\frac{1-\sqrt{a}}{1-a}$	

Question	Answer	Mark	Mark scheme	Additional guidance
14	6	M1	for either $\sin 30 = 0.5$ oe **or** $\sin 45 = \frac{1}{\sqrt{2}}$ or $\frac{\sqrt{2}}{2}$	
		M1	for $\frac{AB}{\sin 45} = \frac{3\sqrt{2}}{\sin 30}$ oe	
		A1	cao	

Question	Answer	Mark	Mark scheme	Additional guidance
16 (a)	$3\sqrt{5}$	M1	for $\frac{15}{\sqrt{5}}\times\frac{\sqrt{5}}{\sqrt{5}}$ or $\frac{15}{\sqrt{5}}\times\frac{-\sqrt{5}}{-\sqrt{5}}$	
		A1	for $3\sqrt{5}$ or $\sqrt{45}$	
(b)	$\frac{32-9\sqrt{3}}{11}$	M1	(indep) for writing $\sqrt{75}$ as $5\sqrt{3}$	This mark can be awarded whenever this is seen, which might be later in the process.
		M1	for method to rationalise the denominator, eg $\frac{\sqrt{75}-2}{1+2\sqrt{3}}\times\frac{1-2\sqrt{3}}{1-2\sqrt{3}}$ **or** $\frac{5\sqrt{3}-2}{1+2\sqrt{3}}\times\frac{1-2\sqrt{3}}{1-2\sqrt{3}}$	
		M1	(dep on previous M1) for expanding terms, condone one error in numerator or denominator eg $\frac{\sqrt{75}-2\sqrt{75}\sqrt{3}-2+4\sqrt{3}}{1-2\sqrt{3}+2\sqrt{3}-4\sqrt{3}\sqrt{3}}$ **or** $\frac{5\sqrt{3}-10\sqrt{3}\sqrt{3}-2+4\sqrt{3}}{1-2\sqrt{3}+2\sqrt{3}-4\sqrt{3}\sqrt{3}}$	
		A1	for $\frac{32-9\sqrt{3}}{11}$ oe eg $\frac{-32+9\sqrt{3}}{-11}$	Accept $a = 32$, $b = 9$, $c = 11$

Paper: 1MA1/1H

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
16 (a)	$5\sqrt{7}$	M1	for $\frac{35}{\sqrt{7}}\times\frac{\sqrt{7}}{\sqrt{7}}$ $(=\frac{35\sqrt{7}}{7})$ or $\frac{35}{\sqrt{7}}\times\frac{-\sqrt{7}}{-\sqrt{7}}$ $(=\frac{-35\sqrt{7}}{-7})$	
		A1	for $5\sqrt{7}$ or $\sqrt{175}$	
(b)	$a = 7$, $b = 5$	B1	for $\sqrt{27}=3\sqrt{3}$ or $2\sqrt{27}=6\sqrt{3}$	B1 can be awarded whenever this is seen, which might be later in the process.
		P1	for process to rationalise the denominator, eg $\frac{\sqrt{27}-1}{2-\sqrt{3}}\times\frac{2+\sqrt{3}}{2+\sqrt{3}}$ **or** $\frac{3\sqrt{3}-1}{2-\sqrt{3}}\times\frac{2+\sqrt{3}}{2+\sqrt{3}}$ oe	
		P1	(dep on previous P1) for expanding terms, condone one error in numerator or denominator, eg $\frac{2\sqrt{27}+\sqrt{27}\sqrt{3}-2-\sqrt{3}}{4+2\sqrt{3}-2\sqrt{3}-\sqrt{3}\sqrt{3}}$ **or** $\frac{6\sqrt{3}+3\sqrt{3}\sqrt{3}-2-\sqrt{3}}{4+2\sqrt{3}-2\sqrt{3}-\sqrt{3}\sqrt{3}}$ **or** $6\sqrt{3}+9-2-\sqrt{3}$ oe	
		A1	for $a = 7$, $b = 5$	Accept $7+5\sqrt{3}$

Paper: 1MA1/1H

edexcel

GCSE

Edexcel GCSE

Mathematics A 1387

July 2004

advancing learning, changing lives

Mark Scheme

Edexcel GCSE

Mathematics A 1387

London Qualifications is one of the leading examining and awarding bodies in the UK and throughout the world. It incorporates all the qualifications previously awarded under the Edexcel and BTEC brand. We provide a wide range of qualifications including general (academic), vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, our centres receive the support they need to help them deliver their education and training programmes to learners.

For further information please call Customer Services on 0870 240 9800, or visit our website at www.edexcel.org.uk

July 2004

Publications Code UG015616

All the material in this publication is copyright
© London Qualifications Ltd 2004

CONTENTS

Notes on Marking Principles....2
Paper 5514....4
Paper 5515....8
Paper 5516.... 11
Paper 5517.... 15
Paper 5518.... 19
Paper 5519.... 23

NOTES ON MARKING PRINCIPLES

NOTES ON MARKING PRINCIPLES

1 Three types of mark are available

M marks - awarded for correct working seen or implied.
A marks - conditional accuracy marks which are awarded for accurate working following the award of M marks.
B marks - unconditional accuracy marks (independent of M).

2 Abbreviations

cao - correct answer only.
ft - follow through.
$\surd$ - Denotes a "follow through" answer.

SC - special case.
isw - ignore subsequent working.
oe - or equivalent (and appropriate).
NB: a candidate cannot benefit from both isw and ft.

3 If no working shown, then

correct answers normally score full marks
incorrect (even though nearly correct) answers score no marks.

4 Marking instructions

Misread loses A marks (and sometimes B marks) on that part but ft can be allowed on subsequent parts. M marks can still be earned. If in doubt contact your team leader.
If there is a wrong answer in the answer space DO CHECK the working in the body of the script.

- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.
- If there is a wrong answer in the answer space DO CHECK the working in the body of the script.
- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.

5 Style of marking

Answer correct: tick and write part mark in margin NEXT TO BRACKETED MARK.
Answer incorrect: cross, but show M, A or B marks if any earned in body of script and transfer the total of these to the margin next to the bracket mark.
Total for each double page at bottom right page (except for back if used), FINAL TOTAL IN RELEVANT BOX ON FRONT COVER.
Nought in margin for fully incorrect question or page AND FOR NO ATTEMPT.
Where no attempt has been made a line should be put in the answer space and zero in the margin next to the bracketed mark
CHOICE OF METHOD – No marks unless one answer is in answer space – then mark that.
CROSSED OUT WORK - if not replaced this should be marked (if legible).
There must always be a mark next to bracketed mark in the margin/

6 Follow Through Marks

Follow throughs are guided by two principles:

(a) Follow throughs which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous, do not award.
(b) Follow throughs which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

7 Probability

Probability answers must be given as fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least two decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.

6 Linear Equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

8 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

PAPER 5501					
No		**Working**	**Answer**	**Mark**	**Notes**
1			K: 1020 L: 8.06	2	B1 for 1020 or 1,020 B1 for 8.06 Accept £8.06p and £8,06
2	(a)		28, 33	2	B1, B1 (B1 ft for "28" + 5 if both numbers >25)
	(b)			1	B1 for add 5, +5, for going up in 5's the difference is 5 oe
	(c)			1	B1 for they end in 3 and 8, it ends in 7; or refers to 383 and 388 are in sequence
3	(a)(i)		$\frac{1}{2}$	2	B2 for ½ accept half (B1 for an equivalent unsimplified fraction eg 4/8 or 50% or 0.5)
	(ii)		2 rectangles shaded	1	B1 for correct shading (any 2 rectangles)
	(b)(i)		40	4	B1 for 40 cao
	(ii)		150 000		B1 for 150 000 cao (accept 150,000 not 150.000)
	(iii)		6.55		B1 for 6.55 cao not $6.5^{½}$
	(iv)		$\frac{3}{8}$		B1 for $\frac{3}{8}$ oe accept 0.375
	(c)		Cross 3cm from *A*	1	B1 mark a cross 3 cm (±2 mm) from *A*
4	(a)(i)		metres	2	B1 for metres (m)
	(ii)		grams		B1 for grams (g)
	(b)		miles	1	B1 for miles
5	(a)		(3, 2)	1	B1 for (3, 2)
	(b)(i)		*Q* at (0, 3)	1	B1 for *Q* plotted correctly on *y*-axis at (0, 3) ±2mm
	(ii)		*R* at (−2, −3)	1	B1 for *R* plotted correctly at (−2, −3) ±2mm
6	(a)(i)		30	2	B1 for 30
	(ii)		25		B1 for 25
	(b)(i)		4 shapes	2	B1 for drawing 4 shapes
	(ii)		1½ shapes		B1 for drawing 1½ shapes

PAPER 5501					
	No	**Working**	**Answer**	**Mark**	**Notes**
7	(a)		30	1	B1 for 30
	(b)	$20 \times 3 + 5$	65	2	M1 for $20 \times 3 + 5$ oe A1 for 65
8	(a)		Draws perp.	1	B1 for correctly drawing perp must touch line or cut line *AB* ± 2mm
	(b)		Sketches a cylinder	1	B1 for sketching cylinder
9	(a)	$\begin{array}{r} 1974 \\ \underline{6580} \\ 8554 \end{array}$	855.4	3	M1 for complete method with relative place value correct, condone 1 error in multiplication A2 cao (A1 for digits 8554 seen or A1 for "855.4" dependent on 1 arithmetic error)
	(b)	$990 \div 69 = 14.3\ldots$ or 14 rem 24	14	4	Method 1 - Everything excluding long division M2 for a valid method with no errors …need to see 966 or 1035 (M1 for a valid method, including estimation, that reaches an answer between 901 and 1099 inclusive. Ignore errors.) A2 for 14 with a fully correct method (A1 (dep on 1st M1) for an answer between 14 and 15 inclusive. Ignore errors. Cannot have used estimation.) Note: Estimation can only score a maximum of M1 unless accompanied by another method or 966 or 1035 see (award M2) Method 2 – long division method M2 for $990 \div 69$ = integer answer between 12 and 19 inclusive and rem 30 seen (M1 for $990 \div 69$ with 1 in tens column.) A2 for 14 with fully correct method (A1 (dep on 1st M1) for an answer between 14 and 15 inclusive.)

PAPER 5501					
	No	**Working**	**Answer**	**Mark**	**Notes**
10	(a)(i)		7	2	B1 for 7
	(ii)		−10		B1 for −10
	(b)(i)		6	2	B1 for 6 (accept −6)
	(ii)		8		B1 for 8 (accept −8)
	(c)		−7	1	B1 for −7 cao
11	(a)		B and D	2	B1, B1 (–1 each extra)
	(b)(i)		A	2	B1 for A
	(ii)		3		B1 for 3
12	(a)	800 − 144	656	2	M1 for at least 1 digit correct and in correct position needs to be 3 digit number A1 for 656
	(b)	144 is less than 200 so Trudy is wrong		2	B1 for 200 B1ft for 'correct' explanation based on cand's "200"
	(c)	$\frac{45\times800}{100}$	360	2	M1 for 45 × 800 ÷ 100 oe A1 for 360
	(d)	$\frac{176}{800}\times100$	22	2	M1 for $\frac{176}{800}\times100$ oe A1 for 22
13	(a)(i)		25	4	B1 for 25
	(ii)		28		B1 for 28
	(iii)		5 and 20		B1 for 5 and 20
	(iv)		26 and 33		B1 for 26 and 33
	(b)	$2^3 = 8$ or $2^3 = 2 \times 2 \times 2$ (which is not 6)		1	B1 for valid 'explanation'
14			10 m^2	2	B1 for 10 B1 indep for m^2.

PAPER 5501					
No		**Working**	**Answer**	**Mark**	**Notes**
15	(a)	$\frac{11}{12}-\frac{10}{12}$	$\frac{1}{12}$	2	M1 for correctly writing both fractions to a common denominator A1 for $\frac{1}{12}$ oe
	(b)	$\frac{70\times400}{200}$	140	2	B2 for 140 (accept 136) (B1 for sight of any two of 70, 400 or 200)
16	(a)		$2y$	1	B1 for $2y$ or $2\times y$, $y2$, $y\times2$
	(b)		$3p^2$.	1	B1 for $3p^2$ or $3\times p^2$ or $p^2\times3$ or $p^2 3$
	(c)		$x(x-3)$	2	B2 for $x(x-3)$ (B1 for x (x…….)) $\frac{25}{60}$ wrongly cancelled gets B1 ISW
17	(a)		2 23 9 34 15 2 9 26 17 25 18 60	3	B3 for all correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
	(b)		$\frac{25}{60}$	1	B1 for $\frac{25}{60}$ or $\frac{5}{12}$ oe $\frac{25}{60}$ wrongly cancelled gets B1 ISW

PAPER 5501					
	No	**Working**	**Answer**	**Mark**	**Notes**
18	(a)(i)		60	2	B1 for 60
	(ii)				B1 for all **angles equal** so equilateral triangle oe
	(b)(i)		130	3	B1 for 130
	(ii)				B1 for isosceles triangle oe or 2 angles equal accept $\angle Q = \angle R$ B1 for angles on a straight **line** add up to **180°**oe (180° could be in working)
	(c)		64	1	B1 for 64
19	(a)		1	1	B1 for 1
	(b)		3	1	B1 for 3
	(c)	$9\times1+3\times2+5\times3+3\times4$ (= 42) "42" ÷ 20	2.1	3	M1 for 9×1, 3×2, 5×3, 3×4 or for 42 seen M1 (dep) for "42" ÷ 20 A1 for 2.1 or $2\frac{1}{10}$ or $2\frac{2}{20}$
20	(a)		09 05	1	B1 for 09 05 oe
	(b)		7	1	B1 for 7
	(c)		10	1	B1 for 10
	(d)	7km in 20 mins	21	3	B1 for 20 (minutes) oe or $\times3$ seen M1 for $\frac{"7"}{"20"}$ A1 cao [SC: "7" × 3 seen gets B1 M1]

PAPER 5501				
No	**Working**	**Answer**	**Mark**	**Notes**
21	Vertices of new triangle at (2, −2) (2, 6) and (6, −2)	Correct triangle drawn	3	B3 cao (B2 for either correct triangle in wrong translated position or for 2 vertices correct) (B1 a right angled triangle with horizontal length 4 or vertical length 8 in the same orientation as the shaded triangle) SC B2 for s + 3 totally correct
22 (a) (b) (c)	 8560 ÷ (10 × 10)	8 Pentagon 85.6	1 1 2	B1 for 8 B1 for pentagon M1 for 8560 ÷ (10 × 10) oe A1 for 85.6
23	$2x + x + 100 + 47 = 360$ $2x + x = 360 - 100 - 47$ $x = 71$ Largest angle = $2x$ =	142	4	M1 for $2x + x + 100 + 47 = 360$ or 360 – 147 or 213 seen M1 dep for correctly separating x-terms and non x terms or "360 – 147" ÷ 3 A1 for $x = 71$provided M2 awarded A1 ft for 142

PAPER 5502					
	No	**Working**	**Answer**	**Mark**	**Notes**
1	(a)		Diagram	1	B1 cao
	(b)		22, 26	1	B1 cao
	(c)		46	1	B1 cao
2	(a)	£5.95 + £1.62 + 25p + 2 × 48p = £8.78	1.22	3	M1 for 3 of £5.95, £1.62, 25p, 48p seen with intention to add or 170 seen A1 for £8.78 A1 ft from " 8.78"
	(b)	160 ÷ 12 = 13.(333…)	14	2	M1 for 160 ÷ 12, or 13 or better or 12×13 or 12×14 A1 14 cao
	(c)	"14" × £12.20 = £170.8**0**	170.80	2	M1 for "14" × 12.20 A1 for 170.8**0** or 158.6**0** if 13 in (b) NB: 170.8 or 158.6 gets M1 A0
3	(a)(i)		1459	1	B1 cao
	(ii)		9541	1	B1 cao
	(b)		9 + 5 = 14	1	B1 cao
	(c)		0	1	B1 cao
4	(a)		1123	1	B1 cao
	(b)		1125	1	B1 cao
	(c)		43	1	B1 cao
	(d)	67 – "43"	24	2	M1 for 67 – "43" A1 ft from "43"
	(e)(i)		80	2	B1 cao
	(ii)		56 – 58		B1 answer in range 56 to 58 inc
	(f)		Completes graph	2	B2 for (Apr , 70) and (May , 60 < p < 65) plotted and joined (B1 for one point plotted)

PAPER 5502					
No		Working	Answer	Mark	Notes
5	(a)(i)		4	1	B1 cao
	(ii)		10	1	B1 cao
	(b)		28	2	B2 for 28 (B1 for 29 or 14×2)
6	(a)		$\frac{3}{4}$	1	B1 accept 0.75, 75%
	(b)	$\frac{3}{4} \times 28 = 21$	15.54	3	M1 for $\frac{3}{4} \times 28$ or 21 seen M1 for "21" $\times$ 74 or 28×74 or 7×74 A1 cao
	(c)(i)		44 – 46	2	B1 44 – 46 inc
	(ii)		5.2 – 5.4		B1 5.2 – 5.4 inc
7	(a)		Correct shape	1	B1
	(b)		Correct shape	1	B1
8	(a)		Hexagon	1	B1
	(b)		Right	1	B1 cao
	(c)		Drawing	2	B2 for 7 additional hexagons with at least 2 points at which 3 hexagons meet (B1 for one point at which 3 hexagons meet)
9	(a)		$\frac{3}{4}$	2	M1 $\frac{80-20}{80}$ or $\frac{60}{80}$ or 0.75 (oe) A1 cao Do not accept equivalents
	(b)		15	1	B1 cao
	(c)		Iron and Cook set	1	B1 for both Accept mixer and fryer
	(d)		Microwave	1	B1 cao
	(e)		Bar chart	2	B1 one column (mixer): 70, 90 B1 second column (fryer): 70, 85 [SC: B1 columns correct but reversed]

PAPER 5502					
No		Working	Answer	Mark	Notes
10	(a)		Acute	1	B1
	(b)		Reflex	1	B1
	(c)		reason	1	B1 reason eg 120 + 230 = 350 ≠ 360
11	(a)		$\frac{1}{2}$	1	B1 for $\frac{1}{2}$ oe
	(b)		reason	1	B1 for reason e.g could get 30 heads
12	(a)	7.5×4	30	2	M1 for 7.5×4 or $7\frac{1}{2} \times 4$ A1 cao
	(b)	$\sqrt{(42.25)}$	6.5	2	B2 for 6.5 (B1 for 42.25)
13	(a)		$6x$	1	B1 accept $6 \times x$, $x6$ oe
	(b)		$x - 4$	1	B1 cao
	(c)		$12(x - 4)$	1	B1 for $12(x - 4)$ or $12x - 48$ or $12 \times x - 48$ oe
	(d)		$18x - 48$	2	M1 for "$6x$" + "$12(x-4)$" or "$6x$" + "$12x - 48$" oe A1 ft $18x - 48$, $2(9x - 24)$, $3(6x - 16)$, $6(3x - 8)$
14	(a)		90	2	B2 for 90 ± 2 (B1 for 9 ± 0.2)
	(b)		317	1	B1 for 317 ± 4
15	(a)	Accurate drawing of triangle		2	B1 cao for 38° (±2°) B1 cao (indep) for BC drawn 7.3 cm (±2mm) and completing the triangle.
	(b)	Measure angle A	63°	1	B1 for 63° (±2°) or ft their diagram (±2°)

PAPER 5502					
	No	Working	Answer	Mark	Notes
16	(a)	0 × 1 = 0 1 × 1 = 1 2 × 4 = 8 3 × 8 = 24 4 × 8 = 32 5 × 7 = 35	100	3	M1 col 1 × col 2 (at least 3 shown) M1 (dep) sum of totals A1 cao SC: M1 M1 A0 for 101
	(b)	Either 35.50 ÷ 1.42 = 25; 26.99 – 25 = 1.99 Cheaper in the USA Or 26.99 × 1.42 = 38.33; 38.33 – 35.50 = 2.83 Cheaper in the USA		3	M1 35.50 ÷ 1.42 A1 25 seen B1 ft (dep on M1) £1.99 with conclusion (must have units) **OR** M1 26.99 × 1.42 A1 38.32 or 38.33… seen B1 ft (dep on M1) $2.83 or $2.82 with conclusion (must have units) must be to 2dp
17	(a)		-1, (1), (3), 5, 7, 9	2	B2 cao (B1 for 2 values)
	(b)		Graph	2	B1 ft for plotting points ±1/2 square B1 cao for line between $x = -2$ and $x = 3$
	(c)(i)		0.4	2	B1 for 0.4 or ft from single straight line with positive gradient
	(ii)		1.2		B1 for 1.2 or ft from single straight line with positive gradient

PAPER 5502					
	No	Working	Answer	Mark	Notes
18	(a)		Plots	1	B1 cao
	(b)		description	1	B1 dynamic relationship or "positive" (correlation)
	(c)		line of best fit	1	Line within overlay region, and to the extent of.
	(d)(i)	280 g →	reading	2	B1 ft from single straight line of positive gradient (±1/2 square)
	(ii)	120 pages →	reading		B1 ft from single straight line of positive gradient (±1/2 square)
19			200 150 225 150 10	3	B3 cao (B2 for three correct, or B1 for one correct)
20	(a)			2	B1 B1 for correct sketch – ignore additional internal lines B1 B0 for rectangle or right-angled triangle or correct shape without line NB 3-D sketch gets B0 B0
	(b)			2	B1 B1 for correct plan – ignore internal lines – accept a rotated plan, not reflected B1 B0 for a single rectangle drawn NB 3-D sketch gets B0 B0
21	(a)	163.25 – 35.50 = 127.75 127.75 ÷ 18.25 = 7	8	3	M1 163.25 – 35.50 (or sight of 127.75) M1 (dep) "127.75" ÷ 18.25 A1 cao SC: M2 for 7 days
	(b)	1.175 × £64 or $\frac{17.5}{100}$ × 64 or £11.20 seen	75.20	2	M1 1.175 × 64, 0.175 × 64 oe or 11.20 or 75.2 seen or 6.40, 3.20 and 1.60 seen A1 cao

PAPER 5502					
	No	Working	Answer	Mark	Notes
22	(a)		6	1	B1 cao
	(b)		$t^2 - 2t$	1	B1 oe
	(c)		$3(y - 4)$	1	B1 Accept $3 \times (y - 4)$ or $3(y - 4$
23	(a)		Reason	1	B1 Makes some mention of bias either directly or making reference to an insufficient or biased range of responses
	(b)		Reason	1	B1 (a) an insufficient range of responses (b) No mention of money (c) No time frame in the question (d) Misunderstanding of "A lot" and "Not much"

PAPER 5503				
No	**Working**	**Answer**	**Mark**	**Notes**
1		10.	1	B1 for 10
2 (a)(i)		$3e + 2f$	2	B1 for $3e + 2f$
(ii)		$3p^2$		B1 for $3p^2$
(b)	$5 \times -3 + 1$	-14	2	M1 for $5 \times -3 + 1$ A1 for -14
3	1974 6580 8554	855.4	3	M1 for complete correct method with relative place value correct, condone 1 error in multiplication. A2 cao (A1 for digits 8554 seen or A1 ft for "855.4" dependent on one arithmetic error only)
4 (a)		2 23 9 34 15 2 9 26 17 25 18 60	3	B3 all correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
(b)		$\frac{25}{60}$	1	B1 for $\frac{25}{60}$ or $\frac{5}{12}$ oe
5 (a)	$\frac{45}{100} \times 800$	360	2	M1 for $\frac{45}{100} \times 800$ oe A1 for 360
(b)	$\frac{176}{800} \times 100$	22	2	M1 for $\frac{176}{800} \times 100$ oe A1 for 22

PAPER 5503					
	No	Working	Answer	Mark	Notes
6	(a)	$\frac{11}{12}-\frac{10}{12}$	$\frac{1}{12}$	2	M1 for correctly writing both fractions to a common denominator A1 for $\frac{1}{12}$ oe
	(b)	$\frac{70\times400}{200}$	140	2	B2 for 140 (accept 136) (B1 for sight of any two of 70, 400 or 200)
7	(a)(i)		60	2	B1 for 60
	(ii)				B1 for all angles equal so equilateral triangle oe
	(b)(i)		130	3	B1 for 130
	(ii)				B1 for isosceles triangle oe B1 for angles on a straight line add up to 180°oe
	(c)		64	1	B1 for 64
8		$9\times1+3\times2+5\times3+3\times4$ (=42) "42" ÷ 20	2.1	3	M1 for 9×1, 3×2, 5×3, 3×4 or for 42 seen M1 (dep) "42" ÷20 A1 for 2.1 or $2\frac{1}{10}$ or $2\frac{2}{20}$
9	(a)		09 05	1	B1 for 09 05 oe
	(b)		7	1	B1 for 7
	(c)		10	1	B1 for 10
	(d)	7 km in 20 minutes	21	3	B1 for 20 (minutes) oe or $\times3$ seen M1 for $\frac{"7"}{"20"}$ A1 cao SC: "7" $\times3$ gets B1 M1

PAPER 5503					
No		**Working**	**Answer**	**Mark**	**Notes**
10		Vertices of new triangle at (2, −2) (2, 6) and (6, −2)	Correct triangle drawn	3	B3 cao (B2 for either correct triangle in wrong translated position or for 2 vertices correct) (B1 for a right-angled triangle with horizontal length 4 or vertical length 8 in the same orientation as shaded triangle) SC B2 for scale factor 3, totally correct
11		$2^2 + 3 =$	7 which is not even	2	B2 (B1 for correctly evaluating $n^2 + 3$ with a prime number value for n.)
12	(a)	360 ÷ 5	72	2	M1 for 360 ÷ 5 oe A1 for 72
	(b)	8560 ÷ (10 × 10)	85.6	2	M1 for 8560 ÷ (10 × 10) oe A1 for 85.6
	(c)(i)	Least length = 100.5	100.5	2	B1 for 100.5
	(ii)	Greatest length = 101.5	101.5		B1 for 101.5 or 101.499 or better
13		$2x + x + 100 + 47 = 360$ $2x + x = 360 - 100 - 47$ $x = 71$ Largest angle = $2x$ =	142	4	M1 for $2x + x + 100 + 47 = 360$ or 360 – 147 or 213 seen M1 (dep) for correctly separating x-terms and non x terms or $\frac{"360-147"}{3}$ A1 for $x = 71$ A1 ft (provided M2 awarded) for "142"
14	(a)(i)		2.21	2	B1 for 2.21
	(ii)		0.013		B1 for 0.013
	(b)	LCM = 3 × 13 × 17 = 3 × 221	663	2	M1 for 3 × 13 × 17 oe A1 for 663

PAPER 5503				
No	**Working**	**Answer**	**Mark**	**Notes**
15		Tick boxes 1,3 & 6	3	B1 for $\frac{\pi abc}{2d}$; B1 for $2a^2$; B1 for $2\left(c^2+d^2\right)$ (−B1 for each additional expression ticked (>3) to a minimum of 0)
16	200×0.2	40	2	M1 for 200×0.2 or $\frac{40}{200}$ seen A1 for 40
17 (a)	eg 2 | 108 2 | 54 3 | 27 3 | 9 3	$2^2 \times 3^3$	3	M2 for full systematic method of at least 4 divisions by prime numbers oe factor trees; condone 1 calculation error (M1 for 108 written as a correct product(s) with only one non-prime or equivalent division or a full process with 2 calculation errors.) A1 for $2^2 \times 3^3$ (accept $2 \times 2 \times 3 \times 3 \times 3$)
(b)		12	1	B1 for 12
18	Perpendicular from P to intersecting arcs (within tramlines); perpendicular at least 2cm long		2	M1 relevant pair of arcs crossing within tramlines A1 SC M1 AO for a full construction of a perpendicular to AB not passing through P.
19	15×10	150 cm^3.	3	M1 for 15×10 A1 for 150 B1 for cm^3.

PAPER 5503				
No	**Working**	**Answer**	**Mark**	**Notes**
20		$y = 2x + 6$	2	B2 for $y = 2x + 6$ (B1 for $y = 2x + k$, $k \neq 1$ or for $y = mx + 6$, m $\neq 0$ or for $2x + 6$)
21	$6\frac{2}{5} = \frac{32}{5}$ Area of triangle = $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ (=2) Length of a side of sq. = $\sqrt{18 \times "2"}$ (=6) Perimeter of square = 4 × 6	24	5	B1 for $6\frac{2}{5} = \frac{32}{5}$ oe or $3\frac{1}{5} = \frac{16}{5}$ or $\frac{30}{8} + \frac{2}{8}$ oe (or implied by area of triangle=2) M1 for $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ oe M1 for area of square = 18 × product of two lengths A1 for $\sqrt{18 \times "2"}$ A1 for 24
22 (a)		$x(x-3)$	2	B2 for $x(x-3)$ (B1 for x (…….))
(b)		k^3.	1	B1 for k^3.
(c)(i)	$4x + 20 + 3x - 21$	$7x - 1$	4	M1 for **three** of 4 terms $4x + 20 + 3x - 21$ (or better) A1 for $7x - 1$
(ii)	$x^2 + 3yx + 2xy + 6y^2$	$x^2 + 5xy + 6y^2$		M1 for **three** of 4 terms $x^2 + 3yx + 2xy + 6y^2$ A1 for $x^2 + 5xy + 6y^2$
(d)		$(p+q)(p+q+5)$	1	B1 for $(p+q)(p+q+5)$

PAPER 5503					
	No	Working	Answer	Mark	Notes
23	(a)		32	1	B1 for 32 (accept 31.5 to 33.5 inclusive)
	(b)			3	B1 for ends of whiskers at 9 and 57, (with a box) B1 for ends of box at 16 and 45 (± 0.5) B1 for median marked at 32 or a complete box and whisker diagram drawn with a median inside the box
	(c)	Median(B)>Median(G); on average boys take longer IQR(B)>IQR(G); times for boys have a greater spread		2	B1 eg for comparison of medians B1 eg for comparison of (interquartile) ranges
24	(a)	$6y = 15 - 5x$	$y = \frac{15-5x}{6}$	2	M1 for either $6y = 15 - 5x$ or -6 $y = 5x - 15$ or for $\frac{5x}{6} + y = \frac{15}{6}$ or a correct ft on sign error to $y=$ A1 for $y = \frac{15-5x}{6}$ oe
	(b)	$6k + 5(-21) = 15$	20	2	M1 for substitution of $x = -21$ (or $x = 21$) into given equation or candidate's answer to (a) A1 for $k = 20$
	(c)(i)		Region R indicated	3	B2 correct region shaded (accept unshaded if R clear) (B1 shaded or R region satisfies 3 of the 4 given inequalities with same boundaries)
	(ii)		(1,1)		B1 for (1,1)

PAPER 5503					
	No	Working	Answer	Mark	Notes
25	(a)		60	1	B1 for 60 B1 for *BDC* = 25
	(b)	eg Angle *BDC* = 25 *ADB* = 60 − 25	35	2	[Award the mark for equivalent approaches] B1 ft for (a) − 25
	(c)	Ben is correct; angle *DAB* = 65 + 25 = 90 and since angle in a semi-circle is 90°, *BD* must be a diameter		1	B1 for full valid justification

PAPER 5504					
	No	Working	Answer	Mark	Notes
1	(a)	Accurate drawing of triangle		2	B1 cao for 38º (±2º) B1 cao (indep) for BC drawn 7.3 cm (±2mm) and completing the triangle.
	(b)	Measure angle A	63º	1	B1 for 63º (±2º) or ft their diagram (±2º)
2	(a)	0 × 1 = 0 1 × 1 = 1 2 × 4 = 8 3 × 8 = 24 4 × 8 = 32 5 × 7 = 35	100	3	M1 col 1 × col 2 (at least 3 shown) could be implied by answers. M1 (dep) sum of totals A1 cao SC: M1 M1 A0 for 101
	(b)	Either $35.50 ÷ 1.42 = £25; £26.99 - £25 = £1.99 Cheaper in the USA Or £26.99 × 1.42= $38.33; $38.33 - $35.50 = $2.83 Cheaper in the USA		3	M1 35.50 ÷ 1.42 A1 25 seen B1 ft (dep on M1) £1.99 with conclusion; units needed OR: M1 26.99 × 1.42 A1 38.33 or 38.32…seen B1 ft (dep on M1) $2.83 or $2.82 with conclusion; units needed
3	(a)		-1, (1), (3), 5, 7, 9	2	B2 cao (B1 for 2 values)
	(b)		Graph	2	B1 ft for plotting points ±1/2 square B1 cao for line between $x = -2$ and $x = 3$
	(c)(i) (ii)		0.4 1.2	2	B1 0.4 or ft single straight line with positive gradient B1 1.2 or ft single straight line with positive gradient

PAPER 5504					
	No	Working	Answer	Mark	Notes
4	(a)			2	B1 Rectangle drawn B1 Triangle attached Ignore additional internal lines; 3D sketch gets B0
	(b)			2	B1 single Rectangle drawn B1 Attached smaller rectangle drawn Accept rotated plan, Ignore internal lines down; Nets get B0
5	(a)	£163.25 - £35.50 = £127.75 £127.75 ÷ £18.25 = 7	8	3	M1 163.25 – £35.50 (or sight of £127.75) M1 (dep) "127.75" ÷ 18.25 A1 cao SC: M2 for 7 days
	(b)	1.175 × £64 or $\frac{17.5}{100}$ × 64 or £11.20 seen	75.20	2	M1 1.175 × 64.00 0.175 × 64 oe or 11.20 or 75.2 seen OR 6.40, 3.20, and 1.60 seen A1 cao
	(c)	54 × £2.38 =	128.52	2	M1 54 × £2.38 A1 cao
6		Tessellation	Drawing	2	B2 seven additional hexagons, with at least 2 points at which 3 hexagons meet (B1 one point at which 3 hexagons meet).
7			200 150 225 150 10	3	B3 cao (B2 for three correct, or B1 for one correct)
8	(a)(i)	Bearing measured	70	2	B1 70 ± 4
	(ii)	Bearing measured	317		B1 317 ± 4
	(b)	Locus (circle) drawn & shaded		2	B1 circle ±2 mm centre Manchester B1 shading with accurate or approximate circle within tolerance

PAPER 5504					
	No	Working	Answer	Mark	Notes
9	(a)		Plots	1	B1 cao
	(b)		Description	1	B1 dynamic relationship or "positive" (correlation)
	(c)		Line of best fit	1	Line within overlay region, and to the extent of.
	(d)(i)	280 g →	Reading	2	B1 ft from single straight line of positive gradient (±1/2 square) B1 ft from single straight line of positive gradient (±1/2 square)
	(ii)	120 pages →	Reading		
10	(a)	$\frac{1}{6}, \frac{3}{8}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}$		2	B2 for all correct (B1 for 1 error or all correct but wrong order, or use of a common denominator decimals).
	(b)	3/5, 65%, 2/3, 0.72, 3/4		2	B2 for all correct (B1 for 1 error or all correct but wrong order or conversions to decimals oe)
11	(a)		182, 178, 180, 184	2	M1 mean of any three consecutive months, eg (147 + 161 + 238) ÷ 3 oe A1 cao
	(b)	Sale price = 80% Fun Friday price = 70% of 80% = 56% (oe)		2	B1 B1 for a fully correct explanation involving a worked example (oe) B1 a partially complete explanation

PAPER 5504					
	No	Working	Answer	Mark	Notes
12	(a)		Reason	1	B1 makes a mention of bias, either directly, or making reference to an insufficient or biased range of responses.
	(b)(i)		Reason	1	B1 make some mention of any one of (a) an insufficient range of responses (b) no mention of money (c) no time frame in the question (d) misunderstanding of "a lot" or "not much"
	(ii)		Question	2	B2 B1 for each of the following upto B2 (a) an improved question eg time frame made clear (b) response boxes (imperfect) (c) response boxes no errors OR For suggesting a generally improved question (a) a question clearly in the context of changes to the canteen (b) at least 3 boxes showing a full range of responses
13	(a)	$20y - 18y = 16 - 9$ oe $2y = 7$	$3\frac{1}{2}$ oe	3	M1 $20y - 18y = 16 - 9$ oe M1 $2y = 7$ A1 cao
	(b)	$40 - x = 3(4 + x)$ $40 - x = 12 + 3x$ $40 - 12 = x + 3x$ $4x = 28$	7	3	M1 multiplying through by 3: $3 \times \frac{40 - x}{3} = 3 \times 4 + 3 \times x$ A1 $40 - 12 = x + 3x$ A1 cao
14	(a)		$12(x\text{-}4)$	1	B1 for $12(x - 4)$ or $12x - 48$ or $12 \times x - 48$ oe $x = 12(x - 4)$ gets B0
	(b)		$18x\text{-}48$	2	M1 $6x +$ "$12(x - 4)$" or $6x +$ "$12x - 48$" A1 ft a linear expression $18x - 48$, $2(9x - 24)$, $3(6x - 16)$, $6(3x - 8)$

PAPER 5504					
No		**Working**	**Answer**	**Mark**	**Notes**
15		x $x^3 - 2x$ 4.1 60.7(21) 4.2 65.6(88) 4.3 70.9(07) Accept answer to 1 dp 4.22 66.7(114) or more 4.23 67.2(269) 4.24 67.7(45) 4.25 68.2(656)	4.2	4	B2 for trial $4.1 \leq x \leq 4.3$ evaluated (B1 for trial $4 < x < 5$ evaluated) B1 for different trial $4.225 \leq x \leq 4.25$ evaluated B1 (dep on at least one previous B1) for 4.2 cao
16	(a)		1.0×10^{-9}	1	B1 accept 1×10^{-9} or just 10^{-9}
	(b)	$1 \div 5 \times 10^{-9}$	2×10^{8}	2	M1 for $1 \div$ ("5×10^{-9}") or digit 2 with zeros only seen Condone omission of bracket for M1. A1 cao
17	(a)	$\frac{6.27 \times 4.52}{4.81 + 9.63} = \frac{28.3404}{14.44} = 1.962631579$	1.9626…	2	B2 for 1.9626… (B1 for 28.34 or 14.44)
	(b)		1.96	1	B1 ft from (a) as rounded to 1dp or 2dp. Do not accept 2, 2.00, but accept 2.0
18			$d = 4n + 6$	2	B2 $d = 4n + 6$ oe (B1 $d = 4n + k$, k an integer or 0, $4n + 6$, $n = 4n + 6$)
19		Vol = $\pi \times 3.8^2 \times 2.5 = \pi \times 14.44 \times 2.5$ = 45.36… × 2.5 = 113.411 Mass = "113" × 1.5 = 170.1165	170	4	M1 for $\pi \times r^2 \times 2.5$ where r = is 3.8 or 7.6 A1 if $r = 3.8$ M1 for "113" × 1.5 A1 for 169.5 – 170.3 cao

PAPER 5504					
	No	Working	Answer	Mark	Notes
20	(a)	$10^2 + 6^2$ or 136 $\sqrt{(100+36)}$ or $\sqrt{136} = 11.66\ldots$	11.7	3	M1 $10^2 + 6^2$ or 136 seen M1 (dep) $\sqrt{100+36}$ or $\sqrt{136}$ A1 11.66 – 11.7
	(b)	$\cos x = \frac{8}{10}$ or 0.8 $x = \cos^{-1} 0.8 = 36.869^\circ$	36.9	3	M1 for $\cos = \frac{8}{10}$, cos = 0.8 (oe) M1 (dep) for $\cos^{-1}$ (oe) A1 for 36.86 – 36.9
21		$18x - 6y = 99$ $\quad 12x - 4y = 66$ $8x + 6y = 18$ $\quad 12x + 9y = 27$ $26x = 117$ $\quad -13y = 39$	$x = 4\frac{1}{2}$ $y = -3$	4	M1 correct process to eliminate either x or y (condone one error) A1 cao for non – eliminated one. M1 (dep on 1st M1) for correct substitution of their found value. A1 cao (need both)
22		$\frac{133}{72} = 1.8472, \frac{160}{82} = 1.9512$ OR $\frac{72}{133} = 0.54135, \frac{82}{160} = 0.5125$ OR $\frac{160}{133} = 1.203...., \frac{82}{72} = 1.1388....$ OR $\frac{133}{160} = 0.83125...., \frac{72}{82} = 0.878$	1.84..≠ 1.95.. 1.20..≠ 1.13..	3	M1 for $\frac{133}{72}$ (= 1.8472…) oe Accept 1.8, 1.85 M1 for $\frac{160}{82}$ (=1.9512…) oe consistent pairing Accept 2.0, 1.9 **OR** M1 for $\frac{160}{133}$ (= 1.203…) oe M1 for $\frac{82}{72}$ (=1.1388) oe A1 for enough decimal places to show that the ratios are not equal; since the scale factors are different the shapes cannot be similar. NB Do Not need conclusion

PAPER 5504					
	No	Working	Answer	Mark	Notes
23	(a)		0.4, 0.6 0.4, 0.6, 0.4	2	B1 for LHS: (0.6), 0.4 B1 for RHS: 0.6, 0.4, 0.6, 0.4
	(b)	(30 × 42) – (25 × 42.8) = 1260 – 1070 = 190 190 ÷ 5 =	38	3	M1 for (30 × 42) – (25 × 42.8) or 1260 – 1070 or 190 seen M1 (dep) for "190" ÷ 5 A1 cao
24	(a)	£12000 × 0.25 = £3000; £12000 – £3000 = £9000 £9000 × 0.25 = £2250; £9000 – £2250 = £6750 £6750 × 0.25 = £1687.50; £6750 – £1687.50 =	£5062.50	3	M1 for 12000 × 0.75 (=9000) oe or £3000 or £23437.50 seen M1 (dep) for at least two further depreciation calculations (complete steps) A1 cao **OR** M2 for $12000 \times (0.75)^3$ or 5062.50 seen (M1 for $12000 \times (0.75)^n$, $n = 2$ or 4)
	(b)	0.8 × 0.8 × 0.8 × 0.8 (oe)	0.4096	2	M1 0.8^4 (oe) A1 cao

PAPER 5505					
No		**Working**	**Answer**	**Mark**	**Notes**
1	(a)(i) (ii) (b)	 LCM = 3 × 13 × 17 = 3 × 221	2.21 0.013 663	2 2	B1 for 2.21 B1 for 0.013 M1 for 3 × 13 × 17 oe A1 for 663
2			Tick boxes 1,3 & 6	3	B1 for $\frac{\pi abc}{2d}$; B1 for $2a^2$; B1 for $2(c^2 + d^2)$ (−B1 for each additional expression ticked (>3) to a minimum of 0)
3	(a) (b)	200 × 0.2 0.2 + 0.4	40 0.6	2 2	M1 for 200 × 0.2 or $\frac{40}{200}$ seen A1 for 40 M1 for 0.2 + 0.4 A1 for 0.6
4	(a) (b)	eg 2 | 108 2 | 54 3 | 27 3 | 9 3 12	$2^2 \times 3^3$ 12	3 1	M2 for full systematic method of at least 4 divisions by prime numbers oe factor trees; condone 1 calculation error (M1 for 108 written as a correct product with only one non-prime or equivalent division or a full process with 2 calculation errors.) A1 for $2^2 \times 3^3$ (accept 2 × 2 × 3 × 3 × 3) B1 for 12
5		Perpendicular from P to intersecting arcs (within tramlines); perpendicular at least 2 cm long		2	M1 relevant pair of arcs crossing within tramlines A1 SC M1A0 for full construction of a line perpendicular to AB not through P
6		15 × 10	150 cm^3.	3	M1 for 15 × 10 A1 for 150 B1 for cm^3

PAPER 5505					
	No	Working	Answer	Mark	Notes
7	(a)		k^3.	1	B1 for k^3.
	(b)(i)	$4x + 20 + 3x - 21$	$7x - 1$	4	M1 for **three** of 4 terms $4x + 20 + 3x - 21$ (or better) A1 for $7x - 1$
	(ii)	$x^2 + 3yx + 2xy + 6y^2$	$x^2 + 5xy + 6y^2$		M1 for **three** of 4 terms $x^2 + 3yx + 2xy + 6y^2$ A1 for $x^2 + 5xy + 6y^2$
	(c)		$(p+q)(p+q+5)$	1	B1 for $(p+q)(p+q+5)$
	(d)		m^8	1	B1 for m^8.
	(e)		$6r^3t^6$	2	B2 for $6r^3t^6$ (B1 for $.....r^3t^6$ or for $6....t^6$)
8	(i)	Least length = 100.5	100.5	2	B1 for 100.5
	(ii)	Greatest length = 101.5	101.5		B1 for 101.5; ACCEPT 101.499 or better
9		$6\frac{2}{5} = \frac{32}{5}$ Area of triangle = $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ (= 2) Length of a side of sq. = $\sqrt{18 \times "2"}$ (=6) Perimeter of square = 4 × 6	24	5	B1 for $6\frac{2}{5} = \frac{32}{5}$ oe or $3\frac{1}{5} = \frac{16}{5}$ oe or $\frac{30}{8} + \frac{2}{8}$ oe (or implied by area of triangle=2) M1 for $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ oe M1 for (area of square) = 18 × product of two lengths A1 = $\sqrt{18 \times "2"}$ A1 for 24

PAPER 5505					
	No	Working	Answer	Mark	Notes
10	(a)	$6y = 15 - 5x$	$y = \frac{15-5x}{6}$	2	M1 for either $6y = 15 - 5x$ or for $\frac{5x}{6} + y = \frac{15}{6}$ or $-6y = 5x - 15$ or a correct ft on sign error to $y =$ A1 for $y = \frac{15-5x}{6}$ oe
	(b)	$6k + 5(-21) = 15$	20	2	M1 for subst. of $x = -21$ (or $x = 21$) into given eq^n or answer to (a) A1 for $k = 20$
	(c)(i)		Region R indicated	3	B2 correct region shaded (accept unshaded if R clear) (B1 shaded (R) region satisfies 3 of the 4 given inequalities with same boundaries)
	(ii)		(1,1)		B1 for (1,1)
11	(a)		$y = 2x + 6$	2	B2 for $y = 2x + 6$ (B1 for $y = 2x + k$, $k \neq 1$ or for $y = mx + 6$, $m \neq 0$ or for $2x+6$)
	(b)	Grad of $AB = 2$; Grad of $BC = \frac{-1}{"2"}$	$y = -\frac{1}{2}x + 6$	2	M1 for Grad of $BC = \frac{-1}{"2"}, -\frac{1}{2}$ or grad of $BC = -\frac{1}{\text{grad of AB}}$ A1 ft for $y = -\frac{1}{2}x + 6$ oe ft on wrong coeff. of x in (a)
	(c)	Eg A rectangle is always a cyclic quadrilateral because the opposite angles of a rectangle always add up to 180° (90+90)		1	B1 for valid explanation. (eg lines from the pt of int. of diagonals of rect to all 4 vertices are equal(radii))
12		Vertices of new triangle at $(-1, -4)$ $(-4, 2)$ and $(2, 2)$	Correct triangle drawn	3	B3 cao (B2 for 3 correct vertices no triangle or triangle with 2 correct vertices (B1 for triangle with two of six co-ordinates correct from using P as centre or any isosceles triangle with base 6 and height 6)

PAPER 5505					
	No	Working	Answer	Mark	Notes
13	(a)		32	1	B1 for 32 (accept 31.5 to 33.5 inclusive)
	(b)			3	B1 for ends of whiskers at 9 and 57 (with a box) B1 for ends of box at 16 and 45/46 ($\pm$ 0.5) B1for median marked at "32"or complete box and whisker diagram drawn with a median inside the box
	(c)	Median(B)>Median(G); on average boys take longer IQR(B)>IQR(G); times for boys have a greater spread		2	B1 eg for comparison of medians (ft on diagrams) B1 eg for comparison of (interquartile) ranges (ft on diagram)
14		Reading top to bottom frequencies are 20; 18; 45; 52		2	B2 all correct (B1 for one frequency correct)
15	(a)	$1600 = p\,q^0$.; $400 = pq^2$ $q^2 = 0.25$	$p = 1600$ $q = 0.5$	3	M1 for either $400 = pq^2$ or $1600 = pq^0$ B1 for $p = 1600$ A1 for $q = 0.5$
	(b)	$V = pq^{-2}$	6400	2	M1 for recognition that t = –2 A1 for 6 400

PAPER 5505					
	No	Working	Answer	Mark	Notes
16	(a)	(I) $AB = AC$ (triangle ABC is isosceles) (II) PB = PC tangents (from a point to a circle are) equal (III) AP = AP (common side) so the 2 triangles are congruent ,SSS.		3	B3 for I, II, III with congruency reason (B2 for any two of I, II or III) (B1 for any one of the I, II or III)
	(b)	$BPC = 20°$ PBC (or PCB) = 90-1/2 "20" (= 80°) $BAC = PBC$ = "80"	50°	4	B4 for 50° (B3 for $BAC = 80°$) (B2 for $PBC = 80°$ or $PCB = 80°$) (B1 for $APC = 10°$ or $BPC = 20°$ or a middle angle = 90°) SC if clear numerical slip seen eg "PBC = 180 – 90 – 10 = 70" then goes on to get correct ft angle $ABC = 55$ deduct 1 from total so this cand would get B4–1 = B3
17	(a)	$\overrightarrow{OP} = \overrightarrow{OA} + \overrightarrow{AP}$ $= \overrightarrow{OA} + \frac{2}{3}(6\mathbf{c} - 6\mathbf{a})$ $= 6\mathbf{a} + 4\mathbf{c} - 4\mathbf{a}$	$2\mathbf{a} + 4\mathbf{c}$	3	M1 for $\overrightarrow{OP} = \overrightarrow{OA} + \overrightarrow{AP}$ or any correct vector journey involving $\overrightarrow{OP}$ M1 for $\overrightarrow{AP} = \frac{2}{3}(6\mathbf{c} - 6\mathbf{a})$ oe or $\overrightarrow{CP} = \frac{1}{3}(-6\mathbf{c} + 6\mathbf{a})$ oe or reverse vectors A1 for $2\mathbf{a} + 4\mathbf{c}$ oe (accept unsimplified)
	(b)	Eg $\overrightarrow{OM} = \overrightarrow{OC} + \overrightarrow{CM} = 6\mathbf{c} + 3\mathbf{a}$ $\overrightarrow{OM} = 1.5\overrightarrow{OP}$	$\overrightarrow{OM} = 1.5\overrightarrow{OP}$ so OPM is a straight line	2	B1 for $\overrightarrow{OM} = 6\mathbf{c} + \frac{1}{2}(6\mathbf{a})$ or $\overrightarrow{PM} = 2\mathbf{c} + \mathbf{a}$ unsimplified or reverse vectors B1 for a fully correct proof.

PAPER 5505					
	No	Working	Answer	Mark	Notes
18	(a)		4	1	B1 for 4 condone $\pm$ 4
	(b)		2	1	B1 for 2 condone ± 2
	(c)	$\sqrt{160} = 4\sqrt{10}$; $\left[\frac{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)-\sqrt{2}\times\sqrt{5}}{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)}\right]\times 100$ $\left[\frac{6\sqrt{10}-\sqrt{10}}{6\sqrt{10}}\right]\times 100$	$\frac{500}{6}$	4	B1 for **either** $\sqrt{160} = 4\sqrt{10}$ or $\sqrt{8} = 2\sqrt{2}$ **or** $\sqrt{20} = 2\sqrt{5}$ M1 for $\left[\frac{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)-\sqrt{2}\times\sqrt{5}}{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)}\right]$ oe $(\times 100)$ B1 for either $6\sqrt{10} - \sqrt{10}$ or $6\sqrt{10}$ A1 for $\frac{500}{6}$ (accept 83.3 if no obvious earlier error)

PAPER 5505					
	No	Working	Answer	Mark	Notes
19	(a)(i)		$(2x-7)(x-14)$	3	M1 x^2 term and constant term (± 98) obtained or $2x(x-14) - 7(x-14)$ or $x(2x-7) - 14(2x-7)$ A1 for $(2x-7)(x-14)$
	(ii)		$x = \frac{7}{2}$; $x = 14$		B1ft ft (i) provided of form $(2x\pm a)(x\pm b)$
	(b)(i)		$\frac{7}{n+7}$	3	B1 for $\frac{7}{n+7}$ oe
	(ii)	$\frac{7}{n+7} = \frac{2}{5} \Rightarrow 2(n+7) = 5 \times 7$ $2n = 21$	n=10.5 is not possible since n has to be an integer		M1 for $2(n+7) = 5\times 7$ or $n+7=5\times 3.5$ (can be implied) ft (b)(i) fractional in terms of n and < 1 A1 ft for n=10.5 not possible (since n not integer) oe
	(c)	$2\times\left(\frac{n}{n+7}\right)\times\left(\frac{7}{n+7}\right) = \frac{4}{9}$ $14n\times 9 = 4(n+7)^2$ $14n\times 9 = 4(n^2+14n+49)$ $4n^2+56n+196-126n = 0$	$2n^2 - 35n + 98 = 0$	5	M1 for $\left(\frac{n}{n+7}\right)\times\left(\frac{7}{n+7}\right)$ seen M1 for $2\times\left(\frac{n}{n+7}\right)\times\left(\frac{7}{n+7}\right)$ oe $(=\frac{4}{9})$ M1(dep on 1st M) elimination of fractions within an equation B1 3 terms correct in expansion of $(n+7)^2 = n^2 + 7n + 7n + 49$ A1 full valid completion to printed answer

PAPER 5505					
	No	Working	Answer	Mark	Notes
	(d)	$\frac{7}{n+7}\times\frac{7}{n+7}=\frac{7}{21}\times\frac{7}{21}=$	$\frac{1}{9}$	2	M1 for $\frac{7}{n+7}\times\frac{7}{n+7}$ or better or ft [answer (b)(i)]^2 or $1-\frac{4}{9}-(\frac{n}{n+7})^2$ A1 for $\frac{1}{9}$ oe cao
20	(a)(i)		$y=1+\sin x$	2	B1 for $y=1+\sin x$
	(ii)		$y=2\sin x$		B1 for $y=2\sin x$ SC both (i) f(x) + 1, (ii) 2f(x) B1
	(b)	Stretch parallel to y-axis scale factor 3 Stretch parallel to x-axis scale factor $\frac{1}{2}$		3	M1 for 'stretch' A1 for Stretch parallel to y-axis scale factor 3 oe A1 for Stretch parallel to x-axis scale factor $\frac{1}{2}$ oe SC if M0 award BI for "sf 3 vertically" **and** "sf $\frac{1}{2}$ horizon."

PAPER 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
1		Question	2	B1 for each of the following upto B2 (a) an improved question eg time frame made clear (b) response boxes (imperfect) (c) response boxes no errors OR For suggesting a generally improved question (a) a question clearly in the context of changes to the canteen (b) at least 3 boxes showing a full range of responses
2	Locus (circle) drawn & shaded		2	B1 circle centre Manchester ±2mm B1 shading with accurate or approximate circle within tolerance
3 (a)		182, 178, 180, 184	2	M1 mean of any three consecutive months, eg (147 + 161 + 238) ÷ 3 oe A1 cao
(b)	Sale price = 80% Fun Friday price = 70% of 80% = 56% (oe)		2	B1 B1 for a fully correct explanation involving a worked example (oe) B1 a partially complete explanation
4	x $x^3 - 2x$ 4.1 60.7(21) 4.2 65.6(88) 4.3 70.9(07) 4.22 66.7(114) 4.23 67.2(269) 4.24 67.7(45) 4.25 68.2(656)	4.2	4	B2 for trial $4.1 \leq x \leq 4.3$ evaluated (B1 for trial $4 < x < 5$ evaluated) B1 for different trial $4.225 \leq x \leq 4.25$ evaluated B1 (dep on at least one previous B1) for 4.2 cao

PAPER 5506					
	No	Working	Answer	Mark	Notes
5	(a)		1.0×10^{-9}	1	B1 accept 1×10^{-9} or just 10^{-9}
	(b)	$1 \div (5 \times 10^{-9})$	2×10^{8}	2	M1 for $1 \div$ ("5×10^{-9}") or digit 2 with zeros only seen Condone omission of bracket for M1. A1 cao
6	(a)	$\frac{6.27 \times 4.52}{4.81 + 9.63} = \frac{28.3404}{14.44} = 1.962631579$	1.9626…	2	B2 for 1.9626… (B1 for 28.34… or 14.44)
	(b)		1.96	1	B1 ft from (a) as rounded to 1dp or 2dp. Do not accept 2, 2.00, but accept 2.0
7			$d = 4n + 6$	2	B2 $d = 4n + 6$ oe (B1 $d = 4n + k$, k an integer $\neq 6$, $4n + 6$, $n = 4n + 6$)
8		Vol = $\pi \times 3.8^2 \times 2.5 = \pi \times 14.44 \times 2.5$ $= 45.36\ldots \times 2.5 = 113.411$ Mass = "113" × 1.5 = 170.1165	170	4	M1 for $\pi \times r^2 \times 2.5$ where r = is 3.8 or 7.6 A1 if $r = 3.8$ M1 for "113" × 1.5 A1 for 169.5 – 170.3 cao

PAPER 5506					
	No	Working	Answer	Mark	Notes
9	(a)	$10^2 + 6^2$ or 136 $\sqrt{(100 + 36)}$ or $\sqrt{136} = 11.66\ldots$	11.7	3	M1 for $10^2 + 6^2$ or 136 seen M1 (dep) $\sqrt{100+36}$ or $\sqrt{136}$ A1 11.66 – 11.7
	(b)	Cos x = 8/10 or 0.8 $x = \cos^{-1} 0.8 = 36.869^o$	36.9	3	M1 for cos = $\frac{8}{10}$, cos = 0.8(oe) M1 (dep) for $\cos^{-1}$ (oe) A1 for 36.86 – 36.9 M1 Use of sine rule and x found M1 for $x = 90 - \sin^{-1}$("0.8") A1 for 36.86 – 36.9
10		$18x - 6y = 99$ $\quad 12x - 4y = 66$ $8x + 6y = 18$ $\quad 12x + 9y = 27$ $26x = 117$ $\quad -13y = 39$	$x = 4\ ½$ $y = –3$	4	M1 correct process to eliminate either x or y (condone one error) A1 cao for non – eliminated one. M1 (dep on 1st M1) for correct substitution of their found value. A1 cao (need both)

PAPER 5506					
	No	Working	Answer	Mark	Notes
11		$\frac{133}{72} = 1.8472\,,\ \frac{160}{82} = 1.9512$ OR $\frac{72}{133} = 0.54135, \frac{82}{160} = 0.5125$ OR $\frac{160}{133} = 1.203...., \frac{82}{72} = 1.1388....$ OR $\frac{133}{160} = 0.83125...., \frac{72}{82} = 0.878$	1.84..≠ 1.95.. 1.20..≠ 1.13..	3	M1 for $\frac{133}{72}$ (= 1.8472…) oe Accept 1.8, 1.85 M1 for $\frac{160}{82}$ (=1.9512…) oe consistent pairing Accept 2.0, 1.9 **OR** M1 for $\frac{160}{133}$ (= 1.203…) oe M1 for $\frac{82}{72}$ (=1.1388) oe A1 for enough decimal places to show that the ratios are not equal; since the scale factors are different the shapes cannot be similar. NB Do Not need conclusion
12	(a)	£12000 × 0.25 = £3000; £12000 – £3000 = £9000 £9000 × 0.25 = £2250; £9000 – £2250 = £6750 £6750 × 0.25 = £1687.50; £6750 – £1687.50 =	£5062.50	3	M1 for 12000 × 0.75 (=9000) oe or £3000 or £23437.50 seen M1 (dep) for at least two further depreciation calculations (complete steps) A1 cao **OR** M2 for 12000 × $(0.75)^3$ or 5062.50 seen (M1 for 12000 × $(0.75)^n$, $n = 2$ or 4)
	(b)	0.8 × 0.8 × 0.8 × 0.8 (oe)	0.4096	2	M1 0.8^4 (oe) A1 cao

PAPER 5506					
No		**Working**	**Answer**	**Mark**	**Notes**
13		$r^2 = \frac{3\times10}{\pi h} = \frac{3\times10}{\pi 1.5} = \frac{3\times10}{4.712..} = 6.36, 6.37$ $r = \sqrt{6.366\ldots}$	2.52-2.54	3	M1 for correct rearrangement to give $r^2 = \frac{3V}{\pi h}$ or $30 \div 4.712..$ or $6.36 - 6.37$ *Allow 0.3, 0.33 for* $\frac{1}{3}$ M1 (dep) for $\sqrt{}$ A1 cao 2.52 – 2.54
14	(a)		0.4 0.6,0.4, 0.6,0.4	2	B1 for LHS: (0.6), 0.4 B1 for RHS: 0.6, 0.4, 0.6, 0.4
	(b)	0.6×0.6	0.36	2	M1 $0.6 \times$ "0.6" [0 < "0.6" < 1] A1 cao
	(c)	$(30 \times 42) - (25 \times 42.8) = (1260 - 1070 = 190$ $190 \div 5 =$	38	3	M1 for $(30 \times 42) - (25 \times 42.8)$ or $1260 - 1070$ or 190 seen M1(dep) for "190" ÷ 5 A1 cao 38

PAPER 5506					
	No	Working	Answer	Mark	Notes
15	(a)	$S = \frac{k}{f^2};\ 125 = \frac{k}{8^2};\ k = 8000$	$S = \frac{8000}{f^2}$	3	M1 for $S = \frac{k}{f^2}$ M1 for $125 = \frac{k}{8^2}$ A1 cao These marks can be awarded if the full formula appears in part (b), rather than in part (a). [SC: $S = \frac{125}{64} f^2$ M1 M0 A0, $S = 1.95(3125) f^2$ M1 M0 A0 $S = \frac{1000}{f}$ M1 M0 A0]
	(b)		500	1	B1 cao

PAPER 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
16	$0.5 \times 3.2 \times 8.4 \times \text{Sin } B = 10$ $\sin B = 0.74404\ldots$ 48.077 $AC^2 = 3.2^2 + 8.4^2 - 2 \times 3.2 \times 8.4 \times \cos B$ $AC^2 = 44.8815\ldots..$ $AC = 6.69\ (936\ldots)$ Perimeter = 18.3 Use the altitude AD, $\frac{h \times 8.4}{2} = 10 \Rightarrow h = (2.381)$ $BD = \sqrt{3.2^2 - h^2} = 2.139$ $DC = 6.261$ $AC = \sqrt{2.38^2 + '6.261'^2} = 6.69(936)$ Perimeter = 18.3	18.3	6	M1 for $0.5 \times 3.2 \times 8.4 \times \sin B$ (= 10) A1 for sin B = 0.74(404…) or B = 47.7 – 48.1 M1 for $3.2^2 + 8.4^2 - 2 \times 3.2 \times 8.4 \times \cos$ "48.077" M1(dep) for AC^2 = "44.8(815)"….. with correct order of evaluation A1 AC = 6.69 – 6.7 A1 18.29 – 18.3 **for** $\frac{h \times 8.4}{2} = 10 \Rightarrow h = (2.381)$ M1 for $BD^2 = 3.2^2 - "2.381"^2$ A1 BD = 2.1 – 2.2 M1 (dep) $AC^2 = "2.381"^2 + "6.261"^2$ A1 AC = 6.69 – 6.7 A1 18.29 – 18.3

PAPER 5506					
	No	Working	Answer	Mark	Notes
17	(a)	$(x + 8)^2 = x^2 + (x + 5)^2$ $x^2 + 16x + 64 = 2x^2 + 10x + 25$		4	B1 for angle $OTA = 90°$ (implied by use of Pythagoras with OA as hypotenuse) M1 $(x + 8)^2 = x^2 + (x + 5)^2$ oe M1 for correct squaring of x+8 or x+5 A1 for completion following correct working
	(b)	$(+6 + \sqrt{(6^2 - - 4\times39)})/2$ $= (6 \pm \sqrt{(36 + 156)})/2$ $= (6 \pm \sqrt{192})/2 = (6 + 13.856)/2$	9.93	3	M1 for substitution into quadratic formula, allow sign errors in b and c M1 for $x = (6 \pm \sqrt{192})/2$ [+ alone will do] A1 for 9.92 – 9.93 M1 for $(x-3)^2-9-39$ M1 for $x = 3 \pm \sqrt{48}$ [+ alone will do it] A1 for 9.92 – 9.93 T&I= 9.93 gets M1,M0,A0
18		$EB = 60 \times \tan 30°$ $BD = \sqrt{(60^2 + 60^2)}$ $\tan BDE = 34.64 \div 84.85$ $EB = 60 \times \tan 30°$ (= 34.64) $ED^2 = 60^2 + (\frac{60}{\cos 30})^2$ $ED = \sqrt{8400} = (91.65)$ $\text{Angle} = \sin^{-1}\left(\frac{EB}{\sqrt{8400}}\right) = 22.2$	22.2°	4	M1 for $EB = 60\times\tan30$ M1 for $BD = \sqrt{(60^2 + 60^2)}$ M1 for $\tan BDE$ = "34.64" ÷ "84.85" A1 22.17 – 22.21 M1 for $EB = 60 \times \tan 30°$ oe M1 for fully correct method for ED M1 for $\sin BDE = \left(\frac{'34.84'}{'\sqrt{8400}'}\right)$ (oe) A1 22.17 – 22.21

PAPER 5506					
	No	Working	Answer	Mark	Notes
19	(a)	$x^2 + 6^2 = 25$ $x^2 = -11$		2	M1 substitutes y=6 to get x^2+36 = 25 A1 deduces x^2=25-36<0; impossible, (so line does not intersect curve)
	(b)	$y^2 = (2x - 2)^2$ $x^2 + (2x - 2)^2 = 25$ $x^2 + 4x^2 - 8x + 4 = 25$ $5x^2 - 8x + 4 - 25 = 0$ $5x^2 - 8x - 21 = 0$ $(5x + 7)(x - 3) = 0$ or $x = \frac{--8 \pm \sqrt{(-8)^2 - 4 \times 5 \times (-21)}}{10} = \frac{8 \pm \sqrt{484}}{10}$ So $x = 3$, or $x = -1.4$ Subs into either equation gives: $y = 4$, or $y = -4.8$	$x = 3$ $y = 4$ $x = -1.4$ $y = -4.8$	6	M1 $y^2=(2x-2)^2$ seen or implied A1 $x^2 + 4x^2 - 8x + 4 = 25$ (oe expanded form) M1 correct attempt to solve 3 term quadratic equation A1 $x = 3$, $x = -1.4$ M1 (dep on previous Ms) Sub one value of x into either equation gives: A1 $y = 4$, $y = -4.8$ NB: incorrectly matched pairs loses the final A1.

PAPER 5506					
	No	Working	Answer	Mark	Notes
20	(a)	400.005 ÷ 59.95 = 6.672310 6.672310 ÷ 1000 × 3600 = 24.02032 399.995 ÷ 60.05 = 6.661032 6.661032 ÷ 1000 × 3600 = 23.97972	24.020 23.980	5	B1 for 400.005 or 59.95 seen oe M1 for "400.005" ÷ "59.95" where "400.005" ∈ [400.005,400.5] and "59.95" ∈ [59.5,59.95] oe B1 for 399.995 or 60.05 seen oe M1 for "399.995" ÷ "60.05" where "399.995" ∈ [399.5,399.95] and "60.05" ∈ [60.05,60.5] oe A1 23.979-23.980 and 24.020-24.0204
	(b)	24.0 because to 1dp the answers are the same		1	B1cao for 24.0 with reason
	(c)	177 × 50/477 = 18.553 → 19 → 18 111 × 50/477 = 11.635 → 12 86 × 50/477 = 9.0147 → 9 82 × 50/477 = 8.595 → 9 21 × 50/477 = 2.201 → 2	18 12 9 9 2	3	M1 for $\frac{50}{"477"} \times 177$ or 111 or 86 or 82 or 21 A1 for all integers or better answers, at least 3 correct A1 cao
21	(a)	$40 - x = 3(4 + x)$ $40 - x = 12 + 3x$ $40 - 12 = x + 3x$ $4x = 28$	7	3	M1 multiplying through by 3: $3 \times \frac{40 - x}{3} = 3 \times 4 + 3 \times x$ A1 $40 - 12 = x + 3x$ A1 cao
	(b)	$\frac{2x(2x - 3)}{(2x - 3)(2x + 3)} = \frac{2x}{2x + 3}$	$\frac{2x}{2x + 3}$	3	B1 for $(2x-3)(2x+3)$ B1 for $2x(2x-3)$ or $(2x + 0)(2x + 3)$ B1 cao

PAPER 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
22	Distance from x axis is y. Distance from (0, 2) is $\sqrt{(x^2+(y-2)^2)}$ $y^2 = x^2 + (y-2)^2$ $y^2 = x^2 + y^2 - 4y + 4$ $0 = x^2 - 4y + 4$ $4y = x^2 + 4$ and finish		4	B1 for $(x-0)^2 + (y-2)^2$ or $\sqrt{((x-0)^2+(y-2)^2)}$ oe seen B1 for $y = \sqrt{(x-0)^2+(y-2)^2}$ or $y^2 = (x-0)^2 + (y-2)^2$ oe B1 $(y-2)^2 = y^2 - 4y + 4$ seen B1 for $4y = x^2 + 4$ and finish

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone: 01623 467467
Fax: 01623 450481
Email: publications@linneydirect.com

Order Code UG015616 July 2004

For more information on Edexcel and BTEC qualifications please contact
Customer Services on 0870 240 9800
or http://enquiries.edexcel.org.uk
or visit our website: www.edexcel.org.uk

London Qualifications Limited, trading as Edexcel. Registered in England and Wales No. 4496750
Registered Office: Stewart House, 32 Russell Square, London WC1B 5DN



edexcel

Edexcel GCSE

Mathematics A 1387

June 2007

advancing learning, changing lives

Mark Scheme

Edexcel GCSE

Mathematics A 1387

Table Of Contents

Notes on Marking Principles	3 - 4
5521 / 01 Mark Scheme	5 - 11
5521 / 02 Mark Scheme	12 – 17
5523 / 03 Mark Scheme	18 – 27
5523 / 04 Mark Scheme	28 – 35
5525 / 05 Mark Scheme	36 – 43
5525 / 06 Mark Scheme	44 - 52

NOTES ON MARKING PRINCIPLES

1 **Types of mark**
M marks: method marks
A marks: accuracy marks
B marks: unconditional accuracy marks (independent of M marks)

2 Abbreviations
cao – correct answer only
ft – follow through
isw – ignore subsequent working
SC: special case
oe – or equivalent (and appropriate)
dep – dependent
indep - independent

3 No working
If no working is shown then correct answers normally score full marks
If no working is shown then incorrect (even though nearly correct) answers score no marks.

4 With working
If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.
If there is no answer on the answer line then check the working for an obvious answer.

5 Follow through marks
Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

6 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

7 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

8 Linear equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

9 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

Remember: if you are having difficulty making a decision on how you should mark a candidate response contact your Team Leader for advice, or send the item to review.

Paper 5521_01					
No		Working	Answer	Mark	Notes
1		Cat |||| /||| 8 Dog |||| | 6 Fish || 2 Hamster |||| 4	8, 6, 2, 4	3	M1 for attempt to tally or one frequency correct in either column A1 for 1 frequency correct or all tallies correct in correct column A1 for all frequencies correct (accept if /20)
2			27, 35, 42, 67, 118	1	B1 cao
3	(a)		Diameter drawn	1	B1 for a diameter
	(b)		Right angle marked	1	B1 *R* marked correctly
	(c)		Rectangle drawn	1	B1 for a rectangle
4	(a)		40	1	B1 cao
	(b)		50	1	B1 cao
	(c)		3 full loaves 4 full loaves + 1 half loaf	2	B1 for 3 full loaves B1 for 4 full loaves + 1 half loaf
5	(a)		7252	1	B1 cao
	(b)		Three thousand and eighty six	1	B1 accept 3 thousand and eighty six (condone 0 hundred)
	(c)		4600	1	B1 accept 4600
	(d)		200	1	B1 for 200 or 2 hundred or 100 or hundred
6	(i)		Cube	2	B1 for 'cube' (accept 'cuboid') ignore spelling
	(ii)		Cylinder		B1 for 'cylinder' ignore spelling
7		5×100	500	2	B2 for 490 or 500 or 510 (B1 for either 5 or 5.0 or 100 seen)

Paper 5521_01					
No		Working	Answer	Mark	Notes
8	(a)		8.4 cm or 84 mm	2	B2 allow ±2mm (B1 for 8.4 or 84, B1 for appropriate unit)
	(b)		37°	1	B1 allow ±2°
9	(a)		Carbon black	1	B1 accept 'black carbon' accept 26%
	(b)		0.1(0)	1	B1 cao
	(c)		0.04	1	B1 cao
	(d)	$\frac{26}{100}$	$\frac{13}{50}$	2	M1 for $\frac{26}{100}$ A1 cao
10	(a)		97	1	B1 cao
	(b)		London Reading	1	B1 cao
	(c)	41 + 57 + 58	156	3	M1 for two of 41, 57, 58 M1(dep) for '41' + '57' + '58' A1 cao
11	(a)		3	1	B1 cao allow ± 0.2
	(b)		–5	1	B1 cao allow ±0.2

Paper 5521_01					
No		**Working**	**Answer**	**Mark**	**Notes**
12	(a)		Centimetres (cm) miles litres (*l*)	3	B3 (B1 for each correct answer) accept abbreviations
	(b)		300	1	B1 cao
	(c)	'1500>1400' or '1.5>1.4'	Reason	1	B1 for No and '1500>1400' or '1.5>1.4'
13	(a)	Height of bars 12, 8, 6 lines drawn between points	Bars drawn	2	B2 for 3 bars correctly drawn (B1 for 2 bars correct)
	(b)		July and August	1	B1 oe
	(c)	24 – 4	20	1	B1 (Accept '4 to 24' oe)
	(d)	The temperatures are rising	Temp's rising oe	1	B1 for reason
14	(a) (i)		$(-4, 3)$	2	B1 cao
	(ii)		$(2, -1)$		B1 cao
	(b) (i)	*D* marked at $(-4, -1)$	Point marked on grid	2	B1 for point marked at $(-4, -1)$ cao
	(ii)		$(-4, -1)$		B1 ft
15	(a)			1	B1 cao
	(b)			1	B1 cao

Paper 5521_01				
No	**Working**	**Answer**	**Mark**	**Notes**
16 (a)		(Pat +) reason	1	B1 correct comment (Pat may be implied)
(b)	21 ÷ 3	7	1	B1 cao
17 (i)		S marked at 1	3	B1 for S within ½ cm of 1
(ii)		P marked at 0		B1 for P marked at 0 cao
(iii)		Q marked at "1/3"		B1 for Q marked at 1/3 ± 1 cm use overlay
18	6 + 6 + 3 or $2\frac{1}{2}\times 6$	15	3	M1 for realizing 6 glasses in one bottle M1 for realizing 3 glasses in ½ a bottle A1 cao (M2 for attempt to find $2\frac{1}{2}\times 6$) oe
19	15 and 16 parts shaded Alternative 1 $\frac{3}{4}=0.75,\ \frac{4}{5}=0.8$ Alternative 2 $\frac{3}{4}=\frac{15}{20},\ \frac{4}{5}=\frac{16}{20}$	$\frac{4}{5}$ + reason	3	M1 for correctly shading 15 parts for 3/4 M1 for correctly shading 16 parts for 4/5 A1 (dependent on M2) for selection of 4/5 Alternative 1 M1 for $\frac{3}{4}=0.75$ M1 for $\frac{4}{5}=0.8$ A1 (dep on M2) for selection of 0.8 or $\frac{4}{5}$ or $\frac{16}{20}$ Alternative 2 M1 for $\frac{3}{4}=\frac{15}{20}$ M1 for $\frac{4}{5}=\frac{16}{20}$ A1(dep on M2) for selection of $\frac{4}{5}$ or $\frac{16}{20}$

Paper 5521_01					
No		**Working**	**Answer**	**Mark**	**Notes**
20	(a)		−5 (−3) (−1) 1 3 5	2	B2 cao (B1 for any 2 or 3 correct)
	(b)	Points plotted	Line	2	B2 for line from (−1, −5) to (4, 5) (B1 ft for plotting at least 5 "points")
21		Shapes shaded on grid	6 tessellating shapes	2	B2 for fully correct with 5 or more additional shapes, no gaps (B1 for 4 shapes tessellating with at least one shape inverted, with or without the given shape ignore extras)
22	(a)		$\frac{7}{20}$	1	B1 for $\frac{7}{20}$ oe
	(b)	7 + 2 (or 20 – 11) are not lime flavour	$\frac{9}{20}$	1	B1 for $\frac{9}{20}$
	(c)		0	1	B1 for 0, zero or nought ($\frac{0}{20}$ gets B0)
23	(a)		$4a$	1	B1 accept $4\times a$, $a \times 4$, $a4$
	(b)		$12b$	1	B1 accept $12\times b$, $b \times 12$, $b12$
	(c)		$2a + 6b$	2	B2 cao (B1 for $2a$ or $6b$ seen)
	(d)		$x(x-6)$	2	B2 cao (B1 for $x(ax+b)$ where a, b are numbers not equal to 0 or x – 6 seen on its own, or as part of an expression)
24	(a)		40	1	B1 cao
	(b)		45	1	B1 for 42 – 48 accept ¾ hour
	(c)	40×2 or $\frac{40}{30}\times 60$ or $40\div\frac{1}{2}$	80	2	M1 for 40×2 or $\frac{40}{30}$ or $40\div\frac{1}{2}$ A1 cao NB $\frac{40}{45}\times 60$ gets M0 A0

Paper 5521_01					
No		**Working**	**Answer**	**Mark**	**Notes**
25	(a)		4560	1	B1 cao
	(b)		45.6	1	B1 cao
	(c)		2.4	1	B1 cao
26		$-7 - 3 = -10$ $2 \times -10 = -20$ $-20 \div 4$	-5	3	M1 for substitution of 2 and –7 into $p(q - 3)$ or sight of -20 or $-14 - 6$ M1dep for '–20' ÷ 4 A1 cao B1 SC for sight of –10 if M0 awarded
27	(a)		Reflection in *y*-axis	1	B1 cao
	(b)		Rotation by half turn about (0,0)	2	B2 cao (B1 for half turn not about (0,0).)
	(c)		Enlargement Scale factor 3 Centre (0,0)	3	B1 for 'enlargement' B1 for "scale factor 3" or 3 seen B1 for 'centre (0,0)
28	(a)		Reason	1	B1 for 'The frequencies are nearly equal' oe
	(b)	$1\times26+2\times26+3\times23+4\times25 = 247$ $247/100$	2.47	3	M1 for *fx* (attempting at least 2 relevant products) M1 for $\sum fx \div 100$ A1 2.47 cao

Paper 5521_01				
No	**Working**	**Answer**	**Mark**	**Notes**
29	5 × 5 × 6	150	4	M1 for attempt at 1 division (e.g. 40 ÷ 8), may be implied by marks or number on one edge of diagram or by 5 or 6 seen M1 for attempt at 3 divisions (40÷8, 40÷8, 60÷10), may be implied by marks or numbers on diagram or by 5,5 and 6 seen. M1 (dep on 1st M1) for "5" × "5" × "6" A1 cao Alternatively M1 for 40 × 40 × 60 **or** 8 × 8 × 10 **or** 96000 or 640 seen M1 for 40 × 40 × 60 **and** 8 × 8 × 10 or 96000 **and** 640 seen M1 (dep on 1st M1) for "(40 × 40 × 60)" ÷ "(8 × 8 × 10)" A1 cao SC:B1 for dividing area of one carton face by area of corresponding box face if M0

Paper 5521_02					
No		Working	Answer	Mark	Notes
1	(a)	Draw diagram.	Diagram	1	B1 cao
	(b)		13,16	1	B1 cao
	(c)		31	1	B1 cao
2	(a)		8	1	B1 cao
	(b)		14	2	B2 for 14 (B1 for 13 or 15)
	(c)		16	2	B2 for 16 (B1 for 15, 17 or 8)
3	(a)		4,7 drawn	2	B2 for car height 4 and bus height 7, (B1 for one correct)
	(b)		6	1	B1 cao
	(c)		Walk	1	B1
	(d)		27	1	B1 cao
4	(a)		40	1	B1 for 40–41 inclusive
	(b)		12	1	B1 for 11.5 – 12.5 inclusive
5	(a)		Row complete	2	B2 for 1+..+11; 36 (B1 for one of the 2 cells complete)
	(b)		Square	1	B1 "square"
6		One line of symmetry		1	B1 within 2mm of centre of base / 2mm of vertex

Paper 5521_02				
No	**Working**	**Answer**	**Mark**	**Notes**
7 (a) (i)		09 06	2	B1 (accept 9 06 oe)
(ii)		39		B1 cao
(b)		06 55	1	B1 (accept 6 55 oe)
(c)		2h 6min	1	B1 cao
(d)		15 min	2	M1 for 0906 – 0645 – "(c)" or 0906 – 0645 – 2hr 6min or 2hr 21min – "(c)" or 2hr 21min – 2hr 6 min or 141 – 126 or 20 – 5 A1 cao SC: B1 for 55 or 75 or 93 seen
8 (i)		8,10,12,20 or 30	3	B1 at least one of 8, 10, 12, 20, 30 (no extras)
(ii)		8,12 or 20		B1 at least one of 8,12, 20 (no extras)
(iii)		3 or 5		B1 3 or 5 or both (no extras)
9 (a)		C or G	1	B1 at least one of C or G (no extras)
(b)		A and F	1	B1 cao
(c)		2	1	B1 (accept –2)
10 (a)		$\frac{7}{10}$	1	B1 7/10 oe
(b)(i)		4 squares	2	B1 4 squares shaded
(ii)		80%		B1 80% or ft from unshaded part (no ft from 0% or 100%)
(c)(i)		2.5	2	B1 2.4–2.6 inclusive
(ii)		1.7		B1 1.6–1.8 inclusive
11 (a)		2	1	B1 for 2 or –2
(b)		14	1	B1 for 14 or –14

Paper 5521_02					
No		Working	Answer	Mark	Notes
12		2 × 8.50 = 17.00 3 × 4.50 = 13.50 Total = 30.50 50.00 – 30.50	19.5(0)(*p*)	3	M1 for adding 5 correct values or 2 × 8.50 + 3 × 4.50 (ignore units) or 30.5(0) or 3050 seen M1 dep for 50 − "30.50" (ignore units) (OR M1for adding at least 1 adult ticket and at least 1 child ticket **and** subtracting from 50) A1 cao SC: B1 for 24 or 37 or 2400 or 3700 seen
13	(a)		Hexagon	1	B1
	(b)(i)		120	2	B1 cao
	(ii)		Str line		B1 reference to a (straight) line and 180°
	(c)		Obtuse	1	B1 Accept "interior"
14	(a)	2, 2, 3, 3, 3, 4, 4, 4, 5, 6	3.5	2	M1 ordering the numbers (condone 1 error or omission) A1 cao
	(b)	36 ÷ 10	3.6	2	M1 sum of numbers ÷ 10 A1 cao SC B1 for 3r 6
	(c)	6 – 2	4	1	B1 cao
15	(a)		Paul	1	B1 cao
	(b)	36 ÷ 2 oe	18	1	B1 cao
	(c)	60/360 =	1/6	2	M1 60/360 oe A1 cao

Paper 5521_02					
No		Working	Answer	Mark	Notes
16		4.7 ÷ 5.9 = 0.796610169	0.7966..	2	B2 for 0.7966 or better (B1 for 0.8, 0.80, 0.79, 0.796, 0.797 or digits 59 seen
17		$6x - 7 + 7 = 38 + 7$ $6x = 45$	7.5	2	M1 $6x = 45$ or +7 both sides A1 7.5 oe; accept 45/6 oe
18		**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	3	B3 all six entries correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
19	(a)	900 × 1.70 =	1530	2	M1 900 × 1.7(0) or digits 153(0) seen A1 cao
	(b)	160 ÷ 1.70 =	£94.12 or £94.11	2	M1 160 ÷ 1.7(0) or digits 941(....) seen A1 cao
20	(a)(i)	180 – 54 (=126) "126" ÷ 2	63	3	M1 for (180–54) ÷ 2 A1 cao
	(ii)		Reason		B1 (indep) angles in triangle add to 180 OR equal angles in isosceles triangle OR equal angles and 2 sides the same (B0 if any incorrect reasoning given eg parallel, equilateral triangle)
	(b)	180 – "x"	117	1	B1 117 or ft 180 – "x" if $x < 90$
21	(a)	3 × 35 + 50	155	2	M1 for 3 × 35 + 50 or digits 155 seen A1 cao
	(b)	260 – 50 = 210 210 ÷ 35 =	6	3	M1 for 260–50 or 210 seen. M1 for "260-50" ÷ 35 or 210÷35 A1 cao SC B1 for starting at a number between 100 and 170 and adding at least two 35's and showing a total between 230 and 290 OR For adding at least three 35's, perhaps with other numbers, and showing a total between 180 and 240 (or between 230 and 290 if 50 is included in the sum)

Paper 5521_02					
No		**Working**	**Answer**	**Mark**	**Notes**
	(c)		P=35h + 50	3	B3 for P=35h+50 or P=35×h+50 oe (B2 for correct RHS or P=h + 50 ×35 or P=35h+k where k is numerical oe) (B1 for P = some other linear expression in h, OR h + 50 ×35 OR 35h seen) NB: P=h scores no marks; ignore £ signs. SC B2 for $h = \frac{P-50}{35}$
22	(a)		Elevation	2	B2 for 4 vertical squares. Accept 4 by 1 rectangle. (B1 for 4 vertical squares with one square added or one parallelogram added at the top, or 3 vertical squares, or 4 horizontal squares)
	(b)		Plan	2	B2 for 2 adjacent squares, vertical or horizontal. Accept 2 by 1 rectangle. (B1 for 3 adjacent horizontal or vertical squares or a rectangle with sides in the ratio 2:1)
23	(i)		5	3	B1 cao
	(ii)		9		B1 cao
	(iii)		6		B1 cao
24		$45.00 + 45.00 \times \frac{15}{100} =$ $45.00 + 6.75 =$	51.75	3	M2 for $45.00 + 45.00 \times \frac{15}{100}$ oe or 45.00×1.15 oe OR $45.00 + 6.75$ OR complete method or 5175 seen. (M1 for $45.00 \times \frac{15}{100}$ oe OR 6.75 seen OR 675 seen OR correct method for calculating 15% of 45) A1 cao SC Award B2 for an answer of 38.25

Paper 5521_02				
No	**Working**	**Answer**	**Mark**	**Notes**
25 (a)		Points	1	B1 all three points ±1 full square
(b)		Negative	1	B1 Negative (ignore additional descriptors unless contradictory)
(c)		lobf	1	B1 A single straight line drawn to cross between (5,30), (5,40) and (40,0), (40,15); accept freehand if considered to be straight.
(d)(i)		18–25	2	B1 18g–25g inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient.
(ii)		30–40		B1 30–40 min inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient
26		300 3 75 150	3	B3 for 4 correct answers (B2 for 2 or 3 correct answers) (B1 for 1 correct answer)
27	$\pi \times 0.65$	2.04–2.05	2	M1 for $\pi \times 0.65$ or 3.14 x 0.65 or 3.142 x 0.65 A1 for 2.04–2.05 SC Award B1 for 2.0 seen (not 2)
28	5 miles = 8 km 70mph÷5×8 = 112 km/h OR 120km/h ÷ 8 × 5 = 75 mph Faster than 70 mph	70mph (Great Britain) (112 km)	3	M1 5 miles = 8 km; OR 70 mph is about 100 km/h OR 1km=0.6(25) miles OR 1mile=1.6km oe M1 70 ÷ 5 × 8 (=112) or 120 ÷ 8 × 5 (=75) A1 (dep on at least M1) GB or 70 mph Refer to both answer line and working. NB GB or 70 mph without working scores 0 marks

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
1		Cuboid drawn	2	B2 for correct isometric drawing in any orientation (ignore points 'behind', mark 7 vertices only); accept lines drawn near to dots as long as there is no ambiguity. (B1 for one of the three faces drawn correctly **or** for an isometric drawing of any cuboid)
2	Different makes of car Tally Frequency	Make of car Tally Frequency	3	B1 for make of car or list of at least 3 different makes B1 for tally or tally marks B1 for frequency or totals
3		6 tessellating shapes	2	B2 for fully correct with 5 or more additional shapes, no gaps (B1 for 4 or more shapes tessellating, with at least one shape inverted, with or without the given shape, ignore extras)
4	24.90 ÷ 3 or 8.30 24.90 – '8.30' or 2 × 8.30	16.6(0)	3	M1 for 24.90 ÷ 3 or 8.30 M1 (dep) for 24.90 – “8.30” or 2 × “8.30” A1 for 16.60 or 16.6

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
5	315 × 24 24 × 315 1260 6300 7560 120 240 7200 7560 **3 1 5** 0 6 | 0 2 | 1 0 **2** 1 2 | 0 4 | 2 0 **4** 7 5 6 0 **300 10 5** 6000 200 100 **20** 1200 40 20 **4** 6000+200+100+1200+40+20 = 7560 **3 0.1 0.05** 60 2 1 **20** 12 0.4 0.2 **4** 60 + 2 + 1 + 12 + 0.4 + 0.2 = 75.6	75.6(0)	3	M1 for a complete method with relative place value correct. Condone 1 multiplication error, addition not necessary. **OR** M1 for a complete grid with not more than 1 multiplication error, addition not necessary. **OR** M1 for sight of a complete partitioning method, condone 1 multiplication error, final addition not necessary. A1 for 7560 or digits 756(0) A1 (dep on M1, but not previous A1) for correct placement of decimal point.

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
6	15 and 16 parts shaded Alternative 1 $\frac{3}{4} = 0.75$ or 75%, $\frac{4}{5} = 0.8$ or 80% Alternative 2 $\frac{3}{4} = \frac{15}{20}$, $\frac{4}{5} = \frac{16}{20}$	$\frac{4}{5}$ + reason	3	M1 for shading 15 parts for $\frac{3}{4}$ M1 for shading 16 parts for $\frac{4}{5}$ A1 (dep on M2) for selection of $\frac{4}{5}$ with correct shading Alternative 1 M1 for $\frac{3}{4} = 0.75$ or 75% M1 for $\frac{4}{5} = 0.8$ or 80% A1 (dep on M2) for selection of 0.8 or 80% or $\frac{4}{5}$ with correct decimals or percentages Alternative 2 M1 for $\frac{3}{4} = \frac{15}{20}$ oe M1 for $\frac{4}{5} = \frac{16}{20}$ oe A1 (dep on M2) for selection of $\frac{4}{5}$ or $\frac{16}{20}$ with equivalent fractions

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
7		$5 \times 5 \times 6$	150	4	M1 for attempt at 1 division (e.g. $40 \div 8$), may be implied by marks or number on one edge of diagram or by 5 or 6 seen M1 for attempt at 3 divisions ($40 \div 8$, $40 \div 8$, $60 \div 10$), may be implied by marks or numbers on diagram or by 5,5 and 6 seen. M1 (dep on 1st M1) for "5" × "5" × "6" A1 cao Alternatively M1 for $40 \times 40 \times 60$ **or** $8 \times 8 \times 10$ **or** 96000 or 640 seen M1 for $40 \times 40 \times 60$ **and** $8 \times 8 \times 10$ or 96000 **and** 640 seen M1 (dep on 1st M1) for "$(40 \times 40 \times 60)$" ÷ "$(8 \times 8 \times 10)$" A1 cao SC:B1 for dividing area of one carton face by area of corresponding box face if M0
8	(a)		$\frac{7}{20}$	1	B1 for $\frac{7}{20}$ oe
	(b)	7 + 2 (or 20 – 11) are not lime flavour	$\frac{9}{20}$	1	B1 for $\frac{9}{20}$ oe
	(c)		0	1	B1 for 0, zero or nought ($\frac{0}{20}$ gets B0)
9	(a)		$80x$	1	B1 for $80x$ (accept $80 \times x$, $x80$, $x \times 80$) seen
	(b)		$95y$	1	B1 for $95y$ (accept $95 \times y$, $y95$, $y \times 95$) seen
	(c)		$80x + 95y$	2	M1ft for adding "$80x$" and "$95y$" (algebraic expressions only) A1 for $80x + 95y$

Paper 5523_03

No	Working	Answer	Mark	Notes
10 (a)		40	1	B1 cao
(b)		45	1	B1 for 42 to 48 (accept 3/4 hour)
(c)	40×2 or $\frac{40}{30}\times60$ or $40\div\frac{1}{2}$	80	2	M1 for 40×2 or $\frac{40}{30}$ or $40\div\frac{1}{2}$ A1 cao NB $\frac{40}{45}\times 60$ gets M0 A0
11 (a)	$3\times3-4\times2$ or $9-8$	1	2	M1 for substitution of 3 and 2 into expression or 9 and 8 seen A1 cao
(b)	$-7-3=-10$ $2\times-10=-20$ $-20\div4$	-5	3	M1 for substitution of 2 and -7 into $p(q-3)$ or sight of -20 or $-14-6$ M1 (dep) for "-20" $\div 4$ A1 cao SC: B1 for -10 seen if M0
12 (a)	6 | 8 9 7 7 | 8 5 9 6 3 8 | 1 3 1 7 1 9 | 0 1	6 | 7 8 9 7 | 3 5 6 8 9 8 | 1 1 1 3 7 9 | 0 1	3	M1 for unordered diagram (condone one error) A1 cao B1 for key (eg 6 | 7 = 67)
(b)(i)		Explanation	2	B1 for '(order numbers and) select middle value' oe
(ii)		79		B1 cao

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
13	(a)		Reflection in y-axis	1	B1 for triangle with vertices at (-1, 1) (-3, 1) and (-1,4)
	(b)		Rotation by half turn about (0, 0)	2	B2 for triangle with vertices (-1, -1) (-3, -1) and (-1, -4) (B1 for half turn not about (0,0))
	(c)		Enlargement Scale factor 3 Centre (0, 0)	3	B1 for 'enlargement' B1 for 'scale factor 3' or 3 seen B1 for 'centre (0,0)' B0 for any combination of transformations
14	(a)		4560	1	B1 cao
	(b)		45.6	1	B1 cao
	(c)		2.4	1	B1 cao
15	(a)	$4a-2a+5b+b$	$2a+6b$	2	B2 cao (B1 for $2a$ or $6b$ seen)
	(b)		$x(x-6)$	2	B2 cao (B1 for $x(ax+b)$ where a, b are numbers not equal to zero or $x-6$ seen on its own, or part of an expression)
	(c)		$3x-2x^3$	2	B2 cao (B1 for $3x$ or $2x^3$)
	(d)		$4x(3y+x)$	2	B2 cao (B1 for $2(6xy+2x^2)$ or $4(3xy+x^2)$ or $x(12y+4x)$ or $2x(6y+2x)$ or $4x(\quad)$)

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
16	(a)	$1 - (0.2 + 0.3 + 0.1)$	0.4	2	M1 for $1 - (0.2 + 0.3 + 0.1)$ A1 for 0.4 oe, accept $\frac{0.4}{1}$
	(b)	0.2×200	40	2	M1 for 0.2×200 A1 cao NB $\frac{40}{200}$ is M1 A0, 40 out of 200 is M1 A1
17	(a) (i)	$180 - 2 \times 25$	130	3	M1 for $180 - 2 \times 25$ A1 cao
	(ii)		Reason		B1 for mentioning isosceles and equal (or base) angles or equal sides and equal (or base) angles
	(b)	$180 - 95$	85	1	B1 cao
18	(a) (i)		7^5	3	B1 cao
	(ii)		7^4		B2 cao (B1 for sight of 7^5 or 7^{2+3} or 7×7^3 or $7^1 \times 7^3$ or $7^2 \times 7^2$ or 7^{2+3-1})
	(b)		$\frac{1}{2}$	1	B1 for $\frac{1}{2}$ or 0.5 or 2^{-1}
19	(a)		3×10^7	1	B1 cao
	(b)		0.002	1	B1 cao

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
20		Box plot	2	3 aspects: 1st aspect – vertical line for median 2nd aspect – box using correct quartiles 3rd aspect – whiskers (could be single line) drawn with correct end points B2 for fully correct box plot (B1 for 1 aspect)
21 (a)	e.g. 2 | 126 3 | 63 3 | 21 7	$2\times3\times3\times7$	2	M1 for a systematic method of at least 2 correct divisions by a prime number oe factor trees; can be implied by digits 2, 3, 3, 7 on answer line. A1 for $2\times3^2\times7$ or $2\times3\times3\times7$
(b)	$2\times3\times7$	42	2	B2 cao (B1 for 6, 14, 21 or $2\times3\times7$)
22	$\frac{8}{3}\times\frac{5}{4}=\frac{8\times5}{3\times4}=\frac{40}{12}$	$3\frac{1}{3}$	3	B1 for $\frac{8}{3}$ oe improper fraction or $\frac{5}{4}$ oe improper fraction M1 (dep on B1) for multiplying numerator and denominator of "$\frac{8}{3}$" and "$\frac{5}{4}$" A1 for $3\frac{1}{3}$ oe mixed number or $\frac{10}{3}$ OR B1 for 1.25 and 2.67 or 2.66(…) M1 (dep on B1) for correct method of multiplication A1 for $3.\dot{3}$

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
23			2	M1 for a relevant pair of intersecting arcs A1 for line drawn within guidelines, at least 3cm in length, accept broken line [SC: B1 for line drawn within guidelines if M0]
24 (a)		$-1, 0, 1, 2, 3$	2	B2 cao (-1 each error or omission)
(b)(i)		$x \geq \frac{7}{2}$	3	M1 for $2x \geq 7$, condone use of = sign or wrong equality A1 for $x \geq \frac{7}{2}$ oe as final answer
(ii)		4		SC:B1 for 3.5 or $\frac{7}{2}$ seen if M0 B1 ft from $x \geq$ "$\frac{7}{2}$"
25	$4x + 2y = 8$ $4x - 10y = 20$ $12y = -12$ $y = -1$ $4x + 2(-1) = 8$ $x = 2.5$	$x = 2.5$ $y = -1$	3	M1 for correct process to eliminate either x or y (condone one arithmetical error) M1 (dep) for substituting found value into either equation A1 for $x = 2.5$, $y = -1$ [SC: B1 for $x = 2.5$ or $y = -1$ if M0]

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
26		Interior angle of hexagon = 180 – (360 ÷ 6) = 120 360 – (90 + 120)	150	4	Alternative 1 M1 for 360 ÷ 6 A1 for 60 M1 (dep on M1) for "60" + 90 A1 cao Alternative 2 M1 for 360 ÷ 6 A1 for 60 M1 (dep on M1) for 360 – (2 × "60" + 90) A1 cao Alternative 3 M1 for (6 – 2) × 180 ÷ 6 A1 for 120 M1 (dep on M1) for 360 – (90 + "120") A1 cao
27	(a)		(16), 50, 82, 96, 100	1	B1 cao
	(b)	Cumulative freq. diag. curve/ segments	Cum. freq graph	2	B1 for 4 or 5 points plotted correctly ± 1 full (2mm) square depending on sensible table (condone 1 addition error) B1 (dep) for points joined by curve or line segments provided no gradient is negative - ignore any part of graph outside range of their points. (SC:B1 if 4 or 5 points plotted not at end but consistent within each interval and joined)
	(c)	100 – 42	58	2	M1 (ft dep on graph being cf) for reading from graph at 18 or 19, can be implied by answer in range 40 to 46 A1 for answer in range 56 to 60 or ft for 100 – '42' ±1 full (2mm) square

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
1	(a)	900 × 1.70 =	1530	2	M1 900 × 1.7(0) or digits 153(0) seen A1 cao
	(b)	160 ÷ 1.70 =	£94.12 or £94.11	2	M1 160 ÷ 1.7(0) or digits 941(....) seen A1 cao
2	(a)	4.7 ÷ 5.9 = 0.796610169	0.7966..	2	B2 for 0.7966 or better (B1 for 0.8, 0.80, 0.79, 0.796, 0.797 or digits 59 seen)
	(b)	0.82, 0.8, 0.85, 0.66, 0.875 0.66, 0.8, 0.82, 0.85, 0.875 2/3, 4/5, 0.82, 85%, 7/8	2/3, 4/5, 0.82, 85%, 7/8	2	B2 correct order (oe decimals in order) (B1 correct order reversed, or one error in ordered listing) with or without decimal equivalents. NB Accept 0.67 or 0.66
3		**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	3	B3 all 6 correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
4	(a)	–3 (–1) (1) 3 5 7	**–3**, 3, **5**, **7**	2	B2 all correct (B1 2,3 correct)
	(b)	(-2,-3),(-1,-1),(0,1),(1,3),(2,5),(3,7)	line	2	B2 cao for line from $x = -2$ to $x = 3$ (B1 plotting at least 5 points correctly or single line passing through (0, 1) or single line of gradient 2) The six possible points are: (-2,-3), (-1,-1), (0,1), (1,3), (2,5), (3,7)
	(c)(i)		–2	2	B1 $y = -2$ or ft from line segment
	(ii)		2.5		B1 $x = 2.5$ or ft from line segment

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
5	(a) (i)	180 – 54 (=126) "126" ÷ 2	63	3	M1 for (180–54) ÷ 2 A1 cao
	(ii)		Reason		B1 (indep) angles in triangle add to 180 OR equal angles in isosceles triangle OR equal angles and 2 sides the same (B0 if any incorrect reasoning given eg parallel, equilateral triangle)
	(b)	180 – "x"	117	1	B1 117 or ft 180 – "x" if $x < 90$
6		15/120×360 = 45 Plain 40/120×360 = 120 Cheese & Onion 55/120×360 = 165 Salt & Vinegar 10/120 × 360 = 30 Beef	45°, 120°, 165°, 30°	4	M1 evidence of method for at least one angle (could be implied by one correct angle of four on pie chart or in the table) A2 All four angles drawn ±2° tolerance, any order (A1 at least 2 angles correctly drawn ±2°, or all 4 angles in the table) B1 (dep on at least 1 angle drawn correctly, and exactly 4 sectors) for labels (flavour or frequency; initials will do) NB: Ignore the table if the pie chart provides the marks.
7	(a)	260 – 50 = 210 210 ÷ 35 =	6	3	M1 for 260–50 or 210 seen. M1 for "260-50" ÷ 35 or 210÷35 A1 cao
	(b)		P=35h + 50	3	B3 for P=35h+50 or P=35×h+50 oe (B2 for correct RHS or P=h + 50 ×35 or P=35h+k where k is numerical oe) (B1 for P = some other linear expression in h, OR h + 50 ×35 OR 35h seen) NB: P=h scores no marks; ignore £ signs. SC B2 for $h = \frac{P-50}{35}$
8	(a)		Elevation	2	B2 for 4 vertical squares. Accept 4 by 1 rectangle. (B1 for 4 vertical squares with one square added or one parallelogram added at the top, or 3 vertical squares, or 4 horizontal squares)

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
(b)		Plan	2	B2 for 2 adjacent squares, vertical or horizontal. Accept 2 by 1 rectangle. (B1 for 3 adjacent horizontal or vertical squares or a rectangle with sides in the ratio 2:1)
9 (a)		Points	1	B1 all three points ±1 full square.
(b)		Negative	1	B1 Negative. Ignore other descriptors unless contradictory.
(c)		lobf	1	B1 A single straight line drawn to cross between (5,30), (5,40) and (40,0), (40,15); accept freehand if considered to be straight.
(d)(i) (ii)		18–25 30–40	2	B1 18g–25g inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient. B1 30–40 min inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient.
10		300, 3, 75, 150	3	B3 for 4 correct (B2 for 2 or 3 correct) (B1 for 1 correct)
11 (a)	$6x - 7 + 7 = 38 + 7$ $6x = 45$	7.5	2	M1 $6x = 45$ or +7 both sides A1 7.5 oe; accept 45/6
(b)	$5y - 2 = 10$ or $20y - 8 = 40$ $5y = 12$ $20y = 48$	$2\frac{2}{5}$	3	M1 $20y–8(=40)$ or $\frac{4(5y-2)}{4} = \frac{40}{4}$ or $5y–2=10$ M1 (indep) for correct rearrangement into the form $ay=b+c$ or better (eg $20y=40+8$ or $5y=10+2$, using own terms) A1 for $2\frac{2}{5}$, 2.4 oe
12	5 miles = 8 km 70mph÷5×8 = 122 km/h OR 120km/h ÷ 8 × 8 = 75 mph Faster than 70 mph	70mph (Great Britain) (112 km)	3	M1 5 miles = 8 km; OR 70 mph is about 100 km/h OR 1km=0.6(25) miles OR 1mile=1.6km oe M1 70 ÷ 5 × 8 (=112) or 120 ÷ 8 × 5 (=75) oe A1 (dep on at least M1) GB or 70 mph Refer to both answer line and working. NB GB or 70 mph without working scores 0 marks.

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
13 (a)	$\pi \times 0.65$	2.04–2.05	2	M1 for $\pi \times 0.65$ or 3.14×0.65 or 3.142×0.65 oe A1 2.04–2.05 SC Award B1 for 2.0 seen (not 2)
(b)	1000 ÷ "(a)"	487–491	2	M1 for 1000 (or 100) ÷ "(a)" A1 for 487–491
14	$45.00 + 45.00 \times \frac{15}{100} =$ $45.00 + 6.75 =$	51.75	3	M2 for $45.00 + 45.00 \times \frac{15}{100}$ oe or 45.00×1.15 oe OR $45.00 + 6.75$ OR complete method or 5175 seen. (M1 for $45.00 \times \frac{15}{100}$ oe OR 6.75 seen OR 675 seen OR correct method for calculating 15% of 45) A1 cao SC Award B2 for an answer of 38.25
15		273	2	M1 for 728 ÷ 8 or 728 ÷ "3+5" or 91 A1 cao SC B1 for 455, or for 273:455
16	3 24 3.7 46.9(53) 4 60 3.8 51.0(72) 3.1 26.6(91) 3.9 55.4(19) 3.2 29.5(68) 3.21 29.8(66…) 3.3 32.6(37) 3.22 30.1(66…) 3.4 35.9(04) 3.23 30.4(68..) 3.5 39.3(75) 3.24 30.7(72...) 3.6 43.0(56) 3.25 31.0(78...) or 31	3.2	4	B2 for trial between 3.2 and 3.3 inclusive (B1 for trial between 3 and 4 inclusive) B1 for different trial between 3.21 and 3.25 inclusive B1 (dep on at least one previous B1) cao for 3.2 as final answer. NB: embedded answers: –B1; award Bs for evaluations rounded or truncated to at least 1 dp or for 31

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
17	$45^2 + 34^2 =$ $2025 + 1156 = 3181$ $\sqrt{3181} = 56.4$	56	4	M1 for $45^2 + 34^2$ M1 (dep) for $\sqrt{(2025 + 1156)}$ A1 for 56.4 … B1 for rounding their diagonal to the nearest integer (dep on evidence from decimal) NB 56 as the final answer gets full marks. NB Scale drawings result in 0 marks.
18	$2000 \times (1.055)^3$ Interest = 2348.48 – 2000 =	348.48	3	M1 for 5.5/100 × 2000 (oe) or 330 or 2330 or 110 or 2110 M1 (dep) for 5.5/100 × (2000 + "110" + "116.05") or 122.4… seen A1 cao (accept only 348.48 or 348.49) **OR** M2 for $2000 \times (1.055)^3$ or 2348.48(…) or 2348.49 seen (M1 for $2000 \times (1.055)^n$, $n \neq 3$) A1 for 348.48 or 348.49 [SC: B2 for 2348.48 - 2348.49]
19		Line	2	B2 line fully within tramlines, crossing AB and CD (B1 a straight line which crosses AB within the tramline, and also crosses CD) NB: Accept dotted or dashed lines, but not curves; accept freehand if considered to be straight. SC B1 for a perpendicular bisector of AB that is at least half way from AB to CD within the tramlines
20 (a)		$90<t\leq100$	1	B1 for $90<t\leq100$; accept 90-100.
(b)	65×12=780 75×22=1650 85×23=1955 95×24=2280 105×19=1995 8660/100=	86.6	4	M1 for use of fx with x consistent within intervals (including end points). Allow one slip. M1 (dep) for use of midpoints M1 (dep on 1st M1) for use of $\sum fx/100$ or $\sum fx/\sum f$ A1 86.6

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
21	(a)	$8 \times \frac{10}{4}$	20	2	M1 $\frac{10}{4}$ or $\frac{4}{10}$ or 0.4 or 2.5 oe seen A1 cao NB ratios get M0 unless of the form 1:*n* OR M1 $\frac{8}{4}, \frac{4}{8}$ oe seen A1 cao
	(b)	$15 \times \frac{4}{10}$	6	2	M1 $15 \times \frac{4}{10}$ oe A1 cao
22	(a)	$x^2 - 4x + 3x - 12 = x^2 - x - 12$	$x^2 - x - 12$	2	M1 for exactly 4 terms correct ignoring signs (eg x^2, 4*x*, 3*x*, 12) or 3 correct terms out of 4 terms with correct signs (eg 3 out of 4 of x^2, –4*x*, +3*x*,–12) A1 cao
	(b)	$(x+2)(x+5)$	$(x+2)(x+5)$	2	B2 cao (B1 for exactly one of $(x+2)$, $(x+5)$)
	(c)	$6=15+4q-20$ $6-15=4(q-5)$ $p-3t=4q-4t$ $6-3\times5=4(q-5)$	2 ¾	3	M1 for correct substitution of *p* and *t*. M1 for correct expansion of $4(q-t)$ oe (eg $4q-20$, $4q-4t$) A1 11/4 or 2 ¾ or 2.75 OR M1 for correct substitution of *p* and *t*. M1 for $\frac{p-3t}{4} = q - t$ oe A1 11/4 or 2 ¾ or 2.75

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
23	(a)	$9.9\times10^8 - 6.0\times10^7$	9.3×10^8	2	M1 for $99\times10^7 - 6\times10^7$ or $9.9\times10^8 - 0.6\times10^8$ or conversion of either to an ordinary number, or 930000000 or 93×10^7 or 9.3×10^n where n is any positive integer A1 cao
	(b)	$\frac{6.0-4.5}{4.5}\times100 = \frac{1.5}{4.5}\times100 =$ or $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}\times100$	33.3%	3	M2 $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}\times100$ oe (M1 for $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}$ or $\frac{6.0\times10^7 - 4.5\times10^7}{6.0\times10^7}\times100$ oe A1 cao OR M2 $\frac{6.0\times10^7}{4.5\times10^7}\times100$ -100 = (M1 $\frac{6.0\times10^7}{4.5\times10^7}\times100$ or 133.33(%)) A1 cao NB Accept any of the above expressions without any reference to 10^7.
24		LCM (40,24) = 120 Bread buns 120÷40 Burgers 120÷24 OR Bread buns: 40 is $2\times2\times2$ (×5) Burgers: 24 is $2\times2\times2$ (×3)	Bread buns 3 Burgers 5	3	M1 attempt to find LCM by eg lists of multiples, or summing of 40 and 24. A1 identify 120 (as LCM) A1 cao (both) OR M1 expansion of either number into its prime factors in a factor tree or 8×5 or 8×3 A1 both expansions correct A1 cao (both) SC B2 if answers given the wrong way around

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
25	8 × 41 = 328 2 × 29 = 58 328 – 58 = 270 270 ÷ 6 = 45	45	3	M1 for either 8 × 41 (=328) or 2 × 29 (=58) M1 (dep) “328” – “58” (=270) A1 cao NB 328 and/or 58 on the answer line gets M1 (implied); 270 on the answer line gets M2 (implied)

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
1 (a)	$1 - (0.2 + 0.3 + 0.1)$	0.4	2	M1 for $1 - (0.2 + 0.3 + 0.1)$ A1 for 0.4 oe, accept $\frac{0.4}{1}$
(b)	0.2×200	40	2	M1 for 0.2×200 A1 cao NB $\frac{40}{200}$ is M1 A0, 40 out of 200 is M1 A1
2	$650 - 430 = 220$ 1 choc ice costs 110p $650 - 5 \times 110 = 100$p	50	3	M1 for $650 - 430$ or 220 or 110 oe seen M1 for $650 - 5 \times \frac{'220'}{2}$ or $430 - 3 \times \frac{'220'}{2}$ oe A1 for 50p or £0.50 or £0.5 *Alternative scheme* $2x + 5y = 650$ $2x + 3y = 430$ oe M1 for subtracting two simultaneous equations to eliminate x (lollies)(2 or 3 terms correct) M1 for $650 - 5 \times 'y'$ or $430 - 3 \times 'y'$ oe A1 for 50p or £0.50 or £0.5 *Alternative scheme* M1 for $3\times (2x + 5y = 650)$ evaluated and $5 \times (2x + 3y = 430)$ evaluated oe (5 or 6 terms correct) M1 for subtraction of equations to eliminate y (choc ices)(2 or 3 terms correct ft) A1 for 50p or £0.50 or £0.5

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
3		question + response boxes oe	2	1st aspect: One question (eg 'how long does it take you to travel to school?' or 'What time did you leave home to get to school?'); ignore other questions. 2nd aspect: Response list (at least two), not overlapping. 3rd aspect: Some mention of units (eg minutes) in either question or responses B2 for all three aspects, or B1 for just one aspects.
4	2 [(3×1) + (4×1)] + (3×6) + (1×6) + (2×6) + (4×6) + (1×6) + (5×6)	110 cm^2	4	M1 for attempt to find the area of one face M1 for at least 6 faces with intention to add A1 cao B1 (indep) for cm^2 (with or without numerical answer)
5	$\left(\frac{-6+4}{2}\right), \left(\frac{5+3}{2}\right)$	(–1, 4)	2	B2 cao [B1 for (–1,*a*) or (b,4) or (4,-1)]
6		Box plot	2	3 aspects: 1st aspect – vertical line for median 2nd aspect – box using correct quartiles 3rd aspect – whiskers (could be single line) drawn with correct end points B2 for fully correct box plot (B1 for 1 aspect)
7			2	M1 for a relevant pair of intersecting arcs A1 for line drawn within guidelines, at least 3cm in length, accept broken line [SC: B1 for line drawn within guidelines if M0]

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
8	(a)	e.g. 2 | 126 3 | 63 3 | 21 7	$2\times3\times3\times7$	2	M1 for a systematic method of at least 2 correct divisions by a prime number oe factor trees; can be implied by digits 2, 3, 3, 7 on answer line. A1 for $2\times3^2\times7$ or $2\times3\times3\times7$
	(b)	$2\times3\times7$	42	2	B2 cao (B1 for 6, 14, 21 or $2\times3\times7$)
9	(a)		$-1,0,1,2,3$	2	B2 cao (-1 each error or omission)
	(b)(i)		$x\geq\frac{7}{2}$	3	M1 for $2x\geq7$, condone use of = sign or wrong equality A1 for $x\geq\frac{7}{2}$ oe as final answer
	(ii)		4		SC:B1 for 3.5 or $\frac{7}{2}$ seen if M0 B1 ft from $x\geq$ "$\frac{7}{2}$" or $x>$"$\frac{7}{2}$"
10	(a)(i)		7^5	3	B1 cao
	(ii)		7^4		B2 cao (B1 for sight of 7^5 or 7^{2+3} or 7×7^3 or $7^1\times7^3$ or $7^2\times7^2$ or 7^{2+3-1})
	(b)		$\frac{1}{2}$	1	B1 for $\frac{1}{2}$ or 0.5 or 2^{-1}
11	(a)	$5n=m+21$	$n=\frac{m+21}{5}$	2	M1 for $5n=m+21$ or for attempt to divide three terms by 5 A1 $n=\frac{m+21}{5}$ oe

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
(b)	$4p-8q=3p+2$ $p-8q=2$ $p=8q+2$	$p=8q+2$	3	M1 for $4p-aq$ or $\frac{3}{4}p+b$ where a is an integer and b is a number M1 (dep) for taking one term correctly to LHS or RHS of expression A1 $p=8q+2$ oe
12		$y=\frac{1}{2}x+3$	2	B2 for $y = ½x + 3$ oe (B1 for $y = ½x + c$, $c \neq 7$ or $y = mx + 3$ oe or $½x + 3$ or $M = ½x + 3$)
13	$\frac{8}{3}\times\frac{5}{4}=\frac{8\times5}{3\times4}=\frac{40}{12}$	$3\frac{1}{3}$	3	B1 for $\frac{8}{3}$ oe or $\frac{5}{4}$ oe M1 (dep on B1) for multiplying numerator and denominator of "$\frac{8}{3}$" and "$\frac{5}{4}$" A1 for $3\frac{1}{3}$ oe mixed number or $\frac{10}{3}$ *OR* B1 for 1.25 and 2.67 or 2.66(…) M1(dep on B1) for correct method of multiplication A1 for $3.\dot{3}$
14	$4x+2y=8$ $4x-10y=20$ $12y=-12$ $y=-1$ $4x+2(-1)=8$ $x=2.5$	$x=2.5$ $y=-1$	3	M1 for correct process to eliminate either x or y (condone one arithmetical error) M1 (dep) for substituting found value into either equation A1 for $x = 2.5$, $y = -1$ [SC: B1 for $x = 2.5$ or $y = -1$ if M0]

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
15		Interior angle of hexagon = 180 – (360 ÷ 6) = 120 360 – (90 + 120)	150	4	Alternative 1 M1 for 360 ÷ 6 A1 for 60 M1 (dep on M1) for "60" + 90 A1 cao Alternative 2 M1 for 360 ÷ 6 A1 for 60 M1(dep on M1) for 360 – (2 × "60" + 90) A1 cao Alternative 3 M1 for (6 – 2) × 180 ÷ 6 A1 for 120 M1(dep on M1) for 360 – (90 + "120") A1 cao
16	(a)		(16), 50, 82, 96, 100	1	B1 cao
	(b)	Cumulative freq. diag. curve/ segments	Cum. freq graph	2	B1 ft for 4 or 5 points plotted correctly ± 1 full (2mm) square depending on sensible table (condone 1 addition error) B1 (dep) for points joined by curve or line segments provided no gradient is negative - ignore any part of graph outside range of their points. (SC:B1 if 4 or 5 points plotted not at end but consistent within each interval and joined)

Paper 5525_05					
No		Working	Answer	Mark	Notes
	(c)	$100 - 42$	58	2	M1 (ft dep on graph being cf) for reading from graph at 18 or 19, can be implied by answer in range 40 to 46 A1 for answer in range 56 to 60 or ft for 100 –'42' ±1 full (2mm) square
17	(a)		0.6 and 0.7,0.3,0.7	2	B1 for 0.6 on LH branch B1 for 0.7, 0.3 and 0.7 on RH branches
	(b)	0.4×0.3	0.12	2	M1 for 0.4×0.3 A1 0.12 oe
	(c)	$0.4\times0.7+0.6\times0.3$	0.46	3	M1 for '0.4×0.7' or '0.6×0.3' M1 for addition of two products from correct branches A1 0.46 oe Alternative M2 for an attempt to evaluate 1-(0.3×0.4 +'0.6×0.7') A1 cao
18		$x = 0.4545...$ so $100x = 45.4545...$ $99x = 45$ $x = \frac{45}{99} = \frac{15}{33}$	proof	3	M1 for $100x = 45.45\ldots$ or $10000x = 4545.45\ldots$ M1 (dep) for subtraction of both sides A1 for $\frac{15}{33}$ from correct proof
19		$\left(\sqrt{3}\right)^2 + \sqrt{3}\sqrt{2} - \sqrt{3}\sqrt{2} - \left(\sqrt{2}\right)^2$ $= 3 - 2$	1	2	B2 cao (B1 for $\sqrt{3}\sqrt{3} + \sqrt{3}\sqrt{2} - \sqrt{3}\sqrt{2} - \sqrt{2}\sqrt{2}$ oe, $\sqrt{3}\sqrt{3} - \sqrt{2}\sqrt{2}$ oe, or for 2,3, $\sqrt{4}$ $\sqrt{6}$ $\sqrt{9}$ seen)
20	(a)	$\frac{96}{24}$ or 4 $\sqrt{4}$ or 2	8	3	M1 for $\frac{96}{24}$ or $\frac{24}{96}$ or 4 or $\frac{1}{4}$ oe M1 for $\sqrt{\frac{96}{24}}$ or $\sqrt{\frac{24}{96}}$ or $\sqrt{'4'}$ or $\frac{1}{\sqrt{'4'}}$ or 2 or $\frac{1}{2}$ oe A1 cao

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
	(b)	12×2^3	96	2	M1 for '2'^{3} or 8 A1 cao
21	(a)	$x\times3-x\times2x^2$	$3x-2x^3$	2	B2 cao (B1 for a two term expression with either $3x$ or $2x^3$)
	(b)		$4x(3y+x)$	2	M1 for taking out a factor of x, $2x$, 2, 4 or $4x$ A1 cao
	(c)	$\frac{5a}{b^2}$	$\frac{5a}{b^2}$	2	B2 for $\frac{5a}{b^2}$ or $5ab^{-2}$ (accept $\frac{5a}{1b^2}$) (B1 for either dealing with the numbers or dealing with the powers of a)
	(d)	$\frac{\cancel{x-3}}{\cancel{(x-3)}(x+3)}$	$\frac{1}{x+3}$	2	M1 for $(x-3)(x+3)$ A1 cao
22	(a)		$\mathbf{a}+\mathbf{b}$	1	B1 $\mathbf{a}+\mathbf{b}$ oe
	(b)	$ED=\mathbf{a}$ $DX=-2\mathbf{b}+2\ AC=2\mathbf{a}$ (So, $DX=2ED$)	$2\mathbf{a}$	3	M1 for $(DX=)\ DA+AX$ A1 for $(DX=)\ -2\boldsymbol{b}+2$ ("$\boldsymbol{a}+\boldsymbol{b}$") A1 $2\boldsymbol{a}$ from fully correct proof
23		$\pi(2x)^2h=\frac{4}{3}\pi(3x)^3$ $h=\frac{\frac{4}{3}\pi(3x)^3}{\pi(2x)^2}$	$9x$	3	M1 for $\pi(2x)^2h=\frac{4}{3}\pi(3x)^3$ (condone absence of brackets) M1 (dep) for valid algebra that gets to $h=ax$ (condone one error in powers of numerical constants) A1 cao

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
24	(i)	$n^2+(n+1)^2=2(n^2+n)+1$	$2n^2+2n+1$	4	M1 for at least 3 terms correct from $n^2+n+n+1$ A1 for $2n^2+2n+1$ oe
	(ii)		$2(n^2+n)$ is always even so $2(n^2+n)+1$ is always odd		M1 for recognizing $2n^2$ is always even A1ft complete proof for their quadratic *Alternative method* M1 for recognizing that if n^2 is odd then $(n+1)^2$ is even or vice versa A1 for complete proof
25			42 318	2	B1 for answer in range 36 – 48 B1 for answer in range 312 – 324
26	(a)	$(x-3)^2-3^2+15$	$p=3, q=6$	2	B2 for $p=3$ and $q=6$ (B1 for $p=3$ OR $q=6$) SC: award B2 for $(x-3)^2+6$ if p and q are not identified
	(b)		Sketch	2	B1 for U shaped curve B1 ft for TP in first quadrant (ft if TP not in first quadrant)
27	(a)	Graph translated 2 units upwards through points (–4, 2), (–2, 4), (0,2) and (3,5)	Sketch	2	M1 for a vertical translation A1 curve through points (–4, 2), (–2, 4), (0,2) and (3,5) ± ½ square
	(b)	Graph reflected in x-axis through points (–4,0), (–2, –2), (0, 0) and (3, -3)	Sketch	2	M1 for reflection in x-axis or y-axis A1 curve through points (–4,0), (–2, –2), (0, 0) and (3, -3) ± ½ square

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
1		273	2	M1 for 728 ÷ 8 or 728 ÷ "3+5" or 91 A1 cao SC B1 for 455 or 273:455
2		$3n - 1$	2	B2 for $3n - 1$ oe (B1 for $3n + k$ where $k \neq -1$ but k could be 0)
3	3 24 3.7 46.9(53) 4 60 3.8 51.0(72) 3.1 26.6(91) 3.9 55.4(19) 3.2 29.5(68) 3.21 29.8(66…) 3.3 32.6(37) 3.22 30.1(66…) 3.4 35.9(04) 3.23 30.4(68..) 3.5 39.3(75) 3.24 30.7(72...) 3.6 43.0(56) 3.25 31.0(78...) or 31	3.2	4	B2 for trial between 3.2 and 3.3 inclusive (B1 for trial between 3 and 4 inclusive) B1 for different trial between 3.21 and 3.25 inclusive B1 (dep on at least one previous B1) cao for 3.2 NB: embedded answers: –B1; award Bs for evaluations rounded or truncated to at least 1 dp or for 31
4	2 is the only even prime number and the product of 2 odd numbers is odd	Yes	2	B2 for 'yes' and '2 is the only even prime number and the product of two odd numbers is odd' oe (B1 for 'yes' and either '2 is the only even prime number' oe **or** 'the product of two odd numbers is odd' oe)
5	$45^2 + 34^2 =$ $2025 + 1156 = 3181$ $\sqrt{3181} = 56.4$	56	4	M1 for $45^2 + 34^2$ M1 for $\sqrt{(2025 + 1156)}$ A1 for 56.4 … B1 for rounding their diagonal to the nearest integer (dep on evidence from decimal) NB 56 with no incorrect working as the final answer gets full marks. NB Scale drawings result in 0 marks.

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
6	$2000 \times (1.055)^3$ Interest = 2348.48 – 2000 =	£348.48	3	M1 for 5.5/100 × 2000 (oe) or 330 or 16.5/100×2000 or 2330 or 110 or 2110 M1 (dep) for 5.5/100 × (2000 + "110" + "116.05") or 122.4… seen A1 cao (accept only 348.48 or 348.49) **OR** M2 for $2000 \times (1.055)^3$ or 2348.48(…) or 2348.49 seen (M1 for $2000 \times (1.055)^n$, $n \neq 3$) A1 for 348.48 or 348.49 [SC: B2 for 2348.48 - 2348.49]
7 (a)		Line	2	B2 line fully within tramlines, crossing AB and CD (B1 a straight line which crosses AB within the tramline, and also crosses CD) NB: Accept dotted or dashed lines, but not curves; accept freehand if considered to be straight. SC B1 for the perpendicular bisector of *AB* reaching halfway or more from *AB*
(b)		Region		B2 correct arc ±2mm and shaded within. Allow dotty or continous arc. (B1 inaccurate arc and shaded or accurate arc unshaded)
8	65×12=780 75×22=1650 85×23=1955 95×24=2280 105×19=1995 8660/100=	86.6	4	M1 for use of fx with x consistent within intervals (including end points). Allow one slip even if outside interval M1 (dep) for use of midpoints M1 (dep on 1st M1) for use of $\sum fx/100$ or $\sum fx/\sum f$ A1 86.6

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
9		Distance ÷ time: 1400 ÷ 2 h 20 min 20 mins is $\frac{1}{3}$ hour 1400 × 3 ÷ 7 =	600 kph	3	B1 20 mins as $\frac{1}{3}$ hour or as 0.33……hour M1 for distance ÷ time eg 1400 ÷ "2h 20 min" A1 cao OR B1 2 hour 20min = 140 (min) M1 Speed = $\frac{1400}{140}$ = (10 km per minute) A1 cao
10	(a)	$(x+18)+2x+(2x+7)=180$	Equation	2	B2 for $(x+18)+2x+(2x+7)=180$ oe (B1 for $(x+18)+2x+(2x+7)$)
	(b)	$5x+25 = 180$ $5x = 155$	31	2	M1 for simplifying to at least $5x+25=180$ or 360 (may be earned in (a)) A1 for $x=31$
11	(a)	(–14) –4 (0) 4 14	–4, 4, 14	2	B2 for all 3 values correct (B1 for 1 or 2 values correct)
	(b)		curve	2	B1 for all 5 points plotted correctly ± ½ square (ft from table if at least B1 awarded in (a)) B1 (indep) ft for any smooth curve through their points

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
12	(a)	$8 \times \frac{10}{4}$	20	2	M1 $\frac{10}{4}$ or $\frac{4}{10}$ or 0.4 or 2.5 oe seen A1 cao NB ratios get M0 unless of the form 1:*n* OR M1 $\frac{8}{4}, \frac{4}{8}$ oe seen A1 cao
	(b)	$15 \times \frac{4}{10}$	6	2	M1 $15 \times \frac{4}{10}$ oe A1 cao
13	(a)	$9.9\times10^8 - 6.0\times10^7$	$\pm 9.3\times10^8$	2	M1 for $99\times10^7 - 6.0\times10^7$ or $9.9\times10^8 - 0.60\times10^8$ or correct conversion of either to an ordinary number or 930000000 or 9.3×10^n where *n* is any positive integer A1 cao
	(b)	$\frac{6.0-4.5}{4.5} \times 100 = \frac{1.5}{4.5} \times 100 =$ or $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7} \times 100$	33.3%	3	M2 $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7} \times 100$ oe (M1 for $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}$ or $\frac{6.0\times10^7 - 4.5\times10^7}{6.0\times10^7} \times 100$ oe A1 33.3 – 33.4 OR M2 $\frac{6.0\times10^7}{4.5\times10^7} \times 100$ -100 or 33.3% (M1 $\frac{6.0\times10^7}{4.5\times10^7} \times 100$ or 133.3%) A1 33.3-33.4 NB Accept any of the above expressions without any reference to 10^7.

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
14	LCM (40,24) = 120 Bread buns 120÷40 Burgers 120÷24 OR Bread buns: 40 is $2 \times 2 \times 2$ (×5) Burgers: 24 is $2 \times 2 \times 2$ (×3)	Bread Buns 3 Burgers 5	3	M1 attempt to find LCM by eg lists of multiples, or summing of 40s and summing of 24s, with at least 3 numbers in each list A1 identify 120 as LCM A1 cao (both) **OR** M1 expansion of either number into its prime factors in a factor tree or 8×5 or 8×3 A1 both expansions correct A1 cao (both) SC B2 if answers given the wrong way around
15	$\sin 32 = \frac{AB}{12}$ $AB = 12 \times \sin 32$ $AB = 6.35903\ldots$	6.36	3	M1 $\sin 32 = \frac{AB}{12}$ (accept $\text{Sin}\ \frac{AB}{12}$) M1 $12 \times \sin 32$ or $12 \times 0.5299..$ A1 accept 6.359 – 6.360 SC Gradians 5.78(1…) Radians 6.62 Get M1M1A0 **OR** Use of Sine Rule $\frac{\sin 32}{AB} = \frac{\sin 90}{12}$ or $\frac{AB}{\sin 32} = \frac{12}{\sin 90}$ M1 $AB = \frac{12 \times \sin 32}{\sin 90}$ M1 AB = 6.359 – 6.36 A1 SC Gradians 5.85(…) Radians 7. 40(…) M1M1A0

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
16		$8 \times 41 = 328$ $2 \times 29 = 58$ $328 - 58 = 270$ $270 \div 6 = 45$	45	3	M1 for either 8×41 (=328) or 2×29 (=58) M1 (dep) "328" – "58" (=270) A1 cao NB 328 and /or 58 on the answer line gets M1 (implied); 270 on the answer line gets M2 (implied)
17	(a)	$x^2 - 4x + 3x - 12 = x^2 - x - 12$	$x^2 - x - 12$	2	M1 for exactly 4 terms correct ignoring signs (x^2, $4x$, $3x$, 12) or 3 out of 4 terms with correct signs (x^2, $-4x$, $+3x$, -12) A1 cao
	(b)	$6x^2 - 8x + 15x - 20 = 6x^2 + 7x - 20$	$6x^2 + 7x - 20$	2	M1 for exactly 4 terms correct ignoring signs ($6x^2$, $8x$, $15x$, 20) or 3 out of 4 terms with correct signs ($6x^2$, $-8x$, $+15x$, -20) A1 cao
	(c)	$(x+2)(x+5)$	$(x+2)(x+5)$	2	B2 cao (B1 for exactly one of $(x + 2)$, $(x + 5)$)
	(d)		$12p^8q^3$	2	B2 cao (B1 for any 2 out of 3 terms correct in a product or 3 terms correct in a sum or part product)
	(e)	$6 = 15 + 4(q - 5)$ $6 = 15 + 4q - 20$ $11 = 4q$	$2\frac{3}{4}$	3	M1 for correct substitution of p and t. M1 for correct expansion of $4(q\text{-}t)$ oe (eg $4q$-20, $4q$-$4t$) A1 11/4 or 2 ¾ or 2.75 OR M1 for correct substitution of p and t. M1 for $\frac{p-3t}{4} = q - t$ oe A1 11/4 or 2 ¾ or 2.75
18	(a)	$T=kx$; $150=6k$; $k=25$	$T=25x$	3	M1 for $T=kx$, k algebraic M1 subs $T=150$ and $x=6$ into $T = kx^n$ $(n \neq 0)$ A1 for $T=25x$ oe SC B1 $T \propto 25x$ oe

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
	(b)	$T = 25 \times 15 =$	375	1	B1 ft on k , $k \neq 1$
	(c)	$600 = 25x$; $x = 600 \div 25 =$	24	1	B1 ft on k , $k \neq 1$
19		$\frac{120}{570} \times 50 = 10.5263$ $\frac{250}{570} \times 50 = 21.9298$ $\frac{200}{570} \times 50 = 17.5438$	10 22 18	3	M1 method shown eg $\frac{120}{120+250+200} \times 50$ or one of 10.5(263) 21.9(298), 17.5(438) **OR** 570 ÷ 50 = 11.4 and one of 120 ÷ 11.4 =10.526…) , 270 ÷ 11.4 (=21.9298…) , 200 ÷ 11.4 (=17.5438…) A1 for 10.5(263), 21.9(298), 17.5(438) or 11, 22, 18 B1 correction to add to 50: 10, 22, 18 or 11, 22, 17
20		$x(x+1) + 4(2x–3) = (2x–3)(x+1)$ $x^2 + x + 8x – 12 = 2x^2 + 2x – 3x – 3$ $x^2 – 10x + 9 = 0$ $(x – 9)(x – 1) = 0$	$x = 1,9$	5	M1 for multiplying through by common denominator $(2x–3)(x+1)$ M1 (dep)for either $x^2 + x + 8x – 12$ or $2x^2 + 2x – 3x – 3$ oe A1 for correct quadratic (= 0) M1 for a correct method to solve 3 term quadratic A1 cao for both solutions
21		$AB^2 = 8^2 + 9^2 – 2\times8\times9\times\cos40$ $AB^2 = 64 + 81 – 144\times\cos40$ $AB^2 = 145 – 144 \times 0.766$ $AB^2 = 145 – 110.31… = 34.6896$ $AB = \sqrt{34.6796} = 5.8897877$	5.89	3	M1 Subs in Cos Rule: $8^2 + 9^2 – 2\times8\times9\times\cos40$ M1 correct order of evaluation of $8^2 + 9^2 – 2\times8\times9\times\cos40$ A1 cao 5.88 – 5.89 SC: Award B2 for one of $AB^2 = 241.03…$ or $AB = 15.525…$ (radians) $AB^2 = 28.50…$ or $AB = 5.33…$ (gradians)
22	(a)		24 30	2	B1 24 cao B1 30 cao
	(b)	Column $60 \leq w < 75$ at 10 Column $75 \leq w < 95$ at 4	10 4	2	B1 10 and correct width (tol $\pm\frac{1}{2}$ small square) B1 4 and correct width (tol $\pm\frac{1}{2}$ small square)

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
23 (a)		100.5	1	B1 cao
(b)		10.515	1	B1 cao
(c)	$\frac{100.5}{10.515}$	9.5577746	2	M1 for greatest distance divided by least time Where 100 < greatest distance ≤ 100.5, 10.51 ≤ least time < 10.52 A1 for 9.555 – 9.56
(d)	$\frac{99.5}{10.525}$	9.45368..	2	M1 for least distance divided by greatest time Where 99.5 ≤ least distance < 100, 10.52 < greatest time ≤10.53 A1 for 9.45 – 9.455
24	$\frac{4}{7}\times\frac{4}{7}\times\frac{3}{7} + \frac{3}{7}\times\frac{3}{7}\times\frac{4}{7} =$ $\frac{48+36}{343} = \frac{84}{343}$ But there are three ways this can be achieved: BBG, BGB, GBB So the probability is $\frac{84}{343}\times 3$ OR $1-\frac{64}{343}-\frac{27}{343}$ NB: 84/343 = 0.244897; 252/343 = 0.73469 $\frac{4}{7} = 0.57(142...)$, $\frac{3}{7} = 0.42(857...)$	$\frac{252}{343}$	3	M1 for $\frac{4}{7}\times\frac{4}{7}\times\frac{3}{7}$ or $\frac{3}{7}\times\frac{3}{7}\times\frac{4}{7}$ oe or $\left(\frac{4}{7}\right)^3$ oe or $\left(\frac{3}{7}\right)^3$ oe or $\frac{91}{343}$ or 0.10(49…) or 0.13(99….) M1 (indep) for identification of all 6 outcomes (M2 for $1-\left[\left(\frac{4}{7}\right)^3+\left(\frac{3}{7}\right)^3\right]$) oe A1 $\frac{252}{343}$, $\frac{36}{49}$, 0.73(469…) oe

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
25	(a)	$\frac{1}{3} \times \pi \times 5^2 \times 8 = \pi \times 25 \times 8 \div 3 =$ 209.4395	209–210	2	M1 for $\frac{1}{3} \times \pi \times 5^2 \times 8$ A1 for 209–210
	(b)	Base radius = $\frac{216}{360} \times 15 = 9$ Height = $\sqrt{(15^2 - 9^2)} =$	12	4	M1 for 216 ÷ 360 A1 for 9 M1 for $\sqrt{(15^2 - \text{“9”}^2)}$, where “9” < 15 A1 cao

PMT

edexcel

Mark Scheme (Results)

June 2011

Modular Mathematics (GCSE)
Unit 2: 5MB2H_01 (Higher)

PMT

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at www.edexcel.com.

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:
http://www.edexcel.com/Aboutus/contact-us/

June 2011
Publications Code UG028428
All the material in this publication is copyright
© Edexcel Ltd 2011

PMT

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labeling conventions.

ii*) select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect canceling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

12 Parts of questions
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers
Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

Guidance on the use of codes within this mark scheme
M1 – method mark A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

PMT

5MB2H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
1		$5 \times 6 \div 2$	15cm^2	3	M1 $5 \times 6 \div 2$ oe seen A1 15 B1 cm^2
2	(a)		$3y + 7x + 3$	1	B1 cao
	(b)		$2x(x - 2)$	2	B2 for $2x(x - 2)$. Accept $2x(x + -2)$. (B1 for $x(2x - 4)$ or $2(x^2 - 2x)$ or $2x$(linear expression in x) or $(x - 2)$(linear expression in x)
	(c)	$11-3x - 6$	$5-3x$	2	M1 for expansion of $-3(x + 2)$ A1 cao
	(d)	$3x^2+7x-18x-42$	$3x^2- 11x - 42$	2	M1 for 4 terms correct with or without signs or 3 out of exactly 4 terms correct (the terms may be in an expression or table) **OR** $x(3x+7)-6(3x+7)$ or $3x(x - 6) + 7(x - 6)$ A1 cao

PMT

5MB2H_01				
Question	**Working**	**Answer**	**Mark**	**Notes**
3 (a)	48=2 × 24 = 2 × 2 × 12 = 2 × 2 × 2 × 6 =2 × 2 × 2× 2 × 3 **OR** Use of factor trees	2 × 2 × 2 × 2 × 3	2	M1 for a systematic method that isolates 2 correct prime numbers correct divisions by a prime number **OR** an equivalent factor tree A1 for 2 × 2 × 2 × 2 × 3 or $2^4 \times 3$
(b)	Exeter coach after 20, 40, 60, 80, 100… Plymouth coach after 16, 32, 48, 64, 80, 96, … **OR** Timetable schedule for the coaches Exeter coach at 8.00, 8.20, 8.40, 9.00, 9.20… Plymouth coach at 8.00, 8.16, 8.32, 8.48, 9.04, 9.20… **OR** LCM of 16 and 20 is 80	9:20(am)	3	M1 for attempt to list multiples of 16 and 20 (at least first 3 of each, condone 1 addition error) M1(dep) for identifying their LCM A1 9:20 am (accept 9:20) **OR** M1 for attempt to draw up a timetable showing when the coaches will run (at least 3 extra times for each, condone 1 addition error) M1(dep) for identifying the first time common to both timetables A1 9:20 oe (do not accept 9:20pm) **OR** M1 attempt to find the LCM M1 for 2 × 2 × 2 × 2 × 5 oe seen A1 9:20 oe (do not accept 9:20pm)
4	10% = 6 6 × 4 = 24	24	2	M1 40 ÷ 100 × 60 oe or any complete method, eg 10% = 6, 6 × 4 A1 cao SC B1 for 36 or 84

5MB2H_01				
Question	**Working**	**Answer**	**Mark**	**Notes**
5 (a)	Table of values x = -1 0 1 2 3 y = 2 2 6 10 14 OR Using $y = mx + c$, gradient = 4, y intercept = 2	Line from (1, 2) to (3,14)	3	**(Table of values)** M1 for at least 2 correct attempts to find points by substituting values of x. M1 ft for plotting at least 2 of their points (any points plotted from their table must be correct) A1 for correct line between 1 and 3 **(No table of values)** M2 for at least 2 correct points (and no incorrect points) plotted OR line segment of $y = 4x + 2$ drawn (ignore any additional incorrect segments) (M1 for at least 3 correct points with no more than 2 incorrect points) A1 for correct line between -1 and 3 **(Use of $y = mx + c$)** M2 for at least 2 correct points (and no incorrect points) plotted OR line segment of $y = 4x + 2$ drawn (ignore any additional incorrect segments) (M1 for line drawn with gradient 4 OR line drawn with a y intercept of 2) A1 for correct line between 1 and 3
(b)(i)		$y = 4x + c$, $c \neq 2$	1	B1 Correct equation given.
(ii)		– 0.25	1	B1 Correct gradient given. Note – 0.25 could be written as $-\frac{1}{4}$ oe

PMT

5MB2H_01				
Question	**Working**	**Answer**	**Mark**	**Notes**
6	$6 \times 2 + 3 \times 2 = 12 + 6 = 18$ or $2 \times 5 + 4 \times 2 = 18$ $18 \div 2.5 = 7.2$ or $2.5 \times 7 = 17.5$ or $2.5 \times 8 = 20$ OR $6 \times 5 - 4 \times 3 = 18$ $18 \div 2.5 = 7.2$ or $2.5 \times 7 = 17.5$ or $2.5 \times 8 = 20$	8 packs	4	M2 for a complete method for finding the total area (can be implied by 18) (M1 for attempt to calculate at least one area, 6×2(=12) or 3×2 or 2×5(=10) or 4×2(=8) or 2×2 or 6×5 (=30) or 3×4(=12)) M1 for their area $\div 2.5$ or repeated addition of 2.5 to within 2.5 of their area C1 for clear communication that 8 full packs are required supported by their calculations, provided at least one of the first two method marks awarded SC B3 for 8 identified as the answer to the problem.
7	$\frac{3+7}{2}, \frac{8+5}{2}$	$(5, 6\frac{1}{2})$	2	M1 for either x or y coordinate correct or $\frac{3+7}{2}, \frac{8+5}{2}$, both seen but not correctly evaluated Al for $(5, 6\frac{1}{2})$ oe
8 (a) (b) (c)		p^8 h^5 x^6	3	B1 for p^8 B1 for h^5 B1 for x^6

PMT

5MB2H_01				
Question	**Working**	**Answer**	**Mark**	**Notes**
9	24 × 2 = 48 56 ÷ 8 × 5 = 35 or 24 × 2 = 48 48 ÷ 5 × 8 = 76.8	Monday	4	M1 for a correct method to obtain two comparable time periods or speeds. eg 24 × 2 (= 48 miles), 56 ÷ 2 (= 28 km) M1 for a correct method to convert between miles and km or mph and km/h eg –, – – – A1 for two correct values (using same units) for comparison C1 for day identified from a clear attempt to use comparable time periods and speeds/distances Note: day alone scores zero
10 (a)		$4n + 2$	2	B2 for oe (B1 for $4n+k$ where $k \neq 2$)
(b)	$2 \times 3^2 - 4$ 18 - 4	14	2	M1 for $2 \times 3^2 - 4$ A1 cao
11	360 ÷ 10 36 ÷ 2	18°	4	M1 for correct method to find the size of an exterior angle eg 360 ÷ 10 A1 for 36 M1 for '36' ÷ 2 or (180 – (180 – 36)) ÷ 2 A1 cao OR M1 for correct method to find the size of an interior angle eg(180 × 8) ÷ 10 A1 144 M1 (180 '144') ÷ 2 A1 cao

5MB2H_01				
Question	**Working**	**Answer**	**Mark**	**Notes**
12	$n^2 - 2n + 1 + n^2 + n^2 + 2n + 1 \equiv 3n^2 + 2$	Proof	2	M1 for correct expansion of either $(n - 1)^2$ or $(n + 1)^2$ or $n^2 - 2n + 1$ or $n^2 + 2n + 1$ seen A1 for $3n^2 + 2$ from correct working
13	$42 \div 2 = 21$ $180 - 90 - 21 = 69$ $69 \times 2 = 138$	138°	3	M1 for 90 seen. A1 for 138 (accept 222) C1 for The tangent to a circle is perpendicular (90°) to the radius (diameter) and Angles in a triangle add up to 180° **OR** The tangent to a circle is perpendicular (90°) to the radius (diameter) and Angles in a quadrilateral (4 sided shape) add up to 360°
14 (a)	$5\sqrt{9\times3}$	$15\sqrt{3}$	2	M1 for sight of $\sqrt{9\times3}$ or $\sqrt{9}\ \sqrt{3}$ or $3\sqrt{3}$ A1 for $15\sqrt{3}$ (accept $n = 15$)
(b)		$7\sqrt{3}$	2	M1 for $\frac{21\sqrt{3}}{\sqrt{3}\sqrt{3}}$ A1 for $7\sqrt{3}$ (accept $\frac{21\sqrt{3}}{3}$)

PMT

5MB2H_01				
Question	**Working**	**Answer**	**Mark**	**Notes**
15 (a)		3	1	B1 cao
(b)		$\frac{1}{5}$	2	M1 $\frac{1}{25^{\frac{1}{2}}}$ or $\sqrt{25} = 5$ A1 $\frac{1}{5}$ or 5^{-1} or 0.2
16 (a)	$\frac{2(y-6)}{(y-6)(y-2)}$	$\frac{2}{y-2}$	3	M1 $2(y-6)$ M1 $(y-6)(y-2)$ A1 cao
(b)	$\frac{3(x+5)-(x-4)}{(x-4)(x+5)}$	$\frac{3(x+5)-(x-4)}{(x-4)(x+5)}$	2	M1 for $3(x+5)$ or $3x+15$ or $3x+5$ or $-(x-4)$ or $-x+4$ or $-x-4$ or $(x-4)(x+5)$ A1 for $\frac{3(x+5)-(x-4)}{(x-4)(x+5)}$ oe

PMT

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481
Email publication.orders@edexcel.com

Order Code UG028428 June 2011

For more information on Edexcel qualifications, please visit
www.edexcel.com/quals

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual
Llywodraeth Cynulliad Cymru
Welsh Assembly Government
Rewarding Learning

PMT

edexcel

Mark Scheme (Results)

June 2012

GCSE Mathematics (2MB01) Higher
5MB2H (Non-Calculator) Paper 01

PMT

Edexcel and BTEC Qualifications
Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our qualifications website at www.edexcel.com. For information about our BTEC qualifications, please call 0844 576 0026, or visit our website at www.btec.co.uk.

If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

Ask The Expert can be accessed online at the following link:

http://www.edexcel.com/Aboutus/contact-us/

Alternatively, you can speak directly to the subject team at Pearson about Edexcel qualifications. Their contact details can be found on this link: www.edexcel.com/teachingservices

Pearson: helping people progress, everywhere
Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

June 2012
Publications Code UG032660
All the material in this publication is copyright
© Pearson Education Ltd 2012

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labeling conventions.

ii) *select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 **With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 **Follow through marks**

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 **Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect canceling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

10 **Probability**

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

PMT

11 Linear equations
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

12 Parts of questions
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers
Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

Guidance on the use of codes within this mark scheme
M1 – method mark A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

PMT

5MB2H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
*1		Angle *BED* = 68° 180 – (68 + 68) OR Angle *BEF* = 180 – 68 = 112 Angle *BED* = 180= 112 = 68 180 – (68 + 68)	x = 44°	4	M1 for Angle *BED* = 68° seen in the diagram or correctly used in the working space M1 for 180 – 2 × "68" A1 for x = 44° (accept angle *BDE* = 44°) [Note: 44° only seen in the body of the script without the "x =" gets M1M1A0] C1 (dep on at least M1) for alternate angles are equal (or equivalent) and one other reason; either angles in a triangle add to 180° or base angles of an isosceles triangle are equal OR for allied angles (or equivalent, eg co-interior) add up to 180° and angles on a straight line add up to 180° and one other reason; either angles in a triangle add to 180° or base angles of an isosceles triangle are equal
2			$4n + 2$	2	B2 for $4n + 2$ oe (eg $4\times n + 2$ or $n4 + 2$, …) or nth = $4n + 2$ (B1 for a linear expression in $4n$ e.g. $4n + a$ ($a \neq 2$) or $n = 4n + 2$) (B0 for $n = 4n$ and $n + 4$)

PMT

5MB2H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
3		35 ÷ (5 + 2) = 5 5 × 5 OR 5 2 10 4 15 6 20 8 25 10	25	2	M1 for 35 ÷ (5 + 2) or 35 ÷ 7 or 35 × $\frac{5}{7}$ A1 for 25 cao OR M1 for listing 5, 10, 15, ….. and 2, 4, 6, ….. to at least 15 and 6 A1 for 25 cao [SC: B1 for an answer of 25:10 if M0 scored]
4		$\frac{16}{40} + \frac{15}{40} = \frac{31}{40}$ OR | | 2 | 5 | | 3 | | 15 | | 8 | 16 | 40 | OR 0.4 + 0.375	$\frac{31}{40}$ or 0.775	2	M1 for attempt to write both fractions with a common denominator (a multiple of 40) with at least one of them correct A1 for $\frac{31}{40}$ oe OR M1 for 40 in the correct cell and 15 or 16 in the correct cell A1 for $\frac{31}{40}$ oe OR M1 for changing both fractions to decimals with both 0.4 and 0.375 seen A1 for 0.775

PMT

5MB2H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
5		$\frac{15}{100} \times 400 = 60$ $400 - 60 =$ OR $\frac{85}{100} \times 400$	340	3	M1 for $\frac{15}{100} \times 400$ (= 60) oe or 40 + 20 or 400 × 0.15 M1 (dep) for 400 – "60" A1 for 340 cao OR M1 for 100–15 (=85) M1 (dep) for $\frac{'85'}{100} \times 400$ or '0.85' × 400 A1 for 340 [SC: B1 for an answer of 460 if M0 scored]
6	(a)		$5(2a + 1)$	1	B1 for $5(2a + 1)$ or $5\times(2a + 1)$ or $(2a + 1)5$ or $5(1 + 2a)$, etc
	(b)	$5x + 35 + 3x - 6$	$8x + 29$	2	M1 for $5x + 35$ or $3x - 6$ or $8x$ or 29 A1 for $8x + 29$
	(c)		$3ab(a + 2b)$	2	B2 for $3ab(a + 2b)$ (B1 for correct partial factorisation $a(3ab + 6b^2)$ or $b(3a^2 + 6ab)$ or $3a(ab + 2b^2)$ or $3b(a^2 + 2ab)$ or $ab(3a + 6b)$ OR $3ab(ma + 2b)$ or $3ab(a + nb)$ where $m \neq 1$, $n \neq 2$) [B0 for partial factorisation using only an integer e.g. $3(a^2b + 2ab^2)$]

PMT

5MB2H_01

Question		Working	Answer	Mark	Notes
7		515 × 35 2575 15450 18025 10 × 515 = 5150 10 × 515 = 5150 10 × 515 = 5150 5 × 515 = 2575 18025 15000 + 2500 + 300 + 50 + 150 + 25 =18025	£180.25	4	M1 for 515 × 0.35 or 515 × 35 This may be implied from an incomplete method of multiplication M1 for a complete method with relative place value correct. Condone one multiplication error, addition not necessary Or for a complete grid, condone one multiplication error, addition not necessary Or for sight of a complete partitioning method. Condone one multiplication error final addition not necessary M1 (dep on the previous M1) for addition of appropriate elements of the calculation A1 for £180.25(p) or 18025p (with '£' sign deleted)

	5	1	5	×
	1 / 5	0 / 3	1 / 5	3
1	2 / 5	0 / 5	2 / 5	5
8	0	2	5	

	500	10	5
30	15000	300	150
5	2500	50	25

PMT

5MB2H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
		OR 3307 × 35 = 115 745 2792 × 35 = 97720 115745 – 97720			OR M1 for 3307 × 0.35 − 2792 × 0.35 or 3307 × 35 − 2792 × 35 M1 for a correct method of multiplication of at least one product, using digits 3307 and 35 or 2792 and 35 Condone one multiplication error, addition not necessary M1 (dep on the previous M1) for addition of appropriate elements of the calculation A1 for £180.25 or 18025p (with '£' sign deleted)

PMT

5MB2H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
8		**Triangular ends** $\frac{1}{2} \times 5 \times 12 = 30$ $\frac{1}{2} \times 5 \times 12 = 30$ **Base** $20 \times 5 = 100$ **Vertical face** $20 \times 12 = 240$ **Slant face** $20 \times 13 = 260$ **Total area** $= 30 + 30 + 100 + 240 + 260$ OR $(5 + 12 + 13) \times 20 + 2 \times \frac{1}{2} \times 5 \times 12$	660	3	M1 for $\frac{1}{2} \times 5 \times 12$ (= 30) or 20×5 (= 100) or 20×12 (= 240) or 20×13 (= 260) M1 (dep) for adding at least 3 areas found from correct methods (of no more than 5 faces) A1 cao OR M1 for $(5 + 12 + 13) \times 20$ or $\frac{1}{2} \times 5 \times 12$ (= 30) M1 (dep) for adding "$(5 + 12 + 13) \times 20$" to at least "$1 \times \frac{1}{2} \times 5 \times 12$" A1 cao Note: Sight of $\frac{1}{2} \times 5 \times 12 \times 20$ or 600 (ie a volume calculation) scores no marks

5MB2H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
9		Table of values x : –2, –1, 0, 1, 2 y : –7, –5, –3, –1, 1 OR Using $y = mx + c$ Gradient 2 intercept –3	Single line drawn from (–2, –7) to (2, 1)	3	**(Table of values)** M1 for at least 2 correct attempts to find points by substituting values of x. M1 (dep) ft for correctly plotting at least 2 of their points (any points plotted from their table must be plotted correctly) A1 for the correct line from (-2, -7) to (2, 1) OR **(No table of values)** M2 for at least 2 correct points (and no incorrect points) correctly plotted or for a line segment of the graph of $y = 2x – 3$ drawn (ignore any additional incorrect line segments) [M1 for at least 3 correct points plotted with no more than 2 incorrect points] A1 for the correct line from (-2, -7) to (2, 1) OR **(Use of $y = mx + c$)** M2 for a single straight line of gradient 2, passing through (0, -3) [M1 for a single straight line of gradient 2 or for a single straight line passing through (0, -3)] A1 for the correct line from (-2, -7) to (2, 1)

PMT

5MB2H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
10		$\frac{1}{2}$ (6 + 4) × 3 OR 6 × 3 – $\frac{1}{2}$ × 2 × 3 OR 4 × 3 + $\frac{1}{2}$ × 2 × 3	15 cm²	4	M2 for $\frac{1}{2}$ (6 + 4) × 3 oe A1 for 15 cao B1 for cm² OR M1 for 6 × 3 (=18) or $\frac{1}{2}$ × 2 × 3 (=3) M1 (dep on "6 × 3" and "$\frac{1}{2}$ × 2 × 3") for "18" – "3" from correct methods A1 for 15 cao B1 for cm² OR M1 for 4×3 (=12) or $\frac{1}{2}$ × 2 × 3 (=3) M1 (dep on "4×3" and "$\frac{1}{2}$ × 2 × 3") for "12" + "3" from correct methods A1 for 15 cao B1 for cm²

PMT

5MB2H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
11		10 000 × 0.8 8000 × 0.9 = 7200 OR 10 000 – $\frac{20}{100}$ × 10 000 = 8000 8000 – $\frac{10}{100}$ × 8 000 OR 10 000 × 0.8 × 0.9	7200	3	M1 for 10 000 × 0.8 (= 8000) M1 (dep) for "8000" × 0.9 (= 7200) A1 for £7200 cao OR M1 for 10 000 – $\frac{20}{100}$ × 10 000 oe M1 (dep) for "8000" – $\frac{10}{100}$ × "8 000" oe A1 7200 cao OR M1 for 0.8 × 0.9 (=0.72) M1 (dep)for 10 000 × "0.72" A1 for 7200 cao [SC: B1 for an answer of 7000 if M0 scored]
12	(a)		$x^2 - 3x - 40$	2	M1 for 3 terms correct (out of no more than 4 terms) from x^2, $5x$, $-8x$ and -40 or 4 terms x^2, $5x$, $8x$ and 40 (ignoring signs) A1 for $x^2 - 3x - 40$ [Note: $x^2 - 3x + 40$ and $x^2 + 3x - 40$ with no working get M0A0]
	(b)		$(x + 4)(x - 4)$	1	B1 for $(x + 4)(x - 4)$ oe
13	(a)		(0, 2, 4)	1	B1 cao
	(b)	$\left(\frac{0+6}{2}, \frac{2+2}{2}, \frac{4+4}{2}\right)$	(3, 2, 4)	2	M1 for an answer of (a, 2, 4) or (3, b, 4) or (3, 2, c) or for[("0" + 6) ÷2, ("2" + 2) ÷2, ("4" + 4) ÷2] This may be implied by their answer with no working seen A1 for (3, 2, 4) cao

PMT

5MB2H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
14	(a)	$x = 0.25555\ldots$ $10x = 2.555\ldots$ $100x = 25.5555\ldots$ $90x = 25 - 2 = 23$ OR $x = 0.25555\ldots$ $10x = 2.555\ldots$ $100x = 25.5555\ldots$ $9x = 2.3$ OR $0.2 + (x = 0.0555\ldots.)$ $10x = 0.555\ldots$ $100x = 5.5555\ldots$ $0.2 + 5/90$	$\frac{23}{90}$	3	M1 for $0.255(5\ldots.)$ or $0.2 + 0.055(5\ldots)$ This can be implied in subsequent working M1 for 2 correct recurring decimals which when subtracted will leave an integer or a terminating decimal number A1 for $\frac{23}{90}$ [Note: $\frac{2.3}{9}$ gets A0] [SC: B1 for an answer of $\frac{25}{99}$ oe $(= 0.\dot{2}\dot{5})$, with or without working if M0 scored]
	(b)	$\frac{12}{\sqrt{6}} \times \frac{\sqrt{6}}{\sqrt{6}} = \frac{12\sqrt{6}}{6}$	$2\sqrt{6}$	2	M1 for multiplying numerator and denominator by $\sqrt{6}$ A1 for $2\sqrt{6}$ (accept $\frac{2\sqrt{6}}{1}$)

PMT

5MB2H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
*15		180 –(90 + 20) = 70 2 × 70 = 140	Angle *TOR* =140º	4	M1 for angle *PTO* (*PRO*) = 90º or seeing it marked on the diagram with a right angle or as 90° M1 (dep) for 180 –(90 + 20) (=70°) or for 360 – (90 + 90 + 40) (=140) A1 for (angle) *TOR* = 140º or for 140º seen in the correct place in the diagram [140º alone without the '*TOR* =' gets A0] C1 (dep on at least M1) for angle between a tangent and a radius = 90° plus at least one other correct reason from: Sum of angles in a triangle is 180º Sum of angles in a quadrilateral is 360º Triangles *PTO* and *PRO* are congruent Tangents from a point are equal in length OR M1 for angle *PTO* (*PRO*) = 90º or seeing it marked on the diagram with a right angle or as 90° M1 for (180 – 40)÷2 (=70) and [180 – 2×(90 –"70")] (=140) A1 for (angle) *TOR* = 140º or for 140º seen in the correct place in the diagram [140º alone without the '*TOR* =' gets A0] C1 (dep on at least M1) for angle between a tangent and a radius = 90° plus at least one other correct reason from: Sum of angles in a triangle is 180º Base angles of an isosceles triangle are equal Triangles *PTO* and *PRO* are congruent Tangents from a point are equal in length
		(180 – 40)÷2 =70 180 – 2×(90 –70) = 140			

PMT

5MB2H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
16		$80 \times \frac{5}{8} = 50$ 50×2.75 OR $80 \times 2.75 = 220$ $220 \times \frac{5}{8}$	136 to 138 inc.	3	M1 for $80 \times \frac{5}{8}$ or $80 \div 1.6$ oe (= 50) M1 (indep) for "50" × 2.75 [accept "50"× 165 (=8250)] or "50" + "50" + "50"÷2 (=25) + "25" ÷2 (=12.5) Note: "50" is what is considered to be their speed in miles per hour calculated using an explicitly stated conversion factor A1 for an answer in the range 136 to 138 inc. OR M1 for 80 × 2.75 (=220) [accept 80 × 165 (=13200)] or 80 + 80 + 80÷2 (=40) + "40"÷2 (=20) M1 (indep) for "220" × $\frac{5}{8}$ Note: "220" is what is considered to be their distance in kilometres calculated using an explicitly stated conversion factor A1 for an answer in the range 136 to 138 inc.
17		$y = -\frac{1}{2}x + 2.5$ *mm'* = –1 shows gradient is 2 $y = 2x + c$ goes through (3, 7) $7 = 2 \times 3 + c$	$y = 2x + 1$	4	M1 for establishing gradient of original line is $-\frac{1}{2}$ or sight of $y = -\frac{1}{2}x + c$ M1 (indep) for (gradient of perpendicular) × "$-\frac{1}{2}$" = –1 This can be implied by a gradient of 2 or ft on "$-\frac{1}{2}$" M1 (dep on the previous M1) for substituting (3, 7) into y = "2"x + c A1 for $y = 2x + 1$ (or equivalent algebraic equation)

PMT

5MB2H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
18		$\frac{2x^2-9x-5}{4x^3+2x^2}$ $\frac{(2x+1)(x-5)}{2x^2(2x+1)}$	$\frac{x-5}{2x^2}$	3	M1 for factorising the numerator correctly M1 for fully factorising the denominator correctly A1 for $\frac{x-5}{2x^2}$ oe eg. $\frac{-5+x}{2x^2}$

PMT

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481
Email publication.orders@edexcel.com

Order Code UG032660 June 2012

For more information on Edexcel qualifications, please visit
www.edexcel.com/quals

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual
Llywodraeth Cynulliad Cymru
Welsh Assembly Government
Rewarding Learning

PMT

Mark Scheme (Results)

November 2015

Pearson Edexcel GCSE
In Mathematics B (2MB01)
Higher (Non-Calculator) Unit 2

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

November 2015
Publications Code UG042910
All the material in this publication is copyright
© Pearson Education Ltd 2015

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Note that in some cases a correct answer alone will not score marks unless supported by working; these situations are made clear in the mark scheme. Examiners should be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will award marks for the quality of written communication (QWC).
The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labelling conventions.

ii*) select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Partial answers shown (usually indicated in the ms by brackets) can be awarded the method mark associated with it (implied).
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks; transcription errors may also gain some credit. Send any such responses to review for the Team Leader to consider.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded (embedded answers).

12 Parts of questions
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers
Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

14 The detailed notes in the mark scheme, and in practice/training material for examiners, should be taken as precedents over the above notes.

Guidance on the use of codes within this mark scheme
M1 – method mark for appropriate method in the context of the question A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
1.			6 525 900 450	3	M1 for × 6 ÷ 4 oe or $\frac{6}{4}$ or 1.5 or 1 correct answer A2 for all correct (A1 for 2 or 3 correct)
2.			$8n + 24$	3	M1 for generating at least 3 terms of sequence 32, 40, 48 or sequence 4, 5, 6 M1 for $8n \pm k$ or $8(n \pm m)$ where $k \neq 24$, $m \neq 3$ A1 for $8n + 24$ or $8(n + 3)$ oe
3.			105 cm^3	4	M1 for writing factor pairs of 35, 21 or 15 M1 for "3" × "5" × "7" using their lengths dep on one at least one correct factor pair A1 ft for 105 (dep on one correct factor pair) B1 for cm^3

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
4.		$y = \frac{1}{2}x + 3$ x | −2 | −1 | 0 | 1 | 2 | 3 | 4 y | 2 | 2.5 | 3 | 3.5 | 4 | 4.5 | 5	Correct line from (−2, 2) to (4, 5)	3	**(Table of values / calculation of values)** M1 for at least 2 correct attempts to find points by substituting values of x. M1 ft for plotting at least 2 of their points (any points plotted from their table must be correctly plotted) A1 for correct line between $x = –2$ and $x = 4$ **(No table of values)** M1 for at least 2 correct points with no more than 2 incorrect points plotted M1 for at least 2 correct points (and no incorrect points) plotted **OR** line segment of $y = \frac{1}{2}x + 3$ drawn A1 for correct line between $x = –2$ and $x = 4$ **(Use of $y = mx + c$)** M1 for line drawn with gradient of $\frac{1}{2}$ **OR** line drawn with a y intercept of 3 M1 for line drawn with gradient of $\frac{1}{2}$ **AND** line drawn with a y intercept of 3 A1 for correct line between $x = –2$ and $x = 4$ **SC** : B2 for correct line from $x = 0$ to $x = 4$

PMT

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
5.			68	5	B1 for correct conversion of units M1 for 250 ÷ 25 (= 10) or 300 ÷ 20 (= 15) M1 for "10" × "15" M1 (dep on M2) for number of tiles ÷ 20, round up, × 8.5(0) A1 for 68 OR B1 for correct conversion of units M1 for 250 ÷ 20 (= 12.5) or 300 ÷ 25 (= 12) M1 for "12.5" × "12" M1 (dep on M2) for number of tiles ÷ 20, round up, × 8.5(0) A1 for 68 OR B1 for correct conversion of units M1 for 3 × 2.5 (= 7 .5) or 20 × 25 (= 500) M1 for "3 × 2.5" ÷ "0.2 × 0.25" or "300 × 250" ÷ "20 × 25" M1 (dep on M2) for number of tiles ÷ 20, round up, × 8.5(0) A1 for 68
*6		1, 4, 7, 10, 13 8, 6, 4, 2, 0	Explanation	2	M1 for listing at least 3 terms of both sequences C1 for Yes and explanation from fully correct working that 4 is in both sequences; numbers in A are increasing; numbers in B are decreasing

PMT

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
7			7.21 (am)	3	M1 for listing multiples 9,18,27,36 and 12,24,36 (condone 1 arithmetic error) or method to find LCM M1 for identifying 36 as LCM A1 cao OR M1 for listing times 6.54, 7.03, 7.12, 7.21 or for listing times 6.57, 7.09, 7.21 (condone one arithmetic error) M1 for listing times 6.54, 7.03, 7.12, 7.21 **and** 6.57, 7.09, 7.21 (condone one arithmetic error) A1 cao
8	(a)		4(3e +1)	1	B1 cao
	(b)		15c – 10d	1	B1 cao
	(c)		$28a^4b^3$	2	B2 for $28a^4b^3$ (B1 for two of 28, a^4 , b^3 as a product)
	(d)		$(x + 7)(x – 7)$	1	B1 cao
	(e)	$2y^2 – 6y + 7y – 21$	$2y^2 + y – 21$	2	M1 for 3 out of no more than 4 terms correct with correct signs or the 4 terms $2y^2$, $6y$, $7y$ and 21 seen, ignoring signs A1 cao

PMT

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
9		360 – 64 140 + 40 + 296 + 40 = 516 540 − 516 = 24	24°	4	M1 for 1 correct relevant angle calculation A1 for 24 cao C2 for all reasons (C1 (dep M1) for reason relating to parallel lines or parallelogram which is relevant to their chosen method) eg angles on a straight line add to 180° angles at a point add to 360° angles in a triangle add to 180° angles in a quadrilateral add to 360° alternate angles are equal corresponding angles are equal allied angles / co-interior angles add up to 180° opposite angles in a parallelogram are equal
10			$4\frac{8}{35}$	3	M1 for converting both fractions to get a common denominator of a multiple of 35 with at least one correctly converted. M1 (dep on M1) for $3+\frac{"28"}{35}+\frac{"15"}{35}\left(=3\frac{43}{35}\right)$ oe A1 for $4\frac{8}{35}$ cao

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
11	(a)		7.8×10^{-3}	1	B1 cao
	(b)		6 710 000	1	B1 cao
	(c)	$-9,\ 0.9,\ 1,\ 3$	$-9,\ 0.9,\ 9^0,\ 9^{\frac{1}{2}}$	2	M1 for either $9^{\frac{1}{2}}$ or 9^0 evaluated correctly A1 for fully correct list from correct working, accept original numbers or evaluated. (SC: B1 for 1 error in position or a correct list in reverse order)
12.	(a)		(0, 5, 5)	1	B1 cao
	(b)		$(2\frac{1}{2}, 5, 2\frac{1}{2})$	1	B1 for $(2\frac{1}{2}, 5, 2\frac{1}{2})$ oe
13.			120	3	M1 for $80\% = 480$ or $\frac{480}{80}(=6)$ or $\frac{480}{80} \times 100\ (=600)$ oe M1 for $600 - 480$ or $\frac{480}{80} \times 20$ oe A1 cao
14.			$A = 9x^2 + 19x - 6$	4	B1 for one of $5x - 2$ or x found M1 for correct method to find area of one relevant rectangle. M1 for complete method to find whole area or simplified expression $9x^2 + 19x - 6$ or correct but not simplified formula A1 for correct, simplified formula $A = 9x^2 + 19x - 6$

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
15.			$y = \frac{3}{2}x - \frac{5}{2}$	4	M1 for attempt to find gradient of AB M1 (dep) for attempt to find gradient of perpendicular line eg use of $-\frac{1}{m}$ M1 for substituting $x = 3, y = 2$ into $y =$ "m" $x + c$ A1 for complete correct equation $y = \frac{3}{2}x - \frac{5}{2}$ oe
16.			$2\sqrt{7}$	3	M1 for multiplying numerator and denominator by $\sqrt{7}$ M1 for correct method to expand $(4+\sqrt{2})(4-\sqrt{2})$ with 3 out of no more than 4 terms correct with correct signs or the 4 terms seen, ignoring signs A1 for $2\sqrt{7}$ (accept $\sqrt{28}$)
17.		$\frac{(x-5)(x+3)}{(x+3)(2x+1)}$	$\frac{x-5}{2x+1}$	3	M1 for correct factorisation of numerator M1 for correct factorisation of denominator A1 for $\frac{x-5}{2x+1}$

PMT

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom

Question 14

Students who gave the exact value of either tan30º or sin60º gained the first mark. Various strategies were used to find these values. Some students worked them out by drawing an appropriate triangle; some used patterns in a table with angles of 0º, 30º, 45º, 60º and 90º and some simply recalled them from memory. Sometimes sin30º ÷ cos30º was used to find the value of tan30º. However, the strategies used were not always successful and incorrect values were very common. When only one of the two values was correct it was more often $\sin 60^\circ = \frac{\sqrt{3}}{2}$. In some responses the values were not labelled. If only one value was correct examiners had to assume from the calculation shown that the order was tan30º×sin60º. Some of the students who

got to $\frac{1}{\sqrt{3}} \times \frac{\sqrt{3}}{2}$ did not go on to get full marks because they gave $\frac{\sqrt{3}}{2\sqrt{3}}$ as the final answer or simplified it to 2. The question required the answer to be given in its simplest form. Some students used incorrect values such as $\tan 30^\circ = 1$ and $\sin 60^\circ = 0.5$ which just happened to give an answer of 0.5. When the "correct" answer clearly came from incorrect working a mark of 0 was awarded.

Question 18

In part (a) many students were able to gain at least one of the two marks for writing $\sqrt{12}$ as $\sqrt{4} \times \sqrt{3}$ or $2\sqrt{3}$. Although most went on to get the correct answer of $3\sqrt{3}$ it was not unusual to see $\sqrt{3} + 2\sqrt{3}$ followed by an answer of $2\sqrt{3}$ or $2\sqrt{6}$. Common mistakes were to write $\sqrt{12}$ as $4\sqrt{3}$ or to write $\sqrt{3} + \sqrt{12}$ as $\sqrt{15}$ and give an answer of $5\sqrt{3}$.

Some students started part (b) by attempting to deal with the power and many were able to obtain $\frac{1}{27\sqrt{3}}$. Different strategies were used to find $\left(\sqrt{3}\right)^7$. Writing out $\sqrt{3}$ seven times and working through the multiplication in a systematic way tended to be a successful approach. The second step for these students was to rationalise the denominator. Instead of multiplying numerator and denominator by $\sqrt{3}$ some students chose to multiply by $27\sqrt{3}$ which resulted in $\frac{27\sqrt{3}}{2187}$. This fraction is not in the required form and the students gained only two of the three marks unless they simplified it to $\frac{\sqrt{3}}{81}$. Students who attempted to deal with the power first but went wrong, obtaining $\frac{7}{7\sqrt{3}}$ for example, sometimes went on to gain a method mark for rationalising the denominator of their fraction. Some students chose to start by rationalising the denominator. Usually this involved multiplying both the numerator and denominator of $\frac{1}{\sqrt{3}}$ by $\sqrt{3}$ to give $\frac{\sqrt{3}}{3}$. Many students could get no further but those who simplified $\left(\frac{\sqrt{3}}{3}\right)^7$ to $\frac{\sqrt{2187}}{2187}$ scored full marks as this fraction is in the required form. Some students started by multiplying the numerator and denominator of $\left(\frac{1}{\sqrt{3}}\right)^7$ by $\left(\sqrt{3}\right)^7$ and frequently got into difficulties. Rationalising the denominator seemed to be much more accessible to students than dealing with the power of 7.

Question 16

It was pleasing to see many students, in part (a), attempting to multiply the given expression by $\frac{\sqrt{11}}{\sqrt{11}}$, some just wrote $\times \sqrt{11}$ but credit was given if the intention was clear. Student should be encouraged to write $\frac{\times\sqrt{11}}{\times\sqrt{11}}$. Many students correctly worked out the surd product but then failed to fully simplify, leaving $\frac{22\sqrt{11}}{11}$ as their answer. This was not awarded the final accuracy mark.

In part (b), students who knew to multiply numerator and denominator by $2\sqrt{3}+1$, gained one mark but were often unable to complete the simplification to gain further credit.

Question 20

Those students who started by writing $\sqrt{180}$ as $6\sqrt{5}$ to simplify the numerator tended to be the more successful because doing so made the subsequent arithmetic easier. Rather surprisingly some students simplified the numerator only as far as $6\sqrt{5}-2\sqrt{5}$ before rationalising the denominator. Sometimes the numerator was simplified but then no attempt was made to rationalise the denominator. Students who started by attempting to rationalise the denominator often made errors when expanding $(\sqrt{180}-2\sqrt{5})(5\sqrt{5}+5)$. Some students used an incorrect method when attempting to rationalise the denominator, e.g. multiplying the numerator and denominator by $5\sqrt{5}-5$, failing to link the process to the difference of two squares. Having got as far as $(100+20\sqrt{5})/100$ some students did not score the final mark because their attempt at simplifying did not lead to $1+\sqrt{5}/5$.

Question 19

There were some excellent responses to this question from the most able students taking this paper but these were few and far between. Able students who were unable to get as far as the given result were sometimes able to score some marks for successfully finding expressions for the area of part of a hexagon or for a whole hexagon, whether it be *ABCDEF* or *FGHIJK*. Where this applied, students usually used trigonometry to find the height of a constituent triangle and then its area by using $\frac{1}{2} \times$ base $\times$ height or they found an area using the formula $\frac{1}{2}ab\sin C$.

Some weaker students used the given result and tried to justify it by using substitution or by algebraic manipulation whilst other students seemed to think that the question revolved around finding angles of a polygon. No marks could be awarded for these approaches.

Question 19

Writing $\sqrt{12}$ as $2\sqrt{3}$ was sometimes seen as the first step and sometimes it was seen later in the process. Some of the students who started by writing $\sqrt{12}$ as $2\sqrt{3}$ and simplified the numerator of the fraction to $8+2\sqrt{3}$ made no further progress. Those students who realised that they needed to

rationalise the denominator were not always able to show a correct method to do so. Attempts at multiplying both the numerator and the denominator by $5+\sqrt{3}$ or by $\sqrt{3}$ were often seen and gained no credit. Those who did appreciate the need to multiply numerator and denominator by $5-\sqrt{3}$ often went on to expand the terms correctly although some students made sign errors. Having got as far as $\frac{(34+2\sqrt{3})}{22}$ some students gave this as the final answer and did not score the final mark because the question asked for an answer in the form $\frac{(a+\sqrt{3})}{b}$.

Question 19

Relatively few students could demonstrate the necessary skills of algebraic manipulation to solve the equation and give the answer in the required form. Many students were able to correctly write the two fractions with a common denominator and gain the first mark but a significant number were then unable to carry the algebraic solution any further. Those that did reduce the equation to a 3 term quadratic often made errors when rearranging and did not get the second mark. Substitution into the quadratic equation formula was generally done well but some students attempted to use completing the square and this was done less well due to not dealing with the coefficient of 2 correctly. Some students who did not get a correct quadratic equation were still able to gain the third mark for dealing correctly with their 3 term quadratic equation. The final step to write $\frac{10 \pm \sqrt{60}}{4}$ in the required form proved difficult with $\frac{5 \pm \sqrt{30}}{2}$ a common incorrect answer.

Question 19

Some students correctly wrote the two fractions with a common denominator and gained the first mark but many were then unable to carry the algebraic solution any further. Those that did reduce the equation to a 3-term quadratic sometimes made errors when rearranging and did not get the second mark. Substitution into the quadratic equation formula was generally done well. Some students who did not get a correct quadratic equation were still able to gain the third mark for dealing correctly with their 3-term quadratic equation. The first of the two accuracy marks was awarded for two correct values of x, e.g. $\frac{-4 \pm \sqrt{32}}{8}$. Some students gave this as the final answer and others made errors when attempting to write it in the required form. Only a few students could demonstrate the necessary skills of algebraic manipulation to solve the equation and give the answer in the required form.

Question 16

Most students started with a process to find the surface area of the sphere. Some made no further progress but many continued with $4\pi r^2 = 200\pi$ or $4r^2 = 200$ and went on to find the radius and gained the first three marks. Inconsistent use of π was a problem for some. It was not uncommon to see a surface area of 200π followed by $4\pi r^2 = 200$. After showing a correct process to find the radius a significant number of students then failed to get the accuracy mark for finding the diameter of the sphere in the required form. Many could not write $\sqrt{50}$ as $5\sqrt{2}$ with $\sqrt{50} = 10\sqrt{5}$ a common mistake and the radius was often given as the final answer . Instead of starting with a process to find the surface area of the sphere some students started by forming the equation $\frac{3}{8} \times 4\pi r^2 = 75\pi$.

Question 23

This question involved writing two fractions with a common denominator, rationalising and simplifying surds. There were many valiant attempts achieving varying degrees of success and not many blank responses. Many students started with a method to write the two fractions with a common denominator and gained the first mark. A common mistake, which prevented the award of the second mark, was incorrectly simplifying $\frac{9}{4\sqrt{3}-3}-\frac{8-2\sqrt{3}}{4\sqrt{3}-3}$ to $\frac{1-2\sqrt{3}}{4\sqrt{3}-3}$. Some students were unable to continue the solution any further and gave their single fraction as the final answer. A good number of students though went on to gain the third mark for a method to rationalise the denominator with some giving a fully correct solution. The alternative approach used by a good proportion of students was to start by rationalising the denominator of each fraction before writing the fractions with a common denominator. The main obstacle to a successful outcome for these students was making errors when manipulating the surds with common mistakes being $\sqrt{3} \times \sqrt{3} = 9$ and $\sqrt{3}(3\sqrt{3}) = 3\sqrt{3} + 3$. Those who used this approach were more likely to give a fully correct answer. Some students used a hybrid approach, rationalising one fraction before trying to achieve a common denominator and then rationalising again. This route had comparable success with the other routes.

Question 24

The final question on the paper was a challenging shape problem requiring candidates to find an expression for the area of an unfamiliar shape. The most common approach was to split the shape into a square and 4 identical triangles. By doing this and correctly identifying an angle inside or outside of the triangle, candidates were able to gain the first mark. To gain the second and third they needed an expression for the area of one of the triangles, or the length of the side of the square. The former was awarded more regularly, as many spotted that the triangles were right-angles and isosceles. The length of the square proved more challenging as it involved using Pythagoras and surds. The final process mark was to put it altogether to form an expression for the total area, before simplifying it to the given form. Although there were some excellent responses, some candidates were let down by the accuracy of their algebra, and we also saw candidates guessing the value of p, but without supporting algebra, this gained no credit. A successful method seen, other than the square and four triangles split, was to split the shape into 8 triangles, 2 trapezia and 1 rectangle. Several responses started with finding the hypotenuse of the small triangle or the area but did not get beyond this. There was evidence that candidates were not very confident at manipulating numbers inside a square root. Some had trouble with

simplifying $\sqrt{2a^2}$ to $a\sqrt{2}$ after using Pythagoras theorem to find the length of the hypotenuse of the small triangle.
If candidates did correctly expand the expression for the area of the square and added on the triangle areas, they were usually able to work backwards from the final answer to rearrange their expression correctly.

Summary

Based on the performance on this paper, students should:

- ensure they are using efficient calculator methods
- take extra care when working in algebra and show each step carefully, and remember the importance of brackets and the length of the lid on root signs
- spend more time interpreting statistical diagrams such as cumulative frequency curves and histograms with unequal class intervals
- read the demand of questions carefully so as to fully understand what the question expects in terms of approach for answering, and also the form in which the answer is expected
- remember that trial and improvement methods should be discouraged
- work on geometrical topics such as the sine rule and ensure that angles are labelled correctly, either using 3 letter notation or on the diagram if introducing a new variable such as x
- ensure candidates are fully prepared for the planned withdrawal of the exam aid for future series

Question 17

Part (a) was generally answered quite well. Most students who knew about rationalising a denominator chose to multiply numerator and denominator by $\sqrt{7}$ although a few used $-\sqrt{7}$ instead. The multiplying was usually carried out correctly. However, $\frac{\sqrt{7}}{7}$ was sometimes incorrectly simplified, often to 1, and the mark could not be awarded. It was common to see $\frac{1\sqrt{7}}{7}$ as the final answer and this was awarded one mark. A common error was to find the reciprocal rather than rationalise. Incorrect manipulation of the surd was seen when students wrote $\sqrt{7}=\sqrt{4+3}=2\sqrt{3}$ or $2+\sqrt{3}$.

In part (b) many students were able to gain at least one of the two marks for writing $\sqrt{80}$ as $\sqrt{16\times5}$ or $\sqrt{16}\times\sqrt{5}$ or $4\sqrt{5}$. Although many went on to get the correct answer of $3\sqrt{5}$ it was not unusual to see $4\sqrt{5}-\sqrt{5}$ followed by an answer of 4 or to see $4\sqrt{5}-\sqrt{5}$ given as the final answer. Writing $\sqrt{80}$ as $\sqrt{10\times8}$ or $\sqrt{4\times20}$ was a first step that often led to no marks being awarded although some students did get to $4\sqrt{5}$. A common mistake was to write $\sqrt{80}-\sqrt{5}$ as $\sqrt{75}$ and this often resulted in an answer of $5\sqrt{3}$.

Question 15

A slightly different style of surds question to the ones we have seen previously, but it was approached well by students. A good number of the stronger students gained the first mark for a method to rationalise, and many of these then gained the second mark. However, some who had the correct algebra leading to the answer, attempted to simplify further often to expressions such as $\frac{\sqrt{a}}{a}$. This further incorrect working was not ignored and lost the A mark. Some students worked with a prime number, such as 3, in place of a. These students were able to gain the M1 if their method to rationalise was correct. The most common error was in trying to rationalise using $\sqrt{a}+1$ rather than $\sqrt{a}-1$.

Question 14

Some learners gained the first mark for stating the value of sin30 or the value of sin45. Those who did not know these values were still able to gain the second mark for a correct sine rule statement such as $\frac{AB}{\sin 45} = \frac{3\sqrt{2}}{\sin 30}$. More than half of those who gained the first two marks did not get a correct final answer, some because they had an incorrect value for sin30 or sin45, some because they could not rearrange their sine rule statement correctly and some because they could not combine the surds correctly. Many learners showed a lack of understanding of the mathematical requirements of the question with some using right-angled trigonometric calculations.

Question 16

Part (a) was answered quite well by those with some understanding of surds and a good proportion of learners gained the method mark for multiplying both the numerator and the denominator by $\sqrt{5}$. Some failed to gain the accuracy mark because they gave $\frac{15\sqrt{5}}{5}$ as the final answer when the question asked for the answer in its simplest form, others because they made errors when trying to write it in its simplest form.

In part (b), writing $\sqrt{75}$ as $5\sqrt{3}$ was sometimes seen as the first step and sometimes it was seen later in the process. Some learners started by writing $\sqrt{75}$ as $5\sqrt{3}$ and simplified the numerator of the fraction to $5\sqrt{3}-2$ but no further progress. Those that realised they needed to rationalise the denominator were not always able to show a correct method to do so. Attempts at multiplying both the numerator and the denominator by $1+2\sqrt{3}$ or by $\sqrt{3}$ were sometimes seen and gained no credit. Some of those that did multiply the numerator and denominator by $1-2\sqrt{3}$ went on to expand the terms correctly but errors were frequently made, often when multiplying $5\sqrt{3}$ by $-2\sqrt{3}$. One error in the numerator or denominator was condoned for the third mark. Those learners who expanded all the terms correctly were usually able to give a correct final answer.

Question 16

Part (a) was answered well by those with some understanding of surds and many responses gained the method mark for multiplying both the numerator and the denominator by $\sqrt{7}$ or, less commonly, by $-\sqrt{7}$. The multiplying was usually carried out correctly. The question asked for the answer in its simplest form and so the accuracy mark was not awarded when $\frac{35\sqrt{7}}{7}$ was given as the final answer, which was very common, or when it was simplified incorrectly.

In part (b), writing $\sqrt{27}$ as $3\sqrt{3}$ gained the first mark. This was sometimes seen as the first step and sometimes it was seen later in the process. Some responses started by writing $\sqrt{27}$ as $3\sqrt{3}$ and simplified the numerator of the fraction to $3\sqrt{3}-1$ but made no further progress. Those that realised they needed to rationalise the denominator were often able to show a correct method to do so. Attempts at multiplying both the numerator and the denominator by $2-\sqrt{3}$ or by $\sqrt{3}$ were often seen and gained no credit. Many of those who did multiply the numerator and denominator by $2+\sqrt{3}$ went on to expand the terms correctly but errors were frequently made, often when multiplying $3\sqrt{3}$ by $\sqrt{3}$. One error in the numerator or denominator was condoned for the third mark. Those responses that expanded all the terms correctly were usually able to give a correct final answer although some lost the accuracy mark due to errors in collecting like terms.