Chhetri Academy
chhetriacademy.com

GCSE MATHS

Algebra

Edexcel 1MA1 | Subtopic collection Topic 15

Laws of indices in algebra

Higher tier

Answers and examiner comments

48	2003-2025	Higher
question groups	available papers	tier

Using this booklet

1. Answer numbers match the companion question booklet.
2. Follow the original question number on marking tables.
3. Examiner comments are included where available.

Name _____ **Date** _____

Some entries use full source marking material in an appendix. Any unavailable mark schemes are identified beside the relevant answer.

Original question 13

The full source mark scheme begins on page 70. Use the original paper code, tier and question number shown above.

Original question 4

No matching mark scheme was available in the supplied archive for this question.

Original question 7

The full source mark scheme begins on page 124. Use the original paper code, tier and question number shown above.

Original question 6

The full source mark scheme begins on page 175. Use the original paper code, tier and question number shown above.

Original question 18

The full source mark scheme begins on page 175. Use the original paper code, tier and question number shown above.

Original question 5

No matching mark scheme was available in the supplied archive for this question.

Original question 16

No matching mark scheme was available in the supplied archive for this question.

14	(a)		12	1	B1 cao
	(b)		24	1	B1 cao
	(c)		49	1	B1 cao

16	(a)		Diagram *(overlay)*	2	B2 within guidelines of the overlay (B1 for exactly one given angle correctly drawn within guidelines of overlay)
	(b)		90	1	B1 for an angle in range 86 to 94 **or** ft 'angle' measured correctly within $\pm 2^o$

1380/1F

Question		Working	Answer	Mark	Notes
15	(a)		m^7	1	B1 for m^7, (accept m^{3+4})
	(b)		p^4	1	B1 for p^4, (accept p^{7-3})
	(c)		$12\,x^3y^5$	2	B2 cao (B1 for two of 12, x^3, y^5, ignore $\times$ signs)

22	(a)	p^{5+4}	p^9	1	B1 (accept p^{5+4})
	(b)	q^{5-2}	q^3	1	B1 (accept q^{5-2})
	(c)		$2u$	2	B2 (accept $2t^0u$, $2t^0u^1$oe) (B1 for 2 correct terms from 2, t^0 and u oe eg u^1)
	(d)		$3wy^3$	2	B2 cao (B1 for 2 correct terms from 3, w and y^3 oe) NB: accept w^1 for w.
	(e)		x^{-2} x^0 $x^{\frac{1}{2}}$ x x^2	2	B2 cao (B1 for any 4 in relative correct order, or all correct but in reverse order)

1380_4H

6	(a)		$6x + 5y$	2	B2 (B1 for either $6x$ or $5y$ seen)
	(b)	$2x = 10 - 3 = 7$ $x = 7 \div 2$	3.5	2	M1 for $2x = 10 - 3$ or $2x = 7$ or $(10 - 3) \div 2$ A1 for 3.5 oe
	(c)(i)		c^{11}	2	B1 accept c^{5+6}
	(ii)		e^{8}		B1 accept e^{12-4}

1380_4H

1380/4H

25	(a)	$(2x + 4y)(4x - 5y)$ $= 8x^2 - 10xy + 16xy - 20y^2$	$8x^2 + 6xy - 20y^2$	2	B2 cao (B1 for 3 out of 4 terms correct or all 4 correct ignoring signs)
	(b)		$x + 10$	1	B1 for $x + 10$ or $(x + 10)$ or $(x + 10)^1$
	(c)	$= \frac{(x+5)(x-5)}{(x+5)(x+2)}$	$\frac{x-5}{x+2}$	3	M1 for $(x+5)(x-5)$ M1 (indep) for $(x+5)(x+2)$ A1 cao
	(d)	$x^2 + 6x - 2 = (x + 3)^2 - 9 - 2$	$p = 3$ $q = -11$	2	M1 for $(x + 3)^2 \pm k$ or $x^2 + 2px + p^2 + q$ oe or $p = 3$ or $q = -11$ A1 cao

1380_4H

1380/4H

12	(a)		1	1	B1 cao
	(b)		$8x^{18}y^{-3}$	2	M1 for $2^3x^{3\times6}y^{3\times-1}$ **or** $8x^{18}y^{k}$ (k ≠ 0) **Or** $8x^{k}y^{-3}$ (k ≠ 0) **or** $kx^{18}y^{-3}$ (k ≠ 0, k ≠ 1) **or** $2x^6y^{-1} \times 2x^6y^{-1} \times 2x^6y^{-1}$ A1 cao

5MB2H_01

5MB2H_01

11	(a)		$3x + 6$	2	M1 for attempted expansion of the bracket eg $3\times x$ and 3×2 seen or $3x + k$ or $kx + 6$ A1 for $3x + 6$
	(b)		$6xy(2x^2 - 3y)$	2	M1 or $6xy$ (two terms involving x and/or y) or correct partial factorisation by taking out two from 6 (or 3 or 2) or x or y A1 cao
	(c)	$2x^2 + 8x - 3x - 12$	$2x^2 + 5x - 12$	2	M1 for 3 out of 4 correct terms with correct signs, or all 4 terms ignoring signs A1 cao
	(d)		$10\,x^7\,y^5$	2	B2 for $10\,x^7\,y^5$ (B1 for product of two of 5×2 oe, x^{4+3}, y^{3+2} ignore $\times$ signs)

5MB2H_01

Question		Working	Answer	Mark	Notes
12	(a)(i)	w^{6+4}	w^{10}	2	B1 accept w^{6+4}
	(ii)	h^{8-3}	h^5		B1 accept h^{8-3}
	(b)	$(12 \div 3)(x \div x^2)(y^3 \div y^3)$	$\frac{4}{x}$	2	B2 for $\frac{4}{x}$ or $4x^{-1}$ (B1 for any one from: $12 \div 3$ in numerator OR x^{1-2} in numerator or x^{2-1} in denominator OR y^{3-3} in numerator OR y^3 cancelled in both numerator and denominator

<table>
<tr><th colspan="2">Question</th><th>Working</th><th>Answer</th><th>Mark</th><th>Notes</th></tr>
<tr><td>18</td><td>(a)</td><td></td><td>c^8k^{20}</td><td>1</td><td>B1</td></tr>
<tr><td></td><td>(b)</td><td>$12x^2 - 3x + 20x - 5$
OR
<table><tr><td>×</td><td>3x</td><td>(+) 5</td></tr><tr><td>4x</td><td>$12x^2$</td><td>(+)$20x$</td></tr><tr><td>−1</td><td>$-3x$</td><td>-5</td></tr></table></td><td>$12x^2 + 17x - 5$</td><td>2</td><td>B2 for fully correct
(B1 for 3 out of 4 terms correct in working including signs OR 4 terms correct, ignore signs. In a grid the $20x$ need not be signed)</td></tr>
<tr><td></td><td>(c)</td><td>$(x - 5)(x + 2) = 0$</td><td>5 and −2</td><td>3</td><td>M1 for $(x \pm 5)(x \pm 2)$
A1 for $(x - 5)(x + 2)$ (= 0)
B1 ft (dep on M1) for $x = 5$ and −2
or
M1 for correct substitution in formula allow sign errors in b and c
M1 for reduction to $\frac{3 \pm \sqrt{49}}{2}$
A1 for 5 and −2
or
M1 for $(x - \frac{3}{2})^2 - \left(\frac{3}{2}\right)^2 - 10 = 0$
M1 for $\frac{3}{2} \pm \sqrt{\frac{49}{4}}$
A1 for 5 and −2

or
T&I B3 both roots
(B1 one root)</td></tr>
</table>

1380_4H

Original question 8

The full source mark scheme begins on page 236. Use the original paper code, tier and question number shown above.

2	(a)		$20\,fg$	1	B1 in any acceptable order
	(b)	$9a + 3(8 - 2a)$ $= 9a + 24 - 6a$	$3a + 24$	2	M1 for 3×8 and $\pm 3 \times 2a$ oe within at most 4 terms seen A1 cao
	(c)		c^8	1	B1 cao
	(d)		x^{15}	1	B1 cao
	(e)		$7(y + 3)$	1	B1 cao

5MB2H_01

Original question 16

The full source mark scheme begins on page 249. Use the original paper code, tier and question number shown above.

15	(a)		m^2	1	B1 for m^2 or m^{5-3}
	(b)		$5x^6y^4$	2	M1 for $x^{4+2}y^a$ or x^by^{3+1} A1 cao

1MA0_1H

9	(a)		a^9	1	B1 for a^{4+5} or a^9
	(b)		$9e^5f^6$	2	B2 cao (B1 for two of 9, e^{6-1}, f^{8-2} as a product)
	(c)		3	1	B1 (accept ± 3 but not just –3)

1MA0_1H					
Question		**Working**	**Answer**	**Mark**	**Notes**
*10		Angle *AED* = 38 alternate angles are equal Angle *ADE* = (180 – 38) ÷ 2 = 71 *x* = 180 – 71 base angles of an isosceles triangle are equal angles in a triangle add up to 180 angles on a straight line sum to 180 **OR** angle *AEF* = 142 allied angles/co-interior angles add up to 180 *ADE* = 142 ÷ 2 = 71 base angles of an isosceles triangle are equal exterior angle of a triangle is equal to the sum of the interior opposite angles , *x* = 180 – 71 angles in a straight line add to 180 **OR** Angle *AED* = 38 alternate angles are equal for angles *BAE* and *AED* **and** *BAD* and *ADC* (*x*) Angle *DAE*= (180 – 38) ÷ 2 = 71 base angles of an isosceles triangle are equal angles in a triangle add up to 180 **Or** Angle *AED* = 38 alternate angles are equal Angle *ADE* = (180 – 38) ÷ 2 = 71 base angles of an isosceles triangle are equal and angles in a triangle sum to 180 *x* = 38 + 71 alternate angles *BAD* and *ADC*(*x*) are equal	*x* = 109	4	B1 for angle *AED* = 38 or *AEF* = 142 M1 for a complete method to find one of the base angles of the isosceles triangle C2 (dep M1) for *x* = 109 with complete reasons (C1 (dep M1) for one reason correctly used and stated)

1MA0_1H

5	(a)		$3x - y$	2	M1 for $3x$ or $-y$ A1 for $3x - y$ or $-y + 3x$
	(b)		$2 - 3x$	1	B1 for $2 - 3x$ or $-3x + 2$
	(c)		t^{11}	1	B1 cao
	(d)		y^4	1	B1 cao
	(e)		c^{12}	1	B1 cao

2	(a)		$5a + 3b$	2	M1 for partial simplification $5a$ **or** $+3b$ A1 cao
	(b)		$5m + 10$	1	B1 cao
	(c)		a^9	1	B1 cao

PAPER: 5MB2H_01

PAPER: 1MA0_2H

Question	Working	Answer	Mark	Notes
11 (a)		x^{10}	1	B1 cao
(b)		m^{12}	1	B1 cao
(c)		$3a^{-4}f^{6}$	2	B2 for $3a^{-4}f^{6}$ **or** $\frac{3f^{6}}{a^{4}}$ (B1 for any two from 3, a^{-4} or $\frac{1}{a^{4}}$, f^{6} in a product)

PAPER: 5MB2H_01

Question		Working	Answer	Mark	Notes
12	(a)		$2a(3b + 5c)$	2	B2 cao (B1 for $a(6b + 10c)$ or $2(3ab + 5ac)$ or $2a$(linear term in b and c))
	(b)		$x^2 + 2x - 35$	2	M1 for 3 terms out of 4 correct including signs or all 4 terms correct ignoring signs A1 cao
	(c)		$\frac{2t^4}{m^2}$	2	B2 for $2m^{-2}t^4$ oe (B1 $\frac{2t^4}{m^n}, n \neq 2$ oe or $\frac{2t^k}{m^2}, k \neq 4$ oe or $m^{-2}t^4$ oe)
	(d)		$(y-4)(y+4)$	1	B1 cao
	(e)		h^{-6}	1	B1 for h^{-6} or $\frac{1}{h^6}$

22	(a)		$27x^6y^{12}$	2	B2 cao (B1 for any two terms correct in a product, e.g. $9x^6y^{12}$)
	(b)		$\frac{x-3}{2x-1}$	3	M1 for $(x-3)(x+3)$ M1 for $(2x-1)(x+3)$ A1 for $\frac{x-3}{2x-1}$ cao

PAPER: 1MA0_1H

PAPER: 5MB2H_01

Question		Working	Answer	Mark	Notes
5	(a)		x^6	1	B1 cao
	(b)		y^2	1	B1 cao
	(c)		t^6	1	B1 cao

PAPER: 1MA0_1H

Question		Working	Answer	Mark	Notes
22	(a)		1	1	B1 cao
	(b)		$\frac{5y}{8x^3}$	2	M1 for correct square root or correct use of reciprocal eg $\frac{8x^3}{5y}$ **or** $\frac{25y^2}{64x^6}$ A1 for $\frac{5y}{8x^3}$ **or** $\frac{5}{8}yx^{-3}$ oe
	(c)		$\frac{x+27}{(x-3)(x+3)}$	3	M1 for denominator $(x-3)(x+3)$ **or** x^2-9 M1 for $\frac{5(x+3)}{(x-3)(x+3)}$ oe **or** $\frac{4(x-3)}{(x-3)(x+3)}$ oe (NB The denominator must be $(x-3)(x+3)$ or x^2-9 **or** another suitable common denominator) A1 for $\frac{x+27}{(x-3)(x+3)}$ **or** $\frac{x+27}{x^2-9}$

PAPER: 1MA0_1H

PAPER: 5MB2H_01

Question		Working	Answer	Mark	Notes
11	(a)		$6x^5y^8$	2	M1 for any two of 6, x^5, y^8 A1 cao
	(b)		$6x^2-11x+3$	2	M1 for 3 out of 4 correct terms with correct signs, or all 4 terms correct ignoring signs A1 cao
	(c)		$4x^3y^2(2y^3-3x)$	2	M1 for $4x^3y^2$ (two terms involving x and/or y) or correct partial factorising taking out two from 4 or x or y A1 cao
	(d)		$(2+x)(e-2f)$	2	M1 for partial factorising of at least two terms eg $2(e-2f)$ or $x(e-2f)$ or $e(2+x)$ or $2f(2+x)$ oe A1 cao

Question		Working	Answer	Mark	Notes
7	(a)		$5y + 4$	2	M1 for $3\times y - 3\times 2$ or $2\times y + 2\times 5$ A1 cao
	(b)		w^2	1	B1 cao
	(c)		$5(x + 4)$	1	B1 cao

11	(a)		±7	2	M1 for intent to divide both sides by 3 as a first step or answer of 7 or -7 A1 ±7
	(b)		0.125 or $\frac{1}{8}$	1	B1 cao
	(c)		$27x^6$	2	M1 for either 27 or x^6 in a two term product A1 cao
	(d)		$p = \frac{w + 16}{4}$	2	M1 for ÷ 4 throughout or adding 16 to both sides as a first step A1 $p = \frac{w+16}{4}$ oe

PAPER: 1MA0_2H

Original question 8

The full source mark scheme begins on page 269. Use the original paper code, tier and question number shown above.

PAPER: 1MA0_2H

Question		Working	Answer	Mark	Notes
18	(a)		$10a^5b^4$	2	B2 cao (B1 for any two of 10, a^5, b^4 in a product)
	(b)		$y = 5p^2 - x$	3	M1 for a correct first step to either remove the square root, eg. $p^2 = \frac{x+y}{5}$ or clear the fraction, eg. $\sqrt{5}p = \sqrt{x+y}$ M1 (dep) for a method to remove the square root **and** a method to clear the fraction , eg. $5p^2 = x + y$ A1 for $y = 5p^2 - x$

10	(a)		a^7	1	B1 cao
	(b)		b^{14}	1	B1 cao
	(c)		1	1	B1 cao
	(d)		$\frac{1}{4}$	1	B1 for $\frac{1}{4}$ oe

Question		Working	Answer	Mark	Notes
8	(a)		$6ab - 7g$	2	M1 for $6ab$ or $-7g$ A1 cao
	(b)		$3(2m - 3)$	1	B1 cao
	(c)		t^5	1	B1 cao
	(d)		$2xy(x + 2y)$	2	B2 cao (B1 for $xy(2x + 4y)$ or $2x(xy + 2y^2)$ or $2y(x^2 + 2xy)$ or $2xy(\ldots)$ where $(\ldots)$ has two terms one of which is correct)
	(e)	$w^2 - 5w - 5w + 25$	$w^2 - 10w + 25$	2	M1 for all 4 terms (and no additional terms) correct with or without signs or 3 out of no more than 4 terms correct with signs A1 cao

Question	Working	Answer	Mark	Notes
6 (a)		6	B1	cao
(b)		5	B1	cao
(c)		Shown	M1	for writing 100^a or 1000^b as a power of 10 ($=10^{2a}$ or 10^{3b}) or 10^{2a+3b} or $100 = 10^2$ and $1000 = 10^3$
			C1	for complete chain of reasoning leading to conclusion

Paper: 1MA1/2H

Question	Answer	Mark	Mark scheme	Additional guidance
1 (a)	m^7	B1	cao	
(b)	$125n^3p^9$	B2	cao	Allow multiplication signs
		(B1	for 2 of 3 terms correct in a single product)	$125n^3p^x$ or $125n^xp^9$ where $x \neq 0$ or an^3p^9 where a is a number
(c)	$8q^6r^3$	B2	cao	Allow multiplication signs
		(B1	for 2 of 3 terms correct in a single product)	$8q^6r^x$ or $8q^xr^3$ where $x \neq 0$ or aq^6r^3 where a is a number

Question	Answer	Mark	Mark scheme	Additional guidance
7 (a)	p^{10}	B1	cao	
(b)	$2x^4y^2$	M1	for any two of $12 \div 6\ (=2)$, $x^{7-3}\ (=x^4)$, $y^{3-1}\ (=y^2)$ in a single product or written as a fraction with complete and correct cancelling of at least two terms	
		A1	cao	

19	$\frac{3}{4}$ oe	P1 P1 A1	for a first step to converting to a common base with one correct conversion, eg. $9^{-\frac{1}{2}} = 3^{-1}$ or $\frac{1}{3}$ or $27^{\frac{1}{4}} = 3^{\frac{3}{4}}$ oe (dep) for $3^{-1} = 3^{\frac{3}{4}} \div 3^{x+1}$ oe cao	$9^{-\frac{1}{2}} = 3^{-1}$ (or $\frac{1}{3}$) oe or $27^{\frac{1}{4}} = 3^{\frac{3}{4}}$ oe seen alone gets the P1

Paper: 1MA1/1H

Paper: 1MA1/2H

Question	Answer	Mark	Mark scheme	Additional guidance
10 (a)	1	B1	cao	
(b)	$\frac{8}{x-4}$	B1	cao	
(c)	$27n^{12}w^6$	B2	cao	
		(B1	for two of 27, n^{12}, w^6 in a product)	

Paper: 1MA1/3H

Question	Answer	Mark	Mark scheme	Additional guidance
1 (a)	n^8	B1	cao	
(b)	cd^3	M1	for partial simplification, eg c or d^3	May be seen as simplification in original fraction
		A1	for cd^3	Accept c^1d^3
(c)	$x > \frac{14}{5}$	M1	for $5x > 14$ or $5x = 14$ or critical value, $\frac{14}{5}$ oe	Must see carried out correctly, ie at least $5x > 7 \times 2$ not just intention seen. Allow other signs for this mark.
		A1	$x > \frac{14}{5}$ or $x > 2\frac{4}{5}$ or $x > 2.8$	

Paper: 1MA1/2H

Question	Answer	Mark	Mark scheme	Additional guidance
1 (a)	x^{15}	B1	cao	
(b)	$40 - 10x$	M1	for method to expand one bracket, or collect like terms eg $4 \times x + 4 \times 3$ $(= 4x + 12)$ **or** $7 \times 4 - 7 \times 2x$ $(= 28 - 14x)$ **or** $4 \times x - 7 \times 2x$ $(= 4x - 14x)$ **and** $4 \times 3 + 7 \times 4$ $(= 12 + 28)$	
		A1	oe	
(c)	$3x^2(5x + y)$	M1	for $3(5x^3 + x^2y)$ **or** $x(15x^2+3xy)$ **or** $3x(5x^2 + xy)$ **or** $x^2(15x + 3y)$ **or** $3x^2(ax + by)$	Where $a \geq 1$ and $b \geq 1$
		A1	cao	

14	(a)	$81x^{20}y^{24}$	B2	cao	
			(B1	for two of 81, x^{20}, y^{24})	
	(b)	$x^3+3x^2-10x-24$	M1	for method to find the product of any two linear expressions (3 out of 4 terms correct or 4 correct terms ignoring signs), eg $x^2 + 2x - 3x - 6$ or $x^2 + 2x + 4x + 8$ or $x^2 + 4x - 3x - 12$	Note that, for example $-x - 6$ in expansion of $(x + 2)(x - 3)$ is regarded as 3 correct terms.
			M1	for a complete method to find all terms, at least half of which are correct (ft their first product), eg $x^3 + 4x^2 + 2x^2 - 3x^2 + 8x - 6x - 12x - 24$	First product must be quadratic with at least 3 terms but need not be simplified or may be simplified incorrectly
			A1	cao	

Question	Answer	Mark	Mark scheme	Additional guidance
11	$3x^7y^2$	M1	for full evaluation of numerator or denominator with at least 2 of 3 terms correct in a product, eg $36x^{10}y^6$ **or** $12x^3y^4$ or full evaluation of $\frac{6x^5y^3}{3x^2y^7}$ or $\frac{6x^5y^3}{4xy^{-3}}$ with at least 2 of 3 terms correct in a product, eg $2x^3y^{-4}$ **or** $1.5x^4y^6$	
		M1	for correct evaluation of numerator and denominator, eg $36x^{10}y^6$ **and** $12x^3y^4$ or for full evaluation of numerator and denominator with no more than one error **and** a final answer of the form ax^by^c with two of a, b and c correct or for correct evaluation of $\frac{6x^5y^3}{3x^2y^7}$ and $\frac{6x^5y^3}{4xy^{-3}}$ eg $2x^3y^{-4}$ **and** $1.5x^4y^6$ or for full evaluation of $\frac{6x^5y^3}{3x^2y^7}$ and $\frac{6x^5y^3}{4xy^{-3}}$ with no more than one error **and** a final answer of the form ax^by^c with two of a, b and c correct	
		A1	for $3x^7y^2$ oe	Accept $a = 3$, $b = 7$, $c = 2$

Paper: 1MA1/1H

Paper: 1MA1/3H

Question	Answer	Mark	Mark scheme	Additional guidance
15 (a)	$\frac{a-3}{5}$	B1	for $\frac{a-3}{5}$ oe	
(b)	$(3k-1)(k+4)$	M1	for $(3k \pm 1)(k \pm 4)$ oe or for brackets which when expanded give 2 out of 3 terms correct	Accept $3k(k+4)-(k+4)$ Accept $k(3k-1)+4(3k-1)$
		A1	cao	Accept $(1-3k)(-k-4)$
(c)	$\frac{2-x}{x}$	M1	for factorisation eg $(4-x^2=)\ (2-x)(2+x)$ **and** $(x^2+3x=)\ x(x+3)$ **or** for inversion and multiplication (condone incorrect factorising) eg $\frac{4-x^2}{x^2+3x}\times\frac{x+3}{x+2}$ oe	
		M1	for factorisation of both quadratics **and** inversion and multiplication, eg $\frac{(2-x)(2+x)}{x(x+3)}\times\frac{x+3}{x+2}$ allow sign errors in one bracket	
		A1	for $\frac{2-x}{x}$ or $\frac{2}{x}-1$	

Paper: 1MA1/3H

9	(a)	$14x^5y^6$	B2	cao	
			(B1	for correct simplification of two terms ax^5y^6 **or** $14x^by^6$ **or** $14x^5y^c$ where $a \neq 14$, $b \neq 5$, $c \neq 6$)	Where a, x^b, y^c can be made up of two products Condone inclusion of multiplication signs for B1
	(b)	m^{-6}	B1	for m^{-6} or $\frac{1}{m^6}$	

Paper: 1MA1/3H

Question	Answer	Mark	Mark scheme	Additional guidance
11 (a)	2	M1	for a correct start, using one rule of indices, eg $3^n \times 3^{12} = 3^{14}$ or $3^n \times 3^{20} = 3^{22}$ or $3^{20} \div 3^8 = 3^{12}$ or $3^{14} \times 3^8 = 3^{22}$ or $\frac{3^{n+20}}{3^8} = 3^{14}$ or $3^{n-8} \times 3^{20} = 3^{14}$ **or** for forming an equation in n, eg $n + 20 - 8 = 14$ oe or (n =) $14 + 8 - 20$	
		A1	cao	Allow an answer of $3^n = 3^2$
			SCB1 for an answer of 3^2 if M0 scored	An answer of 9 or $3^n = 9$, $n \neq 2$ on its own is to be awarded 0 marks
(b)	$16m^8$	M1	for an intention to find the cube root and square, eg $\sqrt[3]{64m^{12}}^2$ or $\sqrt[3]{(64m^{12})^2}$ or $(4m^4)^2$ or $\sqrt[3]{4096m^{24}}$ or for am^8 with $a \neq 16$ or $16m^b$ with $b \neq 8$	Do not condone missing brackets 16 or m^8 imply M1 Allow multiplication sign for M1
		A1	cao	Accept $a = 16$, $b = 8$

Paper: 1MA1/3H

GCSE

Mark Scheme

Mathematics A (1387)

June 2003

Edexcel
Success through qualifications

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information please call our Customer Response Centre on 0870 240 9800, or visit our website at www.edexcel.org.uk.

June 2003

Publications Code UG014129

All the material in this publication is copyright
© London Qualifications Ltd 2003

CONTENTS

Notes on Marking Principles 2
Paper 5501 4
Paper 5502 11
Paper 5503 17
Paper 5504 24
Paper 5505 32
Paper 5506 40

NOTES ON MARKING PRINCIPLES

NOTES ON MARKING PRINCIPLES

1 Three types of mark are available

M marks - awarded for correct working seen or implied.
A marks - conditional accuracy marks which are awarded for accurate working following the award of M marks.
B marks - unconditional accuracy marks (independent of M).

2 Abbreviations

cao - correct answer only.
ft - follow through.
√ - Denotes a "follow through" answer.

SC - special case.
isw - ignore subsequent working.
oe - or equivalent (and appropriate).
NB: a candidate cannot benefit from both isw and ft.

3 If no working shown, then

correct answers normally score full marks
incorrect (even though nearly correct) answers score no marks.

4 Marking instructions

Misread loses A marks (and sometimes B marks) on that part but ft can be allowed on subsequent parts. M marks can still be earned. If in doubt contact your team leader.
If there is a wrong answer in the answer space DO CHECK the working in the body of the script.

- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.
- If there is a wrong answer in the answer space DO CHECK the working in the body of the script.
- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.

5 Style of marking

Answer correct: tick and write part mark in margin NEXT TO BRACKETED MARK.
Answer incorrect: cross, but show M, A or B marks if any earned in body of script and transfer the total of these to the margin next to the bracket mark.
Total for each double page at bottom right page (except for back if used), FINAL TOTAL IN RELEVANT BOX ON FRONT COVER.
Nought in margin for fully incorrect question or page AND FOR NO ATTEMPT.
Where no attempt has been made a line should be put in the answer space and zero in the margin next to the bracketed mark
CHOICE OF METHOD – No marks unless one answer is in answer space – then mark that.
CROSSED OUT WORK - if not replaced this should be marked (if legible).
There must always be a mark next to bracketed mark in the margin/

6 Follow Through Marks

Follow throughs are guided by two principles:

(a) Follow throughs which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous, do not award.
(b) Follow throughs which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

7 Probability

Probability answers must be given as fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least two decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.

8 Linear Equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.



9 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.


Paper 5501

No	Working	Answer	Mark	Notes
1 (a)		46	1	B1 cao
(b)		3.4	1	B1 oe
(c)		Arrow at 430	1	B1 allow ± half graduation
(d)		Arrow at 3.7	1	B1 allow ± half graduation Accept indications other than arrows as long as they are clear
2		1.60 2.05	2	B1) Condone B1) reversal
3 (a)	$\frac{18}{24}$ or $\frac{9}{12}$ or $\frac{6}{8}$	$\frac{3}{4}$	2	B2 for $\frac{3}{4}$ cao (B1 for $\frac{18}{24}$ or $\frac{9}{12}$ or $\frac{6}{8}$) SC B1 for $\frac{1}{4}$ only
(b)		16 squares shaded	1	B1 cao
4 (a)		line	1	B1 within overlay tolerance
(b)		midpoint	1	B1 within overlay tolerance ft from (a) ± 0.2cm
(c)		rectangle	1	B1 for rectangle 6 cm ± 0.2cm by 4 cm ± 0.2cm
5		kilograms, kg litre, *l* or cubic metres, m^3 inches, in	3	B1 B1 B1
6 (a)		parallel lines marked	1	B1
(b)		right angle marked	1	B1
(c)(i)		acute	1	B1
(ii)		reflex	1	B1

Paper 5501

No	Working	Answer	Mark	Notes
7 (i)		sphere	1	B1
(ii)		cylinder	1	B1 Accept circular prism
(iii)		pyramid	1	B1 Condone omission of "triangular" Accept tetrahedron
8 (a)(i)		40	2	B1 cao
(ii)		50		B1 cao
(b)		5 complete symbols	1	B1 cao
(c)		 and oe inc	1	B1
9 (i)		9, 37, 56, 59, 75	5	B1 cao
(ii)		0.067, 0.56, 0.6, 0.605, 0.65		B1 cao Ignore trailing zeros
(iii)		–10, –6, –4, 2, 5		B1 cao
(iv)		$\frac{2}{5}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}$		B2 for all 4 correct (B1 for any 3 in correct order) SC B1 for all 4 in reverse order (applies to(iv) only)
10 (a)			1	B1
(b)		Plotting (4, 24)	1	B1 ft from their matchsticks
(c)		60	1	B1 cao
(d)		$m = 6n$	2	B2 for $m = 6n$ oe (B1 for $6n$ oe or m = multiple of n except $m = n$)
11 (i)		6, 12	4	B1 cao
(ii)		4, 16		B1 cao
(iii)		3, 4, 6 or 3, 4, 6, 12		B1 Condone omission of 12
(iv)		8, 27		B1 cao

Paper 5501

No	Working	Answer	Mark	Notes
12		2.43	4	B1 for 36 or 0.36 B1 for 96 or 0.96 B1 for 125 or 1.25 If none of first 3 B1s awarded then SC B1 for four 24s and five 25s seen OR 4 × 24 and 5 × 25 seen B1 for 2.43 cao
13 (a)(i)		Edinburgh and Plymouth	3	B1 for Edinburgh or −7 B1 for Plymouth or 5
(ii)		12		B1ft from (i) if one positive and one negative
(b)		Cardiff and Belfast London and Plymouth	2	B1 for Cardiff and Belfast OR −6 and −4 B1 for London and Plymouth OR 3 and 5
14		5 9 8	3	B1 cao B1 cao B1 cao
15 (a)			2	✓ / × / 2 / 0 / B2 1 / 0 / B1 1 / 1 / B1 2 / 1 / B1
(b)	$2\times2\times2$	8	2	M1 for $2\times2\times2$ A1 for 8 cao
16 (a)		30	1	B1 cao
(b)		3	1	B1 cao
(c)	$2 \times 29 = 58$ $5 \times 30 = 150$ $2 \times 31 = 62$ $1\times32 = 32$ $\frac{302}{10} = 30.2$	30.2	3	M1 for freq × no pins M1 (dep on 1st M1) for totalling and ÷ 10 A1 for 30.2 cao

Paper 5501

No	Working	Answer	Mark	Notes
17		7 5 5 13 33 52 23	3	B3 all correct (B2 for 4, 5 or 6 correct B1 for 2 or 3 correct)
18 (a)(i) (ii) (iii) (iv) (b)		$4c$ p^4 $8g$ $10pr$ OR $10rp$ $10y - 15$	4 1	B1 oe B1 cao B1 oe B1 B1 cao Accept $10y + - 15$
19		$\frac{2}{3}$	3	M1 for 3 rows (9 squares) shaded M1 for 2 columns (10 squares) shaded A1 for $\frac{2}{3}$
	$\frac{3}{5} = \frac{9}{15}$ $\frac{2}{3} = \frac{10}{15}$	$\frac{2}{3}$	3	M1 for $\frac{3}{5} = \frac{9}{15}$ M1 for $\frac{2}{3} = \frac{10}{15}$ A1 for $\frac{2}{3}$
	$\frac{3}{5} = 0.6$ $\frac{2}{3} = 0.66$ or 0.67 or better	$\frac{2}{3}$	3	M1 for $\frac{3}{5} = 0.6$) Accept M1 for $\frac{2}{3} = 0.66$ or 0.67 or better) percentages A1 for $\frac{2}{3}$


Paper 5501

No	Working	Answer	Mark	Notes
20 (a)	955 × 48 7640 38200 45840 OR 48 × 955 240 2400 43200 45840	458.40	3	M1 for complete correct method (condone one computational error) A2 for 458.40 cao (A1 for digits 4584 OR ft if M1 awarded)
(b)	14. 5 48 | 696. 0 48 216 192 24 0 24 0	14.50	3	M1 for 1 as first digit in answer and remainder 21 M1 (dep) 4 as second digit in answer A1 for 14.50 (Accept 14.5)
21 (a)		$12x$	1	B1 oe
(b)		$12x + 10y$	2	B2 oe ft from (a) (B1 $12x$ + multiple of y or $10y$ seen) SC B1 for $x = 12x + 10y$ OR $y = 12x + 10y$

Paper 5501

No	Working	Answer	Mark	Notes
22	$\frac{1}{3}+\frac{1}{4}=\frac{4}{12}+\frac{3}{12}=\frac{7}{12}$ $1-\frac{7}{12}=\frac{5}{12}$	$\frac{5}{12}$	3	M1 for $\frac{4}{12}$ **and** $\frac{3}{12}$ oe A2 for $\frac{5}{12}$ oe (A1 for $\frac{7}{12}$) B1 for 1–"$\left(\frac{1}{3}+\frac{1}{4}\right)$" correctly evaluated
	$1-\frac{1}{3}=\frac{2}{3}$, $1-\frac{1}{4}=\frac{3}{4}$ $\frac{8}{12}-\frac{3}{12}=\frac{5}{12}$ or $\frac{9}{12}-\frac{4}{12}=\frac{5}{12}$	$\frac{5}{12}$	3	B1 for $\frac{2}{3}$ or $\frac{3}{4}$ seen M1 for $\frac{8}{12}-\frac{3}{12}=\frac{5}{12}$ or $\frac{9}{12}-\frac{4}{12}=\frac{5}{12}$ A1 for $\frac{5}{12}$ oe
		$\frac{5}{12}$	3	M1 for 0.25 and 0.33 or better A1 for 0.58 or better A1 for $0.41\dot{6}$ or recurring
23 (a) (b)(i) (ii)	 180 – (54 + 54)	54 72 Reason	1 3	B1 cao M1 for 180 – (54 + 54) A1 ft from (a) if $x < 90$ B1 for mentioning isosceles and equal or base angles or equal sides and equal or base angles

Paper 5501

No	Working	Answer	Mark	Notes
24 (a)		Bryani	2	M1 for 4×9 or $4 \times 3 \times 3$ or $4 \times x \times x$ or square x first or square 3 first A1 SC 4×3^2 with Bryani scores B2
(b)		64	1	B1 cao
25 (a)		 dotted line may be solid	3	B2 for rectangle base 3 squares and height 4 squares (B1 for rectangle with one correct dimension) B1 for horizontal line 1 cm from top) SC B2 for completely correct elevation on its side
(b)			2	B2 for perspective drawing showing slant face and cutout B1 for perspective drawing with either slant face cutout omitted
26 (a)	20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$	40	2	M1 for 20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$ A1 cao
(b)	$\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes	line from (45, 20) to (65, 0)	2	M1 for $\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes seen A1 for correct line SC If M0, B1 for line from (45, 20) to (t, 0) where $t > 45$ or B1 for a line of the correct gradient.

Paper 5502

No	Working	Answer	Mark	Notes
1 (a)		$\frac{7}{100}$	1	B1 cao accept 0.07
(b)		0.18	1	B1 cao
(c)	20 in 100 oe	40	2	M1 for sight of 20 in 100 or 20 × 2 A1 cao
2 (a)		6cm	2	B1 for 6 ± 0.2 or 60 ± 2 B1 indep for cm or mm consistent with 1st B1
(b)		At centre	1	B1 within overlay
(c)		Circle drawn	1	B1 all within overlay
3		See diagram	3	B3 all correct – see separate sheet (B2 for 3 correct B1 for 2 correct)
4 (a)		$\frac{1}{4}$ oe	1	B1 cao
(b)		0.75	1	B1 cao
(c)		75%	1	B1 cao
(d)(i)		9	2	B1 accept answer in range 9 - 9.2
(ii)		15 - 16		B1 accept answers in range15 - 16
5 (a)		9:30	1	B1 cao
(b)		2hrs 45 min	2	B2 for 2hr 45 min or 2 $\frac{3}{4}$ hr or 165 minutes B1 2:45 or 2.45 or 165 or 45min + 1hr + 1hr oe
(c)		17	1	B1 cao
6 (a)	96 × 4	3.84	2	M1 for 96 × 4 or digits 384 A1 cao
(b)	3 × 96 + 40 = 328	(0).56	2	M1 for 3 × 96 + 40 or digits 328 or digits 56 A1 cao accept 56p

Paper 5502

No	Working	Answer	Mark	Notes
7 (a)		54 000	1	B1 cao accept 54 thousand
(b)		50 000	1	B1 (accept ten thousand or 10 000) oe
8 (a)		14	1	B1 cao
(b)		6	1	B1 cao
(c)		Correct reflection	2	B2 fully correct (B1 correct reflection in a line parallel to the mirror line or condoning 1 block error in shape or position of shape)
9 (a)		Missing horiz label 1 (and 6) missing on vertical scale	2	B1 B1
(b)		Correct graph	2	B1 for bar up to 4 for yellow B1 for bar up to 2 for green
(c)		Blue	1	B1 cao
(d)	3 + 5 + 4 + 2	14	1	B1 ft from (b)
(e)		$\frac{3}{14}$	1	B1 ft on '14'
10	Barry (8) because you double Kath (7) because you add, 1,2,3		2	B1 oe B1 oe SC: B1 for correct rules only
11 (a)		$2n$	1	B1 for $2n$ or $n + n$ OR $2 \times n$ OR $n \times 2$ OR $n2$
(b)		$2n + 15$	1	B1 for "$2n$" + 15 oe
(c)		20q	1	B1 cao
12 (a)	1 + 3 + 5 + 8 + 5	22	2	M1 add frequencies A1 cao
(b)		No, is>no of cups of coffee in the table	1	B1 'average cannot be bigger than 6' oe OR 'Average must be less than 6 oe'
13 (a)		Trapezium	1	B1 cao ignore spelling
(b)		(2, 3)	1	B1 cao
(c)		Isosceles	1	B1 cao ignore spelling
(d)		Q correct	1	B1 cao

Paper 5502

No	Working	Answer	Mark	Notes
14 (a)		250 000	1	B1 cao
(b)	$\frac{28}{4}$	7	2	M1 for $\frac{28}{4}$ oe or "250000" × 28 A1 cao **SC** B1 for 7 000 000
15 (a)		10	1	B1 cao
(b)		5.5	1	B1 ±0.3 pounds
(c)	$\frac{110}{22}$	50	3	M1 for use of graph at 11 or $\frac{110}{22}$ A1 for 5 A1 cao **SC** B2 for 49.5 – 50.6

Paper 5502

No	Working	Answer	Mark	Notes
16 (a)	269.30 – 56.80 = 212.50 $\frac{212.50}{42.50}$	6	2	M1 for $\frac{269.30-56.80}{42.50}$ or 5 seen A1 cao
(b)	5% of £269.30 £269.30 – "£13.465" OR $\frac{95}{100}\times 269.30$	255.83 or 255.84	3	M1 for (5 ÷ 100) × 269.30 M1 for 269.30 – "13.465" A1 cao OR M2 for $\frac{95}{100}\times 269.30$ A1 cao **Alternative Method:** M1 for $\frac{5}{100}\times 56.80$ (= 2.84) and $\frac{5}{100}\times 42.50$ (=2.12(5)) (OR 53.96 AND 40.38 (40.375) seen M1 for 56.80 – "2.84" (= 53.96) 42.50 – "2.12(5)" (= 40.375 or 40.38) "5" × "40.375" + "53.96" A1 cao
17	$4.1^2\times 1.07 = 16.81\times 1.07$	17.9867	2	M1 for ("4.1") followed by squaring, or sight of 16.81 A1 cao SC: *B1 for 18 or better with no working*
18 (a)		60	1	B1 cao
(b)	360 – 60 – 90 – 90	120	2	M1 for 360 – " 60 " – 90 – 90 or 180 – "60" A1 cao
(c)	6 × 2	12	2	M1 for 6 × 2 A1 cao for 12

Paper 5502

No	Working	Answer	Mark	Notes
19 (a)(i)	$240 \times 5 = 1200$	1250	3	B1 cao 1250
(ii)	$\frac{50}{'1250'}$	$\frac{1}{25}$		M1 cao $\frac{50}{'1250'}$ A1 for $\frac{1}{25}$ in its simplest form
(b)	$\frac{60}{100} \times 1000 = 600$	12:5	3	M1 for $\frac{60}{100} \times 1000$ oe A1 for 600 A1 cao
20 (a)		$x + 2$	1	B1 accept $2 + x$ but not $x = x + 2$
(b)	$x + 5 + x + 5 + x + 2 + x + 2$	$4x + 14$	2	M1 adding 4 sides, two of which are '$x + 2$' (all sides to be linear expressions in x) **SC** $x + 5 + x + 2 \times 2$ gets M1 A1 for correct simplified answer or $(20 - 14) \div 4$ oe gets M1
(c)	'$4x + 14$' = 20	1.5 oe	2	M1 for equation A1 cao
21	$\sum f = 90$	Angles drawn, labelled	3	M1 for 1 person = 4°or one angle correct in table or pie chart A1 any 2 correctly drawn angles in pie chart A1 fully correct chart labelled
22 (a)		$2p - q$	2	B1 cao for $2p$ B1 cao for $-q$ accept $(-q + 2p)$, $2p - 1q$ and $2p + -q$
(b)	$5x = 3 + 4$	1.4	2	M1 for either (+3 or sight of 7) or (÷ 5 or sight of 0.8 and 0.6) A1 cao accept $\frac{7}{5}$ or $1\frac{2}{5}$

Paper 5502

No	Working	Answer	Mark	Notes
23 (a)		$\frac{4\times5}{2}$	1	B1 cao
(b)	$1+2+3+4+5+6+7+8$	$\frac{8\times9}{2}$	1	B1 cao
(c)	$\frac{100\times101}{2}$	5050	1	B1 cao
24	0 | 5 7 8 8 1 | 0 0 0 0 2 5 5 5 6 2 | 0 0 0 4 5 3 | 3 5 Key 1 | 2 = 12 (min)	See working	3	B1 for stem 0, 1, 2, 3 or 0, 10, 20, 30 B1 for accurate unordered leaves condone 1 error or omission B1 for key and ordered leaves all correct
25	$3.2\times2.8=8.96$ $2\times4.5\times2.8=25.2$ $2\times4.5\times28.8=28.8$ $\frac{62.96}{2.5}\times2.99$	£75.30	5	M1 for area of any face found correctly M1 for 2 areas found correctly A1 for 62.96 or 54 M1 for $\frac{'62.96'}{2.5}\times2.99$ Al cao Alternate method for candidates who round up $\frac{"62.92"}{2.5}$ M1 for "26" × 2.99 A1 for £77.74 cao **SC**: *for top included B2 for 71.92 m^2 seen or* B3 for £86.02 seen **SC** B4 for £64.58 or £65.78 seen
26	$2.5\times10\,000$	25 000	2	M1 for $2.5\times100\times100$ A1 cao

Paper 5503

No	Working	Answer	Mark	Notes
1 (a)(i)		$8g$	2	B1 oe
(ii)		$10rp$		B1 for $10pr$ or $10\ rp$
(b)		$10y - 15$	1	B1 cao accept $10y$ - + 15
(c)	$6x + 8 - 12x + 15$	$-6x + 23$	2	M1 for 3 correct terms out of 4 A1 cao
2		$m = 6n$	2	B2 for $m = 6n$ oe accept $6 \times n$, n 6 (B1 for $6n$ alone, or $6n$ +1 oe OR m = multiple of n except $m = n$)
3 (i)		0.067, 0.56, 0.6, 0.605, 0.65	1	B1 cao Ignore trailing zeros
(ii)		$-10, -6, -4, 2, 5$	1	B1 cao
(iii)		$\frac{2}{5}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}$	2	B2 all four correct (B1 any three in correct order) SC: B1 all 4 in reverse order
4		7 5 5 13 33 52 23	3	B3 all correct (B2 for 4, 5 or 6 correct entries) (B1 for 2, 3 correct entries)


Paper 5503

No	Working	Answer	Mark	Notes
5		$\frac{2}{3}$	3	M1 for 3 rows (9 squares) shaded M1 for 2 columns (10 squares) shaded A1
	OR $\frac{3}{5}=\frac{9}{15}$ $\frac{2}{3}=\frac{10}{15}$ Therefore $\frac{2}{3}>\frac{3}{5}$	$\frac{2}{3}$		M1 for $\frac{3}{5}=\frac{9}{15}$ M1 for $\frac{2}{3}=\frac{10}{15}$ A1
	OR $\frac{3}{5}=0.6$ or percent $\frac{2}{3}=0.66$ or 0.67 or better Therefore $\frac{2}{3}>\frac{3}{5}$	$\frac{2}{3}$		M1 for $\frac{3}{5}=0.6$ or percent M1 for $\frac{2}{3}=0.66$ or 0.67 or better A1
6	$2\times 29=58$ $5\times 30=150$ $2\times 31=62$ $1\times 32=32$ $\frac{302}{10}=30.2$	30.2	3	M1 for freq × no. pins (at least 3) M1 for totalling and for ÷ 10 (dep on 1st M1) A1 cao
7 (a)			2	B1 for 180° rotation (wrong centre) B1 cao
(b)			3	B1 for any enlargement sf other than 1 B1 for all sides halved B1 for position

Paper 5503

No	Working	Answer	Mark	Notes
8 (a)		$12x$	1	B1 oe
(b)		$12x + 10y$	2	B2 oe or ft from (a) (B1 for $12x$ + a multiple of y or $10y$ seen) **SC** B1 for $x = 12x + 10y$ or $y = 12x + 10y$
9	$\frac{1}{3}+\frac{1}{4}=\frac{4}{12}+\frac{3}{12}=\frac{7}{12}$ $1-\frac{7}{12}=\frac{5}{12}$ OR $1-\frac{1}{3}=\frac{2}{3}, 1-\frac{1}{4}=\frac{3}{4}$ $\frac{9}{12}-\frac{4}{12}=\frac{5}{12}, \frac{8}{12}-\frac{3}{12}=\frac{5}{12}$	$\frac{5}{12}$	3	M1 for $\frac{4}{12}$ and $\frac{3}{12}$ oe A2 for $\frac{5}{12}$ oe (A1 for getting $\frac{7}{12}$) (B1 for $1 - \frac{"1}{3} + \frac{1"}{4}$ correctly evaluated) OR B1 for $\frac{2}{3}$ or $\frac{3}{4}$ seen M1 for $\frac{3}{4}-\frac{1}{3}=\frac{9}{12}-\frac{4}{12}$ or $\frac{2}{3}-\frac{1}{4}=\frac{8}{12}-\frac{3}{12}$ A1 for $\frac{5}{12}$ oe OR M1 0.25 and 0.33 or better A1 for 0.58 or better A1 for 0.416 *recurring*
10 (a)		54	1	B1 cao
(b)(i)	180 – (54 +54)	72	3	M1 for 180 – (“54” + “54”) A1 for 72° ft from (a) if $x < 90°$
(ii)				B1 for mentioning isosceles and equal or base angles or equal sides and equal or base angles

Paper 5503


No	Working	Answer	Mark	Notes
11 (a)	Bryani was correct $4 \times 3^2 = 4 \times 9 = 36$	Bryani	2	M1 for 4×9 or $4 \times 3 \times 3$ or "square the three/x then multiply by four" A1 Bryani **SC** 4×3^2 with Bryani gets B2
(b)			1	B1 cao
12 (a)			3	B2 for rectangle height 4 squares, base 3 squares (B1 for rectangle with one correct dimension) B1for line 1 square from the top **SC** B2 for completely correct elevation on its side
(b)			2	B2 for perspective drawing showing slant face and cut out (B1 for perspective drawing with either slant face or cut out omitted or one aspect incorrect)
13 (a)	20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$	40	2	M1 for 20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$ A1 cao
(b)	$\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes seen	Line from (45, 20) to (65, 0)	2	M1 for $\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes seen A1 for correct line **SC** If M0, B1 for line from (45, 20) to (t, 0) where $t > 45$ or a line of the correct gradient
14 (i)		119.31	3	B1 cao
(ii)		119 310		B1 cao
(iii)		1.23		B1 cao

Paper 5503

No	Working	Answer	Mark	Notes
15	$\frac{10}{100}\times 12000$ 12 000 – 1200 = 10 800 $\frac{10800}{10}=1080$ 10 800 – 1080 = £9720	£9720	3	M1 for $\frac{10}{100}\times 12000$ or sight of 1200 or 2400 or 10 800 or 9600 M1 (dep) for $\frac{10}{100}\times(12\,000-\frac{10}{100}\times 12000)$ or sight of 1080 A1 cao Alternative mark scheme M2 for $12000\times(1-\frac{10}{100})^2$ (M1 for $12000\times(1-\frac{10}{100})$ A1 cao
16 (a)	$2p = 6$	$p = 3$	2	M1 for $7p – 5p = 8 – 2$ or $2p$ or 6 A1 cao
(b)	$7r – 5r = – 20 –2$	–11	2	M1 for $7r + 2 = 5r – 20$ or $\frac{7r}{5}+\frac{2}{5}=r-4$ or $7r – 5r = 20 – 2$ or $\frac{7r}{5}-r=-4-\frac{2}{5}$ A1 cao
17		$5n + 1$	2	B2 oe (B1 for $5n$ seen) NB: n = gets B1 max

Paper 5503

No	Working	Answer	Mark	Notes
18 (a)	–1, 0, 1		2	B2 for –1, 0, 1 (B1 for –1, 0 or 0,1 or –1,1 or –2, -1, 0, 1 only)
(b)	(–1, –1), (0, –1), (1, –1), (0, 0), (1, 0), (1, 1)		3	B3 for 6 points correct B2 for 3 points correct B1 for 1 point correct **NB** –B1 each additional point over six
19 (a)	Triangle with vertices at (0,0) (0,–2) and (3,0)		2	M1 for correct orientation A1 cao
(b)	Rotation, 180°, centre (0,1) Enlargement sf - 1 centre (0,1)		2	B2 for 180° 'rotation' centre (0, 1) B2 for Enlargement sf - 1 centre (0,1) (B1 for any two of the three parts) NB: B0 if additional transformation is included
20	Bisector of $\angle BAC$ Arc around A Region		3	B3 cao (B2 for either two correct boundaries, no shading/ wrong shading or one correct boundary, one incorrect boundary with valid shading) (B1 for either two incorrect boundaries but one drawn from A and one intersection, with valid shading or one correct boundary) Ignore shading outside the triangle
21		Length Volume Area	3	B1 for Length B1 for Volume B1 for Area
22 (a)	Unbiased question with choices		2	B1 for unbiased question B1 for at least 2 choices
(b)			2	Classification 1: A biased question Classification 2: A restricted sample of people Classification 3: Not specifying a range of foods Classification 4: Nothing to do with eating habits B2 reasons which satisfy 2 different classifications (B1 a reason which satisfies one classification)



Paper 5503

No	Working	Answer	Mark	Notes
23 (a)	$6 \times 10^2 \times 8 \times 10^4$ 48×10^6	4.8×10^7	3	M1 for $6 \times 10^a \times 8 \times 10^b$ oe, a and b integers including 0 A1 for 48×10^6 oe A1 cao
(b)	200 000 + 30 000	230000	2	B2 cao (B1 for sight of 200 000 or 30 000 or 2.3×10^5 or 23×10^4)
24 (i)		64	4	B1 cao
(ii)		3		B1 cao
(iii)	$\sqrt{16 \times 9} = \sqrt{144}$	12		B2 cao (B1 for sight of $\sqrt{2^4} \times \sqrt{9}$ or better, or 144 seen)
25 (i)	Tangent 90° to diameter /radius/ line from (through) centre	27°	4	B1 for 27° cao B1 for reason
(ii)	180 –(90+ "27") angle in semicircle (is 90°)/Alternate segments /angle at centre twice at circumference	63°		B1 ft for 90–"27" if not 63° B1 for reason
26 (a)(i)		152	2	B1 cao
(ii)		177		B1 cao
(b)			3	B1 for median marked at 167 B1 ft for postion of box with its ends at "152" and "177" B1 for position of whiskers with ends at 132 and 182 NB: For any points plotted between 141 and 149 give a tolerance of an extra ± 1 square
27	$x^2 + xy + xy + y^2$	$x^2 + 2xy + y^2$	2	M1 for at least 3 of the 4 terms correct A1 cao
		25	2	M1 for recognising 3.47 + 1.53 (= 5) A1 cao

Paper 5504

No	Working	Answer	Mark	Notes
1 (a)	$4.1^2 \times 1.07 = 16.81 \times 1.07$	17.9867	2	M1 for (4.1) followed by squaring, or sight of 16.81 A1 cao SC: B1 for 18 or better with no working
(b)		$(1.6 + 3.8 \times 2.4) \times 4.2$	1	B1 cao Allow additional brackets if they give an expression with value 45.024
2 (a)	269.30 – 56.80 = 212.50	6	2	M1 for $\frac{269.30 - 56.80}{42.50}$ or 5 seen A1 cao
(b)	5% of £269.30 £269.30 – "£13.465" OR $\frac{95}{100} \times 269.30$	255.83 or 255.84	3	M1 for (5 ÷ 100) × 269.30 M1 (dep) for 269.30 – "13.465" A1 cao OR M2 for $\frac{95}{100} \times 269.30$ A1 cao **Alternative Method:** M1 for $\frac{5}{100} \times 56.80$ (= 2.84) and $\frac{5}{100} \times 42.50$ (=2.12(5)) (OR 53.96 AND 40.38 (40.375) seen M1 for 56.80 – "2.84" (= 53.96) 42.50 – "2.12(5)" (= 40.375 or 40.38) "5" × "40.375" + "53.96" A1 cao

Paper 5504

No	Working	Answer	Mark	Notes
3 (a)		60	1	B1 cao
(b)	360 – 60 – 90 – 90	120	2	M1 for 360 – "60" – 90 – 90 or 180 – "60" A1 cao
(c)	6 × 2	12	2	M1 for 6 × 2 A1 cao 12
(d)		Correct drawing	2	B2 for triangle and construction lines (see overlay) (B1 for 1 line of length 4cm and correct arcs crossing OR for correct triangle with either no arcs or incorrect arcs) SC: B1 similar triangle drawn with construction lines
4 a(i)	240 × 5 = 1200	1250	3	B1 cao 1250
(ii)	$\frac{50}{1250}$	$\frac{1}{25}$		M1 $\frac{50}{"1250"}$ A1 oe in its simplest form
(b)	$\frac{60}{100} \times 1000 = 600$ 600: 250	12:5	3	M1 for $\frac{60}{100} \times 1000$ oe A1 for 600 A1 cao
5 (a)		$x + 2$	1	B1
(b)	$x + 5 + x + 5 + x + 2 + x + 2$	$4x + 14$	2	M1 adding 4 sides, two of which must be '$x + 2$' (all sides to be linear expressions in x) A1 for correct simplified answer
(c)	"$4x + 14$" = 20	1.5	2	M1 for equation "$4x + 14$" = 20 OR $\frac{20-14}{4}$ oe A1 cao

Paper 5504

No	Working	Answer	Mark	Notes
6 (a)		$2p - q$	2	B1 cao for $2p$ B1 cao for $-q$ accept $-q + 2p$ and $2p - 1q$
(b)	$5x = 3 + 4$	1.4	2	M1 for either (+3 or sight of 7) or (÷ 5 or sight of 0.8 and 0.6) A1 cao accept $\frac{7}{5}$ or $1\frac{2}{5}$
7 (a)		$\frac{4\times5}{2}$	1	B1 cao
(b)	$1 + 2 + 3 + 4 + 5 + 6 + 7 + 8$	$\frac{8\times9}{2}$	1	B1 cao
(c)	$\frac{100\times101}{2}$	5050	1	B1 cao
(d)		$\frac{n(n+1)}{2}$	2	B2 cao (B1 for any quadratic in n)
8	$3.2 \times 2.8 = 8.96$ $2 \times 4.5 \times 2.8 = 25.2$ $2 \times 4.5 \times 3.2 = 28.8$ $\frac{62.96}{2.5} \times 2.99$	75.30	5	M1 for area of any face found correctly M1 for 2 areas seen A1 for 62.96 or 54 M1 for $\frac{'62.96'}{2.5} \times 2.99$ A1 cao Alternate method for candidates who round up $\frac{"62.96"}{2.5}$ M1 for "26" x 2.99 A1 for £77.74 cao SC: for top included B2 for 71.92 m^2 seen or B3 for £86.02 or £86.71 SC: B4 for £64.58 or £65.78 seen

Paper 5504

No	Working	Answer	Mark	Notes
9	2.5 × 10000	25000	2	M1 for 2.5 × 10000 or 2.5.× 100 ×100 A1 cao
10 (a)	∑f = 90 (88), 144, 32, 96	Angles drawn, labelled	3	M1 for 1 person = 4°or one angle correct in table or pie chart A1 any 2 angles correctly drawn in pie chart A1 fully correct chart labelled
(b)	0.38 + 0.27 + 0.15	0.20	2	M1 1 – sum A1 cao
11	0 | 5 7 8 8 1 | 0 0 0 0 2 5 5 5 6 2 | 0 0 0 4 5 3 | 3 5 Key 1 | 3 = 13 (min)	See working column	3	B1 for stem 0, 1, 2, 3 or 0, 10, 20, 30 B1 for accurate unordered leaves condone 1 error or omission B1 for key and ordered leaves all correct
12 (a)	$V = \pi \times 4^2 \times 10$	502 – 503	2	M1 for $\pi \times 4^2 \times 10$ A1 502 – 503
(b)	$P^2 = 10^2 + 8^2$ $P = \sqrt{164}$	$\sqrt{164} < 13$	3	M1 for sight of a correct right-angled triangle M1 for $10^2 + 8^2$ A1 for conclusion based on a correct calculation or 12.8 seen
13 (a)	2 × 30	2 × 2 × 3 × 5	4	M1 for systematic method, eg division, factor trees (at least one prime) A1 cao
	2 × 48	$2^5 \times 3$		M1 for systematic method, eg division, factor trees (at least one prime) A1 cao
(b)		12	1	B1 cao
(c)	$2^5 \times 3 \times 5$	480	2	B2 cao B1 for $2^5 \times 3 \times 5$ or any correct common multiple

Paper 5504

No	Working	Answer	Mark	Notes
14 (a)		$150 < C \leq 200$	2	M1 use of cum freq to find the cost of the 20^{th} or 20.5^{th} car OR $\frac{1}{2}\Sigma f$ or $\frac{1}{2}(\Sigma f + 1)$ A1 eg 150 to 200
(b)		No, because the 21^{st} value is in the same interval	1	B1 20.5^{th} or 21^{st} in same interval or an alternative correct explanation
(c)	80% = 5200 $\frac{5200}{80} \times 100$	6500	3	M1 for (100 – 20) % = 5200 M1 for $\frac{5200}{"80"} \times 100$ A1 cao
15 (a)	$x^2(x + 1) = 230$	AG	2	M1 for $x \times x \times (x+1)$ or $x \times x \times x + 1$ oe A1 cao from $x \times x \times (x+1)$
(b)	'5 – 150 5.1 – 158.7 5.2 – 167.6 5.3 – 177.0 5.4 – 186.6 5.5 – 196.6 5.6 – 207.0 5.7 – 217.7 5.8 – 228.8 5.9 –240.2 5.85 – 234.4	5.8	4	B 2 for trial between 5.8 and 5.9 inclusive (B1 for different trial between 5 and 6 inclusive) B1 for different trial between 5.8 and 5.85 (not including 5.8) B1 (dep on at least one previous B1) cao for 5.8, 5.81, 5.811

Paper 5504

No	Working	Answer	Mark	Notes
16	$\pi \times \left(\frac{15}{2}\right)^2 = 176.715$	88.4cm^2	3	M1 for $\pi \times \left(\frac{15}{2}\right)^2$ A1 88.3 – 88.4 B1(ind) for cm²
17 (a)	$5 = 0.5x + 1$	8	2	M1 for $5 = 0.5x + 1$ A1 cao
(b)		$y = \frac{1}{2}x + c$	1	B1 $y = \frac{1}{2}x + c$, c ≠ 1, oe
(c)		$x = 2y - 2$ OR $x = 2(y - 1)$	2	M1 for correctly multiplying both sides by 2 or correctly isolating $\frac{x}{2}$ A1 for $x = 2(y - 1)$, $x = \frac{y-1}{0.5}$, $x = \frac{y-1}{\frac{1}{2}}$ oe SC: B1 for $x = 2y - 1$
18	$4x - 6y = 22$ $15x + 6y = 54$ $19x = 76$	$x = 4,\ y = -1$	4	M1 for coefficients of x or y the same followed by correct operation , one arithmetical error A1 cao M1(dep on previous M mark) for sub for other variable A1 cao Trial and improvement –0 unless both x and y correct values found

Paper 5504

No	Working	Answer	Mark	Notes
19 (a)	$SF = \frac{10}{6}$ $\frac{10}{6} \times 4.8 = 8$	8	2	M1 for sight of $\frac{10}{6}$ or $\frac{6}{10}$ or 1.67 or better or $\frac{CD}{10} = \frac{4.8}{6}$ A1 cao
(b)	$\frac{10}{6} \times 4.5 - 4.5 = 3$	19.8	2	M1 for use of SF from (a) to find *BC* or *AC* and adding 4 sides A1 cao
20	$\frac{6 \times 10^{15}}{3.2 \times 10^{8}}$ 1.875×10^{7}	4.3×10^{3}	3	B3 for 4.3×10^{3} to 4.34×10^{3} (B2 for 1.875×10^{7} oe or 4300 to 4340 or final answer of 1.9×10^{7}) (B1 for sight of 6×10^{15} oe or 3.2×10^{8} oe)
21	$8.5 \times \tan 38 = 8.5 \times 0.7813$	6.64	3	M1 for correct use of trig, eg $\tan 38 = \frac{opp}{8.5}$ M1 for 8.5 × tan38 A1 6.64 – 6.641

Paper 5504


No	Working	Answer	Mark	Notes
22 (a)		No, as you would expect about 100. Yes, as it is possible to get 200 sixes with a fair dice	1	B1 for a consistent answer
(b)	$\frac{1}{6}, \frac{5}{6}$ + labels		3	B1 for $\frac{5}{6}$ on the red dice, *not six* branch B1 for a fully complete tree diagram with all branches labelled B1 for $\frac{1}{6}$, $\frac{5}{6}$ on all remaining branches as appropriate

Paper 5505

No	Working	Answer	Mark	Notes
1 (i) (ii) (iii)		119.31 119310 1.23	3	B1 cao B1 cao B1 cao
2	$\frac{10}{100}\times 12000 = 1200$ 12 000 – 1200 = 10 800 10 800 ÷ 10 =1080 10 800 – 1080 = £9720	9720	3	M1 for $\frac{10}{100}\times 12\,000$ or sight of 1200 or 2400 or 10 800 or 9600 M1 (dep) for $\frac{10}{100}\times(12\,000 - \frac{10}{100}\times 12\,000)$ or sight of 1080 A1 cao Alternative markscheme M2 for $12000\times\left(1-\frac{10}{100}\right)^2$ (M1 for $12000\times\left(1-\frac{10}{100}\right)$) A1 cao
3	$7r - 5r = -20 - 2$	–11	2	M1 for $7r + 2 = 5r - 20$ or $\frac{7r}{5} + \frac{2}{5} = r - 4$ or $7r - 5r = -20 - 2$ or $\frac{7r}{5} - r = -4 - \frac{2}{5}$ A1 cao

Paper 5505

No	Working	Answer	Mark	Notes
4 (a)		–1, 0, 1	2	B2 for –1, 0, 1 only (B1 for –1, 0 or 0,1 or –1, 1 or –2, -1, 0, 1 only)
(b)		(–1,–1)(0,–1) (1,–1) (0, 0) (1, 0) (1, 1)	3	B3 for 6 points correct (B2 for 3 points correct) (B1 for 1 point correct) *NB: –B1 for each additional point over six*
5		$5n + 1$	2	B2 oe (B1 for $5n$ seen) NB: n = gets B1 max
6 (a)	Triangle with vertices at (0,0) (0,–2) and (3,0)		2	M1 for correct orientation A1 cao
(b)	Rotation, 180°, centre (0,1) Enlargement sf –1 centre (0,1)		2	B2 for 180° 'rotation' centre (0, 1) or for Enlargement sf - 1 centre (0,1) (B1 for any two of the three parts) NB: B0 if additional transformation is included
7	Bisector of $\angle BAC$ Arc around A Region	See overlay	3	B3 cao (B2 for either two correct boundaries, no shading/ wrong shading or one correct boundary, one incorrect boundary with valid shading) (B1 for either two incorrect boundaries but one drawn from A and one intersection, with valid shading or one correct boundary) Ignore shading outside the triangle
8		Length Volume Area	3	B1 for Length B1 for Volume B1 for Area



Paper 5505

No		Working	Answer	Mark	Notes
9	(a)	Unbiased question with choices		2	B1 for unbiased question B1 for at least 2 choices
	(b)	Leading question and a restricted sample		2	Classification 1: A biased question Classification 2: A restricted sample of people Classification 3: Not specifying a range of foods Classification 4: Nothing to do with eating habits B2 reasons which satisfy 2 different classifications (B1 a reason which satisfies one classification)
10	(a)	$6 \times 10^2 \times 8 \times 10^4$ $48 \times 10^6 = 4.8 \times 10^7$	4.8×10^7	3	M1 for $6 \times 10^a \times 8 \times 10^b$ oe, a and b integers including 0 A1 for 48×10^6 oe A1 cao
	(b)	200 000 + 30 000 = 230 000	230 000	2	B2 cao (B1 for sight of 200 000 or 30 000 or 2.3×10^5 or 23×10^4)
11	(a)	$x^2 + xy + xy + y^2$	$x^2 + 2xy + y^2$	2	M1 for at least 3 of the 4 terms correct A1 cao
	(b)		25	2	M1 for recognising 3.47 + 1.53 (= 5) A1 cao
12	(i)	Tangent 90° to diameter /radius/ line from (through) centre	27°	4	B1 for 27° cao B1 for reason
	(ii)	180 –(90+ "27") angle in semicircle (is 90°)/Alternate segments /angle at centre twice at circumference	63°		B1 ft for 90–"27" if not 63° B1 for reason
13	(i)		p^9	1	B1 cao
	(ii)		$6q^6$	2	B2 for $6q^6$ (B1 for sight of $\frac{6q^9}{q^3}$ or $3q \times 2q^5$ or $3q^4 \times 2q^2$ or $6 \times q \times q \times q \times q \times q \times q$ or final answer of the form kq^6, $k > 0$)

Paper 5505

No	Working	Answer	Mark	Notes
14 (a)(i)		152	2	B1 cao
(ii)		177		B1 cao
(b)			3	B1 for median marked at 167 B1 ft for postion of box with its ends at "152" and "177" B1 for position of whiskers with ends at 132 and 182 NB: For any points plotted between 141 and 149 give a tolerance of an extra ± 1 square
15	(arc =) $\frac{40}{360} \times 2\pi \times 9$ $= 2\pi$	$2\pi + 18$	4	M1 for $\frac{40}{360} \times$ M1 for $2\pi \times 9$ M1 (dep) for $\frac{40}{360} \times 2\pi \times 9$ oe A1 for $\frac{18 \times \pi}{9} + 18$ oe exact form
16 (i)		1	1	B1 cao
(ii)		$\frac{1}{16}$	1	B1 cao accept 0.0625
(iii)		64	1	B1 cao condone ± 64

Paper 5505

No	Working	Answer	Mark	Notes
17 (a)	$F \propto 1/x^2$ $F = k/x^2$ $4 = k / 3^2$ $F = 36/x^2$	$F = \frac{36}{x^2}$	3	M1 for $F = k/x^2$ seen or implied. ($k \neq 1$) M1 (dep) for subst. or sight of $k = 36$ A1 for $F = 36/x^2$
(b)		9	1	B1 ft for 9 (ft on $F = kx^n$, $n \neq 0$)
(c)	$64 = \frac{36}{x^2}$ $x^2 = \frac{36}{64}$ $x = \pm\frac{3}{4}$	$\frac{6}{8}$	2	M1 for $x^2 = \frac{"36"}{64}$ A1 for $\frac{6}{8}$ oe (condone $\pm$) *SC: Use of $F = kx^2$ max M1 M1 A0 B1 ft M0 A0* *SC: Use of* $F = \frac{k}{\sqrt{x}}$ *max M1 M1 A0 B0 M1* ($\sqrt{x} = \frac{4\sqrt{3}}{64}$) *A0*
18	$(5+\sqrt{3})(5-\sqrt{3}) = 5\times5 - 5\sqrt{3} + 5\sqrt{3} - \sqrt{3}\sqrt{3}$ $= 5\times5 - 3$ $\frac{22}{\sqrt{22}} = \frac{22\sqrt{22}}{22}$	$\sqrt{22}$	3	B1 for correct expansion 25 - $5\sqrt{3}$ + $5\sqrt{3}$ - $\sqrt{3}$ $\sqrt{3}$ with 1st three terms reducing to 25 **without** any errors seen B1 (indep) for $\sqrt{3}$ $\sqrt{3}$ = 3 B1 for $\sqrt{22}$ coming from $\frac{22}{\sqrt{22}}$ (S.C $\frac{(5+\sqrt{3})(5-\sqrt{3})\sqrt{22}}{22}$ gets B1)

Paper 5505

No	Working	Answer	Mark	Notes
19 (a)		60 40	2	B1 cao B1 cao
(b)		correct bars	2	B1 for $30 < x \leq 40$ with an area of 2½ squares B1 for $40 < x \leq 70$ with an area of 3 squares *SC:* $\frac{0}{4}$ *give M1 if clearly using area or frequency density*
20 (a)	$6x + 8 - 12x + 15$	$-6x + 23$	2	M1 for 3 of the 4 terms $6x$, $+8$, $-12x$, $+15$ correct A1 cao
(b)		$32x^5y^{15}$	2	B2 cao (B1 for two of 32, x^5, y^{15})
(c)	$\frac{(n+1)(n-1)}{n+1} \times \frac{2}{n-2}$ $\frac{2(n-1)}{n-2}$	$\frac{2(n-1)}{n-2}$	3	M1 for $k(n + 1)(n - 1)$ M1 dep for $\frac{(n+1)(n-1)}{(n+1)} = n - 1$ A1 for $\frac{2(n-1)}{n-2}$
21	Vertices at $(-3, -1\frac{1}{2}), (-4\frac{1}{2}, -1\frac{1}{2}), (-3, -4\frac{1}{2})$		3	B1 for all sides $\times 1\frac{1}{2}$ B1 for correct orientation with 2 vertices almost correct B1 cao

Paper 5505

No	Working	Answer	Mark	Notes
22	Total = 3 + 5 + 2 (=10) $\frac{3}{10}\times\frac{3}{10}\times\frac{5}{10}\left(=\frac{45}{1000}\right), \frac{3}{10}\times\frac{3}{10}\times\frac{2}{10}\left(=\frac{18}{1000}\right)$ $\frac{5}{10}\times\frac{5}{10}\times\frac{3}{10}\left(=\frac{75}{1000}\right), \frac{5}{10}\times\frac{5}{10}\times\frac{2}{10}\left(=\frac{50}{1000}\right)$ $\frac{2}{10}\times\frac{2}{10}\times\frac{3}{10}\left(=\frac{12}{1000}\right), \frac{2}{10}\times\frac{2}{10}\times\frac{5}{10}\left(=\frac{20}{1000}\right)$ $3\times\left(\frac{"45"}{1000}+\frac{"18"}{1000}+\frac{"75"}{1000}+\frac{"50"}{1000}+\frac{"12"}{1000}+\frac{"20"}{1000}\right)$ $\frac{660}{1000}$	$\frac{660}{1000}$ oe	5	M3 for all six expressions seen OR their combined equivalents (M2 for four expressions seen OR their combined equivalents) (M1 for two expressions seen OR their combined equivalents) M1 sum of 18 relevant products condone 1 slip A1 for $\frac{660}{1000}$ oe *SC: without replacement maximum M4 A0* SC: Just 2 beads: Answer either $\frac{38}{100}$ oe OR $\frac{28}{90}$ oe B1

Paper 5505

No	Working	Answer	Mark	Notes
23 (a)(i)		$6\mathbf{b} - 6\mathbf{a}$	2	B1 for $6\mathbf{b} - 6\mathbf{a}$ oe
(ii)		$6\mathbf{a}$		B1 for $6\mathbf{a}$ oe
(b)	$\overrightarrow{EX} = \overrightarrow{EB} + \overrightarrow{BX}$ $= 12\mathbf{b} + \frac{1}{2}\overrightarrow{BC}$	$12\mathbf{b} - 3\mathbf{a}$	2	M1 for $\overrightarrow{EX} = \overrightarrow{EB} + \overrightarrow{BX}$ oe vector journey in a form ready for straightforward substitution A1 for $12\mathbf{b} - 3\mathbf{a}$ oe
(c)	$\overrightarrow{AY} = \frac{5}{3}\overrightarrow{AB}$ or $\overrightarrow{BY} = \frac{2}{3}\overrightarrow{AB}$ $\overrightarrow{EY} = 16\mathbf{b} - 4\mathbf{a}$ or $\overrightarrow{XY} = 4\mathbf{b} - \mathbf{a}$ $\overrightarrow{EY} = 4\overrightarrow{XY}$ or $\overrightarrow{EX} = 3\overrightarrow{XY}$ or $\overrightarrow{EY} = \frac{4}{3}\overrightarrow{EX}$	Printer Answer	3	B1 for either $\overrightarrow{AY} = \frac{5}{3}\overrightarrow{AB}$ or $\overrightarrow{BY} = \frac{2}{3}\overrightarrow{AB}$ oe B1 ft for either $\overrightarrow{EY} = 16\mathbf{b} - 4\mathbf{a}$ or $\overrightarrow{XY} = 4\mathbf{b} - \mathbf{a}$ ft only on parts (*a*) and (*b*) B1 for either $\overrightarrow{EY} = 4\overrightarrow{XY}$ or $\overrightarrow{EX} = 3\overrightarrow{XY}$ or $\overrightarrow{EY} = \frac{4}{3}\overrightarrow{EX}$ oe **plus** conclusion of *E*, *X*, *Y* on the same straight line
24 (a)(i)		$(5, -4)$	4	B1 cao
(ii)		$(2, -9)$		B1 cao
(iii)		$(2, 4)$		B1 cao
(iv)		$(1, -4)$		B1 cao
(b)		$(x-2)^2 - 4$	4	B4 for $(x-2)^2 - 4$ oe eg. $x^2 - 4x$ (B3 for $(x+2)^2 - 4$ or $(x-2)^2 + 4$) (B2 for $x^2 - 4$ or $(x-2)^2$ OR $x^2 + bx$, $b \neq 0$ OR $(x+2)^2 + 4$ OR $f(x-2) - 4$) (B1 for $x^2 + 4$ or $(x+2)^2$ or $ax^2 + bx$ or $x^2 + bx + c$ OR $x - 2 - 4$ or $x^2 - 2 - 4$, $a, b, c \neq 0$)

Paper 5506

No	Working	Answer	Mark	Notes
1 (a)	$V = \pi \times 4^2 \times 10$	502 – 503 cm^3	2	M1 for $\pi \times 4^2 \times 10$ A1 502 – 503
(b)	$P^2 = 10^2 + 8^2$ $P = \sqrt{164}$	$\sqrt{164} < 13$	3	M1 for sight of correct right angled triangle M1 for $10^2 + 8^2$ A1 for conclusion based on a correct calculation Or 12.8 seen
2 (a)(i)	2×30	$2 \times 2 \times 3 \times 5$	4	M1 for systematic method, eg division, factor trees (at least one prime) A1 cao
(ii)	2×48	$2^5 \times 3$		M1 for systematic method, division, factor trees (at least one prime) A1 cao
(b)		12	1	B1 cao
(c)	$2^5 \times 3 \times 5$	480	2	B2 cao B1 for $2^5 \times 3 \times 5$ or any correct common multiple

Paper 5506

No	Working	Answer	Mark	Notes
3 (a)		$150 < C \leq 200$	2	M1 use of cum freq to find the cost of the 20^{th} or 20.5^{th} car OR $\frac{1}{2}\Sigma f$ th or $\frac{1}{2}(\Sigma f + 1)$ th car. A1 eg 150 to 200, 150 – 200
(b)		No, because the 21^{st} value is in the same interval	1	B1 for 20.5^{th} or 21^{st} value in the same internal consistent with 'a' OR Refers to the median value being low in the interval (statement to be mathematically correct) See additional sheet
(c)	80% = 5200 $\frac{5200}{80} \times 100$	6500	3	M1 for (100 – 20)% = 5200 M1 for $\frac{5200}{"80"} \times 100$ A1 cao

Paper 5506

No	Working	Answer	Mark	Notes
4 (a)	$x^2(x+1) = 230$	AG	2	M1 for $x \times x \times (x+1)$ or $x \times x \times x + 1$ oe, $x^2(x+1)$, $x^2 \times x + 1$ A1 cao from $x \times x \times (x+1)$, no need to see 230
(b)	5 – 150 6 – 252 5.1 – 158.7 5.2 – 167.6 5.3 – 177.0 5.4 – 186.6 5.5 – 196.6 5.6 – 207.0 5.7 – 217.7 5.8 – 228.8 5.9 –240.2 5.85 – 234.4	5.8	4	B2 for trial between 5.8 and 5.9 inclusive evaluated (B1 for trial between 5 and 6 inclusive evaluated) B1 for different trial between 5.8 and 5.85 (not including 5.8) B1 dep on at least are previous B1 5.8, 5.81, 5.811
5	$\pi \times \left(\frac{15}{2}\right)^2 = 176.715$	88.4 cm^2	3	M1 for $\pi \times \left(\frac{15}{2}\right)^2$ seen A1 88.3 – 88.4 B1 for cm² (independent)

Paper 5506

No	Working	Answer	Mark	Notes
6 (a)	$5 = 0.5x + 1$	8	2	M1 for $5 = 0.5x + 1$ A1 cao
(b)		$y = \frac{1}{2}x + c$	1	B1for $y = \frac{1}{2}x + c$, $c \neq 1$, oe
(c)		$x = 2y - 2$ OR $x = 2(y - 1)$	2	M1for correctly multiplying both sides by 2 or correctly isolating $\frac{x}{2}$ A1 for $x = 2(y - 1)$, $x = \frac{y-1}{0.5}$, $\frac{y-1}{\frac{1}{2}}$ oe SC B1 for x= 2y -1
7	$4x - 6y = 22$ $15x + 6y = 54$ $19x = 76$	$x = 4$, $y = -1$	4	M1 for coefficients of x or y the same followed by correct operation, allow one arithmetical error A1 cao M1 (dep) for correct sub for other variable A1 cao Trial and improvement 0 marks unless both correct values of x and y found
8 (a)	SF = $\frac{10}{6}$ $\frac{10}{6} \times 4.8 = 8$	8	2	M1 for sight of $\frac{10}{6}$ or $\frac{6}{10}$ or 1.67 or better or $\frac{CD}{10} = \frac{4.8}{6}$ A1 cao
(b)	$\frac{10}{6} \times 4.5 - 4.5 = 3$	19.8	2	M1 for use of SF from "a" to find AC or BC or $\frac{BC}{4.5} = \frac{4}{6}$ and adding 4 sides A1 cao

Paper 5506

No	Working	Answer	Mark	Notes
9	$\frac{6\times10^{15}}{3.2\times10^{8}}$ 1.875×10^{7}	4.3×10^{3}	3	B3 for 4.3×10^{3} to 4.34×10^{3} (B2 for 1.875×10^{7} oe or 4300 to 4340, final answer of 1.9×10^{7} B1 for sight of 6×10^{15} oe or 3.2×10^{8} oe)
10	$8.5\times\tan38$ $=8.5\times0.7813$ $\frac{8.5}{\sin(90-38)}=\frac{AB}{\sin38}$ $AB=\frac{8.5\times\sin38}{\sin(90-38)}$ $=\frac{5.2331}{0.788}=6.64$	6.64	3	M1 for correct use of trig, eg $\tan 38=\frac{opp}{8.5}$ M1 for $8.5\times\tan38$ A1 6.64 – 6.641 OR M1 for correct substitution into the sine rule M1 (dep) for correct rearrangement for AB= A1 6.64 – 6.641
11 (a)		No, as you would expect about 100. Yes, as it is possible to get 200 sixes with a fair dice	1	B1 for a consistent answer See additional sheet

Paper 5506

No	Working	Answer	Mark	Notes
(b)	$\frac{1}{6}, \frac{5}{6}$ + labels		3	B1 for $\frac{5}{6}$ on the red dice, *not six* branch B1 for a fully complete tree diagram with all branches labelled B1 for $\frac{1}{6}$ and $\frac{5}{6}$ on all remaining branches as appropriate
(c)(i)	$\left(\frac{1}{6}\right)^2$	$\frac{1}{36}$	2	M1 $\left(\frac{1}{6}\right)^2$ or $\frac{1}{6}\times\frac{1}{6}$ only or 0.28 A1 $\frac{1}{36}$ or 0.03 or better
(ii)	$1-\left(\frac{5}{6}\right)^2$ OR $\frac{1}{6}\times\frac{5}{6}+\frac{5}{6}\times\frac{1}{6}+\frac{1}{6}\times\frac{1}{6}$	$\frac{11}{36}$	3	M2 for $1-\left(\frac{5}{6}\right)^2$ or $1-\frac{5}{6}\times\frac{5}{6}$ A1 cao OR M1 for $\frac{1}{6}\times\frac{5}{6}$ oe M1 for 2 or 3 only of $\frac{1}{6}\times\frac{5}{6}, \frac{5}{6}\times\frac{1}{6}$, "a" A1 for $\frac{11}{36}$ or 0.31 or better

Paper 5506

No	Working	Answer	Mark	Notes
12 (a)	$\pi \times 30 \times \frac{7.5^2}{3} - \pi \times 10 \times \frac{2.5^2}{3} = 1767 - 65$	1700	3	M1 for either $\pi \times 30 \times \frac{7.5^2}{3}$ or $\pi \times 10 \times \frac{2.5^2}{3}$ M1 (dep) for difference A1 1700 – 1702 **SC** B1 Using d instead of r, 6800 – 6808
(b)	$\frac{S}{2\pi d} = \sqrt{h^2 + d^2}$ $\left(\frac{S}{2\pi d}\right)^2 = h^2 + d^2$	$h = \sqrt{\frac{S^2 - 4\pi^2 d}{4\pi^2 d^2}}$	3	M1 for correctly isolating $\sqrt{h^2 + d^2}$ or $h^2 + d^2$ or $h + d$ or kh^2 or kh M1(indep) squaring both sides A1 $h = \sqrt{\frac{S^2 - 4\pi^2 d^4}{4\pi^2 d^2}}$, $h = \frac{\sqrt{S^2 - 4\pi^2 d^4}}{2\pi d}$ $h = \sqrt{\left(\frac{S}{2\pi d}\right)^2 - d^2}$
(c)	$\left(\frac{30}{20}\right)^2 \times 450$ or $450 \div \left(\frac{20}{30}\right)^2$	1012.5	2	M1 for sight of correct SF^2 including 4:9 A1 1010 to 1013

Paper 5506

No	Working	Answer	Mark	Notes
13 (a)	$\frac{2x(x+20)}{2} = 400$	As given	2	M1 $\frac{2x(x+20)}{2}$ or $\frac{2x \times x + 20}{2}$ or $2x(x + 20) = 800$ A1 cao following correct working, no need for = 400 *SC B1* $2x \times x + \frac{1}{2} \times 2x(10 - \frac{x}{2}) \times 2$
(b)	$\frac{-20 \pm \sqrt{20^2 - 4 \times 1 \times (-400)}}{2}$ $= \frac{-20 \pm 44.721}{2}$	12.361	3	M1 for correct sub, up to signs, in the quad formula A1 for 44.7 or $\sqrt{2000}$ A1 for 12.3606 – 12.361, ignore negative solution T.I B3 for 12.361 OR Completing the square M1 for $(x+10)^2$ seen A1 for $-10 \pm \sqrt{500}$ A1 for 12.3606 – 12.361 ignore negative solution

Paper 5506

No	Working	Answer	Mark	Notes
14 (a)	$0.5\times 8\times 15\times \sin 70^0$	56.4	2	M1 for correct sub into area formula A1 56.38 – 56.4
(b)	$AB^2 = 8^2 + 15^2 - 2\times 8\times 15\times \cos 70^0 = 206.9$ EITHER $0.5\times AB\times CX$ ='56.38' OR $\frac{\sin B}{8} = \frac{\sin 70}{\sqrt{'206.9'}}$ $B = 31.5$ 15 sin '31.5'	7.84	4	M1 for correct sub into cos rule A1 for 206.9 - 207 or 14.38 – 14.4 EITHER M1 for use of area rule to find CX A1 7.83 – 7.84 OR M1 for correct use of sine rule to find sin B or sin A and then sight of 15sin B *or* 8 sin A A1 7.83 – 7.84

Paper 5506

No	Working	Answer	Mark	Notes
15 (a)	$4a^2-4a+1-(4b^2-4b+1)=$ $4(a^2-b^2)-4(a-b)$ $4(a-b)(a+b-1)$ OR $((2a-1)-(2b-1))((2a-1)+(2b-1))$ $(2a-2b)(2a+2b-2)$	AG	3	Expansion Method M1 for a correct expansion of any one of the three terms M1(dep) on an attempt to expand all 3 terms and show LHS = RHS A1 fully correct algebra RHS exp is $4(a^2+ab-a-ba-b^2+b)$ OR Factorisation Method M1 for attempt to use difference of 2 squares on LHS M1 for one bracket correctly simplified A1 fully correct
(b)	Any 2 odd square numbers have the above form If a and b are both even or odd then $a-b$ is even, so $4(a-b)$ is a multiple of 8 If one of a,b is odd, then $a+b-1$ is even, so $4(a+b-1)$ is a multiple of 8		3	B1 'any 2 square nos have the above form' (may be implied by sight of $(2a-1)^2-(2b-1)^2$ in part (b)) B1 first reason B1 second reason SC B1 for $(2r+1)^2-(2r-1)^2$ B1 for 8r
16 (a)	$g_L=\frac{2\times 4.495}{1.35^2\times\sin 30.5}$ $g_u=\frac{2\times 4.505}{1.25^2\times\sin 29.5}$	9.719 11.710	4	B2 for any 4 of 4.505, 1.25, 29.5, 4.495, 1.35, 30.5 seen (B1 for any two or three seen) B1 for 11.710 – 11.7103 B1 cao 9.719 – 9.71904
(b)	Round, until lower and upper bounds agree	10	1	B1 for 10 + reason " they agree to this level of accuracy"

Paper 5506

No	Working	Answer	Mark	Notes
17 (a)(i)		xy	3	B1 cao
(ii)		y^2		B1 for y^2 or $y \times y$
(iii)		$\frac{x}{2}$		B1 for $\frac{x}{2}$ or $0.5x$ or $2^{-1}x$
(b)	Divide to get $2y = 1$	$q = -1$	2	M1 for $2y = 1$ or $\frac{x}{2} = 32$ or $p + q = 5$ or $1 + p + 2q = 5$
		$p = 6$		A1 cao
18 (a)	$x^2 - 2mx + m^2 - k$	$k = m^2$	2	M1 for correct exp of $(x - m)^2$ or correct completion of the square eg $\left(x - \frac{2m}{2}\right)^2 - \left(\frac{2m}{2}\right)^2$
				A1 cao
				SC B1 for $k = -m^2$
(b)(i)	Min value is $-m^2$	$-m^2$	3	M1 for recognition that min value occurs when $(x - m)^2 = 0$ (either (b)(i) or (b)(ii) correct implies this M1)
(ii)	$x = m$	m		A1 ft on 'k', " $-k$" gets M1 A0
				A1 cao
19	$0.06 \times 0.05 = 0.003$	No	2	M1 for 0.06×0.05
				A1 correct conclusion based on 0.003 or 0.06 x 0.05 stated as $\neq 0.0011$
				OR M1 for statement that for the two events to be independent P (BL and CL) = P(BL) × P(CL)

Paper 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
20		50 50 4	3	B1 50 or $\frac{100}{2}$ B1 for 50 or "a" B1 4 or $\frac{360}{90}$ oe

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4LN

Telephone 01623 467467
Fax 01623 450481

Order Code UG014129 June 2003

For more information on Edexcel qualifications please contact our
Customer Response Centre on 0870 240 9800
or email: enquiries@edexcel.org.uk
or visit our website: www.edexcel.org.uk

London Qualifications Limited. Registered in England and Wales no.4496750
Registered Office: Stewart House, 32 Russell Square, London WC1B 5DN



edexcel

GCSE
Edexcel GCSE
Mathematics A 1387

July 2004

advancing learning, changing lives

Mark Scheme

Edexcel GCSE
Mathematics A 1387

London Qualifications is one of the leading examining and awarding bodies in the UK and throughout the world. It incorporates all the qualifications previously awarded under the Edexcel and BTEC brand. We provide a wide range of qualifications including general (academic), vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, our centres receive the support they need to help them deliver their education and training programmes to learners.

For further information please call Customer Services on 0870 240 9800, or visit our website at www.edexcel.org.uk

July 2004

Publications Code UG015616

All the material in this publication is copyright
© London Qualifications Ltd 2004

CONTENTS

Notes on Marking Principles....2
Paper 5514....4
Paper 5515....8
Paper 5516.... 11
Paper 5517.... 15
Paper 5518.... 19
Paper 5519.... 23

NOTES ON MARKING PRINCIPLES

NOTES ON MARKING PRINCIPLES

1 Three types of mark are available

M marks - awarded for correct working seen or implied.
A marks - conditional accuracy marks which are awarded for accurate working following the award of M marks.
B marks - unconditional accuracy marks (independent of M).

2 Abbreviations

cao - correct answer only.
ft - follow through.
$\surd$ - Denotes a "follow through" answer.

SC - special case.
isw - ignore subsequent working.
oe - or equivalent (and appropriate).
NB: a candidate cannot benefit from both isw and ft.

3 If no working shown, then

correct answers normally score full marks
incorrect (even though nearly correct) answers score no marks.

4 Marking instructions

Misread loses A marks (and sometimes B marks) on that part but ft can be allowed on subsequent parts. M marks can still be earned. If in doubt contact your team leader.
If there is a wrong answer in the answer space DO CHECK the working in the body of the script.

- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.
- If there is a wrong answer in the answer space DO CHECK the working in the body of the script.
- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.

5 Style of marking

Answer correct: tick and write part mark in margin NEXT TO BRACKETED MARK.
Answer incorrect: cross, but show M, A or B marks if any earned in body of script and transfer the total of these to the margin next to the bracket mark.
Total for each double page at bottom right page (except for back if used), FINAL TOTAL IN RELEVANT BOX ON FRONT COVER.
Nought in margin for fully incorrect question or page AND FOR NO ATTEMPT.
Where no attempt has been made a line should be put in the answer space and zero in the margin next to the bracketed mark
CHOICE OF METHOD – No marks unless one answer is in answer space – then mark that.
CROSSED OUT WORK - if not replaced this should be marked (if legible).
There must always be a mark next to bracketed mark in the margin/

6 Follow Through Marks

Follow throughs are guided by two principles:

(a) Follow throughs which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous, do not award.
(b) Follow throughs which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

7 Probability

Probability answers must be given as fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least two decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.

6 Linear Equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

8 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

PAPER 5501					
No		**Working**	**Answer**	**Mark**	**Notes**
1			K: 1020 L: 8.06	2	B1 for 1020 or 1,020 B1 for 8.06 Accept £8.06p and £8,06
2	(a)		28, 33	2	B1, B1 (B1 ft for "28" + 5 if both numbers >25)
	(b)			1	B1 for add 5, +5, for going up in 5's the difference is 5 oe
	(c)			1	B1 for they end in 3 and 8, it ends in 7; or refers to 383 and 388 are in sequence
3	(a)(i)		$\frac{1}{2}$	2	B2 for ½ accept half (B1 for an equivalent unsimplified fraction eg 4/8 or 50%
	(ii)		2 rectangles shaded	1	or 0.5) B1 for correct shading (any 2 rectangles)
	(b)(i)		40	4	B1 for 40 cao
	(ii)		150 000		B1 for 150 000 cao (accept 150,000 not 150.000)
	(iii)		6.55		B1 for 6.55 cao not $6.5^{½}$
	(iv)		$\frac{3}{8}$		B1 for $\frac{3}{8}$ oe accept 0.375
	(c)		Cross 3cm from *A*	1	B1 mark a cross 3 cm (±2 mm) from *A*
4	(a)(i)		metres	2	B1 for metres (m)
	(ii)		grams		B1 for grams (g)
	(b)		miles	1	B1 for miles
5	(a)		(3, 2)	1	B1 for (3, 2)
	(b)(i)		*Q* at (0, 3)	1	B1 for *Q* plotted correctly on *y*-axis at (0, 3) ±2mm
	(ii)		*R* at (−2, −3)	1	B1 for *R* plotted correctly at (−2, −3) ±2mm
6	(a)(i)		30	2	B1 for 30
	(ii)		25		B1 for 25
	(b)(i)		4 shapes	2	B1 for drawing 4 shapes
	(ii)		1½ shapes		B1 for drawing 1½ shapes

PAPER 5501					
	No	Working	Answer	Mark	Notes
7	(a)		30	1	B1 for 30
	(b)	$20 \times 3 + 5$	65	2	M1 for $20 \times 3 + 5$ oe A1 for 65
8	(a)		Draws perp.	1	B1 for correctly drawing perp must touch line or cut line *AB* ± 2mm
	(b)		Sketches a cylinder	1	B1 for sketching cylinder
9	(a)	1974 6580 8554	855.4	3	M1 for complete method with relative place value correct, condone 1 error in multiplication A2 cao (A1 for digits 8554 seen or A1 for "855.4" dependent on 1 arithmetic error)
	(b)	990 ÷ 69 = 14.3…. or 14 rem 24	14	4	Method 1 - Everything excluding long division M2 for a valid method with no errors …need to see 966 or 1035 (M1 for a valid method, including estimation, that reaches an answer between 901 and 1099 inclusive. Ignore errors.) A2 for 14 with a fully correct method (A1 (dep on 1st M1) for an answer between 14 and 15 inclusive. Ignore errors. Cannot have used estimation.) Note: Estimation can only score a maximum of M1 unless accompanied by another method or 966 or 1035 see (award M2) Method 2 – long division method M2 for 990 ÷ 69 = integer answer between 12 and 19 inclusive and rem 30 seen (M1 for 990 ÷ 69 with 1 in tens column.) A2 for 14 with fully correct method (A1 (dep on 1st M1) for an answer between 14 and 15 inclusive.)

PAPER 5501					
	No	Working	Answer	Mark	Notes
10	(a)(i)		7	2	B1 for 7
	(ii)		−10		B1 for −10
	(b)(i)		6	2	B1 for 6 (accept −6)
	(ii)		8		B1 for 8 (accept −8)
	(c)		−7	1	B1 for −7 cao
11	(a)		B and D	2	B1, B1 (–1 each extra)
	(b)(i)		A	2	B1 for A
	(ii)		3		B1 for 3
12	(a)	800 − 144	656	2	M1 for at least 1 digit correct and in correct position needs to be 3 digit number A1 for 656
	(b)	144 is less than 200 so Trudy is wrong		2	B1 for 200 B1ft for 'correct' explanation based on cand's "200"
	(c)	$\frac{45\times800}{100}$	360	2	M1 for 45 × 800 ÷ 100 oe A1 for 360
	(d)	$\frac{176}{800}\times100$	22	2	M1 for $\frac{176}{800}\times100$ oe A1 for 22
13	(a)(i)		25	4	B1 for 25
	(ii)		28		B1 for 28
	(iii)		5 and 20		B1 for 5 and 20
	(iv)		26 and 33		B1 for 26 and 33
	(b)	$2^3 = 8$ or $2^3 = 2 \times 2 \times 2$ (which is not 6)		1	B1 for valid 'explanation'
14			10 m^2	2	B1 for 10 B1 indep for m^2.

PAPER 5501					
No		**Working**	**Answer**	**Mark**	**Notes**
15	(a)	$\frac{11}{12}-\frac{10}{12}$	$\frac{1}{12}$	2	M1 for correctly writing both fractions to a common denominator A1 for $\frac{1}{12}$ oe
	(b)	$\frac{70\times400}{200}$	140	2	B2 for 140 (accept 136) (B1 for sight of any two of 70, 400 or 200)
16	(a)		$2y$	1	B1 for $2y$ or $2\times y$, $y2$, $y\times2$
	(b)		$3p^2$.	1	B1 for $3p^2$ or $3\times p^2$ or $p^2\times3$ or $p^2 3$
	(c)		$x(x-3)$	2	B2 for $x(x-3)$ (B1 for x (x…….)) $\frac{25}{60}$ wrongly cancelled gets B1 ISW
17	(a)		2 23 9 34 15 2 9 26 17 25 18 60	3	B3 for all correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
	(b)		$\frac{25}{60}$	1	B1 for $\frac{25}{60}$ or $\frac{5}{12}$ oe $\frac{25}{60}$ wrongly cancelled gets B1 ISW

PAPER 5501					
	No	**Working**	**Answer**	**Mark**	**Notes**
18	(a)(i)		60	2	B1 for 60
	(ii)				B1 for all **angles equal** so equilateral triangle oe
	(b)(i)		130	3	B1 for 130
	(ii)				B1 for isosceles triangle oe or 2 angles equal accept $\angle Q = \angle R$ B1 for angles on a straight **line** add up to **180°**oe (180° could be in working)
	(c)		64	1	B1 for 64
19	(a)		1	1	B1 for 1
	(b)		3	1	B1 for 3
	(c)	$9\times1+3\times2+5\times3+3\times4$ (= 42) "42" ÷ 20	2.1	3	M1 for 9×1, 3×2, 5×3, 3×4 or for 42 seen M1 (dep) for "42" ÷ 20 A1 for 2.1 or $2\frac{1}{10}$ or $2\frac{2}{20}$
20	(a)		09 05	1	B1 for 09 05 oe
	(b)		7	1	B1 for 7
	(c)		10	1	B1 for 10
	(d)	7km in 20 mins	21	3	B1 for 20 (minutes) oe or $\times3$ seen M1 for $\frac{"7"}{"20"}$ A1 cao [SC: "7" × 3 seen gets B1 M1]


PAPER 5501				
No	**Working**	**Answer**	**Mark**	**Notes**
21	Vertices of new triangle at (2, −2) (2, 6) and (6, −2)	Correct triangle drawn	3	B3 cao (B2 for either correct triangle in wrong translated position or for 2 vertices correct) (B1 a right angled triangle with horizontal length 4 or vertical length 8 in the same orientation as the shaded triangle) SC B2 for s + 3 totally correct
22 (a) (b) (c)	 8560 ÷ (10 × 10)	8 Pentagon 85.6	1 1 2	B1 for 8 B1 for pentagon M1 for 8560 ÷ (10 × 10) oe A1 for 85.6
23	$2x + x + 100 + 47 = 360$ $2x + x = 360 - 100 - 47$ $x = 71$ Largest angle = $2x$ =	142	4	M1 for $2x + x + 100 + 47 = 360$ or 360 – 147 or 213 seen M1 dep for correctly separating x-terms and non x terms or "360 – 147" ÷ 3 A1 for $x = 71$provided M2 awarded A1 ft for 142

PAPER 5502					
	No	**Working**	**Answer**	**Mark**	**Notes**
1	(a)		Diagram	1	B1 cao
	(b)		22, 26	1	B1 cao
	(c)		46	1	B1 cao
2	(a)	£5.95 + £1.62 + 25p + 2 × 48p = £8.78	1.22	3	M1 for 3 of £5.95, £1.62, 25p, 48p seen with intention to add or 170 seen A1 for £8.78 A1 ft from " 8.78"
	(b)	160 ÷ 12 = 13.(333…)	14	2	M1 for 160 ÷ 12, or 13 or better or 12 × 13 or 12 × 14 A1 14 cao
	(c)	"14" × £12.20 = £170.8**0**	170.80	2	M1 for "14" × 12.20 A1 for 170.8**0** or 158.6**0** if 13 in (b) NB: 170.8 or 158.6 gets M1 A0
3	(a)(i)		1459	1	B1 cao
	(ii)		9541	1	B1 cao
	(b)		9 + 5 = 14	1	B1 cao
	(c)		0	1	B1 cao
4	(a)		1123	1	B1 cao
	(b)		1125	1	B1 cao
	(c)		43	1	B1 cao
	(d)	67 – "43"	24	2	M1 for 67 – "43" A1 ft from "43"
	(e)(i)		80	2	B1 cao
	(ii)		56 – 58		B1 answer in range 56 to 58 inc
	(f)		Completes graph	2	B2 for (Apr , 70) and (May , 60 < p < 65) plotted and joined (B1 for one point plotted)

PAPER 5502					
	No	Working	Answer	Mark	Notes
5	(a)(i)		4	1	B1 cao
	(ii)		10	1	B1 cao
	(b)		28	2	B2 for 28 (B1 for 29 or 14×2)
6	(a)		$\frac{3}{4}$	1	B1 accept 0.75, 75%
	(b)	$\frac{3}{4} \times 28 = 21$	15.54	3	M1 for $\frac{3}{4} \times 28$ or 21 seen M1 for "21" $\times$ 74 or 28×74 or 7×74 A1 cao
	(c)(i)		44 – 46	2	B1 44 – 46 inc
	(ii)		5.2 – 5.4		B1 5.2 – 5.4 inc
7	(a)		Correct shape	1	B1
	(b)		Correct shape	1	B1
8	(a)		Hexagon	1	B1
	(b)		Right	1	B1 cao
	(c)		Drawing	2	B2 for 7 additional hexagons with at least 2 points at which 3 hexagons meet (B1 for one point at which 3 hexagons meet)
9	(a)		$\frac{3}{4}$	2	M1 $\frac{80-20}{80}$ or $\frac{60}{80}$ or 0.75 (oe) A1 cao Do not accept equivalents
	(b)		15	1	B1 cao
	(c)		Iron and Cook set	1	B1 for both Accept mixer and fryer
	(d)		Microwave	1	B1 cao
	(e)		Bar chart	2	B1 one column (mixer): 70, 90 B1 second column (fryer): 70, 85 [SC: B1 columns correct but reversed]

PAPER 5502					
	No	Working	Answer	Mark	Notes
10	(a)		Acute	1	B1
	(b)		Reflex	1	B1
	(c)		reason	1	B1 reason eg 120 + 230 = 350 ≠ 360
11	(a)		$\frac{1}{2}$	1	B1 for $\frac{1}{2}$ oe
	(b)		reason	1	B1 for reason e.g could get 30 heads
12	(a)	7.5×4	30	2	M1 for 7.5×4 or $7\frac{1}{2} \times 4$ A1 cao
	(b)	$\sqrt{(42.25)}$	6.5	2	B2 for 6.5 (B1 for 42.25)
13	(a)		$6x$	1	B1 accept $6 \times x$, $x6$ oe
	(b)		$x - 4$	1	B1 cao
	(c)		$12(x - 4)$	1	B1 for $12(x - 4)$ or $12x - 48$ or $12 \times x - 48$ oe
	(d)		$18x - 48$	2	M1 for "$6x$" + "$12(x-4)$" or "$6x$" + "$12x - 48$" oe A1 ft $18x - 48$, $2(9x - 24)$, $3(6x - 16)$, $6(3x - 8)$
14	(a)		90	2	B2 for 90 ± 2 (B1 for 9 ± 0.2)
	(b)		317	1	B1 for 317 ± 4
15	(a)	Accurate drawing of triangle		2	B1 cao for 38° (±2°) B1 cao (indep) for BC drawn 7.3 cm (±2mm) and completing the triangle.
	(b)	Measure angle A	63°	1	B1 for 63° (±2°) or ft their diagram (±2°)

PAPER 5502					
	No	Working	Answer	Mark	Notes
16	(a)	0 × 1 = 0 1 × 1 = 1 2 × 4 = 8 3 × 8 = 24 4 × 8 = 32 5 × 7 = 35	100	3	M1 col 1 × col 2 (at least 3 shown) M1 (dep) sum of totals A1 cao SC: M1 M1 A0 for 101
	(b)	Either 35.50 ÷ 1.42 = 25; 26.99 – 25 = 1.99 Cheaper in the USA Or 26.99 × 1.42 = 38.33; 38.33 – 35.50 = 2.83 Cheaper in the USA		3	M1 35.50 ÷ 1.42 A1 25 seen B1 ft (dep on M1) £1.99 with conclusion (must have units) **OR** M1 26.99 × 1.42 A1 38.32 or 38.33… seen B1 ft (dep on M1) \$2.83 or \$2.82 with conclusion (must have units) must be to 2dp
17	(a)		-1, (1), (3), 5, 7, 9	2	B2 cao (B1 for 2 values)
	(b)		Graph	2	B1 ft for plotting points ±1/2 square B1 cao for line between $x = -2$ and $x = 3$
	(c)(i)		0.4	2	B1 for 0.4 or ft from single straight line with positive gradient
	(ii)		1.2		B1 for 1.2 or ft from single straight line with positive gradient

PAPER 5502					
	No	Working	Answer	Mark	Notes
18	(a)		Plots	1	B1 cao
	(b)		description	1	B1 dynamic relationship or "positive" (correlation)
	(c)		line of best fit	1	Line within overlay region, and to the extent of.
	(d)(i)	280 g →	reading	2	B1 ft from single straight line of positive gradient (±1/2 square)
	(ii)	120 pages →	reading		B1 ft from single straight line of positive gradient (±1/2 square)
19			200 150 225 150 10	3	B3 cao (B2 for three correct, or B1 for one correct)
20	(a)			2	B1 B1 for correct sketch – ignore additional internal lines B1 B0 for rectangle or right-angled triangle or correct shape without line NB 3-D sketch gets B0 B0
	(b)			2	B1 B1 for correct plan – ignore internal lines – accept a rotated plan, not reflected B1 B0 for a single rectangle drawn NB 3-D sketch gets B0 B0
21	(a)	163.25 – 35.50 = 127.75 127.75 ÷ 18.25 = 7	8	3	M1 163.25 – 35.50 (or sight of 127.75) M1 (dep) "127.75" ÷ 18.25 A1 cao SC: M2 for 7 days
	(b)	1.175 × £64 or $\frac{17.5}{100}$ × 64 or £11.20 seen	75.20	2	M1 1.175 × 64, 0.175 × 64 oe or 11.20 or 75.2 seen or 6.40, 3.20 and 1.60 seen A1 cao

PAPER 5502					
	No	Working	Answer	Mark	Notes
22	(a)		6	1	B1 cao
	(b)		$t^2 - 2t$	1	B1 oe
	(c)		$3(y - 4)$	1	B1 Accept $3 \times (y - 4)$ or $3(y - 4$
23	(a)		Reason	1	B1 Makes some mention of bias either directly or making reference to an insufficient or biased range of responses
	(b)		Reason	1	B1 (a) an insufficient range of responses (b) No mention of money (c) No time frame in the question (d) Misunderstanding of "A lot" and "Not much"



PAPER 5503				
No	**Working**	**Answer**	**Mark**	**Notes**
1		10.	1	B1 for 10
2 (a)(i)		$3e + 2f$	2	B1 for $3e + 2f$
(ii)		$3p^2$		B1 for $3p^2$
(b)	$5 \times -3 + 1$	-14	2	M1 for $5 \times -3 + 1$ A1 for -14
3	1974 6580 8554	855.4	3	M1 for complete correct method with relative place value correct, condone 1 error in multiplication. A2 cao (A1 for digits 8554 seen or A1 ft for "855.4" dependent on one arithmetic error only)
4 (a)		2 23 9 34 15 2 9 26 17 25 18 60	3	B3 all correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
(b)		$\frac{25}{60}$	1	B1 for $\frac{25}{60}$ or $\frac{5}{12}$ oe
5 (a)	$\frac{45}{100} \times 800$	360	2	M1 for $\frac{45}{100} \times 800$ oe A1 for 360
(b)	$\frac{176}{800} \times 100$	22	2	M1 for $\frac{176}{800} \times 100$ oe A1 for 22

PAPER 5503					
	No	Working	Answer	Mark	Notes
6	(a)	$\frac{11}{12} - \frac{10}{12}$	$\frac{1}{12}$	2	M1 for correctly writing both fractions to a common denominator A1 for $\frac{1}{12}$ oe
	(b)	$\frac{70 \times 400}{200}$	140	2	B2 for 140 (accept 136) (B1 for sight of any two of 70, 400 or 200)
7	(a)(i)		60	2	B1 for 60
	(ii)				B1 for all angles equal so equilateral triangle oe
	(b)(i)		130	3	B1 for 130
	(ii)				B1 for isosceles triangle oe B1 for angles on a straight line add up to 180°oe
	(c)		64	1	B1 for 64
8		$9 \times 1 + 3 \times 2 + 5 \times 3 + 3 \times 4$ (=42) "42" ÷ 20	2.1	3	M1 for 9×1, 3×2, 5×3, 3×4 or for 42 seen M1 (dep) "42" ÷ 20 A1 for 2.1 or $2\frac{1}{10}$ or $2\frac{2}{20}$
9	(a)		09 05	1	B1 for 09 05 oe
	(b)		7	1	B1 for 7
	(c)		10	1	B1 for 10
	(d)	7 km in 20 minutes	21	3	B1 for 20 (minutes) oe or $\times 3$ seen M1 for $\frac{"7"}{"20"}$ A1 cao SC: "7" $\times 3$ gets B1 M1

PAPER 5503					
No		**Working**	**Answer**	**Mark**	**Notes**
10		Vertices of new triangle at (2, −2) (2, 6) and (6, −2)	Correct triangle drawn	3	B3 cao (B2 for either correct triangle in wrong translated position or for 2 vertices correct) (B1 for a right-angled triangle with horizontal length 4 or vertical length 8 in the same orientation as shaded triangle) SC B2 for scale factor 3, totally correct
11		$2^2 + 3 =$	7 which is not even	2	B2 (B1 for correctly evaluating $n^2 + 3$ with a prime number value for n.)
12	(a)	360 ÷ 5	72	2	M1 for 360 ÷ 5 oe A1 for 72
	(b)	8560 ÷ (10 × 10)	85.6	2	M1 for 8560 ÷ (10 × 10) oe A1 for 85.6
	(c)(i)	Least length = 100.5	100.5	2	B1 for 100.5
	(ii)	Greatest length = 101.5	101.5		B1 for 101.5 or 101.499 or better
13		$2x + x + 100 + 47 = 360$ $2x + x = 360 - 100 - 47$ $x = 71$ Largest angle = $2x$ =	142	4	M1 for $2x + x + 100 + 47 = 360$ or 360 – 147 or 213 seen M1 (dep) for correctly separating x-terms and non x terms or $\frac{"360-147"}{3}$ A1 for $x = 71$ A1 ft (provided M2 awarded) for "142"
14	(a)(i)		2.21	2	B1 for 2.21
	(ii)		0.013		B1 for 0.013
	(b)	LCM = 3 × 13 × 17 = 3 × 221	663	2	M1 for 3 × 13 × 17 oe A1 for 663



PAPER 5503					
No		**Working**	**Answer**	**Mark**	**Notes**
15			Tick boxes 1,3 & 6	3	B1 for $\frac{\pi abc}{2d}$; B1 for $2a^2$; B1 for $2\left(c^2+d^2\right)$ (−B1 for each additional expression ticked (>3) to a minimum of 0)
16		200×0.2	40	2	M1 for 200×0.2 or $\frac{40}{200}$ seen A1 for 40
17	(a)	eg $2 \mid \underline{108}$ $2 \mid \underline{54}$ $3 \mid \underline{27}$ $3 \mid \underline{9}$ 3	$2^2 \times 3^3$	3	M2 for full systematic method of at least 4 divisions by prime numbers oe factor trees; condone 1 calculation error (M1 for 108 written as a correct product(s) with only one non-prime or equivalent division or a full process with 2 calculation errors.) A1 for $2^2 \times 3^3$ (accept $2 \times 2 \times 3 \times 3 \times 3$)
	(b)		12	1	B1 for 12
18		Perpendicular from P to intersecting arcs (within tramlines); perpendicular at least 2cm long		2	M1 relevant pair of arcs crossing within tramlines A1 SC M1 AO for a full construction of a perpendicular to AB not passing through P.
19		15×10	150 cm^3.	3	M1 for 15×10 A1 for 150 B1 for cm^3.

PAPER 5503					
No		**Working**	**Answer**	**Mark**	**Notes**
20			$y = 2x + 6$	2	B2 for $y = 2x + 6$ (B1 for $y = 2x + k$, $k \neq 1$ or for $y = mx + 6$, m $\neq 0$ or for $2x + 6$)
21		$6\frac{2}{5} = \frac{32}{5}$ Area of triangle = $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ (=2) Length of a side of sq. = $\sqrt{18 \times "2"}$ (=6) Perimeter of square = 4 × 6	24	5	B1 for $6\frac{2}{5} = \frac{32}{5}$ oe or $3\frac{1}{5} = \frac{16}{5}$ or $\frac{30}{8} + \frac{2}{8}$ oe (or implied by area of triangle=2) M1 for $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ oe M1 for area of square = 18 × product of two lengths A1 for $\sqrt{18 \times "2"}$ A1 for 24
22	(a)		$x(x-3)$	2	B2 for $x(x-3)$ (B1 for x (…….))
	(b)		k^3.	1	B1 for k^3.
	(c)(i)	$4x + 20 + 3x - 21$	$7x - 1$	4	M1 for **three** of 4 terms $4x + 20 + 3x - 21$ (or better) A1 for $7x - 1$
	(ii)	$x^2 + 3yx + 2xy + 6y^2$	$x^2 + 5xy + 6y^2$		M1 for **three** of 4 terms $x^2 + 3yx + 2xy + 6y^2$ A1 for $x^2 + 5xy + 6y^2$
	(d)		$(p+q)(p+q+5)$	1	B1 for $(p+q)(p+q+5)$

PAPER 5503					
	No	Working	Answer	Mark	Notes
23	(a)		32	1	B1 for 32 (accept 31.5 to 33.5 inclusive)
	(b)			3	B1 for ends of whiskers at 9 and 57, (with a box) B1 for ends of box at 16 and 45 (± 0.5) B1 for median marked at 32 or a complete box and whisker diagram drawn with a median inside the box
	(c)	Median(B)>Median(G); on average boys take longer IQR(B)>IQR(G); times for boys have a greater spread		2	B1 eg for comparison of medians B1 eg for comparison of (interquartile) ranges
24	(a)	$6y = 15 - 5x$	$y = \frac{15-5x}{6}$	2	M1 for either $6y = 15 - 5x$ or -6 $y = 5x - 15$ or for $\frac{5x}{6} + y = \frac{15}{6}$ or a correct ft on sign error to $y=$ A1 for $y = \frac{15-5x}{6}$ oe
	(b)	$6k + 5(-21) = 15$	20	2	M1 for substitution of $x = -21$ (or $x = 21$) into given equation or candidate's answer to (a) A1 for $k = 20$
	(c)(i)		Region R indicated	3	B2 correct region shaded (accept unshaded if R clear) (B1 shaded or R region satisfies 3 of the 4 given inequalities with same boundaries)
	(ii)		(1,1)		B1 for (1,1)

PAPER 5503					
	No	Working	Answer	Mark	Notes
25	(a)		60	1	B1 for 60 B1 for *BDC* = 25
	(b)	eg Angle *BDC* = 25 *ADB* = 60 − 25	35	2	[Award the mark for equivalent approaches] B1 ft for (a) − 25
	(c)	Ben is correct; angle *DAB* = 65 + 25 = 90 and since angle in a semi-circle is 90°, *BD* must be a diameter		1	B1 for full valid justification

PAPER 5504					
	No	Working	Answer	Mark	Notes
1	(a)	Accurate drawing of triangle		2	B1 cao for 38º (±2º) B1 cao (indep) for BC drawn 7.3 cm (±2mm) and completing the triangle.
	(b)	Measure angle A	63º	1	B1 for 63º (±2º) or ft their diagram (±2º)
2	(a)	0 × 1 = 0 1 × 1 = 1 2 × 4 = 8 3 × 8 = 24 4 × 8 = 32 5 × 7 = 35	100	3	M1 col 1 × col 2 (at least 3 shown) could be implied by answers. M1 (dep) sum of totals A1 cao SC: M1 M1 A0 for 101
	(b)	Either $35.50 ÷ 1.42 = £25; £26.99 - £25 = £1.99 Cheaper in the USA Or £26.99 × 1.42= $38.33; $38.33 - $35.50 = $2.83 Cheaper in the USA		3	M1 35.50 ÷ 1.42 A1 25 seen B1 ft (dep on M1) £1.99 with conclusion; units needed OR: M1 26.99 × 1.42 A1 38.33 or 38.32…seen B1 ft (dep on M1) $2.83 or $2.82 with conclusion; units needed
3	(a)		-1, (1), (3), 5, 7, 9	2	B2 cao (B1 for 2 values)
	(b)		Graph	2	B1 ft for plotting points ±1/2 square B1 cao for line between $x = -2$ and $x = 3$
	(c)(i)		0.4	2	B1 0.4 or ft single straight line with positive gradient
	(ii)		1.2		B1 1.2 or ft single straight line with positive gradient

PAPER 5504					
	No	Working	Answer	Mark	Notes
4	(a)			2	B1 Rectangle drawn B1 Triangle attached Ignore additional internal lines; 3D sketch gets B0
	(b)			2	B1 single Rectangle drawn B1 Attached smaller rectangle drawn Accept rotated plan, Ignore internal lines down; Nets get B0
5	(a)	£163.25 - £35.50 = £127.75 £127.75 ÷ £18.25 = 7	8	3	M1 163.25 – £35.50 (or sight of £127.75) M1 (dep) "127.75" ÷ 18.25 A1 cao SC: M2 for 7 days
	(b)	1.175 × £64 or $\frac{17.5}{100}$ × 64 or £11.20 seen	75.20	2	M1 1.175 × 64.00 0.175 × 64 oe or 11.20 or 75.2 seen OR 6.40, 3.20, and 1.60 seen A1 cao
	(c)	54 × £2.38 =	128.52	2	M1 54 × £2.38 A1 cao
6		Tessellation	Drawing	2	B2 seven additional hexagons, with at least 2 points at which 3 hexagons meet (B1 one point at which 3 hexagons meet).
7			200 150 225 150 10	3	B3 cao (B2 for three correct, or B1 for one correct)
8	(a)(i)	Bearing measured	70	2	B1 70 ± 4
	(ii)	Bearing measured	317		B1 317 ± 4
	(b)	Locus (circle) drawn & shaded		2	B1 circle ±2 mm centre Manchester B1 shading with accurate or approximate circle within tolerance

PAPER 5504					
	No	Working	Answer	Mark	Notes
9	(a)		Plots	1	B1 cao
	(b)		Description	1	B1 dynamic relationship or "positive" (correlation)
	(c)		Line of best fit	1	Line within overlay region, and to the extent of.
	(d)(i)	280 g →	Reading	2	B1 ft from single straight line of positive gradient (±1/2 square) B1 ft from single straight line of positive gradient (±1/2 square)
	(ii)	120 pages →	Reading		
10	(a)	$\frac{1}{6}, \frac{3}{8}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}$		2	B2 for all correct (B1 for 1 error or all correct but wrong order, or use of a common denominator decimals).
	(b)	3/5, 65%, 2/3, 0.72, 3/4		2	B2 for all correct (B1 for 1 error or all correct but wrong order or conversions to decimals oe)
11	(a)		182, 178, 180, 184	2	M1 mean of any three consecutive months, eg (147 + 161 + 238) ÷ 3 oe A1 cao
	(b)	Sale price = 80% Fun Friday price = 70% of 80% = 56% (oe)		2	B1 B1 for a fully correct explanation involving a worked example (oe) B1 a partially complete explanation

PAPER 5504					
	No	Working	Answer	Mark	Notes
12	(a)		Reason	1	B1 makes a mention of bias, either directly, or making reference to an insufficient or biased range of responses.
	(b)(i)		Reason	1	B1 make some mention of any one of (a) an insufficient range of responses (b) no mention of money (c) no time frame in the question (d) misunderstanding of "a lot" or "not much"
	(ii)		Question	2	B2 B1 for each of the following upto B2 (a) an improved question eg time frame made clear (b) response boxes (imperfect) (c) response boxes no errors OR For suggesting a generally improved question (a) a question clearly in the context of changes to the canteen (b) at least 3 boxes showing a full range of responses
13	(a)	$20y - 18y = 16 - 9$ oe $2y = 7$	$3\frac{1}{2}$ oe	3	M1 $20y - 18y = 16 - 9$ oe M1 $2y = 7$ A1 cao
	(b)	$40 - x = 3(4 + x)$ $40 - x = 12 + 3x$ $40 - 12 = x + 3x$ $4x = 28$	7	3	M1 multiplying through by 3: $3 \times \frac{40 - x}{3} = 3 \times 4 + 3 \times x$ A1 $40 - 12 = x + 3x$ A1 cao
14	(a)		$12(x\text{-}4)$	1	B1 for $12(x - 4)$ or $12x - 48$ or $12 \times x - 48$ oe $x = 12(x - 4)$ gets B0
	(b)		$18x\text{-}48$	2	M1 $6x +$ "$12(x - 4)$" or $6x +$ "$12x - 48$" A1 ft a linear expression $18x - 48$, $2(9x - 24)$, $3(6x - 16)$, $6(3x - 8)$


PAPER 5504				
No	**Working**	**Answer**	**Mark**	**Notes**
15	x $x^3 - 2x$ 4.1 60.7(21) 4.2 65.6(88) 4.3 70.9(07) Accept answer to 1 dp 4.22 66.7(114) or more 4.23 67.2(269) 4.24 67.7(45) 4.25 68.2(656)	4.2	4	B2 for trial $4.1 \leq x \leq 4.3$ evaluated (B1 for trial $4 < x < 5$ evaluated) B1 for different trial $4.225 \leq x \leq 4.25$ evaluated B1 (dep on at least one previous B1) for 4.2 cao
16 (a)		1.0×10^{-9}	1	B1 accept 1×10^{-9} or just 10^{-9}
(b)	$1 \div 5 \times 10^{-9}$	2×10^{8}	2	M1 for $1 \div$ ("5×10^{-9}") or digit 2 with zeros only seen Condone omission of bracket for M1. A1 cao
17 (a)	$\frac{6.27 \times 4.52}{4.81 + 9.63} = \frac{28.3404}{14.44} = 1.962631579$	1.9626…	2	B2 for 1.9626… (B1 for 28.34 or 14.44)
(b)		1.96	1	B1 ft from (a) as rounded to 1dp or 2dp. Do not accept 2, 2.00, but accept 2.0
18		$d = 4n + 6$	2	B2 $d = 4n + 6$ oe (B1 $d = 4n + k$, k an integer or 0, $4n + 6$, $n = 4n + 6$)
19	Vol = $\pi \times 3.8^2 \times 2.5 = \pi \times 14.44 \times 2.5$ = 45.36… × 2.5 = 113.411 Mass = "113" × 1.5 = 170.1165	170	4	M1 for $\pi \times r^2 \times 2.5$ where r = is 3.8 or 7.6 A1 if $r = 3.8$ M1 for "113" × 1.5 A1 for 169.5 – 170.3 cao


PAPER 5504					
	No	Working	Answer	Mark	Notes
20	(a)	$10^2 + 6^2$ or 136 $\sqrt{(100+36)}$ or $\sqrt{136} = 11.66\ldots$	11.7	3	M1 $10^2 + 6^2$ or 136 seen M1 (dep) $\sqrt{100+36}$ or $\sqrt{136}$ A1 11.66 – 11.7
	(b)	$\cos x = \frac{8}{10}$ or 0.8 $x = \cos^{-1} 0.8 = 36.869^\circ$	36.9	3	M1 for $\cos = \frac{8}{10}$, cos = 0.8 (oe) M1 (dep) for $\cos^{-1}$ (oe) A1 for 36.86 – 36.9
21		$18x - 6y = 99$ $\quad 12x - 4y = 66$ $8x + 6y = 18$ $\quad 12x + 9y = 27$ $26x = 117$ $\quad -13y = 39$	$x = 4\frac{1}{2}$ $y = -3$	4	M1 correct process to eliminate either x or y (condone one error) A1 cao for non – eliminated one. M1 (dep on 1st M1) for correct substitution of their found value. A1 cao (need both)
22		$\frac{133}{72} = 1.8472, \frac{160}{82} = 1.9512$ OR $\frac{72}{133} = 0.54135, \frac{82}{160} = 0.5125$ OR $\frac{160}{133} = 1.203...., \frac{82}{72} = 1.1388....$ OR $\frac{133}{160} = 0.83125...., \frac{72}{82} = 0.878$	1.84..≠ 1.95.. 1.20..≠ 1.13..	3	M1 for $\frac{133}{72}$ (= 1.8472…) oe Accept 1.8, 1.85 M1 for $\frac{160}{82}$ (=1.9512…) oe consistent pairing Accept 2.0, 1.9 **OR** M1 for $\frac{160}{133}$ (= 1.203…) oe M1 for $\frac{82}{72}$ (=1.1388) oe A1 for enough decimal places to show that the ratios are not equal; since the scale factors are different the shapes cannot be similar. NB Do Not need conclusion

PAPER 5504					
	No	Working	Answer	Mark	Notes
23	(a)		0.4, 0.6 0.4, 0.6, 0.4	2	B1 for LHS: (0.6), 0.4 B1 for RHS: 0.6, 0.4, 0.6, 0.4
	(b)	(30 × 42) – (25 × 42.8) = 1260 – 1070 = 190 190 ÷ 5 =	38	3	M1 for (30 × 42) – (25 × 42.8) or 1260 – 1070 or 190 seen M1 (dep) for "190" ÷ 5 A1 cao
24	(a)	£12000 × 0.25 = £3000; £12000 – £3000 = £9000 £9000 × 0.25 = £2250; £9000 – £2250 = £6750 £6750 × 0.25 = £1687.50; £6750 – £1687.50 =	£5062.50	3	M1 for 12000 × 0.75 (=9000) oe or £3000 or £23437.50 seen M1 (dep) for at least two further depreciation calculations (complete steps) A1 cao **OR** M2 for $12000 \times (0.75)^3$ or 5062.50 seen (M1 for $12000 \times (0.75)^n$, $n = 2$ or 4)
	(b)	0.8 × 0.8 × 0.8 × 0.8 (oe)	0.4096	2	M1 0.8^4 (oe) A1 cao

PAPER 5505					
	No	Working	Answer	Mark	Notes
1	(a)(i) (ii) (b)	 LCM = 3 × 13 × 17 = 3 × 221	2.21 0.013 663	2 2	B1 for 2.21 B1 for 0.013 M1 for 3 × 13 × 17 oe A1 for 663
2			Tick boxes 1,3 & 6	3	B1 for $\frac{\pi abc}{2d}$; B1 for $2a^2$; B1 for $2(c^2 + d^2)$ (−B1 for each additional expression ticked (>3) to a minimum of 0)
3	(a) (b)	200 × 0.2 0.2 + 0.4	40 0.6	2 2	M1 for 200 × 0.2 or $\frac{40}{200}$ seen A1 for 40 M1 for 0.2 + 0.4 A1 for 0.6
4	(a) (b)	eg 2 | 108 2 | 54 3 | 27 3 | 9 3 12	$2^2 \times 3^3$ 12	3 1	M2 for full systematic method of at least 4 divisions by prime numbers oe factor trees; condone 1 calculation error (M1 for 108 written as a correct product with only one non-prime or equivalent division or a full process with 2 calculation errors.) A1 for $2^2 \times 3^3$ (accept 2 × 2 × 3 × 3 × 3) B1 for 12
5		Perpendicular from P to intersecting arcs (within tramlines); perpendicular at least 2 cm long		2	M1 relevant pair of arcs crossing within tramlines A1 SC M1A0 for full construction of a line perpendicular to AB not through P
6		15 × 10	150 cm^3.	3	M1 for 15 × 10 A1 for 150 B1 for cm^3

PAPER 5505					
	No	Working	Answer	Mark	Notes
7	(a)		k^3.	1	B1 for k^3.
	(b)(i)	$4x + 20 + 3x - 21$	$7x - 1$	4	M1 for **three** of 4 terms $4x + 20 + 3x - 21$ (or better) A1 for $7x - 1$
	(ii)	$x^2 + 3yx + 2xy + 6y^2$	$x^2 + 5xy + 6y^2$		M1 for **three** of 4 terms $x^2 + 3yx + 2xy + 6y^2$ A1 for $x^2 + 5xy + 6y^2$
	(c)		$(p+q)(p+q+5)$	1	B1 for $(p+q)(p+q+5)$
	(d)		m^8	1	B1 for m^8.
	(e)		$6r^3t^6$	2	B2 for $6r^3t^6$ (B1 for $.....r^3t^6$ or for $6....t^6$)
8	(i)	Least length = 100.5	100.5	2	B1 for 100.5
	(ii)	Greatest length = 101.5	101.5		B1 for 101.5; ACCEPT 101.499 or better
9		$6\frac{2}{5} = \frac{32}{5}$ Area of triangle = $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ (= 2) Length of a side of sq. = $\sqrt{18 \times "2"}$ (=6) Perimeter of square = 4 × 6	24	5	B1 for $6\frac{2}{5} = \frac{32}{5}$ oe or $3\frac{1}{5} = \frac{16}{5}$ oe or $\frac{30}{8} + \frac{2}{8}$ oe (or implied by area of triangle=2) M1 for $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ oe M1 for (area of square) = 18 × product of two lengths A1 = $\sqrt{18 \times "2"}$ A1 for 24

PAPER 5505					
	No	Working	Answer	Mark	Notes
10	(a)	$6y = 15 - 5x$	$y = \frac{15-5x}{6}$	2	M1 for either $6y = 15 - 5x$ or for $\frac{5x}{6} + y = \frac{15}{6}$ or $-6y = 5x - 15$ or a correct ft on sign error to $y =$ A1 for $y = \frac{15-5x}{6}$ oe
	(b)	$6k + 5(-21) = 15$	20	2	M1 for subst. of $x = -21$ (or $x = 21$) into given eq^n or answer to (a) A1 for $k = 20$
	(c)(i)		Region R indicated	3	B2 correct region shaded (accept unshaded if R clear) (B1 shaded (R) region satisfies 3 of the 4 given inequalities with same boundaries)
	(ii)		(1,1)		B1 for (1,1)
11	(a)		$y = 2x + 6$	2	B2 for $y = 2x + 6$ (B1 for $y = 2x + k$, $k \neq 1$ or for $y = mx + 6$, $m \neq 0$ or for $2x+6$)
	(b)	Grad of $AB = 2$; Grad of $BC = \frac{-1}{"2"}$	$y = -\frac{1}{2}x + 6$	2	M1 for Grad of $BC = \frac{-1}{"2"}, -\frac{1}{2}$ or grad of $BC = -\frac{1}{\text{grad of AB}}$ A1 ft for $y = -\frac{1}{2}x + 6$ oe ft on wrong coeff. of x in (a)
	(c)	Eg A rectangle is always a cyclic quadrilateral because the opposite angles of a rectangle always add up to 180° (90+90)		1	B1 for valid explanation. (eg lines from the pt of int. of diagonals of rect to all 4 vertices are equal(radii))
12		Vertices of new triangle at $(-1, -4)$ $(-4, 2)$ and $(2, 2)$	Correct triangle drawn	3	B3 cao (B2 for 3 correct vertices no triangle or triangle with 2 correct vertices (B1 for triangle with two of six co-ordinates correct from using P as centre or any isosceles triangle with base 6 and height 6)

PAPER 5505					
	No	Working	Answer	Mark	Notes
13	(a)		32	1	B1 for 32 (accept 31.5 to 33.5 inclusive)
	(b)			3	B1 for ends of whiskers at 9 and 57 (with a box) B1 for ends of box at 16 and 45/46 ($\pm$ 0.5) B1for median marked at "32"or complete box and whisker diagram drawn with a median inside the box
	(c)	Median(B)>Median(G); on average boys take longer IQR(B)>IQR(G); times for boys have a greater spread		2	B1 eg for comparison of medians (ft on diagrams) B1 eg for comparison of (interquartile) ranges (ft on diagram)
14		Reading top to bottom frequencies are 20; 18; 45; 52		2	B2 all correct (B1 for one frequency correct)
15	(a)	$1600 = p\,q^0$.; $400 = pq^2$ $q^2 = 0.25$	$p = 1600$ $q = 0.5$	3	M1 for either $400 = pq^2$ or $1600 = pq^0$ B1 for $p = 1600$ A1 for $q = 0.5$
	(b)	$V = pq^{-2}$	6400	2	M1 for recognition that t = –2 A1 for 6 400

PAPER 5505					
	No	Working	Answer	Mark	Notes
16	(a)	(I) $AB = AC$ (triangle ABC is isosceles) (II) PB = PC tangents (from a point to a circle are) equal (III) AP = AP (common side) so the 2 triangles are congruent ,SSS.		3	B3 for I, II, III with congruency reason (B2 for any two of I, II or III) (B1 for any one of the I, II or III)
	(b)	$BPC = 20°$ PBC (or PCB) = 90-1/2 "20" (= 80°) $BAC = PBC$ = "80"	50°	4	B4 for 50° (B3 for $BAC = 80°$) (B2 for $PBC = 80°$ or $PCB = 80°$) (B1 for $APC = 10°$ or $BPC = 20°$ or a middle angle = 90°) SC if clear numerical slip seen eg "PBC = 180 – 90 – 10 = 70" then goes on to get correct ft angle $ABC = 55$ deduct 1 from total so this cand would get B4–1 = B3
17	(a)	$\overrightarrow{OP} = \overrightarrow{OA} + \overrightarrow{AP}$ $= \overrightarrow{OA} + \frac{2}{3}(6\mathbf{c} - 6\mathbf{a})$ $= 6\mathbf{a} + 4\mathbf{c} - 4\mathbf{a}$	$2\mathbf{a} + 4\mathbf{c}$	3	M1 for $\overrightarrow{OP} = \overrightarrow{OA} + \overrightarrow{AP}$ or any correct vector journey involving $\overrightarrow{OP}$ M1 for $\overrightarrow{AP} = \frac{2}{3}(6\mathbf{c} - 6\mathbf{a})$ oe or $\overrightarrow{CP} = \frac{1}{3}(-6\mathbf{c} + 6\mathbf{a})$ oe or reverse vectors A1 for $2\mathbf{a} + 4\mathbf{c}$ oe (accept unsimplified)
	(b)	Eg $\overrightarrow{OM} = \overrightarrow{OC} + \overrightarrow{CM} = 6\mathbf{c} + 3\mathbf{a}$ $\overrightarrow{OM} = 1.5\overrightarrow{OP}$	$\overrightarrow{OM} = 1.5\overrightarrow{OP}$ so OPM is a straight line	2	B1 for $\overrightarrow{OM} = 6\mathbf{c} + \frac{1}{2}(6\mathbf{a})$ or $\overrightarrow{PM} = 2\mathbf{c} + \mathbf{a}$ unsimplified or reverse vectors B1 for a fully correct proof.

PAPER 5505					
	No	**Working**	**Answer**	**Mark**	**Notes**
18	(a)		4	1	B1 for 4 condone $\pm$ 4
	(b)		2	1	B1 for 2 condone ± 2
	(c)	$\sqrt{160} = 4\sqrt{10}$; $\left[\frac{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)-\sqrt{2}\times\sqrt{5}}{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)}\right]\times 100$ $\left[\frac{6\sqrt{10}-\sqrt{10}}{6\sqrt{10}}\right]\times 100$	$\frac{500}{6}$	4	B1 for **either** $\sqrt{160} = 4\sqrt{10}$ or $\sqrt{8} = 2\sqrt{2}$ **or** $\sqrt{20} = 2\sqrt{5}$ M1 for $\left[\frac{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)-\sqrt{2}\times\sqrt{5}}{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)}\right]$ oe $(\times 100)$ B1 for either $6\sqrt{10} - \sqrt{10}$ or $6\sqrt{10}$ A1 for $\frac{500}{6}$ (accept 83.3 if no obvious earlier error)

PAPER 5505					
	No	Working	Answer	Mark	Notes
19	(a)(i)		$(2x-7)(x-14)$	3	M1 x^2 term and constant term (± 98) obtained or $2x(x-14) - 7(x-14)$ or $x(2x-7) - 14(2x-7)$ A1 for $(2x-7)(x-14)$
	(ii)		$x = \frac{7}{2}$; $x = 14$		B1ft ft (i) provided of form $(2x\pm a)(x\pm b)$
	(b)(i)		$\frac{7}{n+7}$	3	B1 for $\frac{7}{n+7}$ oe
	(ii)	$\frac{7}{n+7} = \frac{2}{5} \Rightarrow 2(n+7) = 5 \times 7$ $2n = 21$	n=10.5 is not possible since n has to be an integer		M1 for $2(n+7) = 5\times 7$ or $n+7=5\times 3.5$ (can be implied) ft (b)(i) fractional in terms of n and < 1 A1 ft for n=10.5 not possible (since n not integer) oe
	(c)	$2\times\left(\frac{n}{n+7}\right)\times\left(\frac{7}{n+7}\right) = \frac{4}{9}$ $14n\times 9 = 4(n+7)^2$ $14n\times 9 = 4(n^2+14n+49)$ $4n^2+56n+196-126n = 0$	$2n^2 - 35n + 98 = 0$	5	M1 for $\left(\frac{n}{n+7}\right)\times\left(\frac{7}{n+7}\right)$ seen M1 for $2\times\left(\frac{n}{n+7}\right)\times\left(\frac{7}{n+7}\right)$ oe $(=\frac{4}{9})$ M1(dep on 1st M) elimination of fractions within an equation B1 3 terms correct in expansion of $(n+7)^2 = n^2 + 7n + 7n + 49$ A1 full valid completion to printed answer

PAPER 5505					
	No	Working	Answer	Mark	Notes
	(d)	$\frac{7}{n+7}\times\frac{7}{n+7}=\frac{7}{21}\times\frac{7}{21}=$	$\frac{1}{9}$	2	M1 for $\frac{7}{n+7}\times\frac{7}{n+7}$ or better or ft [answer (b)(i)]^2 or $1-\frac{4}{9}-(\frac{n}{n+7})^2$ A1 for $\frac{1}{9}$ oe cao
20	(a)(i)		$y=1+\sin x$	2	B1 for $y=1+\sin x$
	(ii)		$y=2\sin x$		B1 for $y=2\sin x$ SC both (i) f(x) + 1, (ii) 2f(x) B1
	(b)	Stretch parallel to y-axis scale factor 3 Stretch parallel to x-axis scale factor $\frac{1}{2}$		3	M1 for 'stretch' A1 for Stretch parallel to y-axis scale factor 3 oe A1 for Stretch parallel to x-axis scale factor $\frac{1}{2}$ oe SC if M0 award BI for "sf 3 vertically" **and** "sf $\frac{1}{2}$ horizon."

PAPER 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
1		Question	2	B1 for each of the following upto B2 (a) an improved question eg time frame made clear (b) response boxes (imperfect) (c) response boxes no errors OR For suggesting a generally improved question (a) a question clearly in the context of changes to the canteen (b) at least 3 boxes showing a full range of responses
2	Locus (circle) drawn & shaded		2	B1 circle centre Manchester ±2mm B1 shading with accurate or approximate circle within tolerance
3 (a)		182, 178, 180, 184	2	M1 mean of any three consecutive months, eg (147 + 161 + 238) ÷ 3 oe A1 cao
(b)	Sale price = 80% Fun Friday price = 70% of 80% = 56% (oe)		2	B1 B1 for a fully correct explanation involving a worked example (oe) B1 a partially complete explanation
4	x $x^3 - 2x$ 4.1 60.7(21) 4.2 65.6(88) 4.3 70.9(07) 4.22 66.7(114) 4.23 67.2(269) 4.24 67.7(45) 4.25 68.2(656)	4.2	4	B2 for trial $4.1 \leq x \leq 4.3$ evaluated (B1 for trial $4 < x < 5$ evaluated) B1 for different trial $4.225 \leq x \leq 4.25$ evaluated B1 (dep on at least one previous B1) for 4.2 cao

PAPER 5506					
	No	Working	Answer	Mark	Notes
5	(a)		1.0×10^{-9}	1	B1 accept 1×10^{-9} or just 10^{-9}
	(b)	$1 \div (5 \times 10^{-9})$	2×10^{8}	2	M1 for $1 \div$ ("5×10^{-9}") or digit 2 with zeros only seen Condone omission of bracket for M1. A1 cao
6	(a)	$\frac{6.27 \times 4.52}{4.81 + 9.63} = \frac{28.3404}{14.44} = 1.962631579$	1.9626…	2	B2 for 1.9626… (B1 for 28.34… or 14.44)
	(b)		1.96	1	B1 ft from (a) as rounded to 1dp or 2dp. Do not accept 2, 2.00, but accept 2.0
7			$d = 4n + 6$	2	B2 $d = 4n + 6$ oe (B1 $d = 4n + k$, k an integer $\neq 6$, $4n + 6$, $n = 4n + 6$)
8		Vol = $\pi \times 3.8^2 \times 2.5 = \pi \times 14.44 \times 2.5$ $= 45.36\ldots \times 2.5 = 113.411$ Mass = "113" × 1.5 = 170.1165	170	4	M1 for $\pi \times r^2 \times 2.5$ where r = is 3.8 or 7.6 A1 if $r = 3.8$ M1 for "113" × 1.5 A1 for 169.5 – 170.3 cao

PAPER 5506					
	No	Working	Answer	Mark	Notes
9	(a)	$10^2 + 6^2$ or 136 $\sqrt{(100 + 36)}$ or $\sqrt{136} = 11.66\ldots$	11.7	3	M1 for $10^2 + 6^2$ or 136 seen M1 (dep) $\sqrt{100+36}$ or $\sqrt{136}$ A1 11.66 – 11.7
	(b)	Cos x = 8/10 or 0.8 $x = \cos^{-1} 0.8 = 36.869^o$	36.9	3	M1 for cos = $\frac{8}{10}$, cos = 0.8(oe) M1 (dep) for $\cos^{-1}$ (oe) A1 for 36.86 – 36.9 M1 Use of sine rule and x found M1 for $x = 90 - \sin^{-1}$("0.8") A1 for 36.86 – 36.9
10		$18x - 6y = 99$ $\quad 12x - 4y = 66$ $8x + 6y = 18$ $\quad 12x + 9y = 27$ $26x = 117$ $\quad -13y = 39$	$x = 4\ ½$ $y = –3$	4	M1 correct process to eliminate either x or y (condone one error) A1 cao for non – eliminated one. M1 (dep on 1st M1) for correct substitution of their found value. A1 cao (need both)

PAPER 5506					
	No	Working	Answer	Mark	Notes
11		$\frac{133}{72} = 1.8472\,,\ \frac{160}{82} = 1.9512$ OR $\frac{72}{133} = 0.54135, \frac{82}{160} = 0.5125$ OR $\frac{160}{133} = 1.203...., \frac{82}{72} = 1.1388....$ OR $\frac{133}{160} = 0.83125...., \frac{72}{82} = 0.878$	1.84..≠ 1.95.. 1.20..≠ 1.13..	3	M1 for $\frac{133}{72}$ (= 1.8472…) oe Accept 1.8, 1.85 M1 for $\frac{160}{82}$ (=1.9512…) oe consistent pairing Accept 2.0, 1.9 **OR** M1 for $\frac{160}{133}$ (= 1.203…) oe M1 for $\frac{82}{72}$ (=1.1388) oe A1 for enough decimal places to show that the ratios are not equal; since the scale factors are different the shapes cannot be similar. NB Do Not need conclusion
12	(a)	£12000 × 0.25 = £3000; £12000 – £3000 = £9000 £9000 × 0.25 = £2250; £9000 – £2250 = £6750 £6750 × 0.25 = £1687.50; £6750 – £1687.50 =	£5062.50	3	M1 for 12000 × 0.75 (=9000) oe or £3000 or £23437.50 seen M1 (dep) for at least two further depreciation calculations (complete steps) A1 cao **OR** M2 for 12000 × $(0.75)^3$ or 5062.50 seen (M1 for 12000 × $(0.75)^n$, $n = 2$ or 4)
	(b)	0.8 × 0.8 × 0.8 × 0.8 (oe)	0.4096	2	M1 0.8^4 (oe) A1 cao

PAPER 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
13	$r^2 = \frac{3\times10}{\pi h} = \frac{3\times10}{\pi 1.5} = \frac{3\times10}{4.712..} = 6.36, 6.37$ $r = \sqrt{6.366\ldots}$	2.52-2.54	3	M1 for correct rearrangement to give $r^2 = \frac{3V}{\pi h}$ or $30 \div 4.712..$ or $6.36 - 6.37$ *Allow 0.3, 0.33 for* $\frac{1}{3}$ M1 (dep) for $\sqrt{}$ A1 cao 2.52 – 2.54
14 (a)		0.4 0.6,0.4, 0.6,0.4	2	B1 for LHS: (0.6), 0.4 B1 for RHS: 0.6, 0.4, 0.6, 0.4
(b)	0.6×0.6	0.36	2	M1 $0.6 \times$ "0.6" [0 < "0.6" < 1] A1 cao
(c)	$(30 \times 42) - (25 \times 42.8) = (1260 - 1070 = 190$ $190 \div 5 =$	38	3	M1 for $(30 \times 42) - (25 \times 42.8)$ or $1260 - 1070$ or 190 seen M1(dep) for "190" ÷ 5 A1 cao 38

PAPER 5506					
	No	Working	Answer	Mark	Notes
15	(a)	$S = \frac{k}{f^2};\ 125 = \frac{k}{8^2};\ k = 8000$	$S = \frac{8000}{f^2}$	3	M1 for $S = \frac{k}{f^2}$ M1 for $125 = \frac{k}{8^2}$ A1 cao These marks can be awarded if the full formula appears in part (b), rather than in part (a). [SC: $S = \frac{125}{64} f^2$ M1 M0 A0, $S = 1.95(3125) f^2$ M1 M0 A0 $S = \frac{1000}{f}$ M1 M0 A0]
	(b)		500	1	B1 cao

PAPER 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
16	$0.5 \times 3.2 \times 8.4 \times \text{Sin } B = 10$ $\sin B = 0.74404\ldots$ 48.077 $AC^2 = 3.2^2 + 8.4^2 - 2 \times 3.2 \times 8.4 \times \cos B$ $AC^2 = 44.8815\ldots..$ $AC = 6.69\ (936\ldots)$ Perimeter = 18.3 Use the altitude AD, $\frac{h \times 8.4}{2} = 10 \Rightarrow h = (2.381)$ $BD = \sqrt{3.2^2 - h^2} = 2.139$ $DC = 6.261$ $AC = \sqrt{2.38^2 + '6.261'^2} = 6.69(936)$ Perimeter = 18.3	18.3	6	M1 for $0.5 \times 3.2 \times 8.4 \times \sin B\ (= 10)$ A1 for sin B = 0.74(404…) or B = 47.7 – 48.1 M1 for $3.2^2 + 8.4^2 - 2 \times 3.2 \times 8.4 \times \cos$ "48.077" M1(dep) for AC^2 = "44.8(815)"….. with correct order of evaluation A1 $AC = 6.69 - 6.7$ A1 18.29 – 18.3 **for** $\frac{h \times 8.4}{2} = 10 \Rightarrow h = (2.381)$ M1 for $BD^2 = 3.2^2 - "2.381"^2$ A1 $BD = 2.1 - 2.2$ M1 (dep) $AC^2 = "2.381"^2 + "6.261"^2$ A1 $AC = 6.69 - 6.7$ A1 18.29 – 18.3

PAPER 5506					
	No	Working	Answer	Mark	Notes
17	(a)	$(x + 8)^2 = x^2 + (x + 5)^2$ $x^2 + 16x + 64 = 2x^2 + 10x + 25$		4	B1 for angle $OTA = 90°$ (implied by use of Pythagoras with OA as hypotenuse) M1 $(x + 8)^2 = x^2 + (x + 5)^2$ oe M1 for correct squaring of x+8 or x+5 A1 for completion following correct working
	(b)	$(+6 + \sqrt{(6^2 - - 4\times39)})/2$ $= (6 \pm \sqrt{(36 + 156)})/2$ $= (6 \pm \sqrt{192})/2 = (6 + 13.856)/2$	9.93	3	M1 for substitution into quadratic formula, allow sign errors in b and c M1 for $x = (6 \pm \sqrt{192})/2$ [+ alone will do] A1 for 9.92 – 9.93
					M1 for $(x-3)^2-9-39$ M1 for $x = 3 \pm \sqrt{48}$ [+ alone will do it] A1 for 9.92 – 9.93 T&I= 9.93 gets M1,M0,A0
18		$EB = 60 \times \tan 30°$ $BD = \sqrt{(60^2 + 60^2)}$ $\tan BDE = 34.64 \div 84.85$	22.2°	4	M1 for $EB = 60\times\tan30$ M1 for $BD = \sqrt{(60^2 + 60^2)}$ M1 for $\tan BDE$ = "34.64" ÷ "84.85" A1 22.17 – 22.21
		$EB = 60 \times \tan 30°$ (= 34.64) $ED^2 = 60^2 + \left(\frac{60}{\cos 30}\right)^2$ $ED = \sqrt{8400} = (91.65)$ $\text{Angle} = \sin^{-1}\left(\frac{EB}{\sqrt{8400}}\right) = 22.2$			M1 for $EB = 60 \times \tan 30°$ oe M1 for fully correct method for ED M1 for $\sin BDE = \left(\frac{'34.84'}{'\sqrt{8400}'}\right)$ (oe) A1 22.17 – 22.21

PAPER 5506					
	No	Working	Answer	Mark	Notes
19	(a)	$x^2 + 6^2 = 25$ $x^2 = -11$		2	M1 substitutes y=6 to get x^2+36 = 25 A1 deduces x^2=25-36<0; impossible, (so line does not intersect curve)
	(b)	$y^2 = (2x - 2)^2$ $x^2 + (2x - 2)^2 = 25$ $x^2 + 4x^2 - 8x + 4 = 25$ $5x^2 - 8x + 4 - 25 = 0$ $5x^2 - 8x - 21 = 0$ $(5x + 7)(x - 3) = 0$ or $x = \frac{--8 \pm \sqrt{(-8)^2 - 4 \times 5 \times (-21)}}{10} = \frac{8 \pm \sqrt{484}}{10}$ So $x = 3$, or $x = -1.4$ Subs into either equation gives: $y = 4$, or $y = -4.8$	$x = 3$ $y = 4$ $x = -1.4$ $y = -4.8$	6	M1 $y^2=(2x-2)^2$ seen or implied A1 $x^2 + 4x^2 - 8x + 4 = 25$ (oe expanded form) M1 correct attempt to solve 3 term quadratic equation A1 $x = 3$, $x = -1.4$ M1 (dep on previous Ms) Sub one value of x into either equation gives: A1 $y = 4$, $y = -4.8$ NB: incorrectly matched pairs loses the final A1.

PAPER 5506					
	No	Working	Answer	Mark	Notes
20	(a)	400.005 ÷ 59.95 = 6.672310 6.672310 ÷ 1000 × 3600 = 24.02032 399.995 ÷ 60.05 = 6.661032 6.661032 ÷ 1000 × 3600 = 23.97972	24.020 23.980	5	B1 for 400.005 or 59.95 seen oe M1 for "400.005" ÷ "59.95" where "400.005" ∈ [400.005,400.5] and "59.95" ∈ [59.5,59.95] oe B1 for 399.995 or 60.05 seen oe M1 for "399.995" ÷ "60.05" where "399.995" ∈ [399.5,399.95] and "60.05" ∈ [60.05,60.5] oe A1 23.979-23.980 and 24.020-24.0204
	(b)	24.0 because to 1dp the answers are the same		1	B1cao for 24.0 with reason
	(c)	177 × 50/477 = 18.553 → 19 → 18 111 × 50/477 = 11.635 → 12 86 × 50/477 = 9.0147 → 9 82 × 50/477 = 8.595 → 9 21 × 50/477 = 2.201 → 2	18 12 9 9 2	3	M1 for $\frac{50}{"477"} \times 177$ or 111 or 86 or 82 or 21 A1 for all integers or better answers, at least 3 correct A1 cao
21	(a)	$40 - x = 3(4 + x)$ $40 - x = 12 + 3x$ $40 - 12 = x + 3x$ $4x = 28$	7	3	M1 multiplying through by 3: $3 \times \frac{40 - x}{3} = 3 \times 4 + 3 \times x$ A1 $40 - 12 = x + 3x$ A1 cao
	(b)	$\frac{2x(2x - 3)}{(2x - 3)(2x + 3)} = \frac{2x}{2x + 3}$	$\frac{2x}{2x + 3}$	3	B1 for $(2x-3)(2x+3)$ B1 for $2x(2x-3)$ or $(2x + 0)(2x + 3)$ B1 cao

PAPER 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
22	Distance from x axis is y. Distance from (0, 2) is $\sqrt{(x^2+(y-2)^2)}$ $y^2 = x^2 + (y-2)^2$ $y^2 = x^2 + y^2 - 4y + 4$ $0 = x^2 - 4y + 4$ $4y = x^2 + 4$ and finish		4	B1 for $(x-0)^2 + (y-2)^2$ or $\sqrt{((x-0)^2+(y-2)^2)}$ oe seen B1 for $y = \sqrt{(x-0)^2+(y-2)^2}$ or $y^2 = (x-0)^2 + (y-2)^2$ oe B1 $(y-2)^2 = y^2 - 4y + 4$ seen B1 for $4y = x^2 + 4$ and finish

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone: 01623 467467
Fax: 01623 450481
Email: publications@linneydirect.com

Order Code UG015616 July 2004

For more information on Edexcel and BTEC qualifications please contact
Customer Services on 0870 240 9800
or http://enquiries.edexcel.org.uk
or visit our website: www.edexcel.org.uk

London Qualifications Limited, trading as Edexcel. Registered in England and Wales No. 4496750
Registered Office: Stewart House, 32 Russell Square, London WC1B 5DN



GCSE

Edexcel GCSE

Mathematics A 1387

Summer 2005
Mark scheme

Edexcel GCSE
Mathematics A 1387

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information please call our Customer Services on 0870 240 9800, or visit our website at www.edexcel.org.uk.

Summer 2005

Publications Code UG016720

All the material in this publication is copyright
© Edexcel Ltd 2005

CONTENTS

Notes on Marking Principles 1

Paper 5521/01 3

Paper 5521/02 12

Paper 5523/03 18

Paper 5523/04 27

Paper 5525/05 37

Paper 5525/06 45

NOTES ON MARKING PRINCIPLES

1 Types of mark

- M marks: method marks
- A marks: accuracy marks
- B marks: unconditional accuracy marks (independent of M marks)

2 Abbreviations

cao – correct answer only
ft – follow through
isw – ignore subsequent working
SC: special case
oe – or equivalent (and appropriate)
dep – dependent
indep - independent

3 No working

If no working is shown then correct answers normally score full marks
If no working is shown then incorrect (even though nearly correct) answers score no marks.

4 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.
If there is no answer on the answer line then check the working for an obvious answer.

5 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

6 **Ignoring subsequent work**
It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.


7 **Probability**
Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

8 **Linear equations**
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

9 **Parts of questions**
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
1	(a)		17252	1	B1 cao
	(b)		5400	1	B1 cao
	(c)		thousands, 1000, 4000	1	B1
2	(a)		grams, g	3	B1 oe spelling
			centimetres, cm		B1 oe spelling
			millilitres, m*l*, cm^3		B1 oe spelling
	(b)		5	1	B1 cao
3	(a)		106, 102	1	B1 cao ignore extras
	(b)		eg take away 4	1	B1 could be indicated on the diagram
	(c)		46	1	B1 cao

Paper 5521/01


No	Working	Answer	Mark	Notes
4	286 × 43 = 858, 11440, 12298 43 × 286 = 258, 3440, 8600, 12298 286×40 = 1140 286×3 = 858 1140+858=12298 x | 40 | 3 | 200 | 8000 | 600 | 8600 80 | 3200 | 240 | 3440 6 | 240 | 18 | 258 | | | 12298  	12298	3	M2 for complete correct method (condone one computational error) (M1 for complete correct method with two computational errors) A1 cao OR B2 inside of grid completed (condone missing zeros and one error) (B1 2 or 3 errors) B1 cao
5 (a) (b)(i) (ii) (c) (d)(i) (ii) (iii)		18, 69 18 or 36 16 or 36 factor 18 11 or 88 69	1 1 1 1 3	B1 B1 B1 B1 B1 cao B1 B1 cao

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
6	(a)		April & May	1	B1 for both
	(b)		Daffodil	1	B1
	(c)		Feb	1	B1
	(d)		Crocus	1	B1
	(e)(i)		$\frac{1}{5}$	2	B1 for $\frac{1}{5}$ oe
	(ii)		× from 56 mm to 64 mm from 0		B1 A single mark on the line, between 56 mm and 64 mm measured from end 0
7	(a)	$\frac{40}{100}$	$\frac{2}{5}$	2	B2 for $\frac{2}{5}$ B1 for $\frac{40}{100}$ or $\frac{4}{10}$ or $\frac{20}{50}$ or $\frac{8}{20}$
	(b)		0.98	1	B1 cao
	(c)		7 500 000	1	B1 cao
	(d)		25	1	B1 cao
	(e)		60	1	B1 cao

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
8	(a)(i)		(0, 2)	2	B1 cao
	(ii)		(4, 1)		B1 cao
	(b)		$(2, 1\frac{1}{2})$ marked	1	B1 Allow 2mm tolerance from $(2, 1\frac{1}{2})$
9	(a)		2.40	1	B1 cao could be indicated on the diagram
	(b)	1.60 + 0.40	2.00	2	M1 for appropriate sum or product in £ or p or 200 seen eg 1.60 + 0.40, 160 + 40, 0.80 + 0.80 + 0.40, 80 + 80 + 40, 0.08×25, 0.80×2.5, 200 A1 cao
	(c)	$1\div0.8$ or $2.50\div2$	1.25	2	M1 for $1.00\div0.8$ or $2.50\div2$ or 125 or appropriate combination eg $1+\frac{1}{2}\times0.50$ A1 cao
10	(a)		hexagon	1	B1 Condone spelling error
	(b)		Sum of angles at a point is 360°	2	B1 for 360 seen B1 for "point", "complete turn" or "a circle" or similar unless accompanied by an incorrect angle SC If neither B1 scored, award B1 for a clear indication that the size of an angle, other than *x*, is 90° or a right angle (may be on diagram)
	(c)	$30\times4+8\times2$	136	2	M1 $30\times4+8\times2$ or attempt to sum 5 or 6 lengths A1 cao

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
11	(a)		13, 67, 76, 103, 130	5	B1 cao
	(b)		$-7, -3, -1, 0, 5$		B1 cao
	(c)		0.07, 0.072,0.7, 0.702, 0.72		B1 cao
	(d)		0.6, $\frac{2}{3}$, 70%, $\frac{3}{4}$		B2 (B1 for any 3 in correct order)
12	(a)		16 30	1	B1 Accept 4 30 pm Do not accept 4 30
	(b)(i)	$33.56 \div 4$ oe	8.39	3	M1 for $33.56 \div 4$ oe eg $3356 \div 4$, division by 2 twice A1 cao
	(ii)		9		B1 ft from "8.39" unless whole number of pounds
13	(a)		6 cm^2	3	B2 for 6 cao for numerical answer (B1 for $5.5 < \text{Area} \leq 7$) then B1 (indep) for cm^2 with or without numerical answer
	(b)	See diagram	Correct shape	2	B2 (B1 for any 2 sides correct or a correct enlargement scale factor $\neq 1$ or 2)

Paper 5521/01				
No	**Working**	**Answer**	**Mark**	**Notes**
14 (a)	$(4+3)\times10$	70	2	M1 for (4 + 3) × 10 A1 cao
(b)	$120\div10-3$	9	2	M1 for $\frac{120}{10}$ or 12 seen eg $12\times10=120$ A1 cao
(c)		$C=10(n+3)$	3	B3 for $C=10(n+3)$ oe such as $C=(n+3)\times10$ (B2 for correct RHS or $C=n+3\times10$, $C=10n+3$ oe B1 for C = some other linear expression in n or for $n+3\times10$, $10n+3$ etc) Note: $C = n$ scores no marks
15		11 13 16 8 21	2	B2 all correct (B1 for 2 correct)

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
16	(a)		$2p + 4q$	2	B2 for $2p + 4q$ (accept $2 \times p$ etc) (B1 for $2p$ or $4q$)
	(b)		$2y^2$	1	B1 accept $2 \times y^2$ oe inc $2 \times y \times y$
	(c)		$3c + 4d$	2	B2 for $3c + 4d$ (accept $3 \times c$ etc) (B1 for $3c$ or $4d$)
	(d)		$8pq$	1	B1 accept in any order but must not include $\times$ sign
17	(a)(i)		60	2	B1 cao
	(ii)		eg top triangle is equilateral		B1 for reason
	(b)		150	2	M1 for $\frac{180-"60"}{2} + 90$ A1 ft from (a)(i) if $x < 90$ SC B1 for "60" + 90 if $x < 90$
18			40	2	M1 for 60×2 or 120 or $60 \div 3$ or 20 or $\frac{120}{180}$ A1 cao
19			correct drawing	2	B2 Condone hidden detail shown with solid lines and missing lines on front face (B1 for a correct sketch with other incorrect sketch(es) or for prism with correct cross section >1 cube wide or for attempt to draw prism with correct cross section or prism with correct plan and side elevation)

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
20		$\frac{600}{3\times10}$ or $\frac{640}{3.2\times10}$	$20\text{-}21\frac{1}{3}$	2	M1 for rounding at least two of the numbers to 1 sf or for sight of 640, 3.2 or 640, 32 or 600, 32 or 30 seen A1 for $20\text{-}21\frac{1}{3}$ Note: 20.3125 scores M0 A0
21	(a)		Points plotted	1	B1 $\pm$ 1 full (2 mm) square
	(b)		positive	1	B1 cao
	(c)		Line of best fit	1	B1 Must pass through (42.5, 1.45), (42.5, 1.55) AND (67.5, 1.75), (67.5, 1.85)
	(d)		~ 1.65	1	B1 ft from single line segment with positive gradient $\pm$ 1 full (2 mm) square
22	(a)	eg $50\times\frac{2000}{500}$	200	2	M1 for $\frac{2000}{500}$ or 4 seen A1 cao
	(b)	eg $400\times\frac{750}{500}$	600	2	M1 for $\frac{750}{500}$ or 1.5 seen or 400 + 200 A1 cao

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
23	(a)	$4\times3-2\times1$ $12-2$	10	3	M1 for 3×4 (=12) or 1×2 or attempt to divide diagram up into rectangles M1 "12"– "2" or sum of areas of rectangles A1 cao
	(b)(i)	$\frac{10}{100}\times680$ or $680\div10$ 680 + 68	748	5	M1 $\frac{10}{100}\times680$ or 680÷10 or 68 seen M1 (dep) 680 + "68" or M2 for 680×1.10 A1 cao
	(ii)	"748"÷50 or 14.96	15		M1 For "748"÷50 or 14.96 Accept "748" rounded up or down to next 50 followed by $\div50$ A1 ft from (b)(i) rounded up SC B1 for 680 (seen) leading to 14

Paper 5521/02					
No		**Working**	**Answer**	**Mark**	**Notes**
1	(a)	Plain llll lll 8 Chicken lll 3 Bovril llll 5 S & Vin llll 4		3	M1 for attempt to tally A1 for 1 frequency correct or all tallies correct A1 for all frequencies correct (accept for /20)
	(b)		4	1	B1 ft
	(c)		Plain or 8	1	B1 ft
2	(a)(i)		11	2	B1 cao
	(ii)		16		B1 cao
	(b)	See diagram	Correct lines	2	B2 cao for both lines correct (B1 for one line correct)
	(c)		12	2	B2 cao (B1 for 11 or 13)
3	(a)		580	1	B1 for 580 (± 2) could be written on line
	(b)		7.2	1	B1 for 7.2 ± 0.02 could be written on line
	(c)		Arrow at 48	1	B1 allow ± half graduation
	(d)		Arrow at 6.7	1	B1 allow ± half graduation
4	(i)		Cylinder	2	B1 ignore spelling
	(ii)		Cuboid		B1 ignore spelling
5	(a)	£10 – (£2.15 + £2.30)	5.55	4	M1 £2.15 + £2.30 A1 for 4.45 M1 £10 – "4.45" A1 cao
	(b)	£60 ÷ £2.80 = 21.42857	21	2	M1 for £60 ÷ 2.80 or sight of digits 214… A1 for 21
	(c)	$120 \times 25 \div 100$	30	2	M1 ¼ of £120 (oe) A1 cao SC B2 for £90

Paper 5521/02					
No		**Working**	**Answer**	**Mark**	**Notes**
6	(a)(i)		143^0	2	B1 for 143 ($\pm 2^0$)
	(ii)		Obtuse		B1 for obtuse (ignore spelling)
	(b)	See diagram	Accurate drawing	1	B1 for accurate drawing ± 2mm
7	(a)(i)		5	2	B1 cao
	(ii)		23		B1 cao
	(b)	$\times 2 - 1$		1	B1 for explaining a suitable method
	(c)	See their diagram		1	B1 for a correct diagram
	(d)		14, 17	2	B2 cao for both (B1 for one only ft from their 14)
8	(a)		90	1	B1 accept –90
	(b)		540	1	B1 accept –540
	(c)		Jupiter	1	B1 accept –150
	(d)		- 230	1	B1 cao

Paper 5521/02					
No		Working	Answer	Mark	Notes
9	(a)	2658 – 2430 = 228 "228" × 32	72.96	4	M1 2658 – 2430 A1 228 M1 "228" × 32 or "228" × 0.32 or digits 7296 seen A1 cao **Or** M1 for 2430 × 32 (or digits 77760 seen) or 2658 × 32 (or digits 85056 seen) A1 if 1 correct M1 for "85056" – "77760" or 7296 seen A1 cao
	(b)	$\frac{2}{5} \times 145 = 58$ 145 – "58"	87	3	M1 $\frac{2}{5} \times 145$ (or M1 $\frac{3}{5}$ seen) A1 58 (or M1 $\frac{3}{5} \times 145$) A1 for 87 ft
	(c)(i) (ii)		80 125	2	B1 for 80 (± 1) B1 125 (± 3)


Paper 5521/02				
No	**Working**	**Answer**	**Mark**	**Notes**
10 (a)		1.5 – 2.0	1	B1 for height between 1.5m – 2.0m inclusive
(b)	Height of man × "2.5"	3 – 6	3	B3 for height between 3m – 6m inclusive (B2 for multiplying (a) by a number between 2 and 3 inclusive) (B1 for multiplying (a) by a number)
11	61 – 19 = 42 42 ÷ 3 = 14	14	2	M1 for – 19 or 42 seen A1 cao
12 (a)		5	1	B1
(b)	4+5+5+5+4+3+2+1+4+5 = 38 mean = 38 ÷ 10 = 3.8	3.8	2	M1 for attempt to add and ÷ 10 or 3.7 or 3.9 seen A1 for 3.8 SC B1 for 33.5 seen
13 (a)		$3x$	1	B1 cao Accept $3 \times x$, $x3$, $x \times 3$, $x + x + x$
(b)		x - 9	1	B1 for x – 9 cao
14 (a)	14.44 – 8.660254038	5.77974(…)	2	M1 for 14.44 seen or 8.66(…) or 5.7 or 5.8 or better rounded or truncated
(b)		6	1	A1 cao B1 ft
15	15 ÷ 24	62.5	2	M1 for 15 ÷ 24 or 1500 ÷ 24 or sight of digits 625 A1 cao
16	2.10 × 450	945	2	M1 for digits 210 × 450 or sight of digits 945 A1 cao
17	See diagram	$2(y + y)$ $2y + 2y$	2	B1 for $2(y + y)$ B1 for $2y + 2y$ (Deduct B1 for each additional tick (>2) to min 0)
18	360° ÷ 18 (=20) Sector angles: G= 60; S= 80; B=220; Correct sectors labelled correctly Use angle measurer	Angles drawn, labelled	4	B4 for fully correct and labelled pie chart (B3 for all angles correct or for a labelled pie chart with 2 correct angles) (B2 for labelled pie chart with 1 correct angle drawn) (B1 for 360° ÷ 18 or 20 seen or implied)


Paper 5521/02					
No		Working	Answer	Mark	Notes
19	(a)		Correct plane	2	B2 for a correct plane defined by showing at least 2 lines. (B1 for a line of symmetry on one face)
	(b)		Correct net	2	B2 cao (B1 for 2 equilateral triangles joined appropriately to at least one rectangle or for 1 equilateral triangle joined appropriately to one of 3 rectangles)
	(c)		Correct drawing	2	B1 for two extra sides of length 6cm (± 2mm) B1 for construction arcs 6cm from each of the ends of the given line.
20	(a)		15	1	B1 for 15 (± 1)
	(b)		15	1	B1 for 15 (± 0.4)
	(c)			2	B1 horiz. line from (2, 20) to (3, 20) B1 line from (3, 20) to (5,0) or horizontal translation of it SC B1 for any journey ending at (5,0)
21	(a)	$x+4+x+x+4+x$	$4x+8$	2	M1 for attempting to add x, $x + 4$, x, $x + 4$ may be implied by $4x + a$ ($a > 0$) A1 for $4x + 8$ or $4(x + 2)$
	(b)	$4x + 8 = 54$ $4x = 46$ $x = 11.5$ Length = "11.5" + 4	15.5	3	M1 for "$4x + 8$" = 54 A1 cao for 11.5 seen B1 ft for "11.5" + 4
22		0.4 + 0.15 1 – "0.55"	0.45	2	M1 for 1 – sum A1 for 0.45 o.e. SC B1 for 0.81
23	(a)		3:1	1	B1 cao
	(b)	$\pi \times 2.45$	7.7	2	M1 for $\pi \times 2.45$ (accept π as 3.1 or better) A1 for 7.59 to 7.70

Paper 5521/02					
No		**Working**	**Answer**	**Mark**	**Notes**
24		7×10000	70 000	2	M1 for 7 X 10 000 or 7 X 100 X 100 A1 cao
25		5.40 ÷ 3 × 7	12.60	3	M1 for 5.40 ÷ 3 or sight of 1.8 M1 dep for "1.80" × 7 A1 for 12.6 or equivalent
26		7.60 x $\frac{17.5}{100}$ = 1.33 7.60 + 1.33 = 8.93 1650 x "8.93"	£14 734.50	4	M1 for 7.60 x $\frac{17.5}{100}$ or 1.33 seen or 7.60 x 1.175 (oe) (Award M1 for 10%, 5% and 2½% correctly calculated) A1 for 8.93 or 893 M1 for 1650 x "8.93" or digits 147345 seen A1 cao Accept 14734.5 **Alternative** M1 for 1650 × 7.6(0) or 12540 seen M1 for "12540" × $\frac{17.5}{100}$ or 2194.5 seen or "12540"× 1.175 (oe) (Award M1 for 10%, 5% and 2½% correctly calculated) M1 for "12540" + "2194.5" (dep on both previous method marks) or digits 147345 seen A1 cao accept 14 734.5

Paper 5523/03				
No	**Working**	**Answer**	**Mark**	**Notes**
1		Correct shape	2	B2 B1 for any 2 sides correct, or a correct enlargement scale factor ≠1 or 2.
2 (a)		11 13 16 8 21	2	B2 all correct (B1 for 2 correct) sign
(b)		$\frac{31}{80}$	1	B1 oe
3 (a)		$2p+3q$	2	B2 for $2p+3q$ (accept $2\times p$ etc) (B1 for $2p$ or $3q$ or $2p3q$)
(b)		$2y^2$	1	B1 accept $2\times y^2$ or $2\times y\times y$
(c)		$3c+4d$	2	B2 for $3c+4d$ (accept $3\times c$ etc) (B1 for $3c$ or $4d$ or $3c4d$)
(d)		$8pq$	1	B1 accept in any order but must not include ×
4 (a)(i)		60	2	B1 cao
(ii)		eg top triangle is equilateral		B1 for reason
(b)		150	2	M1 $\frac{(180-"60")}{2}+90$ A1 ft from (a)(i) if $x < 90°$ SC:B1 for answer from "60" + 90 if $x < 90°$

Paper 5523/03					
No		Working	Answer	Mark	Notes
5	(a)	$4\times3-2\times1$ $12-2$	10	3	M1 for 3×4 (=12) or 1×2 or attempt to divide diagram up into rectangles M1 "12" – "2" A1 cao
	(b)(i)	$\frac{10}{100}\times680$ or $680\div10$ 680 + 68	748	5	M1 $\frac{10}{100}\times680$ or $680\div10$ or 68 seen M1 (dep) 680 +"68" (or M2 for 680×1.10) A1 cao
	(ii)	"748" ÷ 50 or 14.96	15		M1 for "748" $\div50$ or 14.96; accept "748" rounded up or down to next 50 followed by $\div50$ A1 ft from (b)(i) rounded up SC B1 for 680 (seen) leading to 14
6	(a)	$2\times5+5\times-3=10-15=$	-5	2	M1 substitute e.g 2×5 and 5×-3 or 10 and -15 A1 cao
	(b)	$5\times4^2-7$ $5\times16-7$	73	3	M1 substitution e.g $5\times4^2-7$; do not accept 54^2-7 M1 $5\times16-7$ or $5\times4\times4-7$ or $80-7$ (NB 4^2 as 4×4) A1 cao

Paper 5523/03					
No		**Working**	**Answer**	**Mark**	**Notes**
7		679 × 28 = 5432 + 13580 = 19012 or 28 × 679 = 252 + 1960 + 16800 = 19012 or  	190.12	3	M1 for an attempt to multiply the units and tens, or correct partitioning M1 complete correct method (condone one arithmetic error) A1 for 190.12 cao OR M1 for putting the numbers in a grid M1 for multiplying out and addition (condone one error) A1 answer shown with point OR M1 for correct partitioning M1 679×20 and 679×8 calculated oe (condone one error) A1 cao
8	(a)		Points plotted	1	B1 ±1 full (2mm) square
	(b)		positive	1	B1 cao
	(c)		Line of best fit	1	B1 must pass through (42.5, 1.45), (42.5, 1.55) AND (67.5, 1.75), (67.5, 1.85)
	(d)		approx 1.65	1	B1 ft from single line segment with positive gradient ± 1 full (2mm) square

Paper 5523/03					
No		**Working**	**Answer**	**Mark**	**Notes**
9	(a)	eg $50 \times \frac{2000}{500}$	200	2	M1 $2000 \div 500$ or 4 seen A1 cao
	(b)	eg $400 \times \frac{750}{500}$	600	2	M1 $750 \div 500$ or 1.5 seen or 400 + 200 A1 cao
10			$C = 10(n+3)$	3	B3 for $C = 10(n+3)$ oe such as $C = (n+3) \times 10$ (B2 for correct RHS or $C = n + 3 \times 10$, $C = 10n + 3$ etc B1 for C = some other linear expression in n or $n + 3 \times 10$, $10n + 3$ etc) NB: C=n scores no marks
11	(a)		$p(p+6)$	2	B2 for $p(p+6)$ or $p \times (p+6)$ (B1 for $p(ap+b)$ where a,b are numbers or $p+6$ seen on it's own, or part of an expression)
	(b)	$x^2 - 4x + 7x - 28$	$x^2 + 3x - 28$	2	M1 for 4 terms correct ignoring signs (e.g $x^2, 4x, 7x, 28$) or 3 terms with correct signs (e.g $x^2, -4x, 7x, -28$) A1 cao
12			correct drawing	2	B2. Condone hidden detail shown with solid lines, or missing lines on front face. (B1 for : one sketch correct with other sketches incorrect cross-section correct with depth > 1 cube, correct plan and side elevation)

Paper 5523/03					
No		**Working**	**Answer**	**Mark**	**Notes**
13		$\frac{600}{3\times10}$ or $\frac{640}{3.2\times10}$	20 to 21 $\frac{1}{3}$	2	M1 For rounding at least two of the numbers to 1 sf, or for sight of 640, 3.2 or 640, 32 or 600, 32 or 30 seen A1 for 20 to 21 $\frac{1}{3}$ NB: 20.3125 scores M0 A0
14	(a)		correct reflection	2	B2 (B1 reflection in line other than x=3)
	(b)		reflection in $y = x$	2	B2 cao Accept the word "reflected" (B1 any statement including the word "reflection")
15	(a)(i)		5^6	1	B1 accept 15125, 5^{4+2}
	(ii)		5^3	1	B1 accept 125, 5^{9-6}
	(b)	$x + y = 10$ and $x - y = 4$	$x = 7$ $y = 3$	3	M1 for either $x + y = 10$ or $x - y = 4$ A2 for both values correct [(A1 for one value correct) If M0, award B3 for both values correct or B2 for one value correct, otherwise B0] SC B2 for $x = 3$ and $y = 7$

Paper 5523/03				
No	**Working**	**Answer**	**Mark**	**Notes**
16 (a)	$5-3x=2x+2$ $5-2=2x+3x$	$\frac{3}{5}$	3	B1 for $2x+2$ seen OR $2.5-1.5x=x+1$ M1 for correct rearrangement of 4 terms A1 for $\frac{3}{5}$ oe
(b)		$-3,-2,-1,0,1,2$	2	B2 (B1 for 5 correct and not more than one incorrect integers)
17		question + response boxes oe	2	1st aspect: One question with time period (eg each night); ignore other questions. 2nd aspect: Response list (at least two), not overlapping.* 3rd aspect: Some mention of units (eg hours) in either question or responses Award B2 for all three aspects, or B1 for just two aspects. * 0-1, 2-3, 4-5 is OK, but 0-1, 1-2, 2-3 is not OK.

Paper 5523/03					
No		**Working**	**Answer**	**Mark**	**Notes**
18	(a)	$\frac{2}{3}\times\frac{3}{4}=\frac{6}{12}$	$\frac{1}{2}$	2	M1 for $\frac{6}{12}$ or $\frac{3}{6}$ or $\frac{2\times3}{3\times4}$ A1 accept 0.5
	(b)	$1+2+\frac{8}{12}+\frac{9}{12}$	$4\frac{5}{12}$	3	M1 for attempt to convert to fractions with common denominator e.g two fractions, denominator of 12 A1 correct conversion : $\frac{8}{12}$ and $\frac{9}{12}$, or $\frac{20}{12}$ and $\frac{33}{12}$ seen (oe) A1 cao for $4\frac{5}{12}$ OR attempts to convert to decimals: must use at least 2dp M1 0.66+0.75 (or 1.66+2.75) or 0.67+0.75 etc A1 4.41, 4.417, 4.416 A1 4.416 (*recurring*)
19		$2\times\frac{1}{2}\times6\times8$ or 48 $8\times9+6\times9+10\times9$ or 72 +54 + 90	264 cm^2	4	M1 attempt to find the area of one face; $\frac{1}{2}\times6\times8$ or (8×9) or (6×9) or (10×9) or 72 or 54 or 90 or 24 M1 all five faces with an intention to add A1 cao numerical answer of 264 B1 (indep) cm^2 with or without numerical answer

Paper 5523/03					
No		**Working**	**Answer**	**Mark**	**Notes**
20			$\frac{\pi ab^3}{3d}$ $3(c+d)^3$ $3\pi bc^2$	3	B3 (B1 for each one correct) Nb –B1 for each of the 4th,5th,6th tick
21	(a)	$x+0.3+0.2+x=1$	0.25	2	M1 for $x+0.3+0.2+x=1$ oe, or $0.5\div 2$ A1 oe
	(b)	0.3×200	60	2	M1 0.3×200 A1 cao Accept 60 out of 200 (in words) SC B1 for $\frac{60}{200}$
22	(a)		$(-12)-4\quad -2\quad (0)8$	3	B3 for all correct [(B1 for each one correct)
	(b)		points plotted accurately points joined with smooth curve	2	B1 ± 1 full (2mm) square ft table if at least B1 awarded (all 5 points plotted) B1 ft for any smooth curve if previous B1 gained NB: curve must pass within 1 full square of the points
23			$\frac{1}{4}$ on LH branch $\frac{2}{3}$ & $\frac{1}{3}$ & $\frac{2}{3}$ on RH branches	2	B1 B1
24			m=3 n=5	2	B1 for 3 B1 for 5 (B2 for $2^3\times 5$ or $2\times 2\times 2\times 5$) SC: award B1 only if m=5, n=3 or for 8×5 seen

Paper 5523/03					
No		Working	Answer	Mark	Notes
25		$\frac{5--1}{-1-2} = -2$	$y = -2x + 5$	4	M1 for clear attempt to find gradient eg fraction with -1,5 in numerator, 2,-1 in denominator A1 for -2 B2 ft for $y =$"-2"$x + 5$ oe (eg $y = \frac{-6}{3}x + 5$) (B1 for $y = mx + 5$ or , -2x+5 or $y =$"-2"$x + c$)
26	(a)(i)		150	2	B1 accept 150 or 210
	(ii)				B1 for angle at the centre is twice the angle at the circumference
	(b)		30	3	B1 identifies angle between radius and tangent as $90°$ (may be in working or on diagram) M1 $360° - 90 - 90 -$"150" A1 ft from (a)(i) excluding a negative answer OR B1 for 90 M1 for $2 \times (180 - 90 -$"$\frac{150}{2}$"$)$ A1 ft from (a)(i) excluding a negative answer OR B3 for $180 -$(a)
		$360 - 90 - 90 -$"150"			

Paper 5523/04					
No		**Working**	**Answer**	**Mark**	**Notes**
1	(a)	14.44 – 8.660254038	5.77974(….)	2	M1 for 14.44 seen or 8.66(……) seen or 5.7 or 5.8 or better, rounded or truncated A1 cao
	(b)		6	1	B1 ft
2		15 ÷ 24	62.5	2	M1 for 15 ÷ 24 or 1500 ÷ 24 or sight of digits 625 A1 cao
3	(a)	2.10 × 450	945	2	M1 for digits 210 × 450 or sight of digits 945 A1 cao
	(b)	63÷2.10	30	2	M1 for 63 ÷ digits 210 A1 cao
4		See diagram	$2(y + y)$ $2y + 2y$	2	B1 for $2(y + y)$ B1 for $2y + 2y$ (Deduct B1 for each additional tick (>2) to min 0)
5		360° ÷ 18 (=20) Sector angles: G= 60; S= 80; B=220; Correct sectors labelled correctly Use angle measurer	Angles drawn, labelled	4	B4 for fully correct and labelled pie chart (B3 for all angles correct or for a labelled pie chart with two angles correct) (B2 for labelled pie chart with one correct angle drawn) (B1 for 360÷18 or 20 seen or implied)
6	(a)		Correct plane	2	B2 for a correct plane defined by showing at least 2 lines. (B1 for a line of symmetry on one face)
	(b)		Correct net	2	B2 cao (B1 for 2 equilateral triangles joined appropriately to at least one rectangle or for 1 equilateral triangle joined appropriately to one of the three rectangles)
	(c)		Correct drawing	2	B1 for two extra sides of length 6 cm (± 2mm) B1 for construction arcs 6cm from each of the ends of the given line
7		61 – 19 = 42 42 ÷ 3 = 14	14	2	M1 for –19 or 42 seen or $3x + 19$ A1 cao

Paper 5523/04					
No		Working	Answer	Mark	Notes
8	(a) (b) (c)		15 15	1 1 2	B1 cao for $15(\pm 1)$ B1 cao for $15(\pm 0.4)$ B1 horiz. line from (2,20) to (3,20) B1 line from (3,20) to (5,0) or horiz. translation of it SC: B1 for any journey ending at (5,0)
9	(a)	$x+4+x+x+4+x$	$4x+8$	2	M1 for attempting to add $x+4, x, x+4, x$ may be implied by $4x+a$, $a>0$ A1 for $4x+8$ or $4(x+2)$
	(b)	$4x + 8 = 54$ $4x = 46$ $x = 11.5$ Length = "11.5" + 4	15.5	3	M1 for "$4x + 8$" = 54 A1 cao for 11.5 seen B1 ft for "11.5"+ 4
10		0.4 + 0.15 1 – 0.55	0.45	2	M1 for 1 – sum A1 for 0.45 oe SC: B1 for 0.81
11	(a) (b)	 $\pi \times 2.45$	3:1 7.7	1 2	B1 cao M1 for $\pi \times 2.45$ (accept π as 3.1 or better) A1 for 7.59 to 7.70
12		7×10000	70000	2	M1 for 7×10000 or $7 \times 100 \times 100$ A1 cao
13		$5.40 \div 3 \times 7$	12.60	3	M1 for $5.40 \div 3$ or sight of 1.8 M1 (dep) for "1.80" × 7 A1 for 12.6 or equivalent

Paper 5523/04				
No	**Working**	**Answer**	**Mark**	**Notes**
14	$7.60 \times \frac{17.5}{100} = 1.33$ 7.60 + 1.33 = 8.93 1650 × "8.93"	£14734.50	4	M1 for $7.60 \times \frac{17.5}{100}$ or 1.33 seen or 7.60×1.175 (oe) (Award M1 for 10%, 5% and 2½% correctly calculated) A1 for 8.93 or 893 M1 for 1650 × "8.93" or digits 147345 seen A1 cao Accept 14734.5 OR M1 for 1650×7.6 or 12540 seen M1 for "12540" $\times \frac{17.5}{100}$ or 2194.5 seen or "12540" $\times 1.175$ (oe) (Award M1 for 10%, 5%, and 2½% correctly calculated) M1 for "12540" + "2194.5" (dep on both previous M marks) or digits 147345 seen A1 cao accept 14734.5

Paper 5523/04					
No		**Working**	**Answer**	**Mark**	**Notes**
15		$285 \times 1000/(60 \times 60) = 79.1\dot{6}$		3	M2 for $285 \times 1000 \div 60 \div 60$ or $80 \times 60 \times 60 \div 1000$ or for a correct method to obtain two comparable values e.g $80 \times 60 \times 60$ and 285×1000 (M1 for $285 \div 60 \div 60$ or 0.079(….) seen or $80 \times 60 \times 60$ or 288000 seen or for 285×1000 or 285000 seen or $80 \div 1000$ or 0.08 seen) A1 for 288 or 79.(….) or for two correctly calculated comparable values e.g 288000 and 285000
16	(a)	$4x + 12 = 6$ $4x = -6$	-1.5	3	B1 for $4x + 12$ or $x + 3 = \frac{6}{4}$ M1 for a correct re-arrangement of their 3 terms to isolate $4x$ or x A1 for -1.5 oe
	(b)	$v - u = 5t$	$\frac{v-u}{5}$	2	M1 for isolating $\pm 5t$ or $\pm t$ or for dividing through by 5 A1 oe

Paper 5523/04					
No		**Working**	**Answer**	**Mark**	**Notes**
17		3→15 4→48 3.1→17.3(91) 3.2→19.9(68) 3.3→22.7(37) 3.4→25.7(04) 3.5→28.8(75) 3.4→25.7(04) 3.3→22.7(37) 3.35→24.1(95375)	3.3	4	B2 for trial between 3.3 and 3.4 inclusive (B1 for trial between 3 and 4 inclusive) B1 for different trial between 3.3 and 3.4 exclusive B1 (dep on at least one previous B1) for 3.3 NB trials should be evaluated to at least 1 dp truncated or rounded
18	(a)	36 ÷ (7+3+2) "3" × 7	21	3	M1 for 36 ÷ (7+3+2) M1 (dep) for "3" × 7 or 3 or 2 A1 cao
	(b)	$51.5 \times \frac{8.5}{100} = 4.3775$ 51.5 – 4.3775 = 47.1225	47 or 47.1 or 47.12	4	M1 for $51.5 \times \frac{8.5}{100}$ or 4.37(75) seen M1 (dep) for 51.5 – "4.37(75)" A1 for 47 or better B1 (indep) for rounding their answer correctly to the nearest whole number or 1 or 2 d.p OR M1 for $51.5 \times \frac{100-8.5}{100}$ M1 for $51.5 \times$"0.915" or 0.515 × "91.5" A1 for 47 or better B1 (indep) for rounding their answer correct to the nearest whole number or 1 or 2 d.p

Paper 5523/04					
No		**Working**	**Answer**	**Mark**	**Notes**
19	(a)		Angle in a semicircle	1	B1 oe
	(b)	$12^2 + 16^2 = 400$ $\sqrt{400} = 20$	20	3	M1 for $12^2 + 16^2$ M1 for $\sqrt{144+256}$ A1 cao
	(c)	$\pi \times 10^2$	314	3	M1 for $\pi \times \left(\frac{"20"}{2}\right)^2$ M1 (indep) for correct order of evaluation of $\pi \times r^2$ for any r A1 for 314 – 315 inclusive

Paper 5523/04					
No		Working	Answer	Mark	Notes
20	(a)	$(1 \times 10)+(3 \times 15)+(5 \times 30)+(7 \times 35)+$ $(9 \times 25)+(11 \times 5) = 730$ "730" $\div$ 120 = 6.08333	6.08	4	M1 for use of *fx* with *x* consistent within intervals (including end points) M1 (dep) for use of midpoints M1 (dep on 1st M1) for use of $\frac{\sum fx}{\sum f}$ A1 6.08 to 6.085
	(b)		(10),25,55,90,115, 120	1	B1 for all correct
	(c)		graph	2	B1 ft for 5 or 6 points plotted correctly $\pm$ 1 full (2mm) square at the end of interval dep on sensible table (condone 1 addition error) B1(dep) for points joined by curve or line segments provided no gradient is negative – ignore any part of graph outside range of their points. (SC: B1 if 5 or 6 points plotted not at end but consistent within each interval and joined)

Paper 5523/04				
No	**Working**	**Answer**	**Mark**	**Notes**
(d)		72–74	2	M1 (ft dep on graph being cf) for reading from graph at 7 A1 ft ±1 full (2 mm) square Or B2 for $72-74$
21 (a) (b) (c) (d)		a^7 $15x^3y^4$ $x-1$ $(x+3)(x-3)$	1 2 1 1	B1 accept a^{4+3} B2 cao (B1 for two of 15, x^3, y^4 in a product) B1 cao B1 cao
22	80% = 220 $220 \div 80 \times 100$	275	3	M1 for recognising that 80% is equivalent to 220 M1 for $220 \div 80 \times 100$ oe A1 cao

Paper 5523/04					
No		**Working**	**Answer**	**Mark**	**Notes**
23			$x = 3$ $y = 0.5$	3	M1 for coefficients of x or y the same followed by correct operation, condone one arithmetical error M1 (dep) for substituting found value in one equation A1 cao SC: B1 for one correct answer only if M's not awarded
24			1.4×10^{10}	2	B2 for 1.4×10^{10} or 1.44×10^{10} (B1 for 14.4×10^{9} or 14400,000,000 or 14000,000,000 or 14×10^{9})
25	(a)	$\tan x = \frac{1.9}{3.2}$ $x = \tan^{-1}\left(\frac{1.9}{3.2}\right) = 30.7$	30.7	3	M1 for $\tan x = \frac{1.9}{3.2}$ or $\tan \frac{1.9}{3.2}$ M1 for $\tan^{-1}\left(\frac{1.9}{3.2}\right)$ A1 for 30.6 – 30.7
	(b)	90 + "30.7"	121	1	B1 (indep) ft for 90 + "30.7" rounded to 3 or 4 s.f

Paper 5523/04				
No	**Working**	**Answer**	**Mark**	**Notes**
26	SF$=\frac{12}{9}$ $\frac{12}{9}\times 6=8$	2	2	M1 for $\frac{12}{9}$ or $\frac{9}{12}$ or 1.33… seen or 0.75 seen or 8 seen or $\frac{6}{9}$ or $\frac{9}{6}$ or 0.66… or 1.5 or $\frac{1}{3}$ or 3 oe seen A1 cao

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
1	(a)	$x^2-4x+7x-28$	$x^2+3x-28$	2	M1 for 4 terms correct ignoring signs (e.g $x^2,4x,7x,28$) or 3 terms with correct signs (e.g $x^2,-4x,7x,-28$) A1 cao
	(b)		y^4+2y^2	2	B2 cao B1 for y^4 or $2y^2$
	(c)		$p(p+6)$	2	B2 for $p(p+6)$ or $p\times(p+6)$ (B1 for $p(ap+b)$ where a,b are numbers or $p+6$ seen on it's own, or part of an expression)
	(d)		$3x(2x-3y)$	2	B2 (B1 for $3(2x^2-3xy)$ or $x(6x-9y)$ or $3x(...)$)
2			question + response boxes oe	2	1st aspect: One question with time period (eg each night); ignore other questions. 2nd aspect: Response list (at least two), not overlapping.* 3rd aspect: Some mention of units (eg hours) in either question or responses Award B2 for all three aspects, or B1 for just two aspects. * 0-1, 2-3, 4-5 is OK, but 0-1, 1-2, 2-3 is not OK.
3	(b)		reflection in $y=x$	2	B2 cao accept the word "reflected" (B1 any statement including the word "reflection")
4	(a)	$5-3x=2x+2$ $5-2=2x+3x$	$\frac{3}{5}$	3	B1 for $2x+2$ seen OR $2.5-1.5x=x+1$ M1 for correct rearrangement of 4 terms A1 for $\frac{3}{5}$ oe
	(b)		$-3,-2,-1,0,1,2$	2	B2 (B1 for 5 correct and not more than one incorrect integers)

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
5	(a)	$\frac{2}{3}\times\frac{3}{4}=\frac{6}{12}$	$\frac{1}{2}$	2	M1 for $\frac{6}{12}$ or $\frac{3}{6}$ or $\frac{2\times3}{3\times4}$ A1 accept 0.5
	(b)	$1+2+\frac{8}{12}+\frac{9}{12}$	$4\frac{5}{12}$	2	M1 for attempt to convert to fractions with common denominator e.g two fractions, denominator of 12 A1 correct conversion : $\frac{8}{12}$ and $\frac{9}{12}$, or $\frac{20}{12}$ and $\frac{33}{12}$ seen (oe) A1 cao for $4\frac{5}{12}$ OR attempts to convert to decimals: must use at least 2dp M1 0.66+0.75 (or 1.66+2.75) or 0.67+0.75 etc A1 4.41, 4.417, 4.416 or 0.41, 0.417, 0.416 or 0.42, 4.42 A1 4.416 (*recurring*)
6	(a)(i)		5^6	1	B1 accept 15625, 5^{4+2}
	(ii)		5^3	1	B1 accept 125, 5^{9-6}
	(b)	$x+y=10$ and $x-y=4$	$x = 7$ $y = 3$	3	M1 for either $x + y = 10$ or $x-y=4$ A2 for both values correct [(A1 for one value correct) If M0, award B3 for both values correct or B2 for one value correct, otherwise B0] SC B2 for $x = 3$ or $y = 7$

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
7		$2\times\frac{1}{2}\times6\times8$ or 48 $8\times9+6\times9+10\times9$ or 72 +54 + 90	264 cm^2	4	M1 attempt to find the area of one face; $\frac{1}{2}\times6\times8$ or (8×9) or (6×9) or (10×9) or 72 or 54 or 90 or 24 or 48 M1 all five faces with an intention to add A1 cao numerical answer of 264 B1 (indep) cm^2 with or without numerical answer
8			$\frac{\pi ab^3}{3d}$ $3(c+d)^3$ $3\pi bc^2$	3	B3 (B1 for each one correct) Nb –B1 for each of the 4th,5th,6th tick
9	(a)	$x+0.3+0.2+x=1$	0.25	2	M1 for $x+0.3+0.2+x=1$ oe, or 0.5 ÷ 2 A1 oe
	(b)	0.3×200	60	2	M1 0.3×200 A1 cao Accept 60 out of 200 (in words) SC B1 for $\frac{60}{200}$
10	(a)		$(-12)-4\quad-2\quad(0)8$	3	B3 for all correct [(B1 for each one correct)
	(b)		5 points plotted accurately points joined with smooth curve	2	B1 ±1 full (2mm) square ft table if at least B1 awarded (all 5 points plotted) B1 ft for any smooth curve if previous B1 gained NB: curve must pass within 1 full square of the points
11			m=3 n=5	2	B1 for 3 B1 for 5 (B2 for $2^3\times5$ or $2\times2\times2\times5$) SC: award B1 only if m=3, n=3, for 8×5 seen

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
12		$\frac{5--1}{-1-2}$ = -2	$y=-2x+5$	4	M1 for clear attempt to find gradient eg fraction with -1,5 in numerator, 2,-1 in denominator A1 for -2 cao B2 ft for $y=''-2''x+5$ oe (eg $y=\frac{-6}{3}x+5$) (B1 for $y=mx+5$ or , -2x+5 or $y=''-2''x+c$)
13	(a)		$\frac{1}{4}$ on LH branch $\frac{2}{3}$ & $\frac{1}{3}$ & $\frac{2}{3}$ on RH branches	2	B1 cao B1
	(b)	$\frac{3}{4}\times\frac{2}{3}+\frac{1}{4}\times\frac{1}{3}=\frac{6}{12}+\frac{1}{12}$	$\frac{7}{12}$	3	M1 for $\frac{3}{4}\times\frac{2}{3}$ or $\frac{1}{4}\times\frac{1}{3}$ from their tree diagram M1 for sum of two products A1 for $\frac{7}{12}$ oe
	(c)	$n=21\times4$ or $\frac{1}{6}:\frac{1}{4}$ oe $\frac{1}{6}\times84$ or $21\times\frac{2}{3}$	14	3	M1 for either $\frac{1}{3}\times\frac{3}{4}\left(=\frac{1}{4}\right)$ or $\frac{2}{3}\times\frac{1}{4}\left(=\frac{1}{6}\right)$ from their tree diagram M1 for 21×4 (=84) or $\frac{21}{3}\times2$ A1 for 14 cao SC: B2 for 63 seen in fraction or ratio

Paper 5525/05					
No		Working	Answer	Mark	Notes
14	(a)(i)		150	2	B1 accept 150 or 210
	(ii)				B1 for angle at the centre is twice the angle at the circumference
	(b)		30	3	B1 identifies angle between radius and tangent as $90°$ (may be in working or on diagram)
		$360 - 90 - 90 - "150"$ or $180 - "150"$			M1 $360° - 90 - 90 - "150"$ A1 ft from (a)(i) excluding a negative answer Or B1 for 90 M1 for $2 \times (180 - 90 - "\frac{150}{2}")$ A1 ft from (a)(i) excluding a negative answer Or B3 for $180 - (a)$ SC: 180 – "210" can get B1 for $90°$ and/or B1 for "cyclic quadrilateral"
15	(a)		0.2727…	1	B1 for 2.27 recurring or 0.2727…. oe or 0.273
	(b)	eg $x = 0.3939…$ so $100x = 39.3939…$ $99x = 39$ so $x = \frac{39}{99} = \frac{13}{33}$		3	M1 for $100x = 39.39…$ M1 dep for subtraction of both sides A1 for $\frac{13}{33}$ from correct proof **Alternative method** M1 for $13.000 \div 33$ M1 for remainders 31 and 13 A1 for 0.39 recurring SC:B1 for $\frac{39}{99}$

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
16	(a)	$d = kt^2$ $80 = k \times 4^2$	$d = 5t^2$	3	M1 for $d = kt^2$ or $d \propto t^2$ M1 sub d=80 and t=4 into their equation A1 for $d = 5t^2$ oe (cao)
	(b)		245	1	B1 ft from (a) using "k"
	(c)	$45 = 5t^2$	3	2	M1 ft from (a) for substituting d=45 into their equation A1 for 3 cao (condone inclusion of -3)
17	(a)(i) (ii)		(0, 9) (8, 25)	3	B1 cao B1 for $x = 8$ cao B1 for $y = 25$ cao SC: B1 for (25, 8)
	(b)	LHS = $\left(\frac{100-(x^2-16x+64)}{4}\right)$ = $\left(\frac{36+16x-x^2}{4}\right)$ RHS = $\left(\frac{36-2x+18x-x^2}{4}\right)$ = LHS		3	M1 for expansion of either set of brackets with at least 3 of 4 terms correct M1 for common denominator of 4 or multiplying through by 4 or reducing each numerator to a single term A1 for fully correct solution **Alternative method** M1 for $(5-\frac{(x-8)}{2})(5+\frac{(x-8)}{2})$ M1 for $(\frac{2\times5-(x-8)}{2})(\frac{2\times5+(x-8)}{2})$ A1 for $\frac{(18-x)(x+2)}{4}$

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
18	(a)	$\frac{810\pi}{90\pi}$ or 9 $\sqrt{9}$ or 3	12	3	M1 for $\frac{810\pi}{90\pi}$ or 9 or $\frac{1}{9}$ or 1:9 oe M1 for $\sqrt{\frac{810\pi}{90\pi}}$ or $\sqrt{9}$ or 3 or $\frac{1}{3}$ or $\sqrt{9}:\sqrt{1}$ oe A1 cao
	(b)	3^3 or 27 or 2700	2700π	2	SC:B1 for answer of 36 M1 for "3"^{3} or 27 or $\left(\sqrt{9}\right)^3 : (\sqrt{81})^3$ oe or 9^3 or 2700– A1 cao
19	(a)(i)		1	1	B1 cao
	(ii)		8	1	B1 cao
	(iii)	$64^{-\frac{2}{3}} = \frac{1}{64^{\frac{2}{3}}}$ or $64^{-\frac{2}{3}} = (4^2)^{-1}$	$\frac{1}{16}$	2	M1 for knowing negative power is a reciprocal or power of $\frac{1}{3}$ root is a cube root A1 cao for $\frac{1}{16}$
	(b)	$\sqrt{27} = \sqrt{9\times3}$ or $\sqrt{27} = 3\sqrt{3}$ or $\sqrt{27} = 3^{3/2}$	$\frac{5}{2}$ oe	2	M1 for $\sqrt{27} = \sqrt{9\times3}$ or $\sqrt{27} = 3^{3/2}$ A1 for $\frac{5}{2}$ oe (cao) **Alternative method** M1 for $9\times27 = 3^{2n}$ A1 for $\frac{5}{2}$ oe (cao)

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
20	(a)(i) (ii) (b)(i) (ii)		(90, 1) (180, 0) (45, 0) (90, -3)	2 2 	B1 cao could be indicated on diagram B1 cao could be indicated on diagram B1 cao could be indicated on diagram B1 cao could be indicated on diagram
21		$\frac{1}{3}\pi x^2 h = \frac{4}{3}\pi(2x)^3$ $x^2 h = 4\times 8x^3$	$32x$	3	M1 for substitution in correct formulae M1 (dep.) for correct unsimplified expression eg $h = \frac{\frac{4}{3}\pi(2x)^3}{\frac{1}{3}\pi x^2}$ oe or $h = 8x$ oe A1 for $32x$ cao
22	(a)	$\left(\overline{OM} =\right)\ \boldsymbol{a} + 2\boldsymbol{b}$ $\left(\overline{ON} =\right) 3\boldsymbol{a}$ or $\frac{6}{2}\boldsymbol{a}$ $\left(\overline{MN} =\right)\ -\boldsymbol{a} - 2\boldsymbol{b} + 3\boldsymbol{a}$	$2\boldsymbol{a} - 2\boldsymbol{b}$	2	B2 (B1 for either $\overline{OM}$ or $\overline{ON}$ or $-\boldsymbol{a} - 2\boldsymbol{b} + 3\boldsymbol{a}$ SC: B1 for $2\boldsymbol{b} - 2\boldsymbol{a}$
	(b)	$\left(\overline{OX} =\right)\ 2\boldsymbol{a} + \boldsymbol{b}$ $\left(\overline{OY} =\right) \boldsymbol{b} + 4\boldsymbol{a}$ (½ $\overline{QR}$ =) $2\mathbf{a} - \boldsymbol{b}$ or (½ $\overline{RQ}$ =) $\boldsymbol{b} - 2\boldsymbol{a}$	$\overline{XY} = 2\boldsymbol{a}$ (hence parallel)	2	B1 for either $\overline{OX}$ or $\overline{OY}$ or (½ $\overline{QR}$) B1 for $\overline{XY} = 2\boldsymbol{a}$ or $\overline{YX} = -2\boldsymbol{a}$

Paper 5525/06					
No		Working	Answer	Mark	Notes
1	(a)	$v - u = 5t$	$\frac{v-u}{5}$	2	M1 for isolating $\pm 5t$ or $\pm t$ or for dividing through by 5 A1 oe
	(b)	$x - 3 = 5x - 25$ $22 = 4x$	$5\frac{1}{2}$	3	M1 for $x - 3 = 5(x-5)$ or $\frac{x}{5} - \frac{3}{5} = x - 5$ M1 for isolating terms in x correctly from $ax + b = cx + d$ A1 cao accept $5\frac{1}{2}$, $\frac{11}{2}$, 5.5
2	(a)	$36 \div (7+3+2)$ "3" $\times 7$	21	3	M1 for $36 \div (7+3+2)$ M1 (dep) for "3" $\times 7$ or 3 or 2 A1 cao
	(b)	$51.5 \times \frac{8.5}{100} = 4.3775$ $51.5 - 4.3775 = 47.1225$	47 or 47.1 or 47.12	4	M1 for $51.5 \times \frac{8.5}{100}$ or 4.37(75) seen M1 (dep) for 51.5 – "4.37(75)" A1 for 47 or better B1 (indep) for rounding their answer correctly to the nearest whole number or 1 or 2 d.p OR M1 for $51.5 \times \frac{100 - 8.5}{100}$ M1 for $51.5 \times$"0.915" or 0.515 ×"91.5" A1 for 47 or better B1 (indep) for rounding their answer correct to the nearest whole number or 1 or 2 d.p

Paper 5525/06					
No		**Working**	**Answer**	**Mark**	**Notes**
3		3→15 4→48 3.1→17.3(91) 3.2→19.9(68) 3.3→22.7(37) 3.4→25.7(04) 3.5→28.8(75) 3.4→25.7(04) 3.3→22.7(37) 3.35→24.1(95375)	3.3	4	B2 for trial between 3.3 and 3.4 inclusive (B1 for trial between 3 and 4 inclusive) B1 for different trial between 3.3 and 3.4 exclusive B1 (dep on at least one previous B1) for 3.3 NB trials should be evaluated to at least 1 dp truncated or rounded
4	(a)		Angle in a semicircle	1	B1 oe
	(b)	$12^2 + 16^2 = 400$ $\sqrt{400} = 20$	20	3	M1 for $12^2 + 16^2$ M1 for $\sqrt{144+256}$ A1 cao
	(c)	$\pi \times 10^2$	314	3	M1 for $\pi \times \left(\frac{"20"}{2}\right)^2$ M1 (indep) for correct order of evaluation of $\pi \times r^2$ for any r A1 for 314 – 315 inclusive

Paper 5525/06					
No		Working	Answer	Mark	Notes
5	(a)	$(1 \times 10)+(3 \times 15)+(5 \times 30)+(7 \times 35)+$ $(9 \times 25)+(11 \times 5) = 730$ "730" ÷ 120 = 6.08333	6.08	4	M1 for use of *fx* with *x* consistent within intervals (including end points) M1 (dep) for use of midpoints M1 (dep on 1st M1) for use of $\frac{\sum fx}{\sum f}$ A1 6.08 to 6.085
	(b)		(10),25,55,90, 115,120	1	B1 for all correct
	(c)		graph	2	B1 ft for 5 or 6 points plotted correctly ± 1 full (2mm) square at the end of interval dep on sensible table (condone 1 addition error) B1(dep) for points joined by curve or line segments provided no gradient is negative – ignore any part of graph outside range of their points. (SC: B1 if 5 or 6 points plotted not at end but consistent within each interval and joined)
	(d)		72 – 74	2	M1 (ft dep on graph being cf) for reading from graph at 7 A1 ft ±1 full (2 mm) square Or B2 for $72 - 74$

Paper 5525/06				
No	**Working**	**Answer**	**Mark**	**Notes**
6 (a)		a^7	1	B1 accept a^{4+3}
(b)		$15x^3y^4$	2	B2 cao (B1 for two of 15, x^3, y^4 in a product)
(c)		$x-1$	1	B1 cao
(d)		$(a + 3b)(a - 3b)$	2	B2 for $(a + 3b)(a - 3b)$ (B1 for $(a \pm 3b)(a \pm 3b)$
7	80% = 220 $220 \div 80 \times 100$	275	3	M1 for recognising that 80% is equivalent to 220 M1 for $220 \div 80 \times 100$ oe A1 cao
8		$x = 3$ $y = 0.5$	3	M1 for coefficients of x or y the same followed by correct operation, condone one arithmetical error M1 (dep) for substituting found value in one equation A1 cao SC: B1 for one correct answer only if Ms not awarded
9		1.4×10^{10}	2	B2 for 1.4×10^{10} or 1.44×10^{10} (B1 for 14.4×10^9 or 14400,000,000 or 14000,000,000 or 14×10^9)

Paper 5525/06					
No		**Working**	**Answer**	**Mark**	**Notes**
10	(a)	$\tan x = \frac{1.9}{3.2}$ $x = \tan^{-1}\left(\frac{1.9}{3.2}\right) = 30.7$	30.7	3	M1 for $\tan x = \frac{1.9}{3.2}$ or $\tan \frac{1.9}{3.2}$ M1 for $\tan^{-1}\left(\frac{1.9}{3.2}\right)$ A1 for 30.6 – 30.7
	(b)	90 + "30.7"	121	1	B1 (indep) ft for 90 + "30.7" rounded to 3 or 4 s.f
11	(a)	$SF = \frac{12}{9}$ $\frac{12}{9} \times 6 = 8$	2	2	M1 for $\frac{12}{9}$ or $\frac{9}{12}$ or 1.33… seen or 0.75 seen or 8 seen or $\frac{6}{9}$ or $\frac{9}{6}$ or 0.66… or 1.5 or $\frac{1}{3}$ or 3 oe seen A1 cao
	(b)	$SF = \frac{9}{12}, \frac{9}{12} \times 7 = 5.25$	5.25	2	M1 for $\frac{BE}{7} = \frac{9}{12}$ oe A1 cao


Paper 5525/06					
No		**Working**	**Answer**	**Mark**	**Notes**
12	(a)	$84 = 6.7\pi + 2 \times 6.7 + 2a$ $2a + 13.4 = 62.95\ldots$ or $2a + 34.44 = 84$	24.8	3	M1 for substituting correctly, π may be left M1 for correct rearrangement as far as $\pm 2a$ A1 for $24.7 - 24.8$
	(b)	$P = \pi r + 2r + 2a$ $P - 2a = \pi r + 2r$ $P - 2a = (\pi + 2)r$	$\frac{P-2a}{\pi+2}$	3	M1 subtracting $2a$ from each side M1 for factorising to get $(\pi+2)r$ A1 for $\frac{P-2a}{\pi+2}$ oe S.C $\frac{p-2a}{5.14}$ oe is M1 M1 A0
13		Area $\Delta ABC = ½ \times 14 \times 8 \times \sin 106$ (= 53.8)	53.8	3	M1 for $½ \times 14 \times 8 \times \sin 106$ M1 (dep) for $56 \times 0.961(26..)$ or 107.6… A1 $53.8 - 53.9$ SC 107.6 is B2
14	(a)		4.5	1	B1 cao
	(b)	$500 \times 1.045^{20} = 1205.857\ldots\ldots$	1205.86	2	M1 for 500×1.045^{20} A1 for $1205.85 - 1206$ (SC:B1 for $705.85 - 706$ no working)

Paper 5525/06				
No	**Working**	**Answer**	**Mark**	**Notes**
15 (a)	$6x^2 + 11x - 10 + 6x - 4 = 25$ $6x^2 + 17x - 39 = 0$		3	M1 for an expression for the area involving either $(3x-2)(2x+5)+2(3x-2)$ or $3x(3x-2)+(3x-2)(7-x)$ or $3x(2x+5)-2(7-x)$ or $(3x-2)^2 + 2(3x-2) + (3x-2)(7-x)$ where in each case at least one of 2 or 3 product terms must be correct M1 (indep) for one correct expansion involving x^2 A1 for simplification to final answer

Paper 5525/06					
No		Working	Answer	Mark	Notes
15	(b)(i)	$x = \dfrac{-17 \pm \sqrt{17^2 - 4\times 6\times(-39)}}{2\times 6}$ $= \dfrac{-17 \pm \sqrt{289+936}}{12}$ $x = +\dfrac{18}{12}$ or -4.33	1.5, $-\dfrac{13}{3}$	4	M1 for $x = \dfrac{-17 \pm \sqrt{17^2 - 4\times 6\times(-39)}}{2\times 6}$ up to signs in b & c M1 for $x = \dfrac{-17 \pm \sqrt{1225}}{12}$ A1 $x = 1.5$ or -4.33, or better OR M2 for $(3x+13)(2x-3)$ (M1 for $(3x \pm a)(2x \pm b)$ with ab = ±39 A1 x = 1.5 or – 4.33, or better
		$x^2 + \dfrac{17}{6}x - \dfrac{39}{6} = 0$ $\left(x+\dfrac{17}{12}\right)^2 - \left(\dfrac{17}{12}\right)^2 - \dfrac{39}{6} = 0$ $\left(x+\dfrac{17}{12}\right)^2 = \left(\dfrac{17}{12}\right)^2 + \dfrac{39}{6}$			OR M1 for $\left(x+\dfrac{17}{12}\right)^2$ *seen* M1 $\left(x+\dfrac{17}{12}\right)^2 = \left(\dfrac{17}{12}\right)^2 + \dfrac{39}{6}$ A1 x = 1.5 or – 4.33, or better SC:M1 for answer "1.5" with no working or T & I
	(ii)		8		B1 cao length = 8

Paper 5525/06					
No		**Working**	**Answer**	**Mark**	**Notes**
16	(a)	P(win) = $\frac{2}{5}\times\frac{3}{5}+\frac{2}{5}\times\frac{1}{5}(=\frac{8}{25})$	$\frac{8}{25}$	3	M1 for $\frac{2}{5}\times\frac{3}{5}$ or $\frac{2}{5}\times\frac{1}{5}$ or for clearly identifying in P(R) $\times$P(R) + P(B) $\times$ P(B) M1 for P(win) = " $\frac{2}{5}$ " $\times$ " $\frac{3}{5}$ " + " $\frac{2}{5}$ " $\times$ " $\frac{1}{5}$ " A1 for $\frac{8}{25}$, oe
	(b)	$\frac{8}{25}$ $\times$ 100 (=32) 100 $\times$ 20 – "32" $\times$ 50	£4	2	M1 for (" $\frac{8}{25}$ " $\times$100)$\times$50 or $\times$ 0.5 A1 cao

Paper 5525/06					
No		**Working**	**Answer**	**Mark**	**Notes**
17		Lower bound of 1200 is 1150 Upper bound of 60 is 65 $1150 \div 65$	17	4	B1 for 1150 or 1250 seen B1 for 65 or 55 seen M1 (Lower bound of load) ÷ (Upper bound of weight) Where 1150 ≤ LB load < 1200 and 60 < UB Weight ≤ 65 A1 for 17 requires fully correct working OR B1 for 1150 or 1250 seen B1 for 65 or 55 seen M1 (upper bound of load) ÷(lower bound of weight) Where 1200 ‹ UB load ≤ 1250 and 55 ≤ LB weight < 60 A1 for 22 requires fully correct working OR M2 $1200 \div 55$ A1 21.8 A1 21 requires fully correct working OR M2 $1200 \div 65$ A1 18.4(6) A1 18 requires fully correct working
18	(a)	$3^4 x^4 y^8$	$81x^4y^8$	2	B2 for $81x^4y^8$ (B1 for 2 of 81, x^4, y^8)
	(b)	$\frac{x(x-3)}{(x-5)(x-3)}$	$\frac{x}{x-5}$	3	B1 for $x(x-3)$ B1 for $(x-5)(x-3)$ B1 cao

Paper 5525/06					
No		Working	Answer	Mark	Notes
19	(a)	$6^2 - 2^2 = 32$	5.66	2	M1 for $6^2 - 2^2$ (= 32) A1 5.65 – 5.66
	(b)	V  $DVA = 2 \times \sin^{-1}(\frac{2}{6})$ OR $\cos DVA = \frac{6^2 + 6^2 - 16}{2 \times 6 \times 6}$ $= \frac{56}{72}$ $DVA = \cos^{-1}\left(\frac{56}{72}\right) = 38.94$	38.9	3	M1 $\sin x = \frac{2}{6}$ oe M1 for $DVA = 2 \times \sin^{-1}(\frac{2}{6})$ A1 38.9 – 38.95 OR M1 for $(\cos DVA =) \frac{6^2 + 6^2 - 4^2}{2 \times 6 \times 6}$ M1 for $DVA = \cos^{-1}\left(\frac{56}{72}\right)$ A1 38.9 – 38.95

Paper 5525/06				
No	**Working**	**Answer**	**Mark**	**Notes**
19 (c)	$AC^2 = 2^2 + 2^2 - 2 \times 2 \times 2 \times \cos 120^o$ $AC = \sqrt{12}$ **OR** $AN = 2 \times \sin 60 = \sqrt{3}$ **OR** VN = $\sqrt{"32"+1} = \sqrt{33}$ $\cos AVC = \frac{6^2 + 6^2 - 12}{2 \times 6 \times 6}$ $\cos AVC = \frac{60}{72}$ **OR** $AVC = 2 \times \sin^{-1}\frac{\sqrt{3}}{6}$, using AN **OR** $AVC = 2 \times \cos^{-1}\frac{\sqrt{33}}{6}$, using VN	33.6	4	M1 for any valid method for AC or AN or VN where N is the midpoint of AC A1 for $AC^2 = 12$ or $AC = \sqrt{12}$ (= 3.46…) or $AN = \sqrt{3}$ (=1.73…)or $VN = \sqrt{33}$ (=5.74…) M1 (indep) for correct method to find angle AVC A1 33.55 – 33.6

Paper 5525/06					
No		**Working**	**Answer**	**Mark**	**Notes**
20	(a)	Graph translated 1 unit to the right passing through the points (-1,0), (1,2) and (2,0)		2	M1 for translation of $\begin{pmatrix}1\\0\end{pmatrix}$ or $\begin{pmatrix}-1\\0\end{pmatrix}$ A1 for right through the 3 points, $\pm\frac{1}{4}$ sq
	(b)	Graph stretched 2 units parallel to y-axis; passing through the points (-2,0), (0,4) and (1,0)		2	M1 for graph stretched parallel to the y-axis by scale factor 2 A1 through all 3 points; $\pm\frac{1}{4}$ sq not on grid at $x = 2$

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481

Email publications@linneydirect.com

Order Code UG016720 Summer 2005

For more information on Edexcel qualifications, please visit www.edexcel.org.uk/qualifications
Alternatively, you can contact Customer Services at www.edexcel.org.uk/ask or on 0870 240 9800

Edexcel Limited. Registered in England and Wales no.4496750
Registered Office: One90 High Holborn, London, WC1V 7BH

PMT

edexcel

Mark Scheme (Results)

June 2011

Modular Mathematics (GCSE)
Unit 2: 5MB2H_01 (Higher)

PMT

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at www.edexcel.com.

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:
http://www.edexcel.com/Aboutus/contact-us/

June 2011
Publications Code UG028428
All the material in this publication is copyright
© Edexcel Ltd 2011

PMT

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labeling conventions.

ii*) select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect canceling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

12 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers

Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

Guidance on the use of codes within this mark scheme
M1 – method mark A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

PMT

5MB2H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
1		$5 \times 6 \div 2$	15cm^2	3	M1 $5 \times 6 \div 2$ oe seen A1 15 B1 cm^2
2	(a)		$3y + 7x + 3$	1	B1 cao
	(b)		$2x(x - 2)$	2	B2 for $2x(x - 2)$. Accept $2x(x + -2)$. (B1 for $x(2x - 4)$ or $2(x^2 - 2x)$ or $2x$(linear expression in x) or $(x - 2)$(linear expression in x)
	(c)	$11-3x - 6$	$5-3x$	2	M1 for expansion of $-3(x + 2)$ A1 cao
	(d)	$3x^2+7x-18x-42$	$3x^2- 11x - 42$	2	M1 for 4 terms correct with or without signs or 3 out of exactly 4 terms correct (the terms may be in an expression or table) **OR** $x(3x+7)-6(3x+7)$ or $3x(x - 6) + 7(x - 6)$ A1 cao

PMT

5MB2H_01				
Question	**Working**	**Answer**	**Mark**	**Notes**
3 (a)	48=2 × 24 = 2 × 2 × 12 = 2 × 2 × 2 × 6 =2 × 2 × 2× 2 × 3 **OR** Use of factor trees	2 × 2 × 2 × 2 × 3	2	M1 for a systematic method that isolates 2 correct prime numbers correct divisions by a prime number **OR** an equivalent factor tree A1 for 2 × 2 × 2 × 2 × 3 or $2^4 \times 3$
(b)	Exeter coach after 20, 40, 60, 80, 100… Plymouth coach after 16, 32, 48, 64, 80, 96, … **OR** Timetable schedule for the coaches Exeter coach at 8.00, 8.20, 8.40, 9.00, 9.20… Plymouth coach at 8.00, 8.16, 8.32, 8.48, 9.04, 9.20… **OR** LCM of 16 and 20 is 80	9:20(am)	3	M1 for attempt to list multiples of 16 and 20 (at least first 3 of each, condone 1 addition error) M1(dep) for identifying their LCM A1 9:20 am (accept 9:20) **OR** M1 for attempt to draw up a timetable showing when the coaches will run (at least 3 extra times for each, condone 1 addition error) M1(dep) for identifying the first time common to both timetables A1 9:20 oe (do not accept 9:20pm) **OR** M1 attempt to find the LCM M1 for 2 × 2 × 2 × 2 × 5 oe seen A1 9:20 oe (do not accept 9:20pm)
4	10% = 6 6 × 4 = 24	24	2	M1 40 ÷ 100 × 60 oe or any complete method, eg 10% = 6, 6 × 4 A1 cao SC B1 for 36 or 84

5MB2H_01				
Question	**Working**	**Answer**	**Mark**	**Notes**
5 (a)	Table of values x = -1 0 1 2 3 y = 2 2 6 10 14 OR Using $y = mx + c$, gradient = 4, y intercept = 2	Line from (1, 2) to (3,14)	3	**(Table of values)** M1 for at least 2 correct attempts to find points by substituting values of x. M1 ft for plotting at least 2 of their points (any points plotted from their table must be correct) A1 for correct line between 1 and 3 **(No table of values)** M2 for at least 2 correct points (and no incorrect points) plotted OR line segment of $y = 4x + 2$ drawn (ignore any additional incorrect segments) (M1 for at least 3 correct points with no more than 2 incorrect points) A1 for correct line between -1 and 3 **(Use of $y = mx + c$)** M2 for at least 2 correct points (and no incorrect points) plotted OR line segment of $y = 4x + 2$ drawn (ignore any additional incorrect segments) (M1 for line drawn with gradient 4 OR line drawn with a y intercept of 2) A1 for correct line between 1 and 3
(b)(i)		$y = 4x + c$, $c \neq 2$	1	B1 Correct equation given.
(ii)		– 0.25	1	B1 Correct gradient given. Note – 0.25 could be written as $-\frac{1}{4}$ oe

5MB2H_01				
Question	**Working**	**Answer**	**Mark**	**Notes**
6	$6 \times 2 + 3 \times 2 = 12 + 6 = 18$ or $2 \times 5 + 4 \times 2 = 18$ $18 \div 2.5 = 7.2$ or $2.5 \times 7 = 17.5$ or $2.5 \times 8 = 20$ OR $6 \times 5 - 4 \times 3 = 18$ $18 \div 2.5 = 7.2$ or $2.5 \times 7 = 17.5$ or $2.5 \times 8 = 20$	8 packs	4	M2 for a complete method for finding the total area (can be implied by 18) (M1 for attempt to calculate at least one area, 6×2(=12) or 3×2 or 2×5(=10) or 4×2(=8) or 2×2 or 6×5 (=30) or 3×4(=12)) M1 for their area $\div 2.5$ or repeated addition of 2.5 to within 2.5 of their area C1 for clear communication that 8 full packs are required supported by their calculations, provided at least one of the first two method marks awarded SC B3 for 8 identified as the answer to the problem.
7	$\frac{3+7}{2}, \frac{8+5}{2}$	$(5, 6\frac{1}{2})$	2	M1 for either x or y coordinate correct or $\frac{3+7}{2}, \frac{8+5}{2}$, both seen but not correctly evaluated Al for $(5, 6\frac{1}{2})$ oe
8 (a) (b) (c)		p^8 h^5 x^6	3	B1 for p^8 B1 for h^5 B1 for x^6

5MB2H_01				
Question	**Working**	**Answer**	**Mark**	**Notes**
9	24 × 2 = 48 56 ÷ 8 × 5 = 35 or 24 × 2 = 48 48 ÷ 5 × 8 = 76.8	Monday	4	M1 for a correct method to obtain two comparable time periods or speeds. eg 24 × 2 (= 48 miles), 56 ÷ 2 (= 28 km) M1 for a correct method to convert between miles and km or mph and km/h eg –, – – – A1 for two correct values (using same units) for comparison C1 for day identified from a clear attempt to use comparable time periods and speeds/distances Note: day alone scores zero
10 (a)		$4n + 2$	2	B2 for oe (B1 for $4n+k$ where $k \neq 2$)
(b)	$2 \times 3^2 - 4$ 18 - 4	14	2	M1 for $2 \times 3^2 - 4$ A1 cao
11	360 ÷ 10 36 ÷ 2	18°	4	M1 for correct method to find the size of an exterior angle eg 360 ÷ 10 A1 for 36 M1 for '36' ÷ 2 or (180 – (180 – 36)) ÷ 2 A1 cao OR M1 for correct method to find the size of an interior angle eg(180 × 8) ÷ 10 A1 144 M1 (180 '144') ÷ 2 A1 cao

5MB2H_01				
Question	**Working**	**Answer**	**Mark**	**Notes**
12	$n^2 - 2n + 1 + n^2 + n^2 + 2n + 1 \equiv 3n^2 + 2$	Proof	2	M1 for correct expansion of either $(n - 1)^2$ or $(n + 1)^2$ or $n^2 - 2n + 1$ or $n^2 + 2n + 1$ seen A1 for $3n^2 + 2$ from correct working
13	$42 \div 2 = 21$ $180 - 90 - 21 = 69$ $69 \times 2 = 138$	138°	3	M1 for 90 seen. A1 for 138 (accept 222) C1 for The tangent to a circle is perpendicular (90°) to the radius (diameter) and Angles in a triangle add up to 180° **OR** The tangent to a circle is perpendicular (90°) to the radius (diameter) and Angles in a quadrilateral (4 sided shape) add up to 360°
14 (a)	$5\sqrt{9\times3}$	$15\sqrt{3}$	2	M1 for sight of $\sqrt{9\times3}$ or $\sqrt{9}\ \sqrt{3}$ or $3\sqrt{3}$ A1 for $15\sqrt{3}$ (accept $n = 15$)
(b)		$7\sqrt{3}$	2	M1 for $\frac{21\sqrt{3}}{\sqrt{3}\sqrt{3}}$ A1 for $7\sqrt{3}$ (accept $\frac{21\sqrt{3}}{3}$)

PMT

5MB2H_01				
Question	**Working**	**Answer**	**Mark**	**Notes**
15 (a)		3	1	B1 cao
(b)		$\frac{1}{5}$	2	M1 $\frac{1}{25^{\frac{1}{2}}}$ or $\sqrt{25} = 5$ A1 $\frac{1}{5}$ or 5^{-1} or 0.2
16 (a)	$\frac{2(y-6)}{(y-6)(y-2)}$	$\frac{2}{y-2}$	3	M1 $2(y-6)$ M1 $(y-6)(y-2)$ A1 cao
(b)	$\frac{3(x+5)-(x-4)}{(x-4)(x+5)}$	$\frac{3(x+5)-(x-4)}{(x-4)(x+5)}$	2	M1 for $3(x+5)$ or $3x+15$ or $3x+5$ or $-(x-4)$ or $-x+4$ or $-x-4$ or $(x-4)(x+5)$ A1 for $\frac{3(x+5)-(x-4)}{(x-4)(x+5)}$ oe

PMT

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481
Email publication.orders@edexcel.com

Order Code UG028428 June 2011

For more information on Edexcel qualifications, please visit
www.edexcel.com/quals

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual
Llywodraeth Cynulliad Cymru
Welsh Assembly Government
Rewarding Learning

PMT

edexcel

Mark Scheme (Results)

Summer 2012

GCSE Mathematics (Linear) 1MA0
Foundation (Non-Calculator) Paper 1F

PMT

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at www.edexcel.com.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2012
Publications Code UG032619
All the material in this publication is copyright
© Pearson Education Ltd 2012

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labeling conventions.

ii) *select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect canceling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.



12 Parts of questions
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers
Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

Guidance on the use of codes within this mark scheme
M1 – method mark A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working


1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
1	(a)		380	1	B1 cao
	(b)		6.2	1	B1 cao
	(c)		Arrow at 34	1	B1 cao
2	(a)		8	1	B1 for 8 ± 0.2
	(b)		35	1	B1 for 35 ± 2°
	(c)		Circle drawn	1	B1 for all parts within ± 2mm, (use overlay)
3	(a)		4, 7, 4, 3, 2	2	M1 for at least 3 correct tallies **or** at least 3 correct frequencies A1 for all frequencies correct
	(b)		7	1	B1 for 7 **or** ft from frequencies in (a) or tallies if no frequencies
	(c)		Diagram drawn	3	M1 for bar chart or other suitable chart with at least 3 correct heights for their scale (can f.t.) A1 for all 5 bars correctly labelled and vertical axis correctly scaled A1 for fully correct **or** ft frequencies in (a) **OR** M1 for pictogram with at least 3 correct rows (can f.t.) A1 for correct labels on all 5 rows and correctly key A1 for fully correct **or** ft frequencies in (a) **OR** M1 for pie chart with at least 3 correct sectors ± 2° (can f.t.) A1 for all 5 sectors correctly labelled A1 for fully correct **or** ft frequencies in (a)

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
4		£1.18 + 94p = £2.12 £5 – £2.12 – 30p = £2.58 £2.58 ÷ 2 =	1.29	3	M2 for (5 – 1.18 – 0.94 – 0.30) ÷ 2 oe **or** digits 129 (M1 for 1.18 + 0.94 **or** 2.12 seen **or** 1.18 + 0.94 + 0.30 oe **or** 2.42 seen **or** 5 – 1.18 – 0.94 oe **or** 2.88 seen **or** (5 – 1.18 – 0.94) ÷ 2 **or** 1.44 seen **or** 5 – 1.18 – 0.94 – 0.30 oe **or** 2.58 seen) A1 cao **NOTE**: Accept working in £ or pence
5	(a)(i)		(2, 3)	2	B1 cao
	(ii)		(–3, 1)		B1 cao
	(b)		Point plotted at (3, –4)	1	B1 cao
6	(a)		– 5	1	B1 cao
	(b)		6	1	B1 for 6 or – 6
	(c)		3	1	B1 cao
7			(P, B), (P, S), (P, L) (M, B), (M, S), (M, L) (H, B), (H, S), (H, L)	2	M1 for any 3 combinations with no incorrect combinations A1 for all 9 combinations with no duplicates or extras
8	(a)		Walk	1	B1 cao
	(b)	24 ÷ 4 =	6	2	M1 for 24 ÷ 4 oe or ¼ oe seen A1 cao

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
9	(a)		Isosceles triangle	1	B1 for isosceles triangle
	(b)		Rectangle with area 12 cm^2	2	M1 for rectangle drawn A1 cao
10	(a)		A marked at 0	1	B1 for A marked at 0 (within overlay)
	(b)		B marked at 1/4	1	B1 for B marked at 1/4 (within overlay)
11	(a)		9	1	B1 cao
	(b)		33	2	M1 for 5 × 5 **or** 25 seen in the working **or** 2 × 2 × 2 **or** 8 seen in the working A1 cao
12	(a)		20	2	M1 3 × 3 × 3 oe seen or drawn or 27 seen or use of 3 layers A1 cao
	(b)			2	B2 for correct view (B1 for  or )
13	(a)(i)		07 29	2	B1 for 07 29
	(ii)		36		B1 for 36 or ft difference between (i) and 06 53
	(b)		07 51	1	B1 cao
	(c)		09 55	1	B1 for 09 55 or 9 55 or five to ten

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
14		2 + 8 + 2 + 8 = 20 20 ÷ 4 =	5	4	M2 for 2 + 8 + 2 + 8 oe **or** 20 seen **or** (2 + 8) ÷ 2 oe (M1 for the sum of 3 sides of the rectangle) M1 (dep) for the sum of 3 or 4 sides of the rectangle ÷ 4 **or** an attempt to evaluate (2 + 8) ÷ 2 oe to get the length of one side A1 cao **SC**: B1 for an answer of 4 coming from $\sqrt{2\times 8}$ oe
15	(a)		4	1	B1 cao
	(b)	9.5 – 4.75 = **OR** 9.5 ÷ 2 =	4.75	2	M1 for 9.5 – 4.75 **or** 9.5 ÷ 2 **or** 4.75 – 9.5 A1 cao
	(c)		6	1	B1 cao
	(d)	12 × 4 =	48	2	M1 for × 4 seen **or** identifying + 0.5 for every 2 inches **or** 12 + 12 + 12 + 12 oe **or** build up method eg 12, 24, 36, 48 allow one error A1 cao
16	(a)		trapezium	1	B1 for trapezium or isosceles trapezium
	(b)			2	B2 for correct tessellation (at least 5 more shapes) (B1 for at least 4 shapes (including initial shape) correctly tessellating)

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
17*		S: 35 ÷ 100 × 40 = 14 W: 40 ÷ 8 × 3 = 15 **OR** D: 16 ÷ 40 (× 100) = 0.4 (40%) W: 3 ÷ 8 (× 100) = 0.375 (37.5%) **OR** D: $\frac{16}{40} = \frac{80}{200}$ S: $\frac{35}{100} = \frac{70}{200}$ W: $\frac{3}{8} = \frac{75}{200}$	Debbie and correct calculations	4	**Compares Marks out of 40 or fractions with denominator of 40** M1 for 35 ÷ 100 × 40 oe **or** 14 seen (or 14/40 seen) M1 for 40 ÷ 8 × 3 **or** 15 seen (or 15/40 seen) A1 for 14 and 15 **or** $\frac{14}{40}$ and $\frac{15}{40}$ C1 (dep on M1) for correct conclusion for their working QWC with 3 comparable marks: Decision and justification should be clear with working clearly presented and attributable. **OR Decimals (or Percentages)** M1 for 16 ÷ 40 (× 100) oe **or** 0.4 (or 40) seen M1 for 3 ÷ 8 (× 100) oe **or** 0.375 (or 37.5) seen A1 for 0.4 and 0.375 (**or** 40 and 37.5) C1 (dep on M1) for correct conclusion for their working QWC: with 3 comparable decimals (or percentages: Decision and justification should be clear with working clearly presented and attributable. **OR Compares Fractions with denominator other than 40** M1 for attempt to convert all to fractions with a common denominator other than 40 M1 for at least 1 correct A1 for $\frac{80}{200}$ and $\frac{70}{200}$ and $\frac{75}{200}$ oe C1 (dep on M1) for correct conclusion for their working QWC with 3 comparable fractions: Decision and justification should be clear with working clearly presented and attributable.

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
18*	(a)		10	1	B1 cao
	(b)	 **Miles** 0 10 20 30 40 50 **Ed** 0 15 30 45 60 75 **Bill** 10 20 30 40 50 60	Ed is cheaper up to 20 miles, Bill is cheaper for more than 20 miles	3	M1 for correct line for Ed intersecting at (20,30) ±1 sq tolerance **or** $10 + x = 1.5x$ oe C2 (dep on M1) for a correct full statement ft from graph eg. Ed cheaper up to 20 miles **and** Bill cheaper for more than 20 miles (C1 (dep on M1) for a correct conclusion ft from graph eg. cheaper at 10 miles with Ed ; eg. cheaper at 50 miles with Bill eg. same cost at 20 miles; eg for £5 go further with Bill **or** A general statement covering short **and** long distances eg. Ed is cheaper for shorter distances **and** Bill is cheaper for long distances) **OR** (continued on next page)

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
18 (contd)					**OR** M1 for correct method to work out Ed's delivery cost for at least 2 values of n miles where $0 < n \leq 50$ **or** for correct method to work out Ed and Bill's delivery cost for n miles where $0 < n \leq 50$ C2 (dep on M1) for 20 miles linked with £30 for Ed and Bill with correct full statement eg. Ed cheaper up to 20 miles **and** Bill cheaper for more than 20 miles (C1 (dep on M1) for a correct conclusion eg. cheaper at 10 miles with Ed; eg. cheaper at 50 miles with Bill eg. same cost at 20 miles; eg for £5 go further with Bill **or** A general statement covering short **and** long distances eg. Ed is cheaper for shorter distances **and** Bill is cheaper for long distances) SC: B1 for correct full statement seen with no working eg. Ed cheaper up to 20 miles **and** Bill cheaper for more than 20 miles QWC Decision and justification should be clear with working clearly presented and attributable

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
19		1,96 × 2.25 = 4.41 **OR** 4.23 ÷ 9 = 0.47 1.96 ÷ 4 = 0.49 **OR** 4.23 × 4 = 16.92 1.96 × 9 = 17.64 **OR** 4.23 ÷ 9 = 0.47 0.47 × 4 = 1.88 **OR** 1.96 ÷ 4 = 0.49 0.49 × 9 = 4.41 **OR** 9 ÷ 4.23 = 2.12 4 ÷ 1.96 = 2.04	Pack of 9	3	M2 for a fully correct method to enable a conclusion eg 1.96 × 2¼ **OR** M1 for 4.23 ÷ 9 or 423 ÷ 9 **or** 0.47 seen **or** 47 seen M1 for 1.96 ÷ 4 or 196 ÷ 4 or 0.49 seen or 49 seen **OR** M1 for 4.23 × 4 **or** 423 × 4 **or** 16.92 seen **or** 1692 seen M1 for 1.96 × 9 **or** 196 × 9 **or** 17.64 seen **or** 1764 seen **OR** M1 for 4.23 ÷ 9 **or** 423 ÷ 9 **or** 0.47 seen **or** 47 seen M1 for 0.47 × 4 **or** 47 × 4 **or** 1.88 seen **or** 188 seen **OR** M1 for 1.96 ÷ 4 **or** 196 ÷ 4 **or** 0.49 seen **or** 49 seen M1 for 0.49 × 9 **or** 49 × 9 **or** 4.41 seen **or** 441 seen **OR** M1 for 9 ÷ 4.23 **or** 2.12(...) seen **or** 2.13 seen M1 for 4 ÷ 1.96 **or** 2.04(...) seen A1 for Pack of 9 and fully correct calculations **NOTE:** B0 for an answer of 9 not supported by working.
20	(a)		6	1	B1 cao
	(b)		44	1	B1 cao
	(c)		31	2	M1 for 60 – 29 **or** 29 – 60 **or** any correct method that is attempting to find the difference between 29 and 60 (allow 1 arithmetic error) A1 cao

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
21*		Angle *DBC* = (180 – 50) ÷ 2 Base angles of isosceles triangle are equal Angle *ABD* = 180 – 65 Angles on a straight line add up to 180 *x* = 180 – 20 – 115 Angles in a triangle add up to 180 **OR** Angle *DBC* = (180 – 50) ÷ 2 Base angles of isosceles triangle are equal *x* = 65 – 20 Exterior angle of triangle is equal to sum of interior opposite angles **OR** Angle *DCB* = (180 – 50) ÷ 2 Base angles of isosceles triangle are equal *x* = 180 – 50 – 20 – 65 Angles in a triangle add up to 180	45 with reasons	4	M1 for (180 – 50) ÷ 2 oe **or** 65 seen M1 for 180 – 20 – (180 – "65") **or** "65" – 20 or 180 – 50 – 20 –'65' oe C2 for *x* identified as 45 with full reasons QWC: Reasons clearly laid out with correct geometrical language used (C1 (dep on M1) for one reason QWC: Reasons clearly laid out with correct geometrical language used) NOTE: *x* = 45 with no working or without any correct angles marked on the diagram cannot score.

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
22	(a)	360 ÷ 60 = 6 300 ÷ 60 = 5 6 × 5 =	Yes and 30	3	M1 for dividing side of patio by side of paving slab eg 360 ÷ 60 **or** 300 ÷ 60 **or** 3.6 ÷ 0.6 **or** 3 ÷ 0.6 **or** 6 and 5 seen **or** 6 divisions seen on length of diagram **or** 5 divisions seen on width of diagram M1 for correct method to find number of paving slabs eg (360 ÷ 60) × (300 ÷ 60) oe **or** 6 × 5 **or** 30 squares seen on diagram (units may not be consistent) A1 for Yes **and** 30 (or 2 extra) with correct calculations **OR** M1 for correct method to find area of patio **or** paving slab eg 360 × 300 **or** 108000 seen **or** 60 × 60 **or** 3600 seen **or** 3.6 × 3 **or** 10.8 seen **or** 0.6 × 0.6 **or** 0.36 seen M1 for dividing area of patio by area of a paving slab eg. (3.6 × 3) ÷ (0.6 × 0.6) oe (units may not be consistent) A1 for Yes **and** 30 (or 2 extra) with correct calculations **OR** M1 for method to find area of patio and area of 32 slabs eg. 60 × 60 × 32 **or** 360 × 300 M1 for method to find both areas eg. 60 × 60 × 32 **and** 360 × 300 (units may not be consistent) A1 for Yes **and** 115200 **and** 108000 **OR** Yes **and** 11.52 **and** 10.8 **NB :** Throughout the question, candidates could be working in metres or centimetres

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
22	(b)	1726 25890 27616 **8 6 3** **2** 2 4 1 8 9 **3** **7** 1 6 1 2 6 **2** 6 1 6 800 60 3 30 24000 1800 90 2 1600 120 6 24000 + 1800 + 90 + 1600 + 120 + 6 = 27616	276.16	3	M1 for complete correct method with relative place value correct. Condone 1 multiplication error, addition not necessary. **OR** M1 for a complete grid. Condone 1 multiplication error, addition not necessary. **OR** M1 for sight of a complete partitioning method, condone 1 multiplication error. Final addition not necessary. A1 for digits 27616 A1 ft (dep on M1) for correct placement of decimal point after addition (of appropriate values) (SC: B1 for attempting to add 32 lots of 8.63)
23	(a)		30	2	M1 for 25 ÷ 10 **or** 2.5 seen **or** 10 ÷ 25 **or** 0.4 seen **or** 12+12+6 oe **or** a complete method eg. 25×12÷10 oe A1 cao
	(b)	1000 ÷ 200 × 12	60	2	M1 for 500 ÷ 50 **or** 1000 ÷ 200 **or** 500 ÷ 10 **or** correct scale factor clearly linked with one ingredient eg 10 with sugar or 5 with butter or flour or 50 with milk **or** an answer of 120 or 600 A1 cao

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
24		Acton after 24, 48, 72, 96, .. Barton after 20, 40, 60, 80, .. LCM of 20 and 24 is 120 9:00 am + 120 minutes **OR** Acton after 24, 48, 1h 12 min... Barton after 20, 40, 1 h LCM is 2 hours 9:00 am + 2 hours **OR** Times from 9:00 am when each service leaves the bus station Acton at 9:24, 9:48, 10:12.. Barton at 9:20, 9:40, 10:00.. **OR** 20 = 2 × 2 × 5 24 = 2 × 2 × 2 × 3 2 × 2 × 2 × 3 × 5 = 120	11:00 am	3	M1 for listing multiples of 20 and 24 with at least 3 numbers in each list ; multiples could be given in minutes or in hours and minutes (condone one addition error in total in first 3 numbers in lists) A1 identify 120 (mins) **or** 2 (hours) as LCM A1 for 11:00 (am) **or** 11(am) **or** 11 o'clock **OR** M1 for listing times after 9am when each bus leaves the bus station, with at least 3 times in each list (condone one addition error in total in first 3 times after 9am in lists) A1 for correct times in each list up to and including 11:00 A1 for 11:00 (am) **or** 11(am) **or** 11 o'clock **OR** M1 for correct method to write 20 and 24 in terms of their prime factors 2, 2, 5 and 2, 2, 2, 3 (condone one error) A1 identify 120 as LCM A1 for 11:00 (am) **or** 11(am) **or** 11 o'clock

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
25	(a)		$6y - 15$	1	B1 cao
	(b)		$4x(2x + y)$	2	B2 cao (B1 for $x(8x + 4y)$ **or** $2x(4x + 2y)$ **or** $4(2x^2 + xy)$ **or** $4x(ax + by)$ where a, b are positive integers **or** $ax(2x + y)$ where a is a positive integer **or** $4x(2x \quad y)$)
	(c)	$10t = gh$ $h = \frac{10t}{g}$	$\frac{10t}{g}$	2	M1 for clear intention to multiply both sides of the equation by 10 (eg. ×10 seen on both sides of equation) **or** clear intention to divide both sides of the equation by g (eg. ÷ g seen on both sides of equation) **or** $10t = gh$ **or** $\frac{t}{g} = \frac{h}{10}$ **or** fully correct reverse flow diagram eg. ← ×10 ← ÷g ← A1 for $\frac{10t}{g}$ oe

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
26	(a)	$2 \times 5 \times 2 = 20$ $300 \div 20 =$	15	3	M2 for $300 \div (2 \times 5 \times 2)$ oe (M1 for $2 \times 5 \times 2$ **or** 20 seen **or** $300 \div (2 \times 5)$ **or** 30 seen A1 cao
	(b)	$c = \frac{30 \times 40}{150}$ =	8	2	M1 for $\frac{30 \times 40}{150}$ **or** 1200 seen A1 cao
27		$3x - 15 = 2x + 24$ $x = 39$ **OR** $2x + 3x - 15 + 2x + 2x + 24 = 360$ $9x + 9 = 360$ $9x = 351$ $x = 39$ **OR** $2x + 2x + 24 = 180$ $4x + 24 = 180$ $4x = 156$ $x = 39$ **OR** $2x + 3x - 15 = 180$ $5x - 15 = 180$ $5x = 195$ $x = 39$	39	3	M1 for forming an appropriate equation eg $3x - 15 = 2x + 24$ **or** $2x + 3x - 15 + 2x + 2x + 24 = 360$ oe **or** $2x + 2x + 24 = 180$ oe **or** $2x + 3x - 15 = 180$ oe **or** $2x + 3x - 15 = 2x + 2x + 24$ M1 (dep) for correct operation(s) to isolate x and non-x terms in an equation to get $ax = b$ A1 cao **OR** M2 for $\frac{351}{9}$ **or** $\frac{195}{5}$ **or** $\frac{156}{4}$ oe A1 cao

PMT

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481
Email publication.orders@edexcel.com

Order Code UG032619 Summer 2012

For more information on Edexcel qualifications, please visit our website
www.edexcel.com

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual
Llywodraeth Cynulliad Cymru
Welsh Assembly Government
Rewarding Learning

PMT

edexcel

Mark Scheme (Results)

November 2015

Pearson Edexcel GCSE
In Mathematics B (2MB01)
Higher (Non-Calculator) Unit 2

PMT

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

November 2015
Publications Code UG042910
All the material in this publication is copyright
© Pearson Education Ltd 2015

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Note that in some cases a correct answer alone will not score marks unless supported by working; these situations are made clear in the mark scheme. Examiners should be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will award marks for the quality of written communication (QWC).
The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labelling conventions.

ii*) select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Partial answers shown (usually indicated in the ms by brackets) can be awarded the method mark associated with it (implied).
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks; transcription errors may also gain some credit. Send any such responses to review for the Team Leader to consider.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded (embedded answers).

12 Parts of questions
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers
Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

14 The detailed notes in the mark scheme, and in practice/training material for examiners, should be taken as precedents over the above notes.

Guidance on the use of codes within this mark scheme
M1 – method mark for appropriate method in the context of the question A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
1.			6 525 900 450	3	M1 for × 6 ÷ 4 oe or $\frac{6}{4}$ or 1.5 or 1 correct answer A2 for all correct (A1 for 2 or 3 correct)
2.			$8n + 24$	3	M1 for generating at least 3 terms of sequence 32, 40, 48 or sequence 4, 5, 6 M1 for $8n \pm k$ or $8(n \pm m)$ where $k \neq 24$, $m \neq 3$ A1 for $8n + 24$ or $8(n + 3)$ oe
3.			105 cm^3	4	M1 for writing factor pairs of 35, 21 or 15 M1 for "3" × "5" × "7" using their lengths dep on one at least one correct factor pair A1 ft for 105 (dep on one correct factor pair) B1 for cm^3

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
4.		$y = \frac{1}{2}x + 3$ x : −2, −1, 0, 1, 2, 3, 4 y : 2, 2.5, 3, 3.5, 4, 4.5, 5	Correct line from (−2, 2) to (4, 5)	3	**(Table of values / calculation of values)** M1 for at least 2 correct attempts to find points by substituting values of x. M1 ft for plotting at least 2 of their points (any points plotted from their table must be correctly plotted) A1 for correct line between $x = -2$ and $x = 4$ **(No table of values)** M1 for at least 2 correct points with no more than 2 incorrect points plotted M1 for at least 2 correct points (and no incorrect points) plotted **OR** line segment of $y = \frac{1}{2}x + 3$ drawn A1 for correct line between $x = -2$ and $x = 4$ **(Use of $y = mx + c$)** M1 for line drawn with gradient of $\frac{1}{2}$ **OR** line drawn with a y intercept of 3 M1 for line drawn with gradient of $\frac{1}{2}$ **AND** line drawn with a y intercept of 3 A1 for correct line between $x = -2$ and $x = 4$ **SC** : B2 for correct line from $x = 0$ to $x = 4$

PMT

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
5.			68	5	B1 for correct conversion of units M1 for 250 ÷ 25 (= 10) or 300 ÷ 20 (= 15) M1 for "10" × "15" M1 (dep on M2) for number of tiles ÷ 20, round up, × 8.5(0) A1 for 68 OR B1 for correct conversion of units M1 for 250 ÷ 20 (= 12.5) or 300 ÷ 25 (= 12) M1 for "12.5" × "12" M1 (dep on M2) for number of tiles ÷ 20, round up, × 8.5(0) A1 for 68 OR B1 for correct conversion of units M1 for 3 × 2.5 (= 7 .5) or 20 × 25 (= 500) M1 for "3 × 2.5" ÷ "0.2 × 0.25" or "300 × 250" ÷ "20 × 25" M1 (dep on M2) for number of tiles ÷ 20, round up, × 8.5(0) A1 for 68
*6		1, 4, 7, 10, 13 8, 6, 4, 2, 0	Explanation	2	M1 for listing at least 3 terms of both sequences C1 for Yes and explanation from fully correct working that 4 is in both sequences; numbers in A are increasing; numbers in B are decreasing

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
7			7.21 (am)	3	M1 for listing multiples 9,18,27,36 and 12,24,36 (condone 1 arithmetic error) or method to find LCM M1 for identifying 36 as LCM A1 cao OR M1 for listing times 6.54, 7.03, 7.12, 7.21 or for listing times 6.57, 7.09, 7.21 (condone one arithmetic error) M1 for listing times 6.54, 7.03, 7.12, 7.21 **and** 6.57, 7.09, 7.21 (condone one arithmetic error) A1 cao
8	(a)		$4(3e+1)$	1	B1 cao
	(b)		$15c - 10d$	1	B1 cao
	(c)		$28a^4b^3$	2	B2 for $28a^4b^3$ (B1 for two of 28, a^4 , b^3 as a product)
	(d)		$(x + 7)(x - 7)$	1	B1 cao
	(e)	$2y^2 - 6y + 7y - 21$	$2y^2 + y - 21$	2	M1 for 3 out of no more than 4 terms correct with correct signs or the 4 terms $2y^2$, $6y$, $7y$ and 21 seen, ignoring signs A1 cao

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
9		360 – 64 140 + 40 + 296 + 40 = 516 540 − 516 = 24	24°	4	M1 for 1 correct relevant angle calculation A1 for 24 cao C2 for all reasons (C1 (dep M1) for reason relating to parallel lines or parallelogram which is relevant to their chosen method) eg angles on a straight line add to 180° angles at a point add to 360° angles in a triangle add to 180° angles in a quadrilateral add to 360° alternate angles are equal corresponding angles are equal allied angles / co-interior angles add up to 180° opposite angles in a parallelogram are equal
10			$4\frac{8}{35}$	3	M1 for converting both fractions to get a common denominator of a multiple of 35 with at least one correctly converted. M1 (dep on M1) for $3+\frac{"28"}{35}+\frac{"15"}{35}(=3\frac{43}{35})$ oe A1 for $4\frac{8}{35}$ cao

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
11	(a)		7.8×10^{-3}	1	B1 cao
	(b)		6 710 000	1	B1 cao
	(c)	$-9,\ 0.9,\ 1,\ 3$	$-9,\ 0.9,\ 9^0,\ 9^{\frac{1}{2}}$	2	M1 for either $9^{\frac{1}{2}}$ or 9^0 evaluated correctly A1 for fully correct list from correct working, accept original numbers or evaluated. (SC: B1 for 1 error in position or a correct list in reverse order)
12.	(a)		$(0, 5, 5)$	1	B1 cao
	(b)		$(2\frac{1}{2}, 5, 2\frac{1}{2})$	1	B1 for $(2\frac{1}{2}, 5, 2\frac{1}{2})$ oe
13.			120	3	M1 for $80\% = 480$ or $\frac{480}{80} (= 6)$ or $\frac{480}{80} \times 100\ (= 600)$ oe M1 for $600 - 480$ or $\frac{480}{80} \times 20$ oe A1 cao
14.			$A = 9x^2 + 19x - 6$	4	B1 for one of $5x - 2$ or x found M1 for correct method to find area of one relevant rectangle. M1 for complete method to find whole area or simplified expression $9x^2 + 19x - 6$ or correct but not simplified formula A1 for correct, simplified formula $A = 9x^2 + 19x - 6$

5MB2H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
15.			$y = \frac{3}{2}x - \frac{5}{2}$	4	M1 for attempt to find gradient of AB M1 (dep) for attempt to find gradient of perpendicular line eg use of $-\frac{1}{m}$ M1 for substituting $x = 3, y = 2$ into $y =$ "m" $x + c$ A1 for complete correct equation $y = \frac{3}{2}x - \frac{5}{2}$ oe
16.			$2\sqrt{7}$	3	M1 for multiplying numerator and denominator by $\sqrt{7}$ M1 for correct method to expand $(4+\sqrt{2})(4-\sqrt{2})$ with 3 out of no more than 4 terms correct with correct signs or the 4 terms seen, ignoring signs A1 for $2\sqrt{7}$ (accept $\sqrt{28}$)
17.		$\frac{(x-5)(x+3)}{(x+3)(2x+1)}$	$\frac{x-5}{2x+1}$	3	M1 for correct factorisation of numerator M1 for correct factorisation of denominator A1 for $\frac{x-5}{2x+1}$

PMT

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom

Question 7

In general, both parts of this question were answered well. In part (a), p^7 was the modal incorrect answer offered. In part (b), a significant number of students used correct rules of indices to derive expressions of x^4 and/or y^2 but often these were not seen in a product or a fraction. It was also common to see $12 - 6 = 6$ instead of $12 \div 6 = 2$.

Question 19

The correct answer to this question was seen only very occasionally. Some students were able to gain credit for $9^{-\frac{1}{2}} = \frac{1}{3}$ or equivalent. Only a very few were able to write $27^{\frac{1}{4}}$ in any useful format.

Question 10

This question, assessing basic algebra, was not answered particularly well. In part (a) many students gave incorrect answers such as 0 or $\frac{1}{m^2}$ therefore effectively raising to the power 1.

Part (b) was answered slightly better with more students gaining the single mark on offer. However, many tried to expand before simplifying and therefore went wrong. Students should be reminded that the number of marks on offer give a good indication to the amount of work required.

In part (c) with there being two marks available we saw more students gaining some credit. Many students did not understand the 3rd law of indices so didn't score, but those who did typically got the last two terms correct, but struggled with the numerical term, often giving an answer of 3 or 9 rather than 27.

Question 1

This question provided a good start to the paper. Most students scored at least four of the five marks available and nearly all students gained some credit for their answers.

Part (a) of this question required students to use the index law for multiplication of powers of *n*. Nearly all students applied the law correctly. n^{15} was the most commonly seen incorrect answer.

Part (b) of the question was answered less well but nevertheless the great majority of students gained the two marks available. Examiners condoned answers of the form c^1d^3 and $c \times d^3$. It was not unusual for students who did not give a fully correct answer to get one mark for either c or d^3.

In part (c) students could usually find the critical value giving it as either 2.8 or $\frac{14}{5}$.

These students scored at least one mark. A significant number of students gave their final answer as 2.8 or as $x = 2.8$ instead of the correct $x > 2.8$

Question 1

The first question on the paper assessed some basic algebra skills. Part (a) assessed the third law of indices and was generally answered correctly. The most common wrong answer being to add the index numbers rather than multiply.

In part (b) students had to expand a pair of single brackets and simplify. Most students were able to gain at least one mark, for expanding one bracket correctly. Errors tended to come in the collection of terms, and in particular the x terms, often leading to $18x$.

Part (c) was a factorisation question looking for the student to bring two terms out of the expression. A good proportion of the students scored at least one mark for a suitable partial factorisation of the expression, commonly taking out the factor of $3x$ rather than $3x^2$. There was however, a pleasing proportion of students that gained both marks for the correct factorisation.

Question 14

A minority of candidates scored full marks for their responses to this question.
However, most candidates were able to gain some marks and the question was a good discriminator between candidates who could manipulate algebraic expressions with accuracy and those who made errors.
A fully simplified expression was asked for in answering part (a) of this question. It was disappointing to see so few candidates being completely successful and it was much more usual for candidates to gain just one of the two marks available e.g. for $3x^{20}y^{24}$. A common error was for candidates to add the indices 20 and 24 and give their answer in the form cxy^{44}.

Candidates usually scored well in part (b) which required the expansion of a product of three linear expressions to give a fully simplified cubic expression. Though there were some low attaining candidates who attempted to expand the product in one stage, working was usually shown in an organized way or a grid was used to show terms in the products. Errors were usually restricted to incorrect terms or difficulties in dealing with the signs when collecting terms together, rather than a flawed strategy. There were some candidates who omitted terms from their expansion. For candidates who did not give a fully correct answer, it was commonplace for them to earn 2 of the available marks in part (b).

Question15

Most candidates scored full marks for their answers to this question. The most common error seen was where candidates used 7, 13 and 5 in a sum, for example as 7 + 13 + 5, rather than in correct products. Candidates who gained one of the two marks usually did so for giving the correct product leading to 455 but could not complete the question successfully. Some candidates tried unsuccessfully to work backwards from the 555.

Question 11

Many students gained the first mark for simplifying either the numerator or the denominator with at least two of three terms correct in a product. At this stage the most common mistakes were $6x^{10}y^6$ or $36x^{25}y^9$ or ax^7y^5 instead of $36x^{10}y^6$ in the numerator and $12x^2y^4$ instead of $12x^3y^4$ in the denominator. A good proportion of students completed the algebra correctly and arrived at an answer of $3x^7y^2$. Some students made one error in the numerator or denominator and gained two marks for an answer of the form ax^by^c with two of a, b and c correct (of these the error was often an incorrect coefficient). A small number of students used a different approach and instead of simplifying the numerator and the denominator they chose to start by simplifying $6x^5y^3 \div 3x^2y^7$ and $6x^5y^3 \div 4xy^{-3}$. A common error was for students to introduce plus signs into the numerator and then attempt to expand the two brackets, leading to more than one product for the numerator.

Question 15

Part (a) was well answered with a high proportion of students gaining the mark. When the mark was not awarded it was usually as the student had expanded the brackets or for incorrectly simplifying the $\frac{a-3}{5}$. In part (b) most students were able to gain at least one of the available marks. Some were awarded one mark for making an error with the signs or for

giving two pairs of brackets which when expanded gave two out of three terms correct. Students should be encouraged to multiply out their brackets as a final check - there was little evidence of this taking place. A popular error that resulted in no marks being awarded was to factorise the first two terms and give $k(3k + 11) - 4$ as the answer. Some students went on to equate the expression to zero and solve the equation, which on this occasion was condoned and didn't result in loss of marks. It was surprising how many students used the quadratic formula to find the solutions when the expression was equal to zero, this often resulted in no marks being awarded as the students either thought that this was the requirement of the question or didn't know how to use these solutions to help them factorise. A surprising number gave a solution of 3(k-1/3)(k+4) which also scored 1 mark. Students should be aware that factors within brackets should be integer values. Part (c) was a good discriminator and many students were able to show that to divide the algebraic fractions, they needed to multiply the first fraction by the reciprocal of the second fraction. Fewer students realised the need to factorise the numerator and denominator of the first fraction and many spent time expanding the brackets for no gain and could not then simplify the complex expression generated. Of those that realised the need to factorise many struggled to factorise the $4-x^2$ often factorising this to $(x+2)(x-2)$ instead of $(2+x)(2-x)$ as sign errors were condoned in one bracket, these students were able to gain 2 of the 3 marks. Almost all of those who correctly factorised went on to simplify the fraction correctly and gain full marks.

Question 9

A well answered question with common errors being that learners either didn't simplify the 2×7 or added rather than multiplied them. There were some who gave their final answer as a sum and hence scored B0 and the retention of multiplication signs in a final answer scored a maximum of B1.

Question 11

In part (a), many responses demonstrated a secure understanding of index laws, with common approaches including applying the rule for multiplying powers with the same base (e.g. $3^n \times 3^{12} = 3^{14}$ or $3^n \times 3^{20} = 3^{22}$) or forming a linear equation such as $n + 20 - 8 = 14$. Clear and methodical working was often seen in more successful responses. Some responses included errors such as multiplying the bases while adding the powers (e.g. simplifying $3^{14} \times 3^8$ to 9^{22}) or incorrectly multiplying the indices (e.g. $14 \times 8 \div 20$), which led to inaccurate answers such as 5.6.

In part (b), most responses showed understanding of fractional indices and recognised the need to apply both cube root and square operations. Stronger responses often separated the numerical and algebraic components, for example: $(\sqrt{64})2=16$ and $(m12)23=m8$ leading to the final expression $16m^8$. Less successful responses tended to complete only one of the two operations or applied the correct process to only one part of the expression (either 64 or m^{12}), leaving the other unchanged. Partial simplifications such as 16 or m^8 were common and credited where appropriate.

Centres may wish to encourage step-by-step presentation of working, particularly when applying fractional powers. A secure understanding of how powers apply to both numbers and algebraic terms can help improve accuracy and clarity in future responses.