Chhetri Academy

chhetriacademy.com

GCSE MATHS

Algebra

Edexcel 1MA1 | Subtopic collection Topic 27

Iteration

Higher tier

Answers and examiner comments

31	2003-2023	Higher
question groups	available papers	tier

Using this booklet

1. Answer numbers match the companion question booklet.
2. Follow the original question number on marking tables.
3. Examiner comments are included where available.

Name _____ **Date** _____

Some entries use full source marking material in an appendix. Any unavailable mark schemes are identified beside the relevant answer.

Original question 4

The full source mark scheme begins on page 45. Use the original paper code, tier and question number shown above.

Original question 3

The full source mark scheme begins on page 99. Use the original paper code, tier and question number shown above.

9	(a)		8	1	B1 cao
	(b)		C	1	B1 for C or pyramid

14		3→ 33 4 → 72 3.1 → 35.9(91) 3.2 → 39.1(68) 3.3 → 42.5(37) 3.4 → 46.1(04) 3.5 → 49.8(75) 3.6 → 53.8(56) 3.7 → 58.0(53) 3.8 → 62.4(72) 3.9 → 67.1(19) 3.75 → 60.2(34375)	3.7	4	B2 for a trial between 3.7 and 3.8 inclusive (B1 for a trial between 3 and 4 inclusive) B1 for a different trial between 3.7 and 3.8 exclusive B1 (dep on at least one previous B1) for 3.7 NB Trials should be evaluated to at least 1dp truncated or rounded

1380 4H

1380/4H

Question	Working	Answer	Mark	Notes
8	x = 1 gives 11 x = 2 gives 28 x = 1.5, gives 18.(375) x = 1.6, gives 20.(096) x = 1.7, gives 21.(913) x = 1.8, gives 23.(832) x = 1.9, gives 25.(859) x = 1.85, gives 24.8(316..) x = 1.86, gives 25.(03..) x = 1.87, gives 25.2(3..) x = 1.88, gives 25.4(4..) x = 1.89, gives 25.6(5..)	1.9	4	B2 for a trial $1.8 \leq x \leq 1.9$ evaluated (B1 for a trial $1 \leq x \leq 2$ evaluated) B1 for a different trial $1.85 \leq x < 1.9$ evaluated B1 (dep on at least one previous B1) for 1.9 Accept trials correct to the nearest whole number (rounded or truncated) if the value of x is to 1dp but correct to 1dp (rounded or truncated) if the value of x is to 2dp. NB: no working scores no marks even if answer is correct.

Question		Working	Answer	Mark	Notes
12		x | $x^3 + 5x$ 3 | 42 3.4 | 56.(304 3.5 | 60.(375 3.6 | 64.(656 3.7 | 69.(153 3.8 | 73.(872 3.9 | 78.(819 4 | 84 3.65 | 66.8(77 3.66 | 67.3(27 3.67 | 67.7(80 3.68 | 68.2(36 3.69 | 68.6(93	3.7	4	B2 for a trial between 3 and 4 exclusive (B1 for a trial between 3 and 4 inclusive) B1 for a different trial of $3.65 \leq x < 3.7$ B1 (dep on at least one previous B1) for 3.7 NB Trials should be evaluated to at least 2 s.f truncated or rounded for values of x correct to 1 dp. Trials should be evaluated to at least 1 dp for values of x correct to 2 dp truncated or rounded. No working scores 0 marks

11		$x = 4$ gives 40 $x = 5$ gives 95 $x = 4.1$ gives 44.(321) $x = 4.2$ gives 48.(888) $x = 4.3$ gives 53.(707) $x = 4.4$ gives 58.(784) $x = 4.5$ gives 64.(125) $x = 4.6$ gives 69.(736) $x = 4.7$ gives 75.(623) $x = 4.8$ gives 81.(792) $x = 4.9$ gives 88.(249) $x = 4.61$ gives 70.3(12..) $x = 4.62$ gives 70.8(91..) $x = 4.63$ gives 71.4(72..) $x = 4.64$ gives 72.0(57..) $x = 4.65$ gives 72.6(44..)	4.6	4	B2 for a trial $4.6 \leq x \leq 4.7$ evaluated (B1 for a trial $4 \leq x \leq 5$ evaluated) B1 for a different trial $4.6 < x \leq 4.65$ evaluated B1 (dep on at least one previous B1) for 4.6 Accept trials correct to the nearest whole number (rounded or truncated) if the value of x is to 1 dp but correct to 1dp (rounded or truncated) if the value of x is to 2 dp. (Accept 72 for $x = 4.64$) NB : no working scores no marks even if the answer is correct.

1MA0_2H

Question		Working	Answer	Mark	Notes
7		x=1 gives 11 x=2 gives 28 x=1.5 gives 18.(375) x=1.6 gives 20.(096) x=1.7 gives 21.(913..) x=1.8 gives 23.(832..) x=1.9 gives 25.(859..) x=1.75 gives 22.8(59..) x=1.76 gives 23.0(51..) x=1.77 gives 23.2(45..)	1.8	4	B2 for a trial $1.7 \leq x \leq 1.8$ evaluated (B1 for a trial $1 \leq x \leq 2$ evaluated) B1 for a different trial $1.7 < x < 1.8$ evaluated B1 (dependent on at least one previous B1) for 1.8 Accept trials correct to the nearest whole number (rounded or truncated) if the value of x is to 1dp but correct to 1dp (rounded or truncated) if the value of x is to 2dp. NB: no working scores no marks even if answer is correct.

<table>
<tr><th colspan="2">Question</th><th>Working</th><th>Answer</th><th>Mark</th><th>Notes</th></tr>
<tr><td>11</td><td>(a)</td><td></td><td>show</td><td>2</td><td>M1 for $x\times x\times x$ or $2\times5\times x$ or vol of cube = x^3 or vol cuboid = $10x$
A1 correct completion leading to
$x^3-10x=100$</td></tr>
<tr><td></td><td>(b)</td><td><table>
<tr><td>$x = 1$</td><td>−9</td></tr>
<tr><td>$x = 2$</td><td>−2</td></tr>
<tr><td>$x = 3$</td><td>−3</td></tr>
<tr><td>$x = 4$</td><td>24</td></tr>
<tr><td>$x = 5$</td><td>75</td></tr>
<tr><td>$x = 6$</td><td>156</td></tr>
<tr><td>$x = 10$</td><td>900</td></tr>
<tr><td>$x = 5.1$</td><td>81.(651)</td></tr>
<tr><td>$x = 5.2$</td><td>88.(608)</td></tr>
<tr><td>$x = 5.3$</td><td>95.(877)</td></tr>
<tr><td>$x = 5.4$</td><td>103.(464)</td></tr>
<tr><td>$x = 5.5$</td><td>111.(375)</td></tr>
<tr><td>$x = 5.6$</td><td>119.(616)</td></tr>
<tr><td>$x = 5.7$</td><td>128.(193)</td></tr>
<tr><td>$x = 5.8$</td><td>137.(112)</td></tr>
<tr><td>$x = 5.9$</td><td>146.(379)</td></tr>
<tr><td>$x = 5.35$</td><td>99.6(30375)</td></tr>
<tr><td>$x = 5.36$</td><td>100.3(90656)</td></tr>
<tr><td>$x = 5.355$</td><td>100.0(101139)</td></tr>
</table></td><td>5.4</td><td>4</td><td>B2 for a trial $5 \le x \le 6$ evaluated correctly
(B1 for any two trials evaluated correctly for positive values of x)
B1 for a different trial $5.3 < x < 5.4$ evaluated correctly
B1 (dep on at least one previous B1) for 5.4

Accept trials correct to the nearest whole number (rounded or truncated) if the value of x is to 1 d.p., but correct to 1 d.p. (rounded or truncated) if the value of x is to 2 or more d.p.

NB. Allow 100 for a trial of $x = 5.355$</td></tr>
</table>

1MA0_2H

11		x | $x^3 - x$ 3.1 | 26.(691) 3.2 | 29.(568) 3.3 | 32.(637) 3.4 | 35.(904) 3.5 | 39.(375) 3.6 | 43.(056) 3.7 | 46.(953) 3.8 | 51.(072) 3.9 | 55.(419) 3.25 | 31.(078) 3.26 | 31.3(85) 3.27 | 31.6(95) 3.28 | 32.0(07) 3.29 | 32.3(21)	3.3	4	B2 for trial $3.2 \leq x \leq 3.3$ (B1 for trial $3 < x < 4$) B1 for different trial $3.25 \leq x < 3.3$ B1 cao (dep on at least one previous B1) Accept trials correct to the nearest whole number (rounded or truncated) if the value of x is to 1dp but correct to 1dp (rounded or truncated) if the value of x is to 2dp

5MB3H_01

Question		Working	Answer	Mark	Notes
8		x | $x^3 - 3x$ 2 | 2 2.1 | 2.(961) 2.2 | 4.(048) 2.3 | 5.(267) 2.4 | 6.(624) 2.5 | 8.(125) 2.6 | 9.(776) 2.7 | 11.(583) 2.8 | 13.(552) 2.9 | 15.6(89) 3 | 18 2.85 | 14.5(99...) 2.86 | 14.8(13...) 2.87 | 15.0(29...) 2.88 | 15.2(47...) 2.89 | 15.4(67...)	2.9	4	B2 for a trial $2.8 \leq x \leq 2.9$ evaluated correctly (B1 for a trial evaluated correctly for $2 \leq x \leq 3$) B1 for a different trial evaluated correctly for $2.85 \leq x < 2.9$ B1 (dep on at least one previous B1) for 2.9 NB For trials where x has one decimal place: $x \leq 2.6$ trials must be evaluated to at least 1 sf truncated or rounded $2.6 < x < 2.85$ trials must be evaluated to at least 2 sf truncated or rounded $2.85 \leq x \leq 2.9$ trials must be evaluated to at least 3 sf truncated or rounded NB. Accept 15 or 15.0 for trial at x =2.87 No working scores 0 marks. **If candidate is clearly working with $x^3 - 3x - 15 = 0$ then use same scheme as above but subtract 15 from all evaluated values in the table**

1MA0_2H

Original question 10

The full source mark scheme begins on page 160. Use the original paper code, tier and question number shown above.

Original question 10

The full source mark scheme begins on page 175. Use the original paper code, tier and question number shown above.

<table>
<tr><th colspan="2">Question</th><th>Working</th><th>Answer</th><th>Mark</th><th>Notes</th></tr>
<tr><td>11</td><td></td><td>
<table>
<tr><td>4</td><td>40</td></tr>
<tr><td>4.1</td><td>44.(321)</td></tr>
<tr><td>4.2</td><td>48.(888)</td></tr>
<tr><td>4.3</td><td>53.(707)</td></tr>
<tr><td>4.4</td><td>58.(784)</td></tr>
<tr><td>4.5</td><td>64.(125)</td></tr>
<tr><td>4.6</td><td>69.(736)</td></tr>
<tr><td>4.7</td><td>75.(623)</td></tr>
<tr><td>4.8</td><td>81.(792)</td></tr>
<tr><td>4.9</td><td>88.(249)</td></tr>
<tr><td>5</td><td>95</td></tr>
</table>
<table>
<tr><td>4.81</td><td>82.4(2464)</td></tr>
<tr><td>4.82</td><td>83.0(6017)</td></tr>
<tr><td>4.83</td><td>83.6(9859)</td></tr>
<tr><td>4.84</td><td>84.3(399)</td></tr>
<tr><td>4.85</td><td>84.9(8412)</td></tr>
</table>
</td><td>4.8</td><td>4</td><td>B2 for a trial between 4 and 5 exclusive
(B1 for a trial at 4 or 5)
B1 for a different trial of $4.8 < x \leq 4.85$
B1 (dep on at least one previous B1) for 4.8

Trials should be evaluated to at least 2 sf truncated or rounded for values of x correct to 1 decimal place.
Trials should be evaluated to at least 1 dp truncated or rounded for values of x correct to 2 dp.
(For trial at 4.82, accept 83, for trial at 4.85 accept 85)

NB no working scores no marks even if the answer is correct</td></tr>
</table>

3	$x = 3$ gives 39 $x = 4$ gives 80 $x = 3.1$ gives 42.(191) $x = 3.2$ gives 45.(568) $x = 3.3$ gives 49.(137) $x = 3.4$ gives 52.(904) $x = 3.5$ gives 56.(875) $x = 3.6$ gives 61.(056) $x = 3.7$ gives 65.(453) $x = 3.8$ gives 70.(072) $x = 3.9$ gives 74.(919) $x = 3.55$ gives 58.9(38875) $x = 3.56$ gives 59.3(58016) $x = 3.57$ gives 59.7(79293)	3.6	4	B2 for a correct trial in the range $3.5 \leq x \leq 3.6$ evaluated (B1 for a correct trial in the range $3 \leq x \leq 4$ evaluated) B1 for a different correct trial in the range $3.55 \leq x < 3.58$ evaluated B1 (dep on at least one previous B1) for 3.6 Accept trials correct to the nearest whole number (rounded or truncated) if the value of x is to 1 dp but to 1 dp (rounded or truncated) if the value of x is to 2 dp NB: If no working shown, no marks can be awarded, even if the answer is correct.

3	$x = 3$ gives 39 $x = 4$ gives 80 $x = 3.1$ gives 42.(191) $x = 3.2$ gives 45.(568) $x = 3.3$ gives 49.(137) $x = 3.4$ gives 52.(904) $x = 3.5$ gives 56.(875) $x = 3.6$ gives 61.(056) $x = 3.7$ gives 65.(453) $x = 3.8$ gives 70.(072) $x = 3.9$ gives 74.(919) $x = 3.55$ gives 58.9(38875) $x = 3.56$ gives 59.3(58016) $x = 3.57$ gives 59.7(79293)	3.6	4	B2 for a correct trial in the range $3.5 \leq x \leq 3.6$ evaluated (B1 for a correct trial in the range $3 \leq x \leq 4$ evaluated) B1 for a different correct trial in the range $3.55 \leq x < 3.58$ evaluated B1 (dep on at least one previous B1) for 3.6 Accept trials correct to the nearest whole number (rounded or truncated) if the value of x is to 1 dp but to 1 dp (rounded or truncated) if the value of x is to 2 dp NB: If no working shown, no marks can be awarded, even if the answer is correct.

Paper: 5MB3H_01

Question		Working	Answer	Mark	Notes
8		x=2 gives 20 x=2.1 gives 21(.86...) x=2.2 gives 23(.84...) x=2.3 gives 25(.96...) x=2.4 gives 28(.22...) x=2.5 gives 30(.62...) x=2.6 gives 33(.17...) x=2.7 gives 35(.88...) x=2.8 gives 38(.75...) x=2.9 gives 41(.78...) x=3 gives 45 x=2.35 gives 27.0(7...) x=2.36 gives 27.3(0...) x=2.37 gives 27.5(3...)	2.3	4	B2 for a correct trial $2.3 \leq x \leq 2.4$ evaluated (B1 for a correct trial $2 \leq x \leq 3$ evaluated) B1 for a different correct trial $2.3 < x < 2.4$ evaluated B1 (dep on at least one previous B1) for 2.3 Accept trials correct to the nearest whole number (rounded or truncated) if the value of x is to 1 dp but correct to 1 dp (rounded or truncated) if the value of x is to 2 dp NB: no working scores no marks even if answer is correct

PAPER: 1MA0_2H

Question		Working	Answer	Mark	Notes
10		5 115 6 204 5.1 122.(451) 5.2 130.(208) 5.3 138.(277) 5.4 146.(664) 5.5 155.(375) 5.6 164.(416) 5.7 173.(793) 5.8 183.(512) 5.9 193.(579) 5.11 123.2(12..) 5.12 123.9(77..) 5.13 124.7(45..) 5.14 125.5(16..) 5.15 126.2(90..)	5.1	4	B2 for a trial $5.1 \leq x \leq 5.2$ evaluated (B1 for a trial $5 \leq x \leq 6$ evaluated) B1 for a different trial $5.1 < x \leq 5.15$ evaluated B1 (dep on at least one previous B1) for 5.1 Accept trials correct to the nearest whole number (rounded or truncated) if the value of x is to 1 dp but correct to 1dp (rounded or truncated) if the value of x is to 2 dp. (Accept 124 for $x = 5.12$) NB : no working scores no marks even if the answer is correct.
*11			No + reason	4	M1 for intention to find the circumference eg $140 \times \pi$ (= 439.82...) A1 for circumference = 439 - 440 M1 (dep on M1) for a complete method shown that could arrive at two figures that are comparable eg "C"÷60×12 (=87.96..) , 90÷12×60 (=450) , 90×60÷"C" (=12.27), "C"÷90×12 (=58.64..) C1 (dep on both M marks) for No and explanation that shows a correct comparison eg only 84 people could sit around the tables or that 13 tables are needed or that 480 cm is needed.

PAPER: 1MA0_2H

PAPER: 5MB3H_01

Question		Working	Answer	Mark	Notes
9		2 62 2.1 65(.961) 2.2 70(.048) 2.3 74(.267) 2.4 78(.624) 2.5 83(.125) 2.6 87(.776) 2.7 92(.583) 2.8 97(.552) 2.9 102.689 3 108 2.61 88.2(49...) 2.62 88.7(24...) 2.63 89.2(01...) 2.64 89.6(79...) 2.65 90.1(59...) 2.66 90.6(41...) 2.67 91.1(24...) 2.68 91.6(08...) 2.69 92.0(95...)	2.6	4	B2 for a correct trial $2.6 \leq x \leq 2.7$ evaluated (B1 for a correct trial $2 \leq x \leq 3$ evaluated) B1 for a different correct trial $2.6 < x < 2.7$ evaluated B1 (dep on at least one previous B1) for 2.6 Accept trials correct to the nearest whole number (rounded or truncated) if the value of x is to 1 dp but correct to 1 dp (rounded or truncated) if the value of x is to 2 dp NB: no working scores no marks even if answer is correct

2		x	x^3-x^2	3.5	4	B2 for a trial $3.4 \leq x \leq 3.5$ evaluated correctly (B1 for a trial evaluated correctly for $3 \leq x \leq 4$) B1 for a different trial evaluated correctly for $3.45 \leq x < 3.5$ B1 (dep on at least one previous B1) for 3.5 [Note: Trials should be evaluated to at least accuracy shown in table, truncated or rounded] [NB No working scores 0 marks]
		3	18			
		3.1	20.(181)			
		3.2	22.(528)			
		3.3	25.(047)			
		3.4	27.(744)			
		3.5	30.(625)			
		3.6	33.(696)			
		3.7	36.(963)			
		3.8	40.(432)			
		3.9	44.(109)			
		4	48			
		3.45	29.1(61125)			
		3.46	29.4(50136)			
		3.47	29.7(41023)			
		3.48	30.0(33792)			
		3.49	30.3(28449)			

PAPER: 1MA0_2H

7		<table><tr><th>x</th><th>$x^3 - 9x$</th></tr><tr><td>4</td><td>28</td></tr><tr><td>4.1</td><td>32.(021)</td></tr><tr><td>4.2</td><td>36.(288)</td></tr><tr><td>4.3</td><td>40.(807)</td></tr><tr><td>4.4</td><td>45.(584)</td></tr><tr><td>4.5</td><td>50.(625)</td></tr><tr><td>4.6</td><td>55.(936)</td></tr><tr><td>4.7</td><td>61,(523)</td></tr><tr><td>4.8</td><td>67.(392)</td></tr><tr><td>4.9</td><td>73.(549)</td></tr><tr><td>5</td><td>80</td></tr><tr><td>4.41</td><td>46.0(7612)</td></tr><tr><td>4.42</td><td>46.5(7089)</td></tr><tr><td>4.43</td><td>47.0(6831)</td></tr><tr><td>4.44</td><td>47.5(6838)</td></tr><tr><td>4.45</td><td>48.0(7113)</td></tr></table>	4.4	4	B2 for a trial $4.4 \leq x \leq 4.5$ (B1 for a trial evaluated correctly for $4 \leq x \leq 5$) B1 for a different trial evaluated correctly for $4.41 \leq x \leq 4.5$ B1 (dep on at least one previous B1) for 4.4 [Note trials should be evaluated to at least accuracy indicated in the table, truncated or rounded] [NB B0 No working scores 0 marks]

5MB3H_01 November 2015

PAPER: 1MA0_2H

Question		Working	Answer	Mark	Notes
7		x x^3-5x 3 12 4 44 3.1 14.(291) 3.2 16.(768) 3.3 19.(437) 3.4 22.(304) 3.5 25.(375) 3.6 28.(656) 3.7 32.(153) 3.8 35.(872) 3.9 39.(819) 3.75 33.9(84..) 3.76 34.3(57..) 3.77 34.7(32..) 3.78 35.1(10..) 3.79 35.4(89..)	3.8	4	B2 for a trial $3.7 \leq x \leq 3.8$ evaluated (B1 for a trial $3 \leq x \leq 4$ evaluated) B1 for a different trial $3.75 \leq x < 3.8$ evaluated B1 (dep on at least one previous B1) for 3.8 Accept trials correct to the nearest whole number (rounded or truncated) if the value of x is to 1 dp but correct to 1dp (rounded or truncated) if the value of x is to 2 dp. NB: no working scores no marks even if the answer is correct.

Original question 8

The full source mark scheme begins on page 196. Use the original paper code, tier and question number shown above.

PAPER: 5MB3H_01

Question	Working	Answer	Mark	Notes
12 (a)		Equation	2	M1 for $2x \times x \times (x + 4)$ A1 for clear working leading to $x^3 + 4x^2 = 90$
(b)	3 63 4 128 3.1 68.231 3.2 73.728 3.3 79.497 3.4 85.544 3.5 91.875 3.45 88.67363 3.46 89.30814 3.47 89.94552 3.48 90.58579	3.5	4	B2 for trial evaluated using $3 < x < 4$ (B1 for trial evaluated using $3 \leq x \leq 4$) B1 for a different trial evaluated using $3.4 \leq x \leq 3.5$ B1 (dep on at least one previous B1) for 3.5 Accept trials correct to the nearest whole number (rounded or truncated) if the value of x is to 1 dp but correct to 1dp (rounded or truncated) if the value of x is to 2 dp. NB: no working scores no marks even if the answer is correct.

Question	Working	Answer	Mark	Notes
13	$x = 2$ gives 10 $x = 3$ gives 30 $x = 2.1$ gives 11.(361) $x = 2.2$ gives 12.(848) $x = 2.3$ gives 14.(467) $x = 2.4$ gives 16.(224) $x = 2.5$ gives 18.(125) $x = 2.6$ gives 20.(176) $x = 2.7$ gives 22.(383) $x = 2.8$ gives 24.(752) $x = 2.9$ gives 27.(289) $x = 2.61$ gives 20.3(89..) $x = 2.62$ gives 20.6(04..) $x = 2.63$ gives 20.8(21..) $x = 2.64$ gives 21(.039..) $x = 2.65$ gives 21.2(59..)	2.6	4	B2 for a trial $2.6 \leq x \leq 2.7$ evaluated (B1 for a trial $2 \leq x \leq 3$ evaluated) B1 for a different trial $2.6 < x \leq 2.65$ evaluated B1 (dep on at least one previous B1) for 2.6 Accept trials correct to the nearest whole number (rounded or truncated) if the value of x is to 1 dp but correct to 1dp (rounded or truncated) if the value of x is to 2 dp. NB: no working scores no marks even if the answer is correct.
*14		Simple with correct comparable values	4	M1 for a method to calculate 2.15% or 2.3% or 102.15% or 102.3% of 15000 M1 for a complete method to calculate using a compound interest rate of 2.15% for 3 years eg 1.0215^3 (=1.065(89…)) or $1.0215^3 \times 15000$ (=15988.45) M1 for a complete method to calculate using a simple interest rate of 2.3% for 3 years e.g $1+0.023 \times 3$(=1.069) or $15000 + 0.069 \times 15000$ (=16035) or 0.023×3 C1 for a correct decision in a statement based on their two accurate comparable values eg 16035, 15988 to 15988.50 or 1035, 988 to 988.50 or 1.065(89...), 1.069 or 6.6%, 6.9 %

Paper: 1MA0/2H

16	(a)		x_1=-2.64 x_2= -2.57392 x_3=-2.603767255	M1 M1 A1	for substitution of −2.5 into the equation (to get $x_1 = -2.64$) for substitution of "$x_1 = -2.64$" and "$x_2 = -2.57392$" to give x_2 and x_3 for $x_1 = -2.64$ oe, $x_2 = -2.57(392)$ and $x_3 = -2.6(03767255)$ Condone $x_3 = -2.61$ if $x_2 = -2.57$ is used in the substitution
	(b)		Statements	C1 C1	Connection between equation and iterative form in (a) e.g. rearrangement Statement e.g. iteration is an estimation of a solution

Question	Working	Answer	Mark	Notes
15 (a)		Shown	M1	for method to establish at least one root between $x = 0$ and $x = 1$, eg f(0) = – 5 and f(1) = 3
			C1	for correct values and a deduction about the roots eg as there is a sign change there must be at least one root between $x = 0$ and $x = 1$ (as f is continuous)
(b)		Shown	C1	for a correct first step in rearrangement, eg $x(x^2 + 7) – 5 = 0$ or $x^3 + 7x = 5$
			C1	for clear and correct steps showing complete rearrangement
(c)	$x_1 = 0.625$ $x_2 = 0.6765327696$ $x_3 = 0.6704483001$	0.6704(483001)	M1 M1 A1	for substitution of 1 into the formula (to get 0.625) for substitution of "$x_1 = 0.625$" and "$x_2 = 0.6765327696$" to give x_2 and x_3 0.6704(483001)
(d)		Comment	M1 C1	substitutes answer to (c) into expression (to get –0.00549...) appropriate comment, eg accurate as answer is close to 0

Question	Answer	Mark	Mark scheme	Additional guidance
18 (a)	Correct statement	C1	for substituting both 1 and 2 into $x^3 + x$ or into $x^3 + x - 7$	All arithmetic shown must be correct. Ignore any additional trials shown.
		C1	for values 2 and 10 plus explanation that these are above and below 7, or for values -5 and 3 plus explanation that there is a change of sign, thus implying a solution lies between 1 and 2	
(b)	Correct rearrangement	C1	for correct algebraic rearrangement	
(c)	1.74	M1	for substitution of 2 into the formula eg $\sqrt[3]{7-2}$ (= 1.70997…)	x_1 = 1.70997…. x_2 = 1.74241…. x_3 = 1.73884…. Accept an accuracy of 2 dp or more rounded or truncated for values of x_1 and x_2 Award the marks for 1.7 on the answer line provided correct iterations are shown in the working space.
		M1	for a substitution of x_1 to give x_2 (= 1.74241...)	
		A1	for answer in the range 1.738 to 1.74	

Paper: 1MA1/3H

16	(a)	$x_1 = 1.817$ $x_2 = 1.853$ $x_3 = 1.846$	M1 M1 A1	for a correct method to find x_1 eg $\sqrt[3]{10-2\times 2}$ (= 1.8171……) (dep on M1) for substitution of x_1 to give x_2 **and** x_2 to give x_3 for x_1 = 1.81(71…), x_2 = 1.85(33…) and x_3 = 1.84(62.…)	Accept an accuracy of 2dp or more rounded or truncated
	(b)	$a = 2$, $b = -10$	C1	cao	

Paper: 1MA1/2H

Question	Answer	Mark	Mark scheme	Additional guidance
17 (a)	Shows re-arrangement	C1	for showing rearrangement, must see $x^4 = x^2 + 5$ leading to $x = \sqrt[4]{x^2 + 5}$	Can work backwards
(b)	1.669763088	M1	for substitution to find x_1 eg $\sqrt[4]{1.5^2 + 5}$ (= 1.64...)	
		M1	for substitution to find x_2 eg $\sqrt[4]{"1.64..."^2 + 5}$ (= 1.66...)	
		A1	for answer in the range 1.6697 to 1.6698 or 1.67(0)	If a correct value is given and then rounded or rounded incorrectly award full marks

Question	Answer	Mark	Mark scheme	Additional guidance
17 (a)	Shown	M1	For a start to establish at least one root in [1,2], eg substitutes $x = 1$ or $x = 2$ into LHS, eg $1^3 + 2 \times 1 - 6$, or $2^3 + 2 \times 2 - 6$	$f(x) = x^3 + 2x - 6$ $f(1) = -3$ $f(2) = 6$
		C1	For a complete argument, eg evaluates for both values of x and states since there is a sign change there must be at least one root in $1 < x < 2$ (as f is continuous)	
(b)	Shows re-arrangement	C1	for showing re-arrangement; must see $x^3 + 2x - 6 = 0$ leading to $x(x^2 + 2) - 6 = 0$ **or** $x(x^2 + 2) = 6$ or $x = \frac{6}{x^2 + 2}$ leading to $x^3 + 2x = 6$	
(c)	1.4496	M1	for $x_1 = \frac{6}{1.45^2 + 2}$ (=1.462(522851…))	
		M1	for $x_2 = \frac{6}{"1.462(52...)"^2 + 2}$ (=1.449634937)	
		A1	for answer in the range 1.4496 to 1.4497	If a correct value is given and then rounded or rounded incorrectly award full marks.

Paper: 1MA1/2H

GCSE

Mark Scheme

Mathematics A (1387)

June 2003

Edexcel
Success through qualifications

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information please call our Customer Response Centre on 0870 240 9800, or visit our website at www.edexcel.org.uk.

June 2003

Publications Code UG014129

All the material in this publication is copyright
© London Qualifications Ltd 2003

CONTENTS

Notes on Marking Principles 2
Paper 5501 4
Paper 5502 11
Paper 5503 17
Paper 5504 24
Paper 5505 32
Paper 5506 40

NOTES ON MARKING PRINCIPLES

NOTES ON MARKING PRINCIPLES

1 Three types of mark are available

M marks - awarded for correct working seen or implied.
A marks - conditional accuracy marks which are awarded for accurate working following the award of M marks.
B marks - unconditional accuracy marks (independent of M).

2 Abbreviations

cao - correct answer only.
ft - follow through.
$\surd$ - Denotes a "follow through" answer.

SC - special case.
isw - ignore subsequent working.
oe - or equivalent (and appropriate).
NB: a candidate cannot benefit from both isw and ft.

3 If no working shown, then

correct answers normally score full marks
incorrect (even though nearly correct) answers score no marks.

4 Marking instructions

Misread loses A marks (and sometimes B marks) on that part but ft can be allowed on subsequent parts. M marks can still be earned. If in doubt contact your team leader.
If there is a wrong answer in the answer space DO CHECK the working in the body of the script.

- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.
- If there is a wrong answer in the answer space DO CHECK the working in the body of the script.
- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.

5 Style of marking

Answer correct: tick and write part mark in margin NEXT TO BRACKETED MARK.
Answer incorrect: cross, but show M, A or B marks if any earned in body of script and transfer the total of these to the margin next to the bracket mark.
Total for each double page at bottom right page (except for back if used), FINAL TOTAL IN RELEVANT BOX ON FRONT COVER.
Nought in margin for fully incorrect question or page AND FOR NO ATTEMPT.
Where no attempt has been made a line should be put in the answer space and zero in the margin next to the bracketed mark
CHOICE OF METHOD – No marks unless one answer is in answer space – then mark that.
CROSSED OUT WORK - if not replaced this should be marked (if legible).
There must always be a mark next to bracketed mark in the margin/

6 Follow Through Marks

Follow throughs are guided by two principles:

(a) Follow throughs which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous, do not award.

(b) Follow throughs which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

7 Probability

Probability answers must be given as fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least two decimal places (unless tenths).

Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.

8 Linear Equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

9 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

Paper 5501

No	Working	Answer	Mark	Notes
1 (a)		46	1	B1 cao
(b)		3.4	1	B1 oe
(c)		Arrow at 430	1	B1 allow ± half graduation
(d)		Arrow at 3.7	1	B1 allow ± half graduation Accept indications other than arrows as long as they are clear
2		1.60 2.05	2	B1) Condone B1) reversal
3 (a)	$\frac{18}{24}$ or $\frac{9}{12}$ or $\frac{6}{8}$	$\frac{3}{4}$	2	B2 for $\frac{3}{4}$ cao (B1 for $\frac{18}{24}$ or $\frac{9}{12}$ or $\frac{6}{8}$) SC B1 for $\frac{1}{4}$ only
(b)		16 squares shaded	1	B1 cao
4 (a)		line	1	B1 within overlay tolerance
(b)		midpoint	1	B1 within overlay tolerance ft from (a) ± 0.2cm
(c)		rectangle	1	B1 for rectangle 6 cm ± 0.2cm by 4 cm ± 0.2cm
5		kilograms, kg litre, *l* or cubic metres, m^3 inches, in	3	B1 B1 B1
6 (a)		parallel lines marked	1	B1
(b)		right angle marked	1	B1
(c)(i)		acute	1	B1
(ii)		reflex	1	B1

Paper 5501

No	Working	Answer	Mark	Notes
7 (i)		sphere	1	B1
(ii)		cylinder	1	B1 Accept circular prism
(iii)		pyramid	1	B1 Condone omission of "triangular" Accept tetrahedron
8 (a)(i)		40	2	B1 cao
(ii)		50		B1 cao
(b)		5 complete symbols	1	B1 cao
(c)		 and oe inc	1	B1
9 (i)		9, 37, 56, 59, 75	5	B1 cao
(ii)		0.067, 0.56, 0.6, 0.605, 0.65		B1 cao Ignore trailing zeros
(iii)		–10, –6, –4, 2, 5		B1 cao
(iv)		$\frac{2}{5}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}$		B2 for all 4 correct (B1 for any 3 in correct order) SC B1 for all 4 in reverse order (applies to(iv) only)
10 (a)			1	B1
(b)		Plotting (4, 24)	1	B1 ft from their matchsticks
(c)		60	1	B1 cao
(d)		$m = 6n$	2	B2 for $m = 6n$ oe (B1 for $6n$ oe or m = multiple of n except $m = n$)
11 (i)		6, 12	4	B1 cao
(ii)		4, 16		B1 cao
(iii)		3, 4, 6 or 3, 4, 6, 12		B1 Condone omission of 12
(iv)		8, 27		B1 cao

Paper 5501

No	Working	Answer	Mark	Notes
12		2.43	4	B1 for 36 or 0.36 B1 for 96 or 0.96 B1 for 125 or 1.25 If none of first 3 B1s awarded then SC B1 for four 24s and five 25s seen OR 4 × 24 and 5 × 25 seen B1 for 2.43 cao
13 (a)(i)		Edinburgh and Plymouth	3	B1 for Edinburgh or −7 B1 for Plymouth or 5
(ii)		12		B1ft from (i) if one positive and one negative
(b)		Cardiff and Belfast London and Plymouth	2	B1 for Cardiff and Belfast OR −6 and −4 B1 for London and Plymouth OR 3 and 5
14		5 9 8	3	B1 cao B1 cao B1 cao
15 (a)			2	✓ × : 2 0 B2; 1 0 B1; 1 1 B1; 2 1 B1
(b)	$2 \times 2 \times 2$	8	2	M1 for $2 \times 2 \times 2$ A1 for 8 cao
16 (a)		30	1	B1 cao
(b)		3	1	B1 cao
(c)	$2 \times 29 = 58$ $5 \times 30 = 150$ $2 \times 31 = 62$ $1 \times 32 = 32$ $\frac{302}{10} = 30.2$	30.2	3	M1 for freq × no pins M1 (dep on 1st M1) for totalling and ÷ 10 A1 for 30.2 cao

Paper 5501

No	Working	Answer	Mark	Notes
17		7 5 5 13 33 52 23	3	B3 all correct (B2 for 4, 5 or 6 correct B1 for 2 or 3 correct)
18 (a)(i) (ii) (iii) (iv) (b)		$4c$ p^4 $8g$ $10pr$ OR $10rp$ $10y - 15$	4 1	B1 oe B1 cao B1 oe B1 B1 cao Accept $10y + - 15$
19		$\frac{2}{3}$	3	M1 for 3 rows (9 squares) shaded M1 for 2 columns (10 squares) shaded A1 for $\frac{2}{3}$
	$\frac{3}{5} = \frac{9}{15}$ $\frac{2}{3} = \frac{10}{15}$	$\frac{2}{3}$	3	M1 for $\frac{3}{5} = \frac{9}{15}$ M1 for $\frac{2}{3} = \frac{10}{15}$ A1 for $\frac{2}{3}$
	$\frac{3}{5} = 0.6$ $\frac{2}{3} = 0.66$ or 0.67 or better	$\frac{2}{3}$	3	M1 for $\frac{3}{5} = 0.6$) Accept M1 for $\frac{2}{3} = 0.66$ or 0.67 or better) percentages A1 for $\frac{2}{3}$

Paper 5501

No	Working	Answer	Mark	Notes
20 (a)	955 × 48 7640 38200 45840 OR 48 × 955 240 2400 43200 45840	458.40	3	M1 for complete correct method (condone one computational error) A2 for 458.40 cao (A1 for digits 4584 OR ft if M1 awarded)
(b)	14. 5 48 | 696. 0 48 216 192 24 0 24 0	14.50	3	M1 for 1 as first digit in answer and remainder 21 M1 (dep) 4 as second digit in answer A1 for 14.50 (Accept 14.5)
21 (a)		$12x$	1	B1 oe
(b)		$12x + 10y$	2	B2 oe ft from (a) (B1 $12x$ + multiple of y or $10y$ seen) SC B1 for $x = 12x + 10y$ OR $y = 12x + 10y$

Paper 5501

No	Working	Answer	Mark	Notes
22	$\frac{1}{3}+\frac{1}{4}=\frac{4}{12}+\frac{3}{12}=\frac{7}{12}$ $1-\frac{7}{12}=\frac{5}{12}$	$\frac{5}{12}$	3	M1 for $\frac{4}{12}$ **and** $\frac{3}{12}$ oe A2 for $\frac{5}{12}$ oe (A1 for $\frac{7}{12}$) B1 for 1–"$\left(\frac{1}{3}+\frac{1}{4}\right)$" correctly evaluated
	$1-\frac{1}{3}=\frac{2}{3}$, $1-\frac{1}{4}=\frac{3}{4}$ $\frac{8}{12}-\frac{3}{12}=\frac{5}{12}$ or $\frac{9}{12}-\frac{4}{12}=\frac{5}{12}$	$\frac{5}{12}$	3	B1 for $\frac{2}{3}$ or $\frac{3}{4}$ seen M1 for $\frac{8}{12}-\frac{3}{12}=\frac{5}{12}$ or $\frac{9}{12}-\frac{4}{12}=\frac{5}{12}$ A1 for $\frac{5}{12}$ oe
		$\frac{5}{12}$	3	M1 for 0.25 and 0.33 or better A1 for 0.58 or better A1 for $0.41\dot{6}$ or recurring
23 (a) (b)(i) (ii)	 180 – (54 + 54)	54 72 Reason	1 3	B1 cao M1 for 180 – (54 + 54) A1 ft from (a) if $x < 90$ B1 for mentioning isosceles and equal or base angles or equal sides and equal or base angles

Paper 5501

No	Working	Answer	Mark	Notes
24 (a)		Bryani	2	M1 for 4×9 or $4 \times 3 \times 3$ or $4 \times x \times x$ or square x first or square 3 first A1 SC 4×3^2 with Bryani scores B2
(b)		64	1	B1 cao
25 (a)		 dotted line may be solid	3	B2 for rectangle base 3 squares and height 4 squares (B1 for rectangle with one correct dimension) B1 for horizontal line 1 cm from top) SC B2 for completely correct elevation on its side
(b)			2	B2 for perspective drawing showing slant face and cutout B1 for perspective drawing with either slant face cutout omitted
26 (a)	20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$	40	2	M1 for 20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$ A1 cao
(b)	$\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes	line from (45, 20) to (65, 0)	2	M1 for $\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes seen A1 for correct line SC If M0, B1 for line from (45, 20) to (t, 0) where $t > 45$ or B1 for a line of the correct gradient.

Paper 5502

No	Working	Answer	Mark	Notes
1 (a)		$\frac{7}{100}$	1	B1 cao accept 0.07
(b)		0.18	1	B1 cao
(c)	20 in 100 oe	40	2	M1 for sight of 20 in 100 or 20 × 2 A1 cao
2 (a)		6cm	2	B1 for 6 ± 0.2 or 60 ± 2 B1 indep for cm or mm consistent with 1st B1
(b)		At centre	1	B1 within overlay
(c)		Circle drawn	1	B1 all within overlay
3		See diagram	3	B3 all correct – see separate sheet (B2 for 3 correct B1 for 2 correct)
4 (a)		$\frac{1}{4}$ oe	1	B1 cao
(b)		0.75	1	B1 cao
(c)		75%	1	B1 cao
(d)(i)		9	2	B1 accept answer in range 9 - 9.2
(ii)		15 - 16		B1 accept answers in range15 - 16
5 (a)		9:30	1	B1 cao
(b)		2hrs 45 min	2	B2 for 2hr 45 min or 2 $\frac{3}{4}$ hr or 165 minutes B1 2:45 or 2.45 or 165 or 45min + 1hr + 1hr oe
(c)		17	1	B1 cao
6 (a)	96 × 4	3.84	2	M1 for 96 × 4 or digits 384 A1 cao
(b)	3 × 96 + 40 = 328	(0).56	2	M1 for 3 × 96 + 40 or digits 328 or digits 56 A1 cao accept 56p

Paper 5502

No	Working	Answer	Mark	Notes
7 (a)		54 000	1	B1 cao accept 54 thousand
(b)		50 000	1	B1 (accept ten thousand or 10 000) oe
8 (a)		14	1	B1 cao
(b)		6	1	B1 cao
(c)		Correct reflection	2	B2 fully correct (B1 correct reflection in a line parallel to the mirror line or condoning 1 block error in shape or position of shape)
9 (a)		Missing horiz label 1 (and 6) missing on vertical scale	2	B1 B1
(b)		Correct graph	2	B1 for bar up to 4 for yellow B1 for bar up to 2 for green
(c)		Blue	1	B1 cao
(d)	3 + 5 + 4 + 2	14	1	B1 ft from (b)
(e)		$\frac{3}{14}$	1	B1 ft on '14'
10	Barry (8) because you double Kath (7) because you add, 1,2,3		2	B1 oe B1 oe SC: B1 for correct rules only
11 (a)		$2n$	1	B1 for $2n$ or $n + n$ OR $2 \times n$ OR $n \times 2$ OR $n2$
(b)		$2n + 15$	1	B1 for "$2n$" + 15 oe
(c)		20q	1	B1 cao
12 (a)	1 + 3 + 5 + 8 + 5	22	2	M1 add frequencies A1 cao
(b)		No, is>no of cups of coffee in the table	1	B1 'average cannot be bigger than 6' oe OR 'Average must be less than 6 oe'
13 (a)		Trapezium	1	B1 cao ignore spelling
(b)		(2, 3)	1	B1 cao
(c)		Isosceles	1	B1 cao ignore spelling
(d)		Q correct	1	B1 cao

Paper 5502

No	Working	Answer	Mark	Notes
14 (a)		250 000	1	B1 cao
(b)	$\frac{28}{4}$	7	2	M1 for $\frac{28}{4}$ oe or "250000" × 28 A1 cao **SC** B1 for 7 000 000
15 (a)		10	1	B1 cao
(b)		5.5	1	B1 ±0.3 pounds
(c)	$\frac{110}{22}$	50	3	M1 for use of graph at 11 or $\frac{110}{22}$ A1 for 5 A1 cao **SC** B2 for 49.5 – 50.6

Paper 5502

No	Working	Answer	Mark	Notes
16 (a)	$269.30 - 56.80 = 212.50$ $\frac{212.50}{42.50}$	6	2	M1 for $\frac{269.30 - 56.80}{42.50}$ or 5 seen A1 cao
(b)	5% of £269.30 £269.30 – "£13.465" OR $\frac{95}{100} \times 269.30$	255.83 or 255.84	3	M1 for $(5 \div 100) \times 269.30$ M1 for $269.30 - $ "13.465" A1 cao OR M2 for $\frac{95}{100} \times 269.30$ A1 cao **Alternative Method:** M1 for $\frac{5}{100} \times 56.80$ $(= 2.84)$ and $\frac{5}{100} \times 42.50$ (=2.12(5)) (OR 53.96 AND 40.38 (40.375) seen M1 for $56.80 -$ "2.84" (= 53.96) $42.50 -$ "2.12(5)" (= 40.375 or 40.38) "5" × "40.375" + "53.96" A1 cao
17	$4.1^2 \times 1.07 = 16.81 \times 1.07$	17.9867	2	M1 for ("4.1") followed by squaring, or sight of 16.81 A1 cao SC: *B1 for 18 or better with no working*
18 (a)		60	1	B1 cao
(b)	$360 - 60 - 90 - 90$	120	2	M1 for $360 -$ " 60 " $- 90 - 90$ or $180 -$ "60" A1 cao
(c)	6×2	12	2	M1 for 6×2 A1 cao for 12

Paper 5502

No	Working	Answer	Mark	Notes
19 (a)(i)	$240 \times 5 = 1200$	1250	3	B1 cao 1250
(ii)	$\frac{50}{'1250'}$	$\frac{1}{25}$		M1 cao $\frac{50}{'1250'}$ A1 for $\frac{1}{25}$ in its simplest form
(b)	$\frac{60}{100} \times 1000 = 600$	12:5	3	M1 for $\frac{60}{100} \times 1000$ oe A1 for 600 A1 cao
20 (a)		$x + 2$	1	B1 accept $2 + x$ but not $x = x + 2$
(b)	$x + 5 + x + 5 + x + 2 + x + 2$	$4x + 14$	2	M1 adding 4 sides, two of which are '$x + 2$' (all sides to be linear expressions in x) **SC** $x + 5 + x + 2 \times 2$ gets M1 A1 for correct simplified answer or $(20 - 14) \div 4$ oe gets M1
(c)	'$4x + 14$' = 20	1.5 oe	2	M1 for equation A1 cao
21	$\sum f = 90$	Angles drawn, labelled	3	M1 for 1 person = 4°or one angle correct in table or pie chart A1 any 2 correctly drawn angles in pie chart A1 fully correct chart labelled
22 (a)		$2p - q$	2	B1 cao for $2p$ B1 cao for $-q$ accept $(-q + 2p)$, $2p - 1q$ and $2p + -q$
(b)	$5x = 3 + 4$	1.4	2	M1 for either (+3 or sight of 7) or (÷ 5 or sight of 0.8 and 0.6) A1 cao accept $\frac{7}{5}$ or $1\frac{2}{5}$

Paper 5502

No	Working	Answer	Mark	Notes
23 (a)		$\frac{4\times5}{2}$	1	B1 cao
(b)	$1+2+3+4+5+6+7+8$	$\frac{8\times9}{2}$	1	B1 cao
(c)	$\frac{100\times101}{2}$	5050	1	B1 cao
24	0 | 5 7 8 8 1 | 0 0 0 0 2 5 5 5 6 2 | 0 0 0 4 5 3 | 3 5 Key 1 | 2 = 12 (min)	See working	3	B1 for stem 0, 1, 2, 3 or 0, 10, 20, 30 B1 for accurate unordered leaves condone 1 error or omission B1 for key and ordered leaves all correct
25	$3.2\times2.8=8.96$ $2\times4.5\times2.8=25.2$ $2\times4.5\times28.8=28.8$ $\frac{62.96}{2.5}\times2.99$	£75.30	5	M1 for area of any face found correctly M1 for 2 areas found correctly A1 for 62.96 or 54 M1 for $\frac{'62.96'}{2.5}\times2.99$ Al cao Alternate method for candidates who round up $\frac{"62.92"}{2.5}$ M1 for "26" × 2.99 A1 for £77.74 cao **SC**: *for top included B2 for 71.92 m^2 seen or* B3 for £86.02 seen **SC** B4 for £64.58 or £65.78 seen
26	$2.5\times10\,000$	25 000	2	M1 for $2.5\times100\times100$ A1 cao

Paper 5503

No	Working	Answer	Mark	Notes
1 (a)(i)		$8g$	2	B1 oe
(ii)		$10rp$		B1 for $10pr$ or $10\ rp$
(b)		$10y - 15$	1	B1 cao accept $10y$ - + 15
(c)	$6x + 8 - 12x + 15$	$-6x + 23$	2	M1 for 3 correct terms out of 4 A1 cao
2		$m = 6n$	2	B2 for $m = 6n$ oe accept $6 \times n$, n 6 (B1 for $6n$ alone, or $6n$ +1 oe OR m = multiple of n except $m = n$)
3 (i)		0.067, 0.56, 0.6, 0.605, 0.65	1	B1 cao Ignore trailing zeros
(ii)		−10, −6, −4, 2, 5	1	B1 cao
(iii)		$\frac{2}{5}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}$	2	B2 all four correct (B1 any three in correct order) SC: B1 all 4 in reverse order
4		7 5 5 13 33 52 23	3	B3 all correct (B2 for 4, 5 or 6 correct entries) (B1 for 2, 3 correct entries)

Paper 5503

No	Working	Answer	Mark	Notes
5		$\frac{2}{3}$	3	M1 for 3 rows (9 squares) shaded M1 for 2 columns (10 squares) shaded A1
	OR $\frac{3}{5} = \frac{9}{15}$ $\frac{2}{3} = \frac{10}{15}$ Therefore $\frac{2}{3} > \frac{3}{5}$	$\frac{2}{3}$		M1 for $\frac{3}{5} = \frac{9}{15}$ M1 for $\frac{2}{3} = \frac{10}{15}$ A1
	OR $\frac{3}{5} = 0.6$ or percent $\frac{2}{3} = 0.66$ or 0.67 or better Therefore $\frac{2}{3} > \frac{3}{5}$	$\frac{2}{3}$		M1 for $\frac{3}{5} = 0.6$ or percent M1 for $\frac{2}{3} = 0.66$ or 0.67 or better A1
6	$2 \times 29 = 58$ $5 \times 30 = 150$ $2 \times 31 = 62$ $1 \times 32 = 32$ $\frac{302}{10} = 30.2$	30.2	3	M1 for freq × no. pins (at least 3) M1 for totalling and for ÷ 10 (dep on 1st M1) A1 cao
7 (a)			2	B1 for 180° rotation (wrong centre) B1 cao
(b)			3	B1 for any enlargement sf other than 1 B1 for all sides halved B1 for position

Paper 5503

No	Working	Answer	Mark	Notes
8 (a)		$12x$	1	B1 oe
(b)		$12x + 10y$	2	B2 oe or ft from (a) (B1 for $12x$ + a multiple of y or $10y$ seen) **SC** B1 for $x = 12x + 10y$ or $y = 12x + 10y$
9	$\frac{1}{3}+\frac{1}{4}=\frac{4}{12}+\frac{3}{12}=\frac{7}{12}$ $1-\frac{7}{12}=\frac{5}{12}$ OR $1-\frac{1}{3}=\frac{2}{3}, 1-\frac{1}{4}=\frac{3}{4}$ $\frac{9}{12}-\frac{4}{12}=\frac{5}{12}, \frac{8}{12}-\frac{3}{12}=\frac{5}{12}$	$\frac{5}{12}$	3	M1 for $\frac{4}{12}$ and $\frac{3}{12}$ oe A2 for $\frac{5}{12}$ oe (A1 for getting $\frac{7}{12}$) (B1 for $1 - \frac{"1}{3} + \frac{1"}{4}$ correctly evaluated) OR B1 for $\frac{2}{3}$ or $\frac{3}{4}$ seen M1 for $\frac{3}{4}-\frac{1}{3}=\frac{9}{12}-\frac{4}{12}$ or $\frac{2}{3}-\frac{1}{4}=\frac{8}{12}-\frac{3}{12}$ A1 for $\frac{5}{12}$ oe OR M1 0.25 and 0.33 or better A1 for 0.58 or better A1 for 0.416 *recurring*
10 (a)		54	1	B1 cao
(b)(i)	180 – (54 +54)	72	3	M1 for 180 – (“54” + “54”) A1 for 72° ft from (a) if $x < 90°$
(ii)				B1 for mentioning isosceles and equal or base angles or equal sides and equal or base angles

Paper 5503

No	Working	Answer	Mark	Notes
11 (a)	Bryani was correct $4 \times 3^2 = 4 \times 9 = 36$	Bryani	2	M1 for 4×9 or $4 \times 3 \times 3$ or "square the three/x then multiply by four" A1 Bryani **SC** 4×3^2 with Bryani gets B2
(b)			1	B1 cao
12 (a)			3	B2 for rectangle height 4 squares, base 3 squares (B1 for rectangle with one correct dimension) B1for line 1 square from the top **SC** B2 for completely correct elevation on its side
(b)			2	B2 for perspective drawing showing slant face and cut out (B1 for perspective drawing with either slant face or cut out omitted or one aspect incorrect)
13 (a)	20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$	40	2	M1 for 20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$ A1 cao
(b)	$\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes seen	Line from (45, 20) to (65, 0)	2	M1 for $\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes seen A1 for correct line **SC** If M0, B1 for line from (45, 20) to (t, 0) where $t > 45$ or a line of the correct gradient
14 (i)		119.31	3	B1 cao
(ii)		119 310		B1 cao
(iii)		1.23		B1 cao

Paper 5503

No	Working	Answer	Mark	Notes
15	$\frac{10}{100}\times 12000$ 12 000 – 1200 = 10 800 $\frac{10800}{10}=1080$ 10 800 – 1080 = £9720	£9720	3	M1 for $\frac{10}{100}\times 12000$ or sight of 1200 or 2400 or 10 800 or 9600 M1 (dep) for $\frac{10}{100}\times(12\,000-\frac{10}{100}\times 12000)$ or sight of 1080 A1 cao Alternative mark scheme M2 for $12000\times(1-\frac{10}{100})^2$ (M1 for $12000\times(1-\frac{10}{100})$ A1 cao
16 (a)	$2p = 6$	$p = 3$	2	M1 for $7p - 5p = 8 - 2$ or $2p$ or 6 A1 cao
(b)	$7r - 5r = -20 - 2$	–11	2	M1 for $7r + 2 = 5r - 20$ or $\frac{7r}{5}+\frac{2}{5}=r-4$ or $7r - 5r = 20 - 2$ or $\frac{7r}{5}-r=-4-\frac{2}{5}$ A1 cao
17		$5n + 1$	2	B2 oe (B1 for $5n$ seen) NB: n = gets B1 max

Paper 5503

No	Working	Answer	Mark	Notes
18 (a)	–1, 0, 1		2	B2 for –1, 0, 1 (B1 for –1, 0 or 0,1 or –1,1 or –2, -1, 0, 1 only)
(b)	(–1, –1), (0, –1), (1, –1), (0, 0), (1, 0), (1, 1)		3	B3 for 6 points correct B2 for 3 points correct B1 for 1 point correct **NB** –B1 each additional point over six
19 (a)	Triangle with vertices at (0,0) (0,–2) and (3,0)		2	M1 for correct orientation A1 cao
(b)	Rotation, 180°, centre (0,1) Enlargement sf - 1 centre (0,1)		2	B2 for 180° 'rotation' centre (0, 1) B2 for Enlargement sf - 1 centre (0,1) (B1 for any two of the three parts) NB: B0 if additional transformation is included
20	Bisector of $\angle BAC$ Arc around A Region		3	B3 cao (B2 for either two correct boundaries, no shading/ wrong shading or one correct boundary, one incorrect boundary with valid shading) (B1 for either two incorrect boundaries but one drawn from A and one intersection, with valid shading or one correct boundary) Ignore shading outside the triangle
21		Length Volume Area	3	B1 for Length B1 for Volume B1 for Area
22 (a)	Unbiased question with choices		2	B1 for unbiased question B1 for at least 2 choices
(b)			2	Classification 1: A biased question Classification 2: A restricted sample of people Classification 3: Not specifying a range of foods Classification 4: Nothing to do with eating habits B2 reasons which satisfy 2 different classifications (B1 a reason which satisfies one classification)



Paper 5503

No	Working	Answer	Mark	Notes
23 (a)	$6 \times 10^2 \times 8 \times 10^4$ 48×10^6	4.8×10^7	3	M1 for $6 \times 10^a \times 8 \times 10^b$ oe, a and b integers including 0 A1 for 48×10^6 oe A1 cao
(b)	200 000 + 30 000	230000	2	B2 cao (B1 for sight of 200 000 or 30 000 or 2.3×10^5 or 23×10^4)
24 (i)		64	4	B1 cao
(ii)		3		B1 cao
(iii)	$\sqrt{16 \times 9} = \sqrt{144}$	12		B2 cao (B1 for sight of $\sqrt{2^4} \times \sqrt{9}$ or better, or 144 seen)
25 (i)	Tangent 90° to diameter /radius/ line from (through) centre	27°	4	B1 for 27° cao B1 for reason
(ii)	180 –(90+ "27") angle in semicircle (is 90°)/Alternate segments /angle at centre twice at circumference	63°		B1 ft for 90–"27" if not 63° B1 for reason
26 (a)(i)		152	2	B1 cao
(ii)		177		B1 cao
(b)			3	B1 for median marked at 167 B1 ft for postion of box with its ends at "152" and "177" B1 for position of whiskers with ends at 132 and 182 NB: For any points plotted between 141 and 149 give a tolerance of an extra ± 1 square
27	$x^2 + xy + xy + y^2$	$x^2 + 2xy + y^2$	2	M1 for at least 3 of the 4 terms correct A1 cao
		25	2	M1 for recognising 3.47 + 1.53 (= 5) A1 cao

Paper 5504

No	Working	Answer	Mark	Notes
1 (a)	$4.1^2 \times 1.07 = 16.81 \times 1.07$	17.9867	2	M1 for (4.1) followed by squaring, or sight of 16.81 A1 cao SC: B1 for 18 or better with no working
(b)		$(1.6 + 3.8 \times 2.4) \times 4.2$	1	B1 cao Allow additional brackets if they give an expression with value 45.024
2 (a)	269.30 – 56.80 = 212.50	6	2	M1 for $\frac{269.30 - 56.80}{42.50}$ or 5 seen A1 cao
(b)	5% of £269.30 £269.30 – "£13.465" OR $\frac{95}{100} \times 269.30$	255.83 or 255.84	3	M1 for (5 ÷ 100) × 269.30 M1 (dep) for 269.30 – "13.465" A1 cao OR M2 for $\frac{95}{100} \times 269.30$ A1 cao **Alternative Method:** M1 for $\frac{5}{100} \times 56.80$ (= 2.84) and $\frac{5}{100} \times 42.50$ (=2.12(5)) (OR 53.96 AND 40.38 (40.375) seen M1 for 56.80 – "2.84" (= 53.96) 42.50 – "2.12(5)" (= 40.375 or 40.38) "5" × "40.375" + "53.96" A1 cao

Paper 5504

No	Working	Answer	Mark	Notes
3 (a)		60	1	B1 cao
(b)	360 – 60 – 90 – 90	120	2	M1 for 360 – "60" – 90 – 90 or 180 – "60" A1 cao
(c)	6 × 2	12	2	M1 for 6 × 2 A1 cao 12
(d)		Correct drawing	2	B2 for triangle and construction lines (see overlay) (B1 for 1 line of length 4cm and correct arcs crossing OR for correct triangle with either no arcs or incorrect arcs) SC: B1 similar triangle drawn with construction lines
4 a(i)	240 × 5 = 1200	1250	3	B1 cao 1250
(ii)	$\frac{50}{1250}$	$\frac{1}{25}$		M1 $\frac{50}{"1250"}$ A1 oe in its simplest form
(b)	$\frac{60}{100} \times 1000 = 600$ 600: 250	12:5	3	M1 for $\frac{60}{100} \times 1000$ oe A1 for 600 A1 cao
5 (a)		$x + 2$	1	B1
(b)	$x + 5 + x + 5 + x + 2 + x + 2$	$4x + 14$	2	M1 adding 4 sides, two of which must be '$x + 2$' (all sides to be linear expressions in x) A1 for correct simplified answer
(c)	"$4x + 14$" = 20	1.5	2	M1 for equation "$4x + 14$" = 20 OR $\frac{20-14}{4}$ oe A1 cao

Paper 5504

No	Working	Answer	Mark	Notes
6 (a)		$2p - q$	2	B1 cao for $2p$ B1 cao for $-q$ accept $-q + 2p$ and $2p - 1q$
(b)	$5x = 3 + 4$	1.4	2	M1 for either (+3 or sight of 7) or (÷ 5 or sight of 0.8 and 0.6) A1 cao accept $\frac{7}{5}$ or $1\frac{2}{5}$
7 (a)		$\frac{4\times5}{2}$	1	B1 cao
(b)	$1 + 2 + 3 + 4 + 5 + 6 + 7 + 8$	$\frac{8\times9}{2}$	1	B1 cao
(c)	$\frac{100\times101}{2}$	5050	1	B1 cao
(d)		$\frac{n(n+1)}{2}$	2	B2 cao (B1 for any quadratic in n)
8	$3.2 \times 2.8 = 8.96$ $2 \times 4.5 \times 2.8 = 25.2$ $2 \times 4.5 \times 3.2 = 28.8$ $\frac{62.96}{2.5} \times 2.99$	75.30	5	M1 for area of any face found correctly M1 for 2 areas seen A1 for 62.96 or 54 M1 for $\frac{'62.96'}{2.5} \times 2.99$ A1 cao Alternate method for candidates who round up $\frac{"62.96"}{2.5}$ M1 for "26" x 2.99 A1 for £77.74 cao SC: for top included B2 for 71.92 m^2 seen or B3 for £86.02 or £86.71 SC: B4 for £64.58 or £65.78 seen

Paper 5504

No	Working	Answer	Mark	Notes
9	2.5×10000	25000	2	M1 for 2.5×10000 or $2.5 \times 100 \times 100$ A1 cao
10 (a)	$\sum f = 90$ (88), 144, 32, 96	Angles drawn, labelled	3	M1 for 1 person = 4° or one angle correct in table or pie chart A1 any 2 angles correctly drawn in pie chart A1 fully correct chart labelled
(b)	$0.38 + 0.27 + 0.15$	0.20	2	M1 1 – sum A1 cao
11	0 | 5 7 8 8 1 | 0 0 0 0 2 5 5 5 6 2 | 0 0 0 4 5 3 | 3 5 Key 1 | 3 = 13 (min)	See working column	3	B1 for stem 0, 1, 2, 3 or 0, 10, 20, 30 B1 for accurate unordered leaves condone 1 error or omission B1 for key and ordered leaves all correct
12 (a)	$V = \pi \times 4^2 \times 10$	502 – 503	2	M1 for $\pi \times 4^2 \times 10$ A1 502 – 503
(b)	$P^2 = 10^2 + 8^2$ $P = \sqrt{164}$	$\sqrt{164} < 13$	3	M1 for sight of a correct right-angled triangle M1 for $10^2 + 8^2$ A1 for conclusion based on a correct calculation or 12.8 seen
13 (a)	2×30	$2 \times 2 \times 3 \times 5$	4	M1 for systematic method, eg division, factor trees (at least one prime) A1 cao
	2×48	$2^5 \times 3$		M1 for systematic method, eg division, factor trees (at least one prime) A1 cao
(b)		12	1	B1 cao
(c)	$2^5 \times 3 \times 5$	480	2	B2 cao B1 for $2^5 \times 3 \times 5$ or any correct common multiple

Paper 5504

No	Working	Answer	Mark	Notes
14 (a)		$150 < C \leq 200$	2	M1 use of cum freq to find the cost of the 20th or 20.5^{th} car OR $\frac{1}{2}\Sigma f$ or $\frac{1}{2}(\Sigma f + 1)$ A1 eg 150 to 200
(b)		No, because the 21st value is in the same interval	1	B1 20.5^{th} or 21st in same interval or an alternative correct explanation
(c)	80% = 5200 $\frac{5200}{80} \times 100$	6500	3	M1 for (100 – 20) % = 5200 M1 for $\frac{5200}{"80"} \times 100$ A1 cao
15 (a)	$x^2(x + 1) = 230$	AG	2	M1 for $x \times x \times (x+1)$ or $x \times x \times x + 1$ oe A1 cao from $x \times x \times (x+1)$
(b)	‘5 – 150 5.1 – 158.7 5.2 – 167.6 5.3 – 177.0 5.4 – 186.6 5.5 – 196.6 5.6 – 207.0 5.7 – 217.7 5.8 – 228.8 5.9 –240.2 5.85 – 234.4	5.8	4	B 2 for trial between 5.8 and 5.9 inclusive (B1 for different trial between 5 and 6 inclusive) B1 for different trial between 5.8 and 5.85 (not including 5.8) B1 (dep on at least one previous B1) cao for 5.8, 5.81, 5.811

Paper 5504

No	Working	Answer	Mark	Notes
16	$\pi \times \left(\frac{15}{2}\right)^2 = 176.715$	88.4cm^2	3	M1 for $\pi \times \left(\frac{15}{2}\right)^2$ A1 88.3 – 88.4 B1(ind) for cm²
17 (a)	$5 = 0.5x + 1$	8	2	M1 for $5 = 0.5x + 1$ A1 cao
(b)		$y = \frac{1}{2}x + c$	1	B1 $y = \frac{1}{2}x + c$, c ≠ 1, oe
(c)		$x = 2y - 2$ OR $x = 2(y - 1)$	2	M1 for correctly multiplying both sides by 2 or correctly isolating $\frac{x}{2}$ A1 for $x = 2(y - 1)$, $x = \frac{y-1}{0.5}$, $x = \frac{y-1}{\frac{1}{2}}$ oe SC: B1 for $x = 2y - 1$
18	$4x - 6y = 22$ $15x + 6y = 54$ $19x = 76$	$x = 4,\ y = -1$	4	M1 for coefficients of x or y the same followed by correct operation , one arithmetical error A1 cao M1(dep on previous M mark) for sub for other variable A1 cao Trial and improvement –0 unless both x and y correct values found

Paper 5504

No	Working	Answer	Mark	Notes
19 (a)	$SF = \frac{10}{6}$ $\frac{10}{6} \times 4.8 = 8$	8	2	M1 for sight of $\frac{10}{6}$ or $\frac{6}{10}$ or 1.67 or better or $\frac{CD}{10} = \frac{4.8}{6}$ A1 cao
(b)	$\frac{10}{6} \times 4.5 - 4.5 = 3$	19.8	2	M1 for use of SF from (a) to find *BC* or *AC* and adding 4 sides A1 cao
20	$\frac{6 \times 10^{15}}{3.2 \times 10^{8}}$ 1.875×10^{7}	4.3×10^{3}	3	B3 for 4.3×10^{3} to 4.34×10^{3} (B2 for 1.875×10^{7} oe or 4300 to 4340 or final answer of 1.9×10^{7}) (B1 for sight of 6×10^{15} oe or 3.2×10^{8} oe)
21	$8.5 \times \tan 38 = 8.5 \times 0.7813$	6.64	3	M1 for correct use of trig, eg $\tan 38 = \frac{opp}{8.5}$ M1 for 8.5 × tan38 A1 6.64 – 6.641

Paper 5504

No	Working	Answer	Mark	Notes
22 (a)		No, as you would expect about 100. Yes, as it is possible to get 200 sixes with a fair dice	1	B1 for a consistent answer
(b)	$\frac{1}{6}, \frac{5}{6}$ + labels		3	B1 for $\frac{5}{6}$ on the red dice, *not six* branch B1 for a fully complete tree diagram with all branches labelled B1 for $\frac{1}{6}$, $\frac{5}{6}$ on all remaining branches as appropriate

Paper 5505

No	Working	Answer	Mark	Notes
1 (i) (ii) (iii)		119.31 119310 1.23	3	B1 cao B1 cao B1 cao
2	$\frac{10}{100}\times 12000 = 1200$ 12 000 – 1200 = 10 800 10 800 ÷ 10 =1080 10 800 – 1080 = £9720	9720	3	M1 for $\frac{10}{100}\times 12\,000$ or sight of 1200 or 2400 or 10 800 or 9600 M1 (dep) for $\frac{10}{100}\times(12\,000 - \frac{10}{100}\times 12\,000)$ or sight of 1080 A1 cao Alternative markscheme M2 for $12000\times\left(1-\frac{10}{100}\right)^2$ (M1 for $12000\times\left(1-\frac{10}{100}\right)$) A1 cao
3	$7r - 5r = -20 - 2$	–11	2	M1 for $7r + 2 = 5r - 20$ or $\frac{7r}{5} + \frac{2}{5} = r - 4$ or $7r - 5r = -20 - 2$ or $\frac{7r}{5} - r = -4 - \frac{2}{5}$ A1 cao

Paper 5505

No	Working	Answer	Mark	Notes
4 (a)		–1, 0, 1	2	B2 for –1, 0, 1 only (B1 for –1, 0 or 0,1 or –1, 1 or –2, -1, 0, 1 only)
(b)		(–1,–1)(0,–1) (1,–1) (0, 0) (1, 0) (1, 1)	3	B3 for 6 points correct (B2 for 3 points correct) (B1 for 1 point correct) *NB: –B1 for each additional point over six*
5		$5n + 1$	2	B2 oe (B1 for $5n$ seen) NB: n = gets B1 max
6 (a)	Triangle with vertices at (0,0) (0,–2) and (3,0)		2	M1 for correct orientation A1 cao
(b)	Rotation, 180°, centre (0,1) Enlargement sf –1 centre (0,1)		2	B2 for 180° 'rotation' centre (0, 1) or for Enlargement sf - 1 centre (0,1) (B1 for any two of the three parts) NB: B0 if additional transformation is included
7	Bisector of $\angle BAC$ Arc around A Region	See overlay	3	B3 cao (B2 for either two correct boundaries, no shading/ wrong shading or one correct boundary, one incorrect boundary with valid shading) (B1 for either two incorrect boundaries but one drawn from A and one intersection, with valid shading or one correct boundary) Ignore shading outside the triangle
8		Length Volume Area	3	B1 for Length B1 for Volume B1 for Area



Paper 5505

No		Working	Answer	Mark	Notes
9	(a)	Unbiased question with choices		2	B1 for unbiased question B1 for at least 2 choices
	(b)	Leading question and a restricted sample		2	Classification 1: A biased question Classification 2: A restricted sample of people Classification 3: Not specifying a range of foods Classification 4: Nothing to do with eating habits B2 reasons which satisfy 2 different classifications (B1 a reason which satisfies one classification)
10	(a)	$6 \times 10^2 \times 8 \times 10^4$ $48 \times 10^6 = 4.8 \times 10^7$	4.8×10^7	3	M1 for $6 \times 10^a \times 8 \times 10^b$ oe, a and b integers including 0 A1 for 48×10^6 oe A1 cao
	(b)	200 000 + 30 000 = 230 000	230 000	2	B2 cao (B1 for sight of 200 000 or 30 000 or 2.3×10^5 or 23×10^4)
11	(a)	$x^2 + xy + xy + y^2$	$x^2 + 2xy + y^2$	2	M1 for at least 3 of the 4 terms correct A1 cao
	(b)		25	2	M1 for recognising 3.47 + 1.53 (= 5) A1 cao
12	(i)	Tangent 90° to diameter /radius/ line from (through) centre	27°	4	B1 for 27° cao B1 for reason
	(ii)	180 –(90+ "27") angle in semicircle (is 90°)/Alternate segments /angle at centre twice at circumference	63°		B1 ft for 90–"27" if not 63° B1 for reason
13	(i)		p^9	1	B1 cao
	(ii)		$6q^6$	2	B2 for $6q^6$ (B1 for sight of $\frac{6q^9}{q^3}$ or $3q \times 2q^5$ or $3q^4 \times 2q^2$ or $6 \times q \times q \times q \times q \times q \times q$ or final answer of the form kq^6, $k > 0$)

Paper 5505

No	Working	Answer	Mark	Notes
14 (a)(i)		152	2	B1 cao
(ii)		177		B1 cao
(b)			3	B1 for median marked at 167 B1 ft for postion of box with its ends at "152" and "177" B1 for position of whiskers with ends at 132 and 182 NB: For any points plotted between 141 and 149 give a tolerance of an extra ± 1 square
15	(arc =) $\frac{40}{360} \times 2\pi \times 9$ $= 2\pi$	$2\pi + 18$	4	M1 for $\frac{40}{360} \times$ M1 for $2\pi \times 9$ M1 (dep) for $\frac{40}{360} \times 2\pi \times 9$ oe A1 for $\frac{18 \times \pi}{9} + 18$ oe exact form
16 (i)		1	1	B1 cao
(ii)		$\frac{1}{16}$	1	B1 cao accept 0.0625
(iii)		64	1	B1 cao condone ± 64

Paper 5505

No	Working	Answer	Mark	Notes
17 (a)	$F \propto 1/x^2$ $F = k/x^2$ $4 = k / 3^2$ $F = 36/x^2$	$F = \frac{36}{x^2}$	3	M1 for $F = k/x^2$ seen or implied. ($k \neq 1$) M1 (dep) for subst. or sight of $k = 36$ A1 for $F = 36/x^2$
(b)		9	1	B1 ft for 9 (ft on $F = kx^n$, $n \neq 0$)
(c)	$64 = \frac{36}{x^2}$ $x^2 = \frac{36}{64}$ $x = \pm\frac{3}{4}$	$\frac{6}{8}$	2	M1 for $x^2 = \frac{"36"}{64}$ A1 for $\frac{6}{8}$ oe (condone $\pm$) *SC: Use of $F = kx^2$ max M1 M1 A0 B1 ft M0 A0* *SC: Use of* $F = \frac{k}{\sqrt{x}}$ *max M1 M1 A0 B0 M1* ($\sqrt{x} = \frac{4\sqrt{3}}{64}$) *A0*
18	$(5+\sqrt{3})(5-\sqrt{3}) = 5\times5 - 5\sqrt{3} + 5\sqrt{3} - \sqrt{3}\sqrt{3}$ $= 5\times5 - 3$ $\frac{22}{\sqrt{22}} = \frac{22\sqrt{22}}{22}$	$\sqrt{22}$	3	B1 for correct expansion 25 - $5\sqrt{3}$ + $5\sqrt{3}$ - $\sqrt{3}$ $\sqrt{3}$ with 1st three terms reducing to 25 **without** any errors seen B1 (indep) for $\sqrt{3}$ $\sqrt{3}$ = 3 B1 for $\sqrt{22}$ coming from $\frac{22}{\sqrt{22}}$ (S.C $\frac{(5+\sqrt{3})(5-\sqrt{3})\sqrt{22}}{22}$ gets B1)

Paper 5505

No	Working	Answer	Mark	Notes
19 (a)		60 40	2	B1 cao B1 cao
(b)		correct bars	2	B1 for $30 < x \leq 40$ with an area of 2½ squares B1 for $40 < x \leq 70$ with an area of 3 squares *SC:* $\frac{0}{4}$ *give M1 if clearly using area or frequency density*
20 (a)	$6x + 8 - 12x + 15$	$-6x + 23$	2	M1 for 3 of the 4 terms $6x$, $+8$, $-12x$, $+15$ correct A1 cao
(b)		$32x^5y^{15}$	2	B2 cao (B1 for two of 32, x^5, y^{15})
(c)	$\frac{(n+1)(n-1)}{n+1} \times \frac{2}{n-2}$ $\frac{2(n-1)}{n-2}$	$\frac{2(n-1)}{n-2}$	3	M1 for $k(n + 1)(n - 1)$ M1 dep for $\frac{(n+1)(n-1)}{(n+1)} = n - 1$ A1 for $\frac{2(n-1)}{n-2}$
21	Vertices at $(-3, -1\frac{1}{2}), (-4\frac{1}{2}, -1\frac{1}{2}), (-3, -4\frac{1}{2})$		3	B1 for all sides $\times 1\frac{1}{2}$ B1 for correct orientation with 2 vertices almost correct B1 cao

Paper 5505

No	Working	Answer	Mark	Notes
22	Total = 3 + 5 + 2 (=10) $\frac{3}{10}\times\frac{3}{10}\times\frac{5}{10}\left(=\frac{45}{1000}\right), \frac{3}{10}\times\frac{3}{10}\times\frac{2}{10}\left(=\frac{18}{1000}\right)$ $\frac{5}{10}\times\frac{5}{10}\times\frac{3}{10}\left(=\frac{75}{1000}\right), \frac{5}{10}\times\frac{5}{10}\times\frac{2}{10}\left(=\frac{50}{1000}\right)$ $\frac{2}{10}\times\frac{2}{10}\times\frac{3}{10}\left(=\frac{12}{1000}\right), \frac{2}{10}\times\frac{2}{10}\times\frac{5}{10}\left(=\frac{20}{1000}\right)$ $3\times\left(\frac{"45"}{1000}+\frac{"18"}{1000}+\frac{"75"}{1000}+\frac{"50"}{1000}+\frac{"12"}{1000}+\frac{"20"}{1000}\right)$ $\frac{660}{1000}$	$\frac{660}{1000}$ oe	5	M3 for all six expressions seen OR their combined equivalents (M2 for four expressions seen OR their combined equivalents) (M1 for two expressions seen OR their combined equivalents) M1 sum of 18 relevant products condone 1 slip A1 for $\frac{660}{1000}$ oe *SC: without replacement maximum M4 A0* SC: Just 2 beads: Answer either $\frac{38}{100}$ oe OR $\frac{28}{90}$ oe B1

Paper 5505

No	Working	Answer	Mark	Notes
23 (a)(i)		$6\mathbf{b} - 6\mathbf{a}$	2	B1 for $6\mathbf{b} - 6\mathbf{a}$ oe
(ii)		$6\mathbf{a}$		B1 for $6\mathbf{a}$ oe
(b)	$\overrightarrow{EX} = \overrightarrow{EB} + \overrightarrow{BX}$ $= 12\mathbf{b} + ½\ \overrightarrow{BC}$	$12\mathbf{b} - 3\mathbf{a}$	2	M1 for $\overrightarrow{EX} = \overrightarrow{EB} + \overrightarrow{BX}$ oe vector journey in a form ready for straightforward substitution A1 for $12\mathbf{b} - 3\mathbf{a}$ oe
(c)	$\vec{AY} = \frac{5}{3}\vec{AB}$ or $\vec{BY} = \frac{2}{3}\vec{AB}$ $\vec{EY} = 16\mathbf{b} - 4\mathbf{a}$ or $\vec{XY} = 4\mathbf{b} - \mathbf{a}$ $\vec{EY} = 4\vec{XY}$ or $\vec{EX} = 3\vec{XY}$ or $\vec{EY} = \frac{4}{3}\vec{EX}$	Printer Answer	3	B1 for either $\vec{AY} = \frac{5}{3}\vec{AB}$ or $\vec{BY} = \frac{2}{3}\vec{AB}$ oe B1 ft for either $\vec{EY} = 16\mathbf{b} - 4\mathbf{a}$ or $\vec{XY} = 4\mathbf{b} - \mathbf{a}$ ft only on parts (*a*) and (*b*) B1 for either $\vec{EY} = 4\vec{XY}$ or $\vec{EX} = 3\vec{XY}$ or $\vec{EY} = \frac{4}{3}\vec{EX}$ oe **plus** conclusion of *E*, *X*, *Y* on the same straight line
24 (a)(i)		(5, – 4)	4	B1 cao
(ii)		(2, – 9)		B1 cao
(iii)		(2, 4)		B1 cao
(iv)		(1, – 4)		B1 cao
(b)		$(x-2)^2 - 4$	4	B4 for $(x-2)^2 - 4$ oe eg. $x^2 - 4x$ (B3 for $(x+2)^2 - 4$ or $(x-2)^2 + 4$) (B2 for $x^2 - 4$ or $(x-2)^2$ OR $x^2 + bx,\ b \neq 0$ OR $(x+2)^2 + 4$ OR $f(x-2) - 4$) (B1 for $x^2 + 4$ or $(x+2)^2$ or $ax^2 + bx$ or $x^2 + bx + c$ OR $x - 2 - 4$ or $x^2 - 2 - 4$, $a, b, c \neq 0$)

Paper 5506

No	Working	Answer	Mark	Notes
1 (a)	$V = \pi \times 4^2 \times 10$	502 – 503 cm^3	2	M1 for $\pi \times 4^2 \times 10$ A1 502 – 503
(b)	$P^2 = 10^2 + 8^2$ $P = \sqrt{164}$	$\sqrt{164} < 13$	3	M1 for sight of correct right angled triangle M1 for $10^2 + 8^2$ A1 for conclusion based on a correct calculation Or 12.8 seen
2 (a)(i)	2×30	$2 \times 2 \times 3 \times 5$	4	M1 for systematic method, eg division, factor trees (at least one prime) A1 cao
(ii)	2×48	$2^5 \times 3$		M1 for systematic method, division, factor trees (at least one prime) A1 cao
(b)		12	1	B1 cao
(c)	$2^5 \times 3 \times 5$	480	2	B2 cao B1 for $2^5 \times 3 \times 5$ or any correct common multiple

Paper 5506

No	Working	Answer	Mark	Notes
3 (a)		$150 < C \leq 200$	2	M1 use of cum freq to find the cost of the 20th or 20.5^{th} car OR $\frac{1}{2}\Sigma f$ th or $\frac{1}{2}(\Sigma f + 1)$ th car. A1 eg 150 to 200, 150 – 200
(b)		No, because the 21st value is in the same interval	1	B1 for 20.5^{th} or 21st value in the same internal consistent with 'a' OR Refers to the median value being low in the interval (statement to be mathematically correct) See additional sheet
(c)	80% = 5200 $\frac{5200}{80} \times 100$	6500	3	M1 for (100 – 20)% = 5200 M1 for $\frac{5200}{"80"} \times 100$ A1 cao

Paper 5506

No	Working	Answer	Mark	Notes
4 (a)	$x^2(x+1) = 230$	AG	2	M1 for $x \times x \times (x+1)$ or $x \times x \times x + 1$ oe, $x^2(x+1)$, $x^2 \times x + 1$ A1 cao from $x \times x \times (x+1)$, no need to see 230
(b)	5 – 150 6 – 252 5.1 – 158.7 5.2 – 167.6 5.3 – 177.0 5.4 – 186.6 5.5 – 196.6 5.6 – 207.0 5.7 – 217.7 5.8 – 228.8 5.9 –240.2 5.85 – 234.4	5.8	4	B2 for trial between 5.8 and 5.9 inclusive evaluated (B1 for trial between 5 and 6 inclusive evaluated) B1 for different trial between 5.8 and 5.85 (not including 5.8) B1 dep on at least are previous B1 5.8, 5.81, 5.811
5	$\pi \times \left(\frac{15}{2}\right)^2 = 176.715$	88.4 cm^2	3	M1 for $\pi \times \left(\frac{15}{2}\right)^2$ seen A1 88.3 – 88.4 B1 for cm² (independent)

Paper 5506

No	Working	Answer	Mark	Notes
6 (a)	$5 = 0.5x + 1$	8	2	M1 for $5 = 0.5x + 1$ A1 cao
(b)		$y = \frac{1}{2}x + c$	1	B1for $y = \frac{1}{2}x + c$, $c \neq 1$, oe
(c)		$x = 2y - 2$ OR $x = 2(y - 1)$	2	M1for correctly multiplying both sides by 2 or correctly isolating $\frac{x}{2}$ A1 for $x = 2(y - 1)$, $x = \frac{y-1}{0.5}$, $\frac{y-1}{\frac{1}{2}}$ oe SC B1 for x= 2y -1
7	$4x - 6y = 22$ $15x + 6y = 54$ $19x = 76$	$x = 4,\ y = -1$	4	M1 for coefficients of x or y the same followed by correct operation, allow one arithmetical error A1 cao M1 (dep) for correct sub for other variable A1 cao Trial and improvement 0 marks unless both correct values of x and y found
8 (a)	SF = $\frac{10}{6}$ $\frac{10}{6} \times 4.8 = 8$	8	2	M1 for sight of $\frac{10}{6}$ or $\frac{6}{10}$ or 1.67 or better or $\frac{CD}{10} = \frac{4.8}{6}$ A1 cao
(b)	$\frac{10}{6} \times 4.5 - 4.5 = 3$	19.8	2	M1 for use of SF from "a" to find AC or BC or $\frac{BC}{4.5} = \frac{4}{6}$ and adding 4 sides A1 cao

Paper 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
9	$\frac{6\times10^{15}}{3.2\times10^{8}}$ 1.875×10^{7}	4.3×10^{3}	3	B3 for 4.3×10^{3} to 4.34×10^{3} (B2 for 1.875×10^{7} oe or 4300 to 4340, final answer of 1.9×10^{7} B1 for sight of 6×10^{15} oe or 3.2×10^{8} oe)
10	$8.5\times\tan38$ $=8.5\times0.7813$ $\frac{8.5}{\sin(90-38)}=\frac{AB}{\sin38}$ $AB=\frac{8.5\times\sin38}{\sin(90-38)}$ $=\frac{5.2331}{0.788}=6.64$	6.64	3	M1 for correct use of trig, eg $\tan 38=\frac{opp}{8.5}$ M1 for $8.5\times\tan38$ A1 6.64 – 6.641 OR M1 for correct substitution into the sine rule M1 (dep) for correct rearrangement for AB= A1 6.64 – 6.641
11 (a)		No, as you would expect about 100. Yes, as it is possible to get 200 sixes with a fair dice	1	B1 for a consistent answer See additional sheet

Paper 5506

No	Working	Answer	Mark	Notes
(b)	$\frac{1}{6}, \frac{5}{6}$ + labels		3	B1 for $\frac{5}{6}$ on the red dice, *not six* branch B1 for a fully complete tree diagram with all branches labelled B1 for $\frac{1}{6}$ and $\frac{5}{6}$ on all remaining branches as appropriate
(c)(i)	$\left(\frac{1}{6}\right)^2$	$\frac{1}{36}$	2	M1 $\left(\frac{1}{6}\right)^2$ or $\frac{1}{6}\times\frac{1}{6}$ only or 0.28 A1 $\frac{1}{36}$ or 0.03 or better
(ii)	$1-\left(\frac{5}{6}\right)^2$ OR $\frac{1}{6}\times\frac{5}{6}+\frac{5}{6}\times\frac{1}{6}+\frac{1}{6}\times\frac{1}{6}$	$\frac{11}{36}$	3	M2 for $1-\left(\frac{5}{6}\right)^2$ or $1-\frac{5}{6}\times\frac{5}{6}$ A1 cao OR M1 for $\frac{1}{6}\times\frac{5}{6}$ oe M1 for 2 or 3 only of $\frac{1}{6}\times\frac{5}{6}, \frac{5}{6}\times\frac{1}{6}$, "a" A1 for $\frac{11}{36}$ or 0.31 or better

Paper 5506

No	Working	Answer	Mark	Notes
12 (a)	$\pi \times 30 \times \frac{7.5^2}{3} - \pi \times 10 \times \frac{2.5^2}{3} = 1767 - 65$	1700	3	M1 for either $\pi \times 30 \times \frac{7.5^2}{3}$ or $\pi \times 10 \times \frac{2.5^2}{3}$ M1 (dep) for difference A1 1700 – 1702 **SC** B1 Using d instead of r, 6800 – 6808
(b)	$\frac{S}{2\pi d} = \sqrt{h^2 + d^2}$ $\left(\frac{S}{2\pi d}\right)^2 = h^2 + d^2$	$h = \sqrt{\frac{S^2 - 4\pi^2 d}{4\pi^2 d^2}}$	3	M1 for correctly isolating $\sqrt{h^2 + d^2}$ or $h^2 + d^2$ or $h + d$ or kh^2 or kh M1(indep) squaring both sides A1 $h = \sqrt{\frac{S^2 - 4\pi^2 d^4}{4\pi^2 d^2}}$, $h = \frac{\sqrt{S^2 - 4\pi^2 d^4}}{2\pi d}$ $h = \sqrt{\left(\frac{S}{2\pi d}\right)^2 - d^2}$
(c)	$\left(\frac{30}{20}\right)^2 \times 450$ or $450 \div \left(\frac{20}{30}\right)^2$	1012.5	2	M1 for sight of correct SF^2 including 4:9 A1 1010 to 1013

Paper 5506

No	Working	Answer	Mark	Notes
13 (a)	$\frac{2x(x+20)}{2} = 400$	As given	2	M1 $\frac{2x(x+20)}{2}$ or $\frac{2x \times x + 20}{2}$ or $2x(x + 20) = 800$ A1 cao following correct working, no need for = 400 *SC B1* $2x \times x + \frac{1}{2} \times 2x(10 - \frac{x}{2}) \times 2$
(b)	$\frac{-20 \pm \sqrt{20^2 - 4 \times 1 \times (-400)}}{2}$ $= \frac{-20 \pm 44.721}{2}$	12.361	3	M1 for correct sub, up to signs, in the quad formula A1 for 44.7 or $\sqrt{2000}$ A1 for 12.3606 – 12.361, ignore negative solution T.I B3 for 12.361 OR Completing the square M1 for $(x+10)^2$ seen A1 for $-10 \pm \sqrt{500}$ A1 for 12.3606 – 12.361 ignore negative solution

Paper 5506

No	Working	Answer	Mark	Notes
14 (a)	$0.5\times 8\times 15\times \sin 70^0$	56.4	2	M1 for correct sub into area formula A1 56.38 – 56.4
(b)	$AB^2 = 8^2 + 15^2 - 2\times 8\times 15\times \cos 70^0 = 206.9$ EITHER $0.5\times AB\times CX$ ='56.38' OR $\frac{\sin B}{8} = \frac{\sin 70}{\sqrt{'206.9'}}$ $B = 31.5$ 15 sin '31.5'	7.84	4	M1 for correct sub into cos rule A1 for 206.9 - 207 or 14.38 – 14.4 EITHER M1 for use of area rule to find CX A1 7.83 – 7.84 OR M1 for correct use of sine rule to find sin B or sin A and then sight of 15sin B *or* 8 sin A A1 7.83 – 7.84

Paper 5506

No	Working	Answer	Mark	Notes
15 (a)	$4a^2 - 4a + 1 - (4b^2 - 4b + 1) =$ $4(a^2 - b^2) - 4(a - b)$ $4(a - b)(a + b - 1)$ OR $((2a - 1) - (2b - 1))((2a - 1) + (2b - 1))$ $(2a - 2b)(2a + 2b - 2)$	AG	3	Expansion Method M1 for a correct expansion of any one of the three terms M1(dep) on an attempt to expand all 3 terms and show LHS = RHS A1 fully correct algebra RHS exp is $4(a^2 + ab - a - ba - b^2 + b)$ OR Factorisation Method M1 for attempt to use difference of 2 squares on LHS M1 for one bracket correctly simplified A1 fully correct
(b)	Any 2 odd square numbers have the above form If a and b are both even or odd then $a - b$ is even, so $4(a - b)$ is a multiple of 8 If one of a,b is odd, then $a + b - 1$ is even, so $4(a + b - 1)$ is a multiple of 8		3	B1 'any 2 square nos have the above form' (may be implied by sight of $(2a - 1)^2 - (2b - 1)^2$ in part (b)) B1 first reason B1 second reason SC B1 for $(2r + 1)^2 - (2r - 1)^2$ B1 for 8r
16 (a)	$g_L = \frac{2 \times 4.495}{1.35^2 \times \sin 30.5}$ $g_u = \frac{2 \times 4.505}{1.25^2 \times \sin 29.5}$	9.719 11.710	4	B2 for any 4 of 4.505, 1.25, 29.5, 4.495, 1.35, 30.5 seen (B1 for any two or three seen) B1 for 11.710 – 11.7103 B1 cao 9.719 – 9.71904
(b)	Round, until lower and upper bounds agree	10	1	B1 for 10 + reason " they agree to this level of accuracy"

Paper 5506

No	Working	Answer	Mark	Notes
17 (a)(i)		xy	3	B1 cao
(ii)		y^2		B1 for y^2 or $y \times y$
(iii)		$\frac{x}{2}$		B1 for $\frac{x}{2}$ or $0.5x$ or $2^{-1}x$
(b)	Divide to get $2y = 1$	$q = -1$	2	M1 for $2y = 1$ or $\frac{x}{2} = 32$ or $p + q = 5$ or $1 + p + 2q = 5$
		$p = 6$		A1 cao
18 (a)	$x^2 - 2mx + m^2 - k$	$k = m^2$	2	M1 for correct exp of $(x - m)^2$ or correct completion of the square eg $\left(x - \frac{2m}{2}\right)^2 - \left(\frac{2m}{2}\right)^2$
				A1 cao
				SC B1 for $k = -m^2$
(b)(i)	Min value is $-m^2$	$-m^2$	3	M1 for recognition that min value occurs when $(x - m)^2 = 0$ (either (b)(i) or (b)(ii) correct implies this M1)
(ii)	$x = m$	m		A1 ft on 'k', " $-k$" gets M1 A0
				A1 cao
19	$0.06 \times 0.05 = 0.003$	No	2	M1 for 0.06×0.05
				A1 correct conclusion based on 0.003 or 0.06 x 0.05 stated as $\neq 0.0011$
				OR M1 for statement that for the two events to be independent P (BL and CL) = P(BL) × P(CL)

Paper 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
20		50 50 4	3	B1 50 or $\frac{100}{2}$ B1 for 50 or "a" B1 4 or $\frac{360}{90}$ oe

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4LN

Telephone 01623 467467
Fax 01623 450481

Order Code UG014129 June 2003

For more information on Edexcel qualifications please contact our
Customer Response Centre on 0870 240 9800
or email: enquiries@edexcel.org.uk
or visit our website: www.edexcel.org.uk

London Qualifications Limited. Registered in England and Wales no.4496750
Registered Office: Stewart House, 32 Russell Square, London WC1B 5DN



GCSE

Edexcel GCSE

Mathematics A 1387

Summer 2005
Mark scheme

Edexcel GCSE
Mathematics A 1387

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information please call our Customer Services on 0870 240 9800, or visit our website at www.edexcel.org.uk.

Summer 2005

Publications Code UG016720

All the material in this publication is copyright
© Edexcel Ltd 2005

CONTENTS

Notes on Marking Principles 1

Paper 5521/01 3

Paper 5521/02 12

Paper 5523/03 18

Paper 5523/04 27

Paper 5525/05 37

Paper 5525/06 45

NOTES ON MARKING PRINCIPLES

1 Types of mark

- M marks: method marks
- A marks: accuracy marks
- B marks: unconditional accuracy marks (independent of M marks)

2 Abbreviations

cao – correct answer only
ft – follow through
isw – ignore subsequent working
SC: special case
oe – or equivalent (and appropriate)
dep – dependent
indep - independent

3 No working

If no working is shown then correct answers normally score full marks
If no working is shown then incorrect (even though nearly correct) answers score no marks.

4 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.
If there is no answer on the answer line then check the working for an obvious answer.

5 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

6 **Ignoring subsequent work**
It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

7 **Probability**
Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

8 **Linear equations**
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

9 **Parts of questions**
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

Paper 5521/01					
No		Working	Answer	Mark	Notes
1	(a)		17252	1	B1 cao
	(b)		5400	1	B1 cao
	(c)		thousands, 1000, 4000	1	B1
2	(a)		grams, g	3	B1 oe spelling
			centimetres, cm		B1 oe spelling
			millilitres, m*l*, cm^3		B1 oe spelling
	(b)		5	1	B1 cao
3	(a)		106, 102	1	B1 cao ignore extras
	(b)		eg take away 4	1	B1 could be indicated on the diagram
	(c)		46	1	B1 cao

Paper 5521/01

No	Working	Answer	Mark	Notes
4	286 × 43 = 11440 + 12298 = 12298 43 × 286 = 258 + 3440 + 8600 = 12298 286 × 40 = 1140 286 × 3 = 858 1140+858=12298 x | 40 | 3 | 8600 200 | 8000 | 600 80 | 3200 | 240 | 3440 6 | 240 | 18 | 258 12298 2 8 6 1 | 0 8 | 3 2 | 2 4 | 4 2 | 0 6 | 2 4 | 1 8 | 3 2 9 8	12298	3	M2 for complete correct method (condone one computational error) (M1 for complete correct method with two computational errors) A1 cao OR B2 inside of grid completed (condone missing zeros and one error) (B1 2 or 3 errors) B1 cao
5 (a)		18, 69	1	B1
(b)(i)		18 or 36	1	B1
(ii)		16 or 36	1	B1
(c)		factor	1	B1
(d)(i)		18	3	B1 cao
(ii)		11 or 88		B1
(iii)		69		B1 cao

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
6	(a)		April & May	1	B1 for both
	(b)		Daffodil	1	B1
	(c)		Feb	1	B1
	(d)		Crocus	1	B1
	(e)(i)		$\frac{1}{5}$	2	B1 for $\frac{1}{5}$ oe
	(ii)		× from 56 mm to 64 mm from 0		B1 A single mark on the line, between 56 mm and 64 mm measured from end 0
7	(a)	$\frac{40}{100}$	$\frac{2}{5}$	2	B2 for $\frac{2}{5}$ B1 for $\frac{40}{100}$ or $\frac{4}{10}$ or $\frac{20}{50}$ or $\frac{8}{20}$
	(b)		0.98	1	B1 cao
	(c)		7 500 000	1	B1 cao
	(d)		25	1	B1 cao
	(e)		60	1	B1 cao

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
8	(a)(i)		(0, 2)	2	B1 cao
	(ii)		(4, 1)		B1 cao
	(b)		$(2, 1\frac{1}{2})$ marked	1	B1 Allow 2mm tolerance from $(2, 1\frac{1}{2})$
9	(a)		2.40	1	B1 cao could be indicated on the diagram
	(b)	1.60 + 0.40	2.00	2	M1 for appropriate sum or product in £ or p or 200 seen eg 1.60 + 0.40, 160 + 40, 0.80 + 0.80 + 0.40, 80 + 80 + 40, 0.08×25, 0.80×2.5, 200 A1 cao
	(c)	$1\div0.8$ or $2.50\div2$	1.25	2	M1 for $1.00\div0.8$ or $2.50\div2$ or 125 or appropriate combination eg $1+\frac{1}{2}\times0.50$ A1 cao
10	(a)		hexagon	1	B1 Condone spelling error
	(b)		Sum of angles at a point is 360°	2	B1 for 360 seen B1 for "point", "complete turn" or "a circle" or similar unless accompanied by an incorrect angle SC If neither B1 scored, award B1 for a clear indication that the size of an angle, other than *x*, is 90° or a right angle (may be on diagram)
	(c)	$30\times4+8\times2$	136	2	M1 $30\times4+8\times2$ or attempt to sum 5 or 6 lengths A1 cao

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
11	(a)		13, 67, 76, 103, 130	5	B1 cao
	(b)		$-7, -3, -1, 0, 5$		B1 cao
	(c)		0.07, 0.072,0.7, 0.702, 0.72		B1 cao
	(d)		0.6, $\frac{2}{3}$, 70%, $\frac{3}{4}$		B2 (B1 for any 3 in correct order)
12	(a)		16 30	1	B1 Accept 4 30 pm Do not accept 4 30
	(b)(i)	$33.56 \div 4$ oe	8.39	3	M1 for $33.56 \div 4$ oe eg $3356 \div 4$, division by 2 twice A1 cao
	(ii)		9		B1 ft from "8.39" unless whole number of pounds
13	(a)		6 cm^2	3	B2 for 6 cao for numerical answer (B1 for $5.5 < \text{Area} \leq 7$) then B1 (indep) for cm^2 with or without numerical answer
	(b)	See diagram	Correct shape	2	B2 (B1 for any 2 sides correct or a correct enlargement scale factor $\neq 1$ or 2)

Paper 5521/01				
No	**Working**	**Answer**	**Mark**	**Notes**
14 (a)	$(4+3)\times10$	70	2	M1 for (4 + 3) × 10 A1 cao
(b)	$120\div10-3$	9	2	M1 for $\frac{120}{10}$ or 12 seen eg $12\times10=120$ A1 cao
(c)		$C=10(n+3)$	3	B3 for $C=10(n+3)$ oe such as $C=(n+3)\times10$ (B2 for correct RHS or $C=n+3\times10$, $C=10n+3$ oe B1 for C = some other linear expression in n or for $n+3\times10$, $10n+3$ etc) Note: $C = n$ scores no marks
15		11 13 16 8 21	2	B2 all correct (B1 for 2 correct)

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
16	(a)		$2p + 4q$	2	B2 for $2p + 4q$ (accept $2 \times p$ etc) (B1 for $2p$ or $4q$)
	(b)		$2y^2$	1	B1 accept $2 \times y^2$ oe inc $2 \times y \times y$
	(c)		$3c + 4d$	2	B2 for $3c + 4d$ (accept $3 \times c$ etc) (B1 for $3c$ or $4d$)
	(d)		$8pq$	1	B1 accept in any order but must not include $\times$ sign
17	(a)(i)		60	2	B1 cao
	(ii)		eg top triangle is equilateral		B1 for reason
	(b)		150	2	M1 for $\frac{180-"60"}{2} + 90$ A1 ft from (a)(i) if $x < 90$ SC B1 for "60" + 90 if $x < 90$
18			40	2	M1 for 60×2 or 120 or $60 \div 3$ or 20 or $\frac{120}{180}$ A1 cao
19			correct drawing	2	B2 Condone hidden detail shown with solid lines and missing lines on front face (B1 for a correct sketch with other incorrect sketch(es) or for prism with correct cross section >1 cube wide or for attempt to draw prism with correct cross section or prism with correct plan and side elevation)

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
20		$\frac{600}{3\times10}$ or $\frac{640}{3.2\times10}$	$20\text{-}21\frac{1}{3}$	2	M1 for rounding at least two of the numbers to 1 sf or for sight of 640, 3.2 or 640, 32 or 600, 32 or 30 seen A1 for $20\text{-}21\frac{1}{3}$ Note: 20.3125 scores M0 A0
21	(a)		Points plotted	1	B1 $\pm$ 1 full (2 mm) square
	(b)		positive	1	B1 cao
	(c)		Line of best fit	1	B1 Must pass through (42.5, 1.45), (42.5, 1.55) AND (67.5, 1.75), (67.5, 1.85)
	(d)		~ 1.65	1	B1 ft from single line segment with positive gradient $\pm$ 1 full (2 mm) square
22	(a)	eg $50\times\frac{2000}{500}$	200	2	M1 for $\frac{2000}{500}$ or 4 seen A1 cao
	(b)	eg $400\times\frac{750}{500}$	600	2	M1 for $\frac{750}{500}$ or 1.5 seen or 400 + 200 A1 cao

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
23	(a)	$4\times3-2\times1$ $12-2$	10	3	M1 for 3×4 (=12) or 1×2 or attempt to divide diagram up into rectangles M1 "12"– "2" or sum of areas of rectangles A1 cao
	(b)(i)	$\frac{10}{100}\times680$ or $680\div10$ 680 + 68	748	5	M1 $\frac{10}{100}\times680$ or 680÷10 or 68 seen M1 (dep) 680 + "68" or M2 for 680×1.10 A1 cao
	(ii)	"748"÷50 or 14.96	15		M1 For "748"÷50 or 14.96 Accept "748" rounded up or down to next 50 followed by $\div50$ A1 ft from (b)(i) rounded up SC B1 for 680 (seen) leading to 14

Paper 5521/02					
No		**Working**	**Answer**	**Mark**	**Notes**
1	(a)	Plain llll lll 8 Chicken lll 3 Bovril llll 5 S & Vin llll 4		3	M1 for attempt to tally A1 for 1 frequency correct or all tallies correct A1 for all frequencies correct (accept for /20)
	(b)		4	1	B1 ft
	(c)		Plain or 8	1	B1 ft
2	(a)(i)		11	2	B1 cao
	(ii)		16		B1 cao
	(b)	See diagram	Correct lines	2	B2 cao for both lines correct (B1 for one line correct)
	(c)		12	2	B2 cao (B1 for 11 or 13)
3	(a)		580	1	B1 for 580 (± 2) could be written on line
	(b)		7.2	1	B1 for 7.2 ± 0.02 could be written on line
	(c)		Arrow at 48	1	B1 allow ± half graduation
	(d)		Arrow at 6.7	1	B1 allow ± half graduation
4	(i)		Cylinder	2	B1 ignore spelling
	(ii)		Cuboid		B1 ignore spelling
5	(a)	£10 – (£2.15 + £2.30)	5.55	4	M1 £2.15 + £2.30 A1 for 4.45 M1 £10 – "4.45" A1 cao
	(b)	£60 ÷ £2.80 = 21.42857	21	2	M1 for £60 ÷ 2.80 or sight of digits 214… A1 for 21
	(c)	$120 \times 25 \div 100$	30	2	M1 ¼ of £120 (oe) A1 cao SC B2 for £90

Paper 5521/02					
No		Working	Answer	Mark	Notes
6	(a)(i)		143^0	2	B1 for 143 ($\pm 2^0$)
	(ii)		Obtuse		B1 for obtuse (ignore spelling)
	(b)	See diagram	Accurate drawing	1	B1 for accurate drawing ± 2mm
7	(a)(i)		5	2	B1 cao
	(ii)		23		B1 cao
	(b)	$\times 2 - 1$		1	B1 for explaining a suitable method
	(c)	See their diagram		1	B1 for a correct diagram
	(d)		14, 17	2	B2 cao for both (B1 for one only ft from their 14)
8	(a)		90	1	B1 accept –90
	(b)		540	1	B1 accept –540
	(c)		Jupiter	1	B1 accept –150
	(d)		- 230	1	B1 cao

Paper 5521/02					
No		Working	Answer	Mark	Notes
9	(a)	2658 – 2430 = 228 "228" × 32	72.96	4	M1 2658 – 2430 A1 228 M1 "228" × 32 or "228" × 0.32 or digits 7296 seen A1 cao **Or** M1 for 2430 × 32 (or digits 77760 seen) or 2658 × 32 (or digits 85056 seen) A1 if 1 correct M1 for "85056" – "77760" or 7296 seen A1 cao
	(b)	$\frac{2}{5} \times 145 = 58$ 145 – "58"	87	3	M1 $\frac{2}{5} \times 145$ (or M1 $\frac{3}{5}$ seen) A1 58 (or M1 $\frac{3}{5} \times 145$) A1 for 87 ft
	(c)(i) (ii)		80 125	2	B1 for 80 (± 1) B1 125 (± 3)

Paper 5521/02					
No		**Working**	**Answer**	**Mark**	**Notes**
10	(a)		1.5 – 2.0	1	B1 for height between 1.5m – 2.0m inclusive
	(b)	Height of man × "2.5"	3 – 6	3	B3 for height between 3m – 6m inclusive (B2 for multiplying (a) by a number between 2 and 3 inclusive) (B1 for multiplying (a) by a number)
11		61 – 19 = 42 42 ÷ 3 = 14	14	2	M1 for – 19 or 42 seen A1 cao
12	(a)		5	1	B1
	(b)	4+5+5+5+4+3+2+1+4+5 = 38 mean = 38 ÷ 10 = 3.8	3.8	2	M1 for attempt to add and ÷ 10 or 3.7 or 3.9 seen A1 for 3.8 SC B1 for 33.5 seen
13	(a)		$3x$	1	B1 cao Accept $3 \times x$, $x3$, $x \times 3$, $x + x + x$
	(b)		x - 9	1	B1 for x – 9 cao
14	(a)	14.44 – 8.660254038	5.77974(…)	2	M1 for 14.44 seen or 8.66(…) or 5.7 or 5.8 or better rounded or truncated
	(b)		6	1	A1 cao B1 ft
15		15 ÷ 24	62.5	2	M1 for 15 ÷ 24 or 1500 ÷ 24 or sight of digits 625 A1 cao
16		2.10 × 450	945	2	M1 for digits 210 × 450 or sight of digits 945 A1 cao
17		See diagram	$2(y + y)$ $2y + 2y$	2	B1 for $2(y + y)$ B1 for $2y + 2y$ (Deduct B1 for each additional tick (>2) to min 0)
18		360° ÷ 18 (=20) Sector angles: G= 60; S= 80; B=220; Correct sectors labelled correctly Use angle measurer	Angles drawn, labelled	4	B4 for fully correct and labelled pie chart (B3 for all angles correct or for a labelled pie chart with 2 correct angles) (B2 for labelled pie chart with 1 correct angle drawn) (B1 for 360° ÷ 18 or 20 seen or implied)

Paper 5521/02					
No		Working	Answer	Mark	Notes
19	(a)		Correct plane	2	B2 for a correct plane defined by showing at least 2 lines. (B1 for a line of symmetry on one face)
	(b)		Correct net	2	B2 cao (B1 for 2 equilateral triangles joined appropriately to at least one rectangle or for 1 equilateral triangle joined appropriately to one of 3 rectangles)
	(c)		Correct drawing	2	B1 for two extra sides of length 6cm (± 2mm) B1 for construction arcs 6cm from each of the ends of the given line.
20	(a)		15	1	B1 for 15 (± 1)
	(b)		15	1	B1 for 15 (± 0.4)
	(c)			2	B1 horiz. line from (2, 20) to (3, 20) B1 line from (3, 20) to (5,0) or horizontal translation of it SC B1 for any journey ending at (5,0)
21	(a)	$x+4+x+x+4+x$	$4x+8$	2	M1 for attempting to add x, $x + 4$, x, $x + 4$ may be implied by $4x + a$ ($a > 0$) A1 for $4x + 8$ or $4(x + 2)$
	(b)	$4x + 8 = 54$ $4x = 46$ $x = 11.5$ Length = "11.5" + 4	15.5	3	M1 for "$4x + 8$" = 54 A1 cao for 11.5 seen B1 ft for "11.5" + 4
22		0.4 + 0.15 1 – "0.55"	0.45	2	M1 for 1 – sum A1 for 0.45 o.e. SC B1 for 0.81
23	(a)		3:1	1	B1 cao
	(b)	$\pi \times 2.45$	7.7	2	M1 for $\pi \times 2.45$ (accept π as 3.1 or better) A1 for 7.59 to 7.70

Paper 5521/02					
No		**Working**	**Answer**	**Mark**	**Notes**
24		7×10000	70 000	2	M1 for 7 X 10 000 or 7 X 100 X 100 A1 cao
25		5.40 ÷ 3 × 7	12.60	3	M1 for 5.40 ÷ 3 or sight of 1.8 M1 dep for "1.80" × 7 A1 for 12.6 or equivalent
26		7.60 x $\frac{17.5}{100}$ = 1.33 7.60 + 1.33 = 8.93 1650 x "8.93"	£14 734.50	4	M1 for 7.60 x $\frac{17.5}{100}$ or 1.33 seen or 7.60 x 1.175 (oe) (Award M1 for 10%, 5% and 2½% correctly calculated) A1 for 8.93 or 893 M1 for 1650 x "8.93" or digits 147345 seen A1 cao Accept 14734.5 **Alternative** M1 for 1650 × 7.6(0) or 12540 seen M1 for "12540" × $\frac{17.5}{100}$ or 2194.5 seen or "12540"× 1.175 (oe) (Award M1 for 10%, 5% and 2½% correctly calculated) M1 for "12540" + "2194.5" (dep on both previous method marks) or digits 147345 seen A1 cao accept 14 734.5

Paper 5523/03				
No	**Working**	**Answer**	**Mark**	**Notes**
1		Correct shape	2	B2 B1 for any 2 sides correct, or a correct enlargement scale factor ≠1 or 2.
2 (a)		11 13 16 8 21	2	B2 all correct (B1 for 2 correct) sign
(b)		$\frac{31}{80}$	1	B1 oe
3 (a)		$2p+3q$	2	B2 for $2p+3q$ (accept $2\times p$ etc) (B1 for $2p$ or $3q$ or $2p3q$)
(b)		$2y^2$	1	B1 accept $2\times y^2$ or $2\times y\times y$
(c)		$3c+4d$	2	B2 for $3c+4d$ (accept $3\times c$ etc) (B1 for $3c$ or $4d$ or $3c4d$)
(d)		$8pq$	1	B1 accept in any order but must not include ×
4 (a)(i)		60	2	B1 cao
(ii)		eg top triangle is equilateral		B1 for reason
(b)		150	2	M1 $\frac{(180-"60")}{2}+90$ A1 ft from (a)(i) if $x < 90°$ SC:B1 for answer from "60" + 90 if $x < 90°$

Paper 5523/03					
No		Working	Answer	Mark	Notes
5	(a)	$4\times3-2\times1$ $12-2$	10	3	M1 for 3×4 (=12) or 1×2 or attempt to divide diagram up into rectangles M1 "12" – "2" A1 cao
	(b)(i)	$\frac{10}{100}\times680$ or $680\div10$ 680 + 68	748	5	M1 $\frac{10}{100}\times680$ or $680\div10$ or 68 seen M1 (dep) 680 +"68" (or M2 for 680×1.10) A1 cao
	(ii)	"748" ÷ 50 or 14.96	15		M1 for "748" $\div50$ or 14.96; accept "748" rounded up or down to next 50 followed by $\div50$ A1 ft from (b)(i) rounded up SC B1 for 680 (seen) leading to 14
6	(a)	$2\times5+5\times-3=10-15=$	-5	2	M1 substitute e.g 2×5 and 5×-3 or 10 and -15 A1 cao
	(b)	$5\times4^2-7$ $5\times16-7$	73	3	M1 substitution e.g $5\times4^2-7$; do not accept 54^2-7 M1 $5\times16-7$ or $5\times4\times4-7$ or $80-7$ (NB 4^2 as 4×4) A1 cao

Paper 5523/03					
No		**Working**	**Answer**	**Mark**	**Notes**
7		679 × 28 = 5432 + 13580 = 19012 or 28 × 679 = 252 + 1960 + 16800 = 19012 or  	190.12	3	M1 for an attempt to multiply the units and tens, or correct partitioning M1 complete correct method (condone one arithmetic error) A1 for 190.12 cao OR M1 for putting the numbers in a grid M1 for multiplying out and addition (condone one error) A1 answer shown with point OR M1 for correct partitioning M1 679×20 and 679×8 calculated oe (condone one error) A1 cao
8	(a)		Points plotted	1	B1 ±1 full (2mm) square
	(b)		positive	1	B1 cao
	(c)		Line of best fit	1	B1 must pass through (42.5, 1.45), (42.5, 1.55) AND (67.5, 1.75), (67.5, 1.85)
	(d)		approx 1.65	1	B1 ft from single line segment with positive gradient ± 1 full (2mm) square

Paper 5523/03					
No		**Working**	**Answer**	**Mark**	**Notes**
9	(a)	eg $50 \times \frac{2000}{500}$	200	2	M1 $2000 \div 500$ or 4 seen A1 cao
	(b)	eg $400 \times \frac{750}{500}$	600	2	M1 $750 \div 500$ or 1.5 seen or 400 + 200 A1 cao
10			$C = 10(n+3)$	3	B3 for $C = 10(n+3)$ oe such as $C = (n+3) \times 10$ (B2 for correct RHS or $C = n + 3 \times 10$, $C = 10n + 3$ etc B1 for C = some other linear expression in n or $n + 3 \times 10$, $10n + 3$ etc) NB: C=n scores no marks
11	(a)		$p(p+6)$	2	B2 for $p(p+6)$ or $p \times (p+6)$ (B1 for $p(ap+b)$ where a,b are numbers or $p+6$ seen on it's own, or part of an expression)
	(b)	$x^2 - 4x + 7x - 28$	$x^2 + 3x - 28$	2	M1 for 4 terms correct ignoring signs (e.g $x^2, 4x, 7x, 28$) or 3 terms with correct signs (e.g $x^2, -4x, 7x, -28$) A1 cao
12			correct drawing	2	B2. Condone hidden detail shown with solid lines, or missing lines on front face. (B1 for : one sketch correct with other sketches incorrect cross-section correct with depth > 1 cube, correct plan and side elevation)

Paper 5523/03					
No		**Working**	**Answer**	**Mark**	**Notes**
13		$\frac{600}{3\times10}$ or $\frac{640}{3.2\times10}$	20 to 21 $\frac{1}{3}$	2	M1 For rounding at least two of the numbers to 1 sf, or for sight of 640, 3.2 or 640, 32 or 600, 32 or 30 seen A1 for 20 to 21 $\frac{1}{3}$ NB: 20.3125 scores M0 A0
14	(a)		correct reflection	2	B2 (B1 reflection in line other than x=3)
	(b)		reflection in $y = x$	2	B2 cao Accept the word "reflected" (B1 any statement including the word "reflection")
15	(a)(i)		5^6	1	B1 accept 15125, 5^{4+2}
	(ii)		5^3	1	B1 accept 125, 5^{9-6}
	(b)	$x + y = 10$ and $x - y = 4$	$x = 7$ $y = 3$	3	M1 for either $x + y = 10$ or $x - y = 4$ A2 for both values correct [(A1 for one value correct) If M0, award B3 for both values correct or B2 for one value correct, otherwise B0] SC B2 for $x = 3$ and $y = 7$

Paper 5523/03				
No	**Working**	**Answer**	**Mark**	**Notes**
16 (a)	$5-3x=2x+2$ $5-2=2x+3x$	$\frac{3}{5}$	3	B1 for $2x+2$ seen OR $2.5-1.5x=x+1$ M1 for correct rearrangement of 4 terms A1 for $\frac{3}{5}$ oe
(b)		$-3,-2,-1,0,1,2$	2	B2 (B1 for 5 correct and not more than one incorrect integers)
17		question + response boxes oe	2	1st aspect: One question with time period (eg each night); ignore other questions. 2nd aspect: Response list (at least two), not overlapping.* 3rd aspect: Some mention of units (eg hours) in either question or responses Award B2 for all three aspects, or B1 for just two aspects. * 0-1, 2-3, 4-5 is OK, but 0-1, 1-2, 2-3 is not OK.

Paper 5523/03				
No	**Working**	**Answer**	**Mark**	**Notes**
18 (a)	$\frac{2}{3}\times\frac{3}{4}=\frac{6}{12}$	$\frac{1}{2}$	2	M1 for $\frac{6}{12}$ or $\frac{3}{6}$ or $\frac{2\times3}{3\times4}$ A1 accept 0.5
(b)	$1+2+\frac{8}{12}+\frac{9}{12}$	$4\frac{5}{12}$	3	M1 for attempt to convert to fractions with common denominator e.g two fractions, denominator of 12 A1 correct conversion : $\frac{8}{12}$ and $\frac{9}{12}$, or $\frac{20}{12}$ and $\frac{33}{12}$ seen (oe) A1 cao for $4\frac{5}{12}$ OR attempts to convert to decimals: must use at least 2dp M1 0.66+0.75 (or 1.66+2.75) or 0.67+0.75 etc A1 4.41, 4.417, 4.416 A1 4.416 (*recurring*)
19	$2\times\frac{1}{2}\times6\times8$ or 48 $8\times9+6\times9+10\times9$ or 72 +54 + 90	264 cm^2	4	M1 attempt to find the area of one face; $\frac{1}{2}\times6\times8$ or (8×9) or (6×9) or (10×9) or 72 or 54 or 90 or 24 M1 all five faces with an intention to add A1 cao numerical answer of 264 B1 (indep) cm^2 with or without numerical answer

Paper 5523/03					
No		**Working**	**Answer**	**Mark**	**Notes**
20			$\frac{\pi ab^3}{3d}$ $3(c+d)^3$ $3\pi bc^2$	3	B3 (B1 for each one correct) Nb –B1 for each of the 4th,5th,6th tick
21	(a)	$x+0.3+0.2+x=1$	0.25	2	M1 for $x+0.3+0.2+x=1$ oe, or $0.5\div 2$ A1 oe
	(b)	0.3×200	60	2	M1 0.3×200 A1 cao Accept 60 out of 200 (in words) SC B1 for $\frac{60}{200}$
22	(a)		$(-12)-4$ -2 $(0)8$	3	B3 for all correct [(B1 for each one correct)
	(b)		points plotted accurately points joined with smooth curve	2	B1 ± 1 full (2mm) square ft table if at least B1 awarded (all 5 points plotted) B1 ft for any smooth curve if previous B1 gained NB: curve must pass within 1 full square of the points
23			$\frac{1}{4}$ on LH branch $\frac{2}{3}$ & $\frac{1}{3}$ & $\frac{2}{3}$ on RH branches	2	B1 B1
24			m=3 n=5	2	B1 for 3 B1 for 5 (B2 for $2^3\times 5$ or $2\times 2\times 2\times 5$) SC: award B1 only if m=5, n=3 or for 8×5 seen

Paper 5523/03					
No		Working	Answer	Mark	Notes
25		$\frac{5--1}{-1-2} = -2$	$y = -2x + 5$	4	M1 for clear attempt to find gradient eg fraction with -1,5 in numerator, 2,-1 in denominator A1 for -2 B2 ft for $y =$"-2"$x + 5$ oe (eg $y = \frac{-6}{3}x + 5$) (B1 for $y = mx + 5$ or , $-2x+5$ or $y =$"-2"$x + c$)
26	(a)(i)		150	2	B1 accept 150 or 210
	(ii)				B1 for angle at the centre is twice the angle at the circumference
	(b)	$360 - 90 - 90 -$"150"	30	3	B1 identifies angle between radius and tangent as $90°$ (may be in working or on diagram) M1 $360° - 90 - 90 -$"150" A1 ft from (a)(i) excluding a negative answer OR B1 for 90 M1 for $2 \times (180 - 90 -$"$\frac{150}{2}$"$)$ A1 ft from (a)(i) excluding a negative answer OR B3 for $180 -$(a)

Paper 5523/04					
No		**Working**	**Answer**	**Mark**	**Notes**
1	(a)	14.44 – 8.660254038	5.77974(….)	2	M1 for 14.44 seen or 8.66(……) seen or 5.7 or 5.8 or better, rounded or truncated A1 cao
	(b)		6	1	B1 ft
2		15 ÷ 24	62.5	2	M1 for 15 ÷ 24 or 1500 ÷ 24 or sight of digits 625 A1 cao
3	(a)	2.10 × 450	945	2	M1 for digits 210 × 450 or sight of digits 945 A1 cao
	(b)	63÷2.10	30	2	M1 for 63 ÷ digits 210 A1 cao
4		See diagram	2(y + y) 2y + 2y	2	B1 for 2(y + y) B1 for 2y + 2y (Deduct B1 for each additional tick (>2) to min 0)
5		360° ÷ 18 (=20) Sector angles: G= 60; S= 80; B=220; Correct sectors labelled correctly Use angle measurer	Angles drawn, labelled	4	B4 for fully correct and labelled pie chart (B3 for all angles correct or for a labelled pie chart with two angles correct) (B2 for labelled pie chart with one correct angle drawn) (B1 for 360÷18 or 20 seen or implied)
6	(a)		Correct plane	2	B2 for a correct plane defined by showing at least 2 lines. (B1 for a line of symmetry on one face)
	(b)		Correct net	2	B2 cao (B1 for 2 equilateral triangles joined appropriately to at least one rectangle or for 1 equilateral triangle joined appropriately to one of the three rectangles)
	(c)		Correct drawing	2	B1 for two extra sides of length 6 cm (± 2mm) B1 for construction arcs 6cm from each of the ends of the given line
7		61 – 19 = 42 42 ÷ 3 = 14	14	2	M1 for –19 or 42 seen or 3x + 19 A1 cao

Paper 5523/04					
No		Working	Answer	Mark	Notes
8	(a) (b) (c)		15 15	1 1 2	B1 cao for 15(±1) B1 cao for 15(±0.4) B1 horiz. line from (2,20) to (3,20) B1 line from (3,20) to (5,0) or horiz. translation of it SC: B1 for any journey ending at (5,0)
9	(a)	x+4+x+x+4+x	4x+8	2	M1 for attempting to add $x+4, x, x+4, x$ may be implied by 4x+a, a>0 A1 for 4x+8 or 4(x + 2)
	(b)	4 x + 8 = 54 4 x = 46 x = 11.5 Length = "11.5" + 4	15.5	3	M1 for "4x + 8" = 54 A1 cao for 11.5 seen B1 ft for "11.5"+ 4
10		0.4 + 0.15 1 – 0.55	0.45	2	M1 for 1 – sum A1 for 0.45 oe SC: B1 for 0.81
11	(a) (b)	 π × 2.45	3:1 7.7	1 2	B1 cao M1 for π × 2.45 (accept π as 3.1 or better) A1 for 7.59 to 7.70
12		7 × 10000	70000	2	M1 for 7 × 10000 or 7 × 100 × 100 A1 cao
13		5.40 ÷ 3 × 7	12.60	3	M1 for 5.40 ÷ 3 or sight of 1.8 M1 (dep) for "1.80" × 7 A1 for 12.6 or equivalent

Paper 5523/04				
No	**Working**	**Answer**	**Mark**	**Notes**
14	$7.60 \times \frac{17.5}{100} = 1.33$ 7.60 + 1.33 = 8.93 1650 × "8.93"	£14734.50	4	M1 for $7.60 \times \frac{17.5}{100}$ or 1.33 seen or 7.60 ×1.175 (oe) (Award M1 for 10%, 5% and 2½% correctly calculated) A1 for 8.93 or 893 M1 for 1650 × "8.93" or digits 147345 seen A1 cao Accept 14734.5 OR M1 for 1650×7.6 or 12540 seen M1 for "12540" $\times \frac{17.5}{100}$ or 2194.5 seen or "12540"×1.175 (oe) (Award M1 for 10%, 5%, and 2½% correctly calculated) M1 for "12540" + "2194.5" (dep on both previous M marks) or digits 147345 seen A1 cao accept 14734.5

Paper 5523/04					
No		**Working**	**Answer**	**Mark**	**Notes**
15		$285 \times 1000/(60 \times 60) = 79.1\dot{6}$		3	M2 for $285 \times 1000 \div 60 \div 60$ or $80 \times 60 \times 60 \div 1000$ or for a correct method to obtain two comparable values e.g $80 \times 60 \times 60$ and 285×1000 (M1 for $285 \div 60 \div 60$ or 0.079(….) seen or $80 \times 60 \times 60$ or 288000 seen or for 285×1000 or 285000 seen or $80 \div 1000$ or 0.08 seen) A1 for 288 or 79.(….) or for two correctly calculated comparable values e.g 288000 and 285000
16	(a)	$4x + 12 = 6$ $4x = -6$	-1.5	3	B1 for $4x + 12$ or $x + 3 = \frac{6}{4}$ M1 for a correct re-arrangement of their 3 terms to isolate $4x$ or x A1 for -1.5 oe
	(b)	$v - u = 5t$	$\frac{v-u}{5}$	2	M1 for isolating $\pm 5t$ or $\pm t$ or for dividing through by 5 A1 oe

Paper 5523/04					
No		**Working**	**Answer**	**Mark**	**Notes**
17		3→15 4→48 3.1→17.3(91) 3.2→19.9(68) 3.3→22.7(37) 3.4→25.7(04) 3.5→28.8(75) 3.4→25.7(04) 3.3→22.7(37) 3.35→24.1(95375)	3.3	4	B2 for trial between 3.3 and 3.4 inclusive (B1 for trial between 3 and 4 inclusive) B1 for different trial between 3.3 and 3.4 exclusive B1 (dep on at least one previous B1) for 3.3 NB trials should be evaluated to at least 1 dp truncated or rounded
18	(a)	36 ÷ (7+3+2) "3" × 7	21	3	M1 for 36 ÷ (7+3+2) M1 (dep) for "3" × 7 or 3 or 2 A1 cao
	(b)	$51.5 \times \frac{8.5}{100} = 4.3775$ 51.5 – 4.3775 = 47.1225	47 or 47.1 or 47.12	4	M1 for $51.5 \times \frac{8.5}{100}$ or 4.37(75) seen M1 (dep) for 51.5 – "4.37(75)" A1 for 47 or better B1 (indep) for rounding their answer correctly to the nearest whole number or 1 or 2 d.p OR M1 for $51.5 \times \frac{100-8.5}{100}$ M1 for $51.5 \times$"0.915" or 0.515 × "91.5" A1 for 47 or better B1 (indep) for rounding their answer correct to the nearest whole number or 1 or 2 d.p

Paper 5523/04					
No		Working	Answer	Mark	Notes
19	(a)		Angle in a semicircle	1	B1 oe
	(b)	$12^2 + 16^2 = 400$ $\sqrt{400} = 20$	20	3	M1 for $12^2 + 16^2$ M1 for $\sqrt{144+256}$ A1 cao
	(c)	$\pi \times 10^2$	314	3	M1 for $\pi \times \left(\frac{"20"}{2}\right)^2$ M1 (indep) for correct order of evaluation of $\pi \times r^2$ for any r A1 for 314 – 315 inclusive

Paper 5523/04					
No		Working	Answer	Mark	Notes
20	(a)	$(1 \times 10)+(3 \times 15)+(5 \times 30)+(7 \times 35)+$ $(9 \times 25)+(11 \times 5) = 730$ "730" $\div$ 120 = 6.08333	6.08	4	M1 for use of *fx* with *x* consistent within intervals (including end points) M1 (dep) for use of midpoints M1 (dep on 1st M1) for use of $\frac{\sum fx}{\sum f}$ A1 6.08 to 6.085
	(b)		(10),25,55,90,115, 120	1	B1 for all correct
	(c)		graph	2	B1 ft for 5 or 6 points plotted correctly $\pm$ 1 full (2mm) square at the end of interval dep on sensible table (condone 1 addition error) B1(dep) for points joined by curve or line segments provided no gradient is negative – ignore any part of graph outside range of their points. (SC: B1 if 5 or 6 points plotted not at end but consistent within each interval and joined)

Paper 5523/04				
No	**Working**	**Answer**	**Mark**	**Notes**
(d)		72–74	2	M1 (ft dep on graph being cf) for reading from graph at 7 A1 ft ±1 full (2 mm) square Or B2 for $72-74$
21 (a) (b) (c) (d)		a^7 $15x^3y^4$ $x-1$ $(x+3)(x-3)$	1 2 1 1	B1 accept a^{4+3} B2 cao (B1 for two of 15, x^3, y^4 in a product) B1 cao B1 cao
22	80% = 220 $220 \div 80 \times 100$	275	3	M1 for recognising that 80% is equivalent to 220 M1 for $220 \div 80 \times 100$ oe A1 cao

Paper 5523/04					
No		**Working**	**Answer**	**Mark**	**Notes**
23			$x = 3$ $y = 0.5$	3	M1 for coefficients of x or y the same followed by correct operation, condone one arithmetical error M1 (dep) for substituting found value in one equation A1 cao SC: B1 for one correct answer only if M's not awarded
24			1.4×10^{10}	2	B2 for 1.4×10^{10} or 1.44×10^{10} (B1 for 14.4×10^{9} or 14400,000,000 or 14000,000,000 or 14×10^{9})
25	(a)	$\tan x = \frac{1.9}{3.2}$ $x = \tan^{-1}\left(\frac{1.9}{3.2}\right) = 30.7$	30.7	3	M1 for $\tan x = \frac{1.9}{3.2}$ or $\tan \frac{1.9}{3.2}$ M1 for $\tan^{-1}\left(\frac{1.9}{3.2}\right)$ A1 for 30.6 – 30.7
	(b)	90 + "30.7"	121	1	B1 (indep) ft for 90 + "30.7" rounded to 3 or 4 s.f

Paper 5523/04				
No	**Working**	**Answer**	**Mark**	**Notes**
26	SF$=\frac{12}{9}$ $\frac{12}{9}\times 6=8$	2	2	M1 for $\frac{12}{9}$ or $\frac{9}{12}$ or 1.33… seen or 0.75 seen or 8 seen or $\frac{6}{9}$ or $\frac{9}{6}$ or 0.66… or 1.5 or $\frac{1}{3}$ or 3 oe seen A1 cao

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
1	(a)	$x^2-4x+7x-28$	$x^2+3x-28$	2	M1 for 4 terms correct ignoring signs (e.g $x^2,4x,7x,28$) or 3 terms with correct signs (e.g $x^2,-4x,7x,-28$) A1 cao
	(b)		y^4+2y^2	2	B2 cao B1 for y^4 or $2y^2$
	(c)		$p(p+6)$	2	B2 for $p(p+6)$ or $p\times(p+6)$ (B1 for $p(ap+b)$ where a,b are numbers or $p+6$ seen on it's own, or part of an expression)
	(d)		$3x(2x-3y)$	2	B2 (B1 for $3(2x^2-3xy)$ or $x(6x-9y)$ or $3x(...)$)
2			question + response boxes oe	2	1st aspect: One question with time period (eg each night); ignore other questions. 2nd aspect: Response list (at least two), not overlapping.* 3rd aspect: Some mention of units (eg hours) in either question or responses Award B2 for all three aspects, or B1 for just two aspects. * 0-1, 2-3, 4-5 is OK, but 0-1, 1-2, 2-3 is not OK.
3	(b)		reflection in $y=x$	2	B2 cao accept the word "reflected" (B1 any statement including the word "reflection")
4	(a)	$5-3x=2x+2$ $5-2=2x+3x$	$\frac{3}{5}$	3	B1 for $2x+2$ seen OR $2.5-1.5x=x+1$ M1 for correct rearrangement of 4 terms A1 for $\frac{3}{5}$ oe
	(b)		$-3,-2,-1,0,1,2$	2	B2 (B1 for 5 correct and not more than one incorrect integers)

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
5	(a)	$\frac{2}{3}\times\frac{3}{4}=\frac{6}{12}$	$\frac{1}{2}$	2	M1 for $\frac{6}{12}$ or $\frac{3}{6}$ or $\frac{2\times3}{3\times4}$ A1 accept 0.5
	(b)	$1+2+\frac{8}{12}+\frac{9}{12}$	$4\frac{5}{12}$	2	M1 for attempt to convert to fractions with common denominator e.g two fractions, denominator of 12 A1 correct conversion : $\frac{8}{12}$ and $\frac{9}{12}$, or $\frac{20}{12}$ and $\frac{33}{12}$ seen (oe) A1 cao for $4\frac{5}{12}$ OR attempts to convert to decimals: must use at least 2dp M1 0.66+0.75 (or 1.66+2.75) or 0.67+0.75 etc A1 4.41, 4.417, 4.416 or 0.41, 0.417, 0.416 or 0.42, 4.42 A1 4.416 (*recurring*)
6	(a)(i)		5^6	1	B1 accept 15625, 5^{4+2}
	(ii)		5^3	1	B1 accept 125, 5^{9-6}
	(b)	$x+y=10$ and $x-y=4$	$x = 7$ $y = 3$	3	M1 for either $x + y = 10$ or $x-y=4$ A2 for both values correct [(A1 for one value correct) If M0, award B3 for both values correct or B2 for one value correct, otherwise B0] SC B2 for $x = 3$ or $y = 7$

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
7		$2\times\frac{1}{2}\times6\times8$ or 48 $8\times9+6\times9+10\times9$ or 72 +54 + 90	264 cm^2	4	M1 attempt to find the area of one face; $\frac{1}{2}\times6\times8$ or (8×9) or (6×9) or (10×9) or 72 or 54 or 90 or 24 or 48 M1 all five faces with an intention to add A1 cao numerical answer of 264 B1 (indep) cm^2 with or without numerical answer
8			$\frac{\pi ab^3}{3d}$ $3(c+d)^3$ $3\pi bc^2$	3	B3 (B1 for each one correct) Nb –B1 for each of the 4th,5th,6th tick
9	(a)	$x+0.3+0.2+x=1$	0.25	2	M1 for $x+0.3+0.2+x=1$ oe, or 0.5 ÷ 2 A1 oe
	(b)	0.3×200	60	2	M1 0.3×200 A1 cao Accept 60 out of 200 (in words) SC B1 for $\frac{60}{200}$
10	(a)		$(-12)-4\quad-2\quad(0)8$	3	B3 for all correct [(B1 for each one correct)
	(b)		5 points plotted accurately points joined with smooth curve	2	B1 ±1 full (2mm) square ft table if at least B1 awarded (all 5 points plotted) B1 ft for any smooth curve if previous B1 gained NB: curve must pass within 1 full square of the points
11			m=3 n=5	2	B1 for 3 B1 for 5 (B2 for $2^3\times5$ or $2\times2\times2\times5$) SC: award B1 only if m=3, n=3, for 8×5 seen

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
12		$\frac{5--1}{-1-2} = -2$	$y = -2x + 5$	4	M1 for clear attempt to find gradient eg fraction with -1,5 in numerator, 2,-1 in denominator A1 for -2 cao B2 ft for $y =$"-2"$x + 5$ oe (eg $y = \frac{-6}{3}x + 5$) (B1 for $y = mx + 5$ or , $-2x+5$ or $y =$"-2"$x + c$)
13	(a)		$\frac{1}{4}$ on LH branch $\frac{2}{3}$ & $\frac{1}{3}$ & $\frac{2}{3}$ on RH branches	2	B1 cao B1
	(b)	$\frac{3}{4} \times \frac{2}{3} + \frac{1}{4} \times \frac{1}{3} = \frac{6}{12} + \frac{1}{12}$	$\frac{7}{12}$	3	M1 for $\frac{3}{4} \times \frac{2}{3}$ or $\frac{1}{4} \times \frac{1}{3}$ from their tree diagram M1 for sum of two products A1 for $\frac{7}{12}$ oe
	(c)	$n = 21 \times 4$ or $\frac{1}{6} : \frac{1}{4}$ oe $\frac{1}{6} \times 84$ or $21 \times \frac{2}{3}$	14	3	M1 for either $\frac{1}{3} \times \frac{3}{4} \left(= \frac{1}{4}\right)$ or $\frac{2}{3} \times \frac{1}{4} \left(= \frac{1}{6}\right)$ from their tree diagram M1 for 21×4 (=84) or $\frac{21}{3} \times 2$ A1 for 14 cao SC: B2 for 63 seen in fraction or ratio

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
14	(a)(i)		150	2	B1 accept 150 or 210
	(ii)				B1 for angle at the centre is twice the angle at the circumference
	(b)		30	3	B1 identifies angle between radius and tangent as $90°$ (may be in working or on diagram)
		$360 - 90 - 90 - "150"$ or $180 - "150"$			M1 $360° - 90 - 90 - "150"$ A1 ft from (a)(i) excluding a negative answer Or B1 for 90 M1 for $2 \times (180 - 90 - "\frac{150}{2}")$ A1 ft from (a)(i) excluding a negative answer Or B3 for $180 - (a)$ SC: 180 – "210" can get B1 for $90°$ and/or B1 for "cyclic quadrilateral"
15	(a)		0.2727…	1	B1 for 2.27 recurring or 0.2727…. oe or 0.273
	(b)	eg $x = 0.3939…$ so $100x = 39.3939…$ $99x = 39$ so $x = \frac{39}{99} = \frac{13}{33}$		3	M1 for $100x = 39.39…$ M1 dep for subtraction of both sides A1 for $\frac{13}{33}$ from correct proof **Alternative method** M1 for $13.000 \div 33$ M1 for remainders 31 and 13 A1 for 0.39 recurring SC:B1 for $\frac{39}{99}$

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
16	(a)	$d = kt^2$ $80 = k \times 4^2$	$d = 5t^2$	3	M1 for $d = kt^2$ or $d \propto t^2$ M1 sub d=80 and t=4 into their equation A1 for $d = 5t^2$ oe (cao)
	(b)		245	1	B1 ft from (a) using "k"
	(c)	$45 = 5t^2$	3	2	M1 ft from (a) for substituting d=45 into their equation A1 for 3 cao (condone inclusion of -3)
17	(a)(i) (ii)		(0, 9) (8, 25)	3	B1 cao B1 for $x = 8$ cao B1 for $y = 25$ cao SC: B1 for (25, 8)
	(b)	LHS = $\left(\frac{100-(x^2-16x+64)}{4}\right)$ = $\left(\frac{36+16x-x^2}{4}\right)$ RHS = $\left(\frac{36-2x+18x-x^2}{4}\right)$ = LHS		3	M1 for expansion of either set of brackets with at least 3 of 4 terms correct M1 for common denominator of 4 or multiplying through by 4 or reducing each numerator to a single term A1 for fully correct solution **Alternative method** M1 for $(5-\frac{(x-8)}{2})(5+\frac{(x-8)}{2})$ M1 for $(\frac{2\times5-(x-8)}{2})(\frac{2\times5+(x-8)}{2})$ A1 for $\frac{(18-x)(x+2)}{4}$

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
18	(a)	$\frac{810\pi}{90\pi}$ or 9 $\sqrt{9}$ or 3	12	3	M1 for $\frac{810\pi}{90\pi}$ or 9 or $\frac{1}{9}$ or 1:9 oe M1 for $\sqrt{\frac{810\pi}{90\pi}}$ or $\sqrt{9}$ or 3 or $\frac{1}{3}$ or $\sqrt{9}:\sqrt{1}$ oe A1 cao SC:B1 for answer of 36
	(b)	3^3 or 27 or 2700	2700π	2	M1 for "3"^{3} or 27 or $\left(\sqrt{9}\right)^3 : (\sqrt{81})^3$ oe or 9^3 or 2700– A1 cao
19	(a)(i)		1	1	B1 cao
	(ii)		8	1	B1 cao
	(iii)	$64^{-\frac{2}{3}} = \frac{1}{64^{\frac{2}{3}}}$ or $64^{-\frac{2}{3}} = (4^2)^{-1}$	$\frac{1}{16}$	2	M1 for knowing negative power is a reciprocal or power of $\frac{1}{3}$ root is a cube root A1 cao for $\frac{1}{16}$
	(b)	$\sqrt{27} = \sqrt{9\times3}$ or $\sqrt{27} = 3\sqrt{3}$ or $\sqrt{27} = 3^{3/2}$	$\frac{5}{2}$ oe	2	M1 for $\sqrt{27} = \sqrt{9\times3}$ or $\sqrt{27} = 3^{3/2}$ A1 for $\frac{5}{2}$ oe (cao) **Alternative method** M1 for $9\times27 = 3^{2n}$ A1 for $\frac{5}{2}$ oe (cao)

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
20	(a)(i) (ii) (b)(i) (ii)		(90, 1) (180, 0) (45, 0) (90, -3)	2 2 	B1 cao could be indicated on diagram B1 cao could be indicated on diagram B1 cao could be indicated on diagram B1 cao could be indicated on diagram
21		$\frac{1}{3}\pi x^2 h = \frac{4}{3}\pi(2x)^3$ $x^2 h = 4\times 8x^3$	$32x$	3	M1 for substitution in correct formulae M1 (dep.) for correct unsimplified expression eg $h = \frac{\frac{4}{3}\pi(2x)^3}{\frac{1}{3}\pi x^2}$ oe or $h = 8x$ oe A1 for $32x$ cao
22	(a)	$\left(\overline{OM} =\right)\ \boldsymbol{a} + 2\boldsymbol{b}$ $\left(\overline{ON} =\right) 3\boldsymbol{a}$ or $\frac{6}{2}\boldsymbol{a}$ $\left(\overline{MN} =\right)\ -\boldsymbol{a} - 2\boldsymbol{b} + 3\boldsymbol{a}$	$2\boldsymbol{a} - 2\boldsymbol{b}$	2	B2 (B1 for either $\overline{OM}$ or $\overline{ON}$ or $-\boldsymbol{a} - 2\boldsymbol{b} + 3\boldsymbol{a}$ SC: B1 for $2\boldsymbol{b} - 2\boldsymbol{a}$
	(b)	$\left(\overline{OX} =\right)\ 2\boldsymbol{a} + \boldsymbol{b}$ $\left(\overline{OY} =\right) \boldsymbol{b} + 4\boldsymbol{a}$ (½ $\overline{QR}$ =) $2\mathbf{a} - \boldsymbol{b}$ or (½ $\overline{RQ}$ =) $\boldsymbol{b} - 2\boldsymbol{a}$	$\overline{XY} = 2\boldsymbol{a}$ (hence parallel)	2	B1 for either $\overline{OX}$ or $\overline{OY}$ or (½ $\overline{QR}$) B1 for $\overline{XY} = 2\boldsymbol{a}$ or $\overline{YX} = -2\boldsymbol{a}$

Paper 5525/06					
No		**Working**	**Answer**	**Mark**	**Notes**
1	(a)	$v - u = 5t$	$\frac{v-u}{5}$	2	M1 for isolating $\pm 5t$ or $\pm t$ or for dividing through by 5 A1 oe
	(b)	$x - 3 = 5x - 25$ $22 = 4x$	$5\frac{1}{2}$	3	M1 for $x - 3 = 5(x-5)$ or $\frac{x}{5} - \frac{3}{5} = x - 5$ M1 for isolating terms in x correctly from $ax + b = cx + d$ A1 cao accept $5\frac{1}{2}$, $\frac{11}{2}$, 5.5
2	(a)	$36 \div (7+3+2)$ "3" $\times 7$	21	3	M1 for $36 \div (7+3+2)$ M1 (dep) for "3" $\times 7$ or 3 or 2 A1 cao
	(b)	$51.5 \times \frac{8.5}{100} = 4.3775$ $51.5 - 4.3775 = 47.1225$	47 or 47.1 or 47.12	4	M1 for $51.5 \times \frac{8.5}{100}$ or 4.37(75) seen M1 (dep) for 51.5 – "4.37(75)" A1 for 47 or better B1 (indep) for rounding their answer correctly to the nearest whole number or 1 or 2 d.p OR M1 for $51.5 \times \frac{100 - 8.5}{100}$ M1 for $51.5 \times$"0.915" or 0.515 ×"91.5" A1 for 47 or better B1 (indep) for rounding their answer correct to the nearest whole number or 1 or 2 d.p

Paper 5525/06					
No		**Working**	**Answer**	**Mark**	**Notes**
3		3→15 4→48 3.1→17.3(91) 3.2→19.9(68) 3.3→22.7(37) 3.4→25.7(04) 3.5→28.8(75) 3.4→25.7(04) 3.3→22.7(37) 3.35→24.1(95375)	3.3	4	B2 for trial between 3.3 and 3.4 inclusive (B1 for trial between 3 and 4 inclusive) B1 for different trial between 3.3 and 3.4 exclusive B1 (dep on at least one previous B1) for 3.3 NB trials should be evaluated to at least 1 dp truncated or rounded
4	(a)		Angle in a semicircle	1	B1 oe
	(b)	$12^2 + 16^2 = 400$ $\sqrt{400} = 20$	20	3	M1 for $12^2 + 16^2$ M1 for $\sqrt{144+256}$ A1 cao
	(c)	$\pi \times 10^2$	314	3	M1 for $\pi \times \left(\frac{"20"}{2}\right)^2$ M1 (indep) for correct order of evaluation of $\pi \times r^2$ for any r A1 for 314 – 315 inclusive

Paper 5525/06					
No		Working	Answer	Mark	Notes
5	(a)	$(1 \times 10)+(3 \times 15)+(5 \times 30)+(7 \times 35)+$ $(9 \times 25)+(11 \times 5) = 730$ "730" ÷ 120 = 6.08333	6.08	4	M1 for use of *fx* with *x* consistent within intervals (including end points) M1 (dep) for use of midpoints M1 (dep on 1st M1) for use of $\frac{\sum fx}{\sum f}$ A1 6.08 to 6.085
	(b)		(10),25,55,90, 115,120	1	B1 for all correct
	(c)		graph	2	B1 ft for 5 or 6 points plotted correctly ± 1 full (2mm) square at the end of interval dep on sensible table (condone 1 addition error) B1(dep) for points joined by curve or line segments provided no gradient is negative – ignore any part of graph outside range of their points. (SC: B1 if 5 or 6 points plotted not at end but consistent within each interval and joined)
	(d)		72 – 74	2	M1 (ft dep on graph being cf) for reading from graph at 7 A1 ft ±1 full (2 mm) square Or B2 for $72 - 74$

Paper 5525/06					
No		Working	Answer	Mark	Notes
6	(a)		a^7	1	B1 accept a^{4+3}
	(b)		$15x^3y^4$	2	B2 cao (B1 for two of 15, x^3, y^4 in a product)
	(c)		$x-1$	1	B1 cao
	(d)		$(a + 3b)(a - 3b)$	2	B2 for $(a + 3b)(a - 3b)$ (B1 for $(a \pm 3b)(a \pm 3b)$
7		80% = 220 $220 \div 80 \times 100$	275	3	M1 for recognising that 80% is equivalent to 220 M1 for $220 \div 80 \times 100$ oe A1 cao
8			$x = 3$ $y = 0.5$	3	M1 for coefficients of x or y the same followed by correct operation, condone one arithmetical error M1 (dep) for substituting found value in one equation A1 cao SC: B1 for one correct answer only if Ms not awarded
9			1.4×10^{10}	2	B2 for 1.4×10^{10} or 1.44×10^{10} (B1 for 14.4×10^9 or 14400,000,000 or 14000,000,000 or 14×10^9)

Paper 5525/06					
No		Working	Answer	Mark	Notes
10	(a)	$\tan x = \frac{1.9}{3.2}$ $x = \tan^{-1}\left(\frac{1.9}{3.2}\right) = 30.7$	30.7	3	M1 for $\tan x = \frac{1.9}{3.2}$ or $\tan \frac{1.9}{3.2}$ M1 for $\tan^{-1}\left(\frac{1.9}{3.2}\right)$ A1 for 30.6 – 30.7
	(b)	90 + "30.7"	121	1	B1 (indep) ft for 90 + "30.7" rounded to 3 or 4 s.f
11	(a)	$SF = \frac{12}{9}$ $\frac{12}{9} \times 6 = 8$	2	2	M1 for $\frac{12}{9}$ or $\frac{9}{12}$ or 1.33… seen or 0.75 seen or 8 seen or $\frac{6}{9}$ or $\frac{9}{6}$ or 0.66… or 1.5 or $\frac{1}{3}$ or 3 oe seen A1 cao
	(b)	$SF = \frac{9}{12}, \frac{9}{12} \times 7 = 5.25$	5.25	2	M1 for $\frac{BE}{7} = \frac{9}{12}$ oe A1 cao

Paper 5525/06					
No		Working	Answer	Mark	Notes
12	(a)	$84 = 6.7\pi + 2 \times 6.7 + 2a$ $2a + 13.4 = 62.95\ldots$ or $2a + 34.44 = 84$	24.8	3	M1 for substituting correctly, π may be left M1 for correct rearrangement as far as $\pm 2a$ A1 for $24.7 - 24.8$
	(b)	$P = \pi r + 2r + 2a$ $P - 2a = \pi r + 2r$ $P - 2a = (\pi + 2)r$	$\frac{P-2a}{\pi+2}$	3	M1 subtracting $2a$ from each side M1 for factorising to get $(\pi+2)r$ A1 for $\frac{P-2a}{\pi+2}$ oe S.C $\frac{p-2a}{5.14}$ oe is M1 M1 A0
13		Area $\Delta ABC = \frac{1}{2} \times 14 \times 8 \times \sin 106$ (= 53.8)	53.8	3	M1 for $\frac{1}{2} \times 14 \times 8 \times \sin 106$ M1 (dep) for $56 \times 0.961(26..)$ or $107.6\ldots$ A1 $53.8 - 53.9$ SC 107.6 is B2
14	(a)		4.5	1	B1 cao
	(b)	$500 \times 1.045^{20} = 1205.857\ldots\ldots$	1205.86	2	M1 for 500×1.045^{20} A1 for $1205.85 - 1206$ (SC:B1 for $705.85 - 706$ no working)

Paper 5525/06				
No	**Working**	**Answer**	**Mark**	**Notes**
15 (a)	$6x^2 + 11x - 10 + 6x - 4 = 25$ $6x^2 + 17x - 39 = 0$		3	M1 for an expression for the area involving either $(3x-2)(2x+5)+2(3x-2)$ or $3x(3x-2)+(3x-2)(7-x)$ or $3x(2x+5)-2(7-x)$ or $(3x-2)^2 + 2(3x-2) + (3x-2)(7-x)$ where in each case at least one of 2 or 3 product terms must be correct M1 (indep) for one correct expansion involving x^2 A1 for simplification to final answer

Paper 5525/06					
No		Working	Answer	Mark	Notes
15	(b)(i)	$x = \frac{-17 \pm \sqrt{17^2 - 4\times 6\times(-39)}}{2\times 6}$ $= \frac{-17 \pm \sqrt{289+936}}{12}$ $x = +\frac{18}{12}$ or -4.33	1.5, $-\frac{13}{3}$	4	M1 for $x = \frac{-17 \pm \sqrt{17^2 - 4\times 6\times(-39)}}{2\times 6}$ up to signs in b & c M1 for $x = \frac{-17 \pm \sqrt{1225}}{12}$ A1 $x = 1.5$ or -4.33, or better OR M2 for $(3x+13)(2x-3)$ (M1 for $(3x \pm a)(2x \pm b)$ with ab = ±39 A1 x = 1.5 or – 4.33, or better OR M1 for $\left(x+\frac{17}{12}\right)^2$ *seen* M1 $\left(x+\frac{17}{12}\right)^2 = \left(\frac{17}{12}\right)^2 + \frac{39}{6}$ A1 x = 1.5 or – 4.33, or better SC:M1 for answer "1.5" with no working or T & I
		$x^2 + \frac{17}{6}x - \frac{39}{6} = 0$ $\left(x+\frac{17}{12}\right)^2 - \left(\frac{17}{12}\right)^2 - \frac{39}{6} = 0$ $\left(x+\frac{17}{12}\right)^2 = \left(\frac{17}{12}\right)^2 + \frac{39}{6}$			
	(ii)		8		B1 cao length = 8

Paper 5525/06					
No		**Working**	**Answer**	**Mark**	**Notes**
16	(a)	P(win) = $\frac{2}{5}\times\frac{3}{5}+\frac{2}{5}\times\frac{1}{5}(=\frac{8}{25})$	$\frac{8}{25}$	3	M1 for $\frac{2}{5}\times\frac{3}{5}$ or $\frac{2}{5}\times\frac{1}{5}$ or for clearly identifying in P(R) $\times$P(R) + P(B) $\times$ P(B) M1 for P(win) = " $\frac{2}{5}$ " $\times$ " $\frac{3}{5}$ " + " $\frac{2}{5}$ " $\times$ " $\frac{1}{5}$ " A1 for $\frac{8}{25}$, oe
	(b)	$\frac{8}{25}$ $\times$ 100 (=32) 100 $\times$ 20 – "32" $\times$ 50	£4	2	M1 for (" $\frac{8}{25}$ " $\times$100)$\times$50 or $\times$ 0.5 A1 cao

Paper 5525/06					
No		**Working**	**Answer**	**Mark**	**Notes**
17		Lower bound of 1200 is 1150 Upper bound of 60 is 65 $1150 \div 65$	17	4	B1 for 1150 or 1250 seen B1 for 65 or 55 seen M1 (Lower bound of load) ÷ (Upper bound of weight) Where 1150 ≤ LB load < 1200 and 60 < UB Weight ≤ 65 A1 for 17 requires fully correct working OR B1 for 1150 or 1250 seen B1 for 65 or 55 seen M1 (upper bound of load) ÷(lower bound of weight) Where 1200 ‹ UB load ≤ 1250 and 55 ≤ LB weight < 60 A1 for 22 requires fully correct working OR M2 $1200 \div 55$ A1 21.8 A1 21 requires fully correct working OR M2 $1200 \div 65$ A1 18.4(6) A1 18 requires fully correct working
18	(a)	$3^4 x^4 y^8$	$81x^4y^8$	2	B2 for $81x^4y^8$ (B1 for 2 of 81, x^4, y^8)
	(b)	$\frac{x(x-3)}{(x-5)(x-3)}$	$\frac{x}{x-5}$	3	B1 for $x(x-3)$ B1 for $(x-5)(x-3)$ B1 cao

Paper 5525/06					
No		Working	Answer	Mark	Notes
19	(a)	$6^2 - 2^2 = 32$	5.66	2	M1 for $6^2 - 2^2$ (= 32) A1 5.65 – 5.66
	(b)	V  $DVA = 2 \times \sin^{-1}(\frac{2}{6})$ OR $\cos DVA = \frac{6^2 + 6^2 - 16}{2 \times 6 \times 6}$ $= \frac{56}{72}$ $DVA = \cos^{-1}\left(\frac{56}{72}\right) = 38.94$	38.9	3	M1 $\sin x = \frac{2}{6}$ oe M1 for $DVA = 2 \times \sin^{-1}(\frac{2}{6})$ A1 38.9 – 38.95 OR M1 for $(\cos DVA =) \frac{6^2 + 6^2 - 4^2}{2 \times 6 \times 6}$ M1 for $DVA = \cos^{-1}\left(\frac{56}{72}\right)$ A1 38.9 – 38.95

Paper 5525/06				
No	**Working**	**Answer**	**Mark**	**Notes**
19 (c)	$AC^2 = 2^2 + 2^2 - 2 \times 2 \times 2 \times \cos 120^o$ $AC = \sqrt{12}$ **OR** $AN = 2 \times \sin 60 = \sqrt{3}$ **OR** $VN = \sqrt{"32"+1} = \sqrt{33}$ $\cos AVC = \frac{6^2+6^2-12}{2\times 6\times 6}$ $\cos AVC = \frac{60}{72}$ **OR** $AVC = 2 \times \sin^{-1}\frac{\sqrt{3}}{6}$, using AN **OR** $AVC = 2\times\cos^{-1}\frac{\sqrt{33}}{6}$, using VN	33.6	4	M1 for any valid method for AC or AN or VN where N is the midpoint of AC A1 for $AC^2 = 12$ or $AC = \sqrt{12}$ (= 3.46…) or $AN = \sqrt{3}$ (=1.73…)or $VN = \sqrt{33}$ (=5.74…) M1 (indep) for correct method to find angle AVC A1 33.55 – 33.6

Paper 5525/06					
No		**Working**	**Answer**	**Mark**	**Notes**
20	(a)	Graph translated 1 unit to the right passing through the points (-1,0), (1,2) and (2,0)		2	M1 for translation of $\begin{pmatrix}1\\0\end{pmatrix}$ or $\begin{pmatrix}-1\\0\end{pmatrix}$ A1 for right through the 3 points, $\pm\frac{1}{4}$ sq
	(b)	Graph stretched 2 units parallel to y-axis; passing through the points (-2,0), (0,4) and (1,0)		2	M1 for graph stretched parallel to the y-axis by scale factor 2 A1 through all 3 points; $\pm\frac{1}{4}$ sq not on grid at $x = 2$

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481

Email publications@linneydirect.com

Order Code UG016720 Summer 2005

For more information on Edexcel qualifications, please visit www.edexcel.org.uk/qualifications
Alternatively, you can contact Customer Services at www.edexcel.org.uk/ask or on 0870 240 9800

Edexcel Limited. Registered in England and Wales no.4496750
Registered Office: One90 High Holborn, London, WC1V 7BH

PMT

edexcel

Mark Scheme (Results)

March 2013

GCSE Mathematics (2MB01) Higher
5MB3H (Calculator) Paper 01

PMT

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk for our BTEC qualifications.
Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

If you have any subject specific questions about this specification that require the help of a subject specialist, you can speak directly to the subject team at Pearson.
Their contact details can be found on this link: www.edexcel.com/teachingservices.

You can also use our online Ask the Expert service at www.edexcel.com/ask. You will need an Edexcel username and password to access this service.

Pearson: helping people progress, everywhere
Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

March 2013
Publications Code UG035055
All the material in this publication is copyright
© Pearson Education Ltd 2013

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labeling conventions.

ii) *select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect canceling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

12 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers

Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

Guidance on the use of codes within this mark scheme
M1 – method mark A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
*1			small and correct answers	3	M1 for 4.2 ÷ 1.75 (=2.4(0)) **or** 8.68 ÷ 3.5 (=2.48) M1 for 4.2 ÷ 1.75 (=2.4(0)) **and** 8.68 ÷ 3.5 (=2.48) C1 (dep on at least M1) for small and (£)2.4(0) and (£)2.48 **OR** M1 for 1.75 ÷ 4.2 (=0.41(6)) **or** 3.5 ÷ 8.68 (=0.40(3)) M1 for 1.75 ÷ 4.2 (=0.41(6)) **and** 3.5 ÷ 8.68 (=0.40(3)) C1 (dep on at least M1) for small and 0.41(6)(kg) and 0.40(3)(kg) **OR** M1 for 4.2 ÷ 1.75 (=2.4(0)) **or** 8.68 ÷ 3.5 (=2.48) M1 for (4.2 ÷ 1.75) × 3.5 (=8.4(0)) or (8.68 ÷ 3.5) × 1.75 (=4.34) C1 (dep on at least M1) for small and (£)8.4(0) or for small and (£)4.34 **OR** M1 for 1.75 ÷ 4.2 (=0.41(6)) or 3.5 ÷ 8.68 (=0.40(3)) M1 for (1.75 ÷ 4.2) × 8.68 (=3.61(6)) or (3.5 ÷ 8.68) × 4.2 (=1.69(3)) C1 (dep on at least M1) for small and 3.61(6)(kg) or 1.69(3)(kg) **OR** M2 for 8.68 ÷ 2 (=4.34) or 4.20 × 2 (=8.40) C1 (dep on at least M1) for small and (£)4.34 or (£8.40) **OR** Any other method considered equivalent. Figures may be truncated or rounded as long as their method is clear.

PMT

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
2		$7 \times 8^2 + 320 = 448 + 320$ 768p = 7.68	£7.68	3	M1 for 7×8^2 or 448 seen A1 for 768 B1 for correct monetary answer written correctly eg £7.68 or 768p
3			correct shaded region	2	B1 for arc of circle, centre oak tree, radius 4cm(±2mm) B1 for shaded region where the tree can be planted, bounded by sides of garden and arc of circle centre oak tree
4	(a)		−3, −2, −1,0,1	2	B2 for all 5 values and no others (B1 for 4 correct values and no others or -4,-3,-2,-1,0,1 or -3,-2,-1,0,1,2
	(b)		$-2 \leq x < 4$	2	B2 for $-2 \leq x < 4$ (B1 for $-2 \leq x$ or $x < 4$ or $-2 < x \leq 4$) [Note: accept the use of any letter other than x throughout and ignore any attempt to list integer values]
5			25.1-25.2	2	M1 for $2 \times 3.142 \times 4$ or 8π oe A1 for 25.1 – 25. 2
6	(a)		5, −4, −3	2	B2 for 5, −4 and −3 (B1 for 5 or −4 or −3)
	(b)		correct curve	2	B2 for fully correct curve (B1 ft for at least 5 points plotted correctly)

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
7			enlargement, scale factor 2, centre (6, −5)	3	B1 for enlargement B1 for (scale) factor 2 or ×2 or sf 2 B1 for (6, −5) NB: award 0 marks for an explanation that includes reference to more than one transformation.
*8		Cost of wages, rent and other expenses last year 92 000+10 800+7000=109 800 Cost of wages, rent and other expenses this year $1.075\times92\,000 + \frac{7}{9} \times 10$ 800 + 7000÷2 = 98 900+8400+3500 = 110 800 Costs are more this year, so she cannot increase the amount she spends on goods	no, with working and reason	4	M1 for an attempt to calculate using 7.5% and 92000 (eg 7.5% of 92000 or increase of 7.5%) eg 92000 + 6900 or 98900 M1 for an attempt to find a fractional amount of 10800 (eg $\frac{7}{9}$ × 10800 or 8400 or $\frac{2}{9}$ × 10800 or 2400) M1 for complete method of increasing 92000 by 7.5%, finding $\frac{7}{9}$ of 10800, and halving 7000, with at least all of these three added together. C1 for statement eg "no" with both 110800 & 109800 seen, OR "no" with "1000 more" OR "no" and compares 181800 with 182800, oe.
9	(a) (b)	$\sqrt{30} = 5.4772255\ldots$ $2.5^2 = 6.25$ 5.4772255...÷ 6.25 = 4.5 × 1000 × 1000	0.876 4 500 000	2 2	M1 for $\sqrt{30} \div 6.25$ or $5.4(7\ldots) \div 2.5^2$ A1 for any answer in the range 0.876 to 0.877 M1 for complete method equivalent to 4.5 × 1000 × 1000 A1 for 4 500 000 oe

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
10		3 → 21 4 → 56 3.1 → 23.5(9) 3.2 → 26.3(6) 3.3 → 29.3(3) 3.4 → 32.5(0) 3.5 → 35.8(7) 3.6 → 39.4(5) 3.7 → 43.2(5) 3.8 → 47.2(7) 3.9 → 51.(5) 3.35 → 30.8(9)	3.3	4	B2 for a trial between 3.3 and 3.4 inclusive (B1 for a trial between 3 and 4 inclusive) B1 for a different trial between 3.3 and 3.4 exclusive B1 (dep on at least one previous B1) for 3.3 NB trials should be evaluated to at least 1dp truncated or rounded
11			45	3	M1 for 60000 × 0.75 or ÷1000 or digits 45 or the numbers 60 or 0.00075) M1 for the complete method 60 000 × 0.75 ÷ 1000 oe A1 cao
12	(a)		−7	2	M1 for an attempt to get *x*s on one side or number terms on one side A1 cao
	(b)		8.5	2	M1 for an attempt to divide each side by 4 or an attempt to expand the bracket eg $4 \times x - 4 \times 5$ A1 for 8.5 oe

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
*13		2500×0.04×2 =£200 2500×$(1.039)^2$ = £2698.80 £2698.80 – £2500 = £198.80 Scheme A gives £1.20 more	statement	4	M1 for a correct method to find 4% of 2500 or 3.9% of 2500 A1 for a correctly calculated amount 2700 or 2698.8(0) or 200 or 198.8(0); or percentage rate calculated over the 2 years for comparison: 3.93% or 3.976% oe M1 for a correct compound interest method using 3.9% and 2500 over 2 years C1 for statement of scheme A, with two correct comparable figures.
14		$3x - 2y = 7$ $7x + 2y = 13$ $10x = 20$ $x = 2$ $3\times2 - 2y = 7$ $-2y = 1$ $y = -0.5$	2, −0.5	3	M1 for a correct process to eliminate either x or y (allow one arithmetic error) M1 (dep) for correct substitution of their found variable or an otherwise correct method to eliminate the other variable A1 for 2 and −0.5 oe
15	(a)		3.7×10^{-3}	1	B1 cao
	(b)		49 000	1	B1 cao
	(c)	500 ÷ 250 000 [= 0.002] OR $(5 \times 10^2) \div (2.5 \times 10^5)$ [= $(5\div2.5)\times10^{2-5}$] OR $\frac{500}{250} \times \frac{1}{10^3}$ [= 2×10^{-3}] OR $\frac{500\div100}{250000\div100} = \frac{5}{2500}$ [= 0.002]	2×10^{-3}	2	M1 for an attempt to divide using two numbers that are in consistent form eg following an attempt to convert to ordinary numbers, or standard form numbers. A1 for 2×10^{-3}

PMT

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
16			12.7	3	M1 for 3.142 × 5 × 5 oe or 3.142 × 5 × 5 × '*h*' (=78.5 – 78.55) M1 for 1000 ÷ (3.142 × 5 × 5) A1 for 12.7 – 12.8 NB: multiples of π acceptable for M marks
17		$4x - 3 = 2x + 2y$ $2x - 3 = 2y$ $2x = 2y + 3$ $x = y + 1.5$	$x = y + 1.5$	3	M1 for attempt to expand brackets eg $2\times x+2\times y$ or divide through by 2 (each term). M1 for attempt to get x on one side of equation. A1 for $(x =) y + 1.5$, $\frac{2y + 3}{2}$, oe
18			(11, 21)	3	M1 for 5 – 2 or 9 – 3 OR 3 or 6 as long as these are not related to the 3 from *A* or multiple 3 or 9,18 M1 for 2 + 3 × "3" and 3 + 3 × "6" OR for 5 + 2 × "3" and 9 + 2 × "6" OR for 2 + 3 × "3" (=11) and (y=) 2 × '11' – 1 A1 cao SC: B1 if no method shown and answer shows x coordinate as 11 or y coordinate as 21.

PMT

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
19			4.6	5	M1 for $AC = \sqrt{(3^2+4^2)}$ (=5) or $DA = \sqrt{(5^2+2^2)}$ M1 for tan BAC = 0.75 oe or (BAC=) 36.8(6) or 36.9 M1 for tan DAC = 0.4 oe or (DAC=) 21.8(0) M1 for $\sqrt{'5'^2+2^2}$ × sin("36.8(6)"+"21.8(0)") or 5.385 × sin 58.7 A1 for 4.59 – 4.61 OR M1 for $AC = \sqrt{3^2+4^2}(=5)$ or $DP = \sqrt{(29 - AP^2)}$ M1 for cos BAC = 0.8 oe or (BAC=) 36.8(6) oe M1 for angle CDP = angle BAC M1 for (DP=) 3 + 2×cos CDP or (AP=) 4 – 2×sin CDP A1 for 4.59 – 4.61
20			0.5, −3	3	M1 for $(2x \pm 1)(x \pm 3)$ M1 for $(2x - 1)(x + 3)$ A1 cao OR M1 for $\frac{-5 \pm \sqrt{5^2 - 4 \times 2 \times -3}}{2 \times 2}$ allow sub. of $c = \pm 3$ M1 for $\frac{-5 \pm \sqrt{49}}{4}$ A1 cao NB: any other method put in review.

PMT

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
21		$1180.5 \div \frac{4}{3}\pi(6.15)^3$	1.21	4	M1 for 1180 ± 0.5 or 6.2 ± 0.05 M1 for 'max. mass' ÷ 'min. volume' where "min volume" is defined as any expression involving 6.15 M1 for $1180.5 \div \frac{4}{3}\pi(6.15)^3$ A1 for 1.21 – 1.22
22			proof	3	B1 for *AD* = *AB* and *AE* = *AG* B1 for angle *EAB* = angle *DAG* (= 90 + angle *DAE*) B1 (dep on previous B marks) for 2 sides and included angle (SAS) oe
23	(a)		$3\boldsymbol{a} + \boldsymbol{b}$	1	B1 oe
	(b)		3**c**	4	M1 for writing *QB* as a combination of vectors eg $\frac{1}{2}$ ('3**a** + **b**') + **b** or 3**a** + 2**b** – $\frac{1}{2}$ ('3**a** + **b**') or 2**a** + 2**b** − **c** oe M1 for equating **a** + **b** = 2**c** or writing *PQ* in terms of vectors **a** and **b** eg −**a** + $\frac{1}{2}$ ('3**a** + **b**') or 2**a** + **b** – $\frac{1}{2}$ ('3**a** + **b**') A1 for (*PQ*=) $\frac{1}{2}$**a** + $\frac{1}{2}$**b or** (*QB*=) $\frac{3}{2}$**a** + $\frac{3}{2}$**b** or (*QB*=) 2**c**−**a** + **a** + **c** A1 cao NB: accept equivalent expressions throughout

PMT

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481
Email publication.orders@edexcel.com
Order Code UG035055 March 2013

For more information on Edexcel qualifications, please visit our website
www.edexcel.com

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual
Llywodraeth Cynulliad Cymru
Welsh Assembly Government
Rewarding Learning

PMT

edexcel

Mark Scheme (Results)

Summer 2013

GCSE Mathematics (Linear) 1MA0
Higher (Calculator) Paper 2H

PMT

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk for our BTEC qualifications.
Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

If you have any subject specific questions about this specification that require the help of a subject specialist, you can speak directly to the subject team at Pearson.
Their contact details can be found on this link: www.edexcel.com/teachingservices.

You can also use our online Ask the Expert service at www.edexcel.com/ask. You will need an Edexcel username and password to access this service.

Pearson: helping people progress, everywhere
Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2013
Publications Code UG037224
All the material in this publication is copyright
© Pearson Education Ltd 2013

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labeling conventions.

ii*) select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect canceling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

12 Parts of questions
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers
Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

Guidance on the use of codes within this mark scheme
M1 – method mark A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

PAPER: 1MA0_2H					
Question		**Working**	**Answer**	**Mark**	**Notes**
1			40.5	3	M1 for 1.5×6 or 1.5 ×1.5 M1 for adding area of 5 or 6 faces provided at least 3 are the correct area A1 cao NB: anything that leads to a volume calculation 0 marks.
*2			Not enough mincemeat since 600<700 OR Only able to make 38 mince pies since insufficient mincemeat	4	M1 for 45 ÷ 18 (= 2.5) M1 for 2.5 used as factor or divisor A1 for ingredients as 562.5 and 875 and 250 and 700 and 2.5 (accept 2 or 3) OR for availables as 400, 400, 200 240, 2.4 (accept 2 or 3) C1 ft (dep on at least M1) for identifying and stating which ingredient is insufficient for the recipe (with some supportive evidence) OR M1 for a correct method to determine the number of pies one ingredient could produce M1 for a correct method to determine the number of pies **all** ingredient could produce A1 for 80 and 51 and 90 and 38 and 108 C1 ft (dep on at least M1) for identifying and stating which ingredient is insufficient for the recipe. (with some supportive evidence)

PAPER: 1MA0_2H					
Question		**Working**	**Answer**	**Mark**	**Notes**
3	(a)		Points plotted at (1,8200) and (3.5,5000)	1	B1 for points accurately plotted ±1/2 square tolerance
	(b)		'the older the car the lower the value' 'the greater the value the newer the car'	1	B1 for an acceptable relationship eg. 'the older the car the lower the value' (accept 'negative correlation' but not just 'negative')
	(c)		5200 to 6600	2	M1 for a single line segment with negative gradient that could be used as a line of best fit or a vertical line from 2.5 or a point at (2.5,*y*) where *y* is from 5200 to 6600 A1 for given answer in the range 5200 − 6600
4			126	3	M1 for 1 – 0.05 – 0.32 (= 0.63) M1 for '0.63' × 200 A1 cao OR M1 for 0.05 × 200 (= 10) or 0.32 × 200 (= 64) or 0.37 × 200 (=74) M1 for 200 – '10' – '64' A1 cao OR M1 for 100 – 5 – 32 (= 63) M1 for $\frac{"63"}{100} \times 200$ A1 cao SC: B2 for $\frac{126}{200}$ as the answer.

PAPER: 1MA0_2H					
Question		**Working**	**Answer**	**Mark**	**Notes**
5	(a)		Response boxes overlap and are not exhaustive	2	B2 for **TWO** aspects from: No time frame given Non-exhaustive responses Response boxes over-lapping (B1 for ONE correct aspect)
	(b)		How many magazines do you buy each month? 0-4 5-8 over 8	2	B1 for a question with a time frame B1 for at least 3 correctly labelled response boxes (non-overlapping, need not be exhaustive) **or** for a set of response boxes that are exhaustive (could be overlapping) [Do not allow inequalities in response boxes]
	(c)		One reason	1	B1 for ONE reason Eg. All the same age, may all be males, may all like same types of magazines, sample too small, biased
6			4.8	4	M1 for 60 × 60 (=3600) M1 for 15000÷ 20 (=750) or 20÷15000 (=0.00133..) or "3600"÷15000 (=0.24) or 15000÷"3600" (=4.16..) M1 for "3600" ÷ (15000÷20) or "3600"×20÷15000 oe A1 cao

PMT

PAPER: 1MA0_2H					
Question		**Working**	**Answer**	**Mark**	**Notes**
7			28% or $\frac{14}{50}$	4	M1 for 100 30 (= 70) or $1 - \frac{3}{10}$ $(= \frac{7}{10})$ M1 for "70" ÷ (3 + 2) (= 14) or $"\frac{7}{10}" \div (3+2)$ $(= \frac{7}{50})$ M1 for "14" × 2 or $\frac{7}{50} \times 2$ A1 for 28% or $\frac{14}{50}$ oe **OR** M1 for a correct method to find (100-30)% of any actual sum of money M1 for "350" ÷ (3 + 2) (= 70) M1 for "70" × 2 A1 for 28% or $\frac{14}{50}$ oe **OR** M1 for starting with two numbers in ratio 3:2, eg 21 and 14 M1 for equating sum of their numbers to 100 – 30 (=70%), eg '21' + '14' (=35) M1 for scaling sum of their numbers to 100%, eg '35'÷70×100 (=50) A1 for 28% or $\frac{14}{50}$ oe SC: award B3 for oe answers expressed in an incorrect form eg $\frac{2.8}{10}$

PAPER: 1MA0_2H					
Question		**Working**	**Answer**	**Mark**	**Notes**
8			10752	4	M1 for splitting the pentagon (or show the recognition of the "absent" triangle) and using a correct method to find the area of one shape M1 for a complete and correct method to find the total area M1 (dep on at least one prev M1) for multiplying their total area by 2.56 (where total area is a calculation involving at least two areas) A1 cao
9			55	4	M1 for a correct method to find a different angle using 35° M1 for setting up a complete process to calculate angle x A1 cao B1 states one of the following reasons relating to their chosen method: Alternate angles are equal; Corresponding angles are equal; Allied angles / Co-interior angles add up to 180; the exterior angle of a triangle is equal to the sum of the interior opposite angles.

<table>
<tr><th colspan="5">PAPER: 1MA0_2H</th></tr>
<tr><th colspan="2">Question</th><th>Working</th><th>Answer</th><th>Mark</th><th>Notes</th></tr>
<tr><td>10</td><td></td><td>
<table>
<tr><th>x</th><th>$x^3 + 2x$</th></tr>
<tr><td>4</td><td>72</td></tr>
<tr><td>4.1</td><td>77.(121)</td></tr>
<tr><td>4.2</td><td>82.(488)</td></tr>
<tr><td>4.3</td><td>88.(107)</td></tr>
<tr><td>4.4</td><td>93.(984)</td></tr>
<tr><td>4.5</td><td>100.(125)</td></tr>
<tr><td>4.6</td><td>106.(536)</td></tr>
<tr><td>4.7</td><td>113.(223)</td></tr>
<tr><td>4.8</td><td>120.(192)</td></tr>
<tr><td>4.9</td><td>127.(449)</td></tr>
<tr><td>5</td><td>135</td></tr>
<tr><td></td><td></td></tr>
<tr><td>4.65</td><td>109.8(44625)</td></tr>
<tr><td>4.66</td><td>110.5(14696)</td></tr>
<tr><td>4.67</td><td>111.1(87563)</td></tr>
<tr><td>4.68</td><td>111.8(63232)</td></tr>
<tr><td>4.69</td><td>112.5(41709)</td></tr>
</table>
</td><td>4.7</td><td>4</td><td>B2 for a trial $4.6 \leq x \leq 4.7$ evaluated correctly
(B1 for a trial evaluated correctly for $4 \leq x \leq 5$)
B1 for a different trial evaluated correctly for $4.65 \leq x < 4.7$
B1 (dep on at least one previous B1) for 4.7

[Note: Trials should be evaluated to at least accuracy shown in table, truncated or rounded]

No working scores 0 marks</td></tr>
<tr><td>11</td><td></td><td></td><td>3.52</td><td>3</td><td>M1 for $1.35^2 + 3.25^2$
M1 (dep) for $\sqrt{(1.35^2 + 3.25^2)}$ $(= \sqrt{12.385})$
A1 for answer in the range 3.51 to 3.52</td></tr>
</table>

PMT

PAPER: 1MA0_2H					
Question		**Working**	**Answer**	**Mark**	**Notes**
12	(a)	$3x - 6 = x + 7$ $2x = 13$	6.5	3	M1 for $3\times x - 3\times 2$ ($=3x - 6$) or $\frac{x}{3} + \frac{7}{3}$ seen M1 for correct method to isolate the terms in x or the number terms on opposite sides of an equation A1 for 6.5 oe
	(b)	$2 - y = 1 \times 5$	-3	2	M1 for intention to multiply both sides by 5 (to give $2 - y = 1 \times 5$) A1 cao
13	(a)		(3, 3.5) oe	2	M1 for a correct method to find the value of either the x coordinate or the y coordinate of the midpoint **or** $x = 3$ **or** $y = 3.5$ A1 cao
	(b)		-1.8 oe	2	M1 for correct method to find the gradient OR (+)1.8 A1 for -1.8 oe

PAPER: 1MA0_2H					
Question		**Working**	**Answer**	**Mark**	**Notes**
*14			The Friendly Bank	4	M1 for a correct method to find interest for the first year for either bank **OR** correct method to find the value of investment after one year for either bank **OR** use of the multiplier 1.04 or 1.05 M1 for a correct full method to find the value of the investment (or the value of the total interest) at the end of 2 years in either bank A1 for 2100.8(0) and 2110.5(0) (accept 100.8(0) and 110.5(0)) C1 (dep on M1) ft for a correct comparison of *their* total amounts, identifying the bank from their calculations **OR** M1 for either 1.04 × 1.01 or 1.05 × 1.005 M1 for 1.04 × 1.01 and 1.05 × 1.005 A1 for 1.0504 and 1.05525 C1 (dep on M1) ft for a correct comparison of *their* total multiplying factors identifying the bank from their calculations

PMT

PAPER: 1MA0_2H					
Question		**Working**	**Answer**	**Mark**	**Notes**
15	(a)		-2 -1 0 1 2 3 4 **8** 3 0 **-1** **0** 3 **8**	2	B2 for 8, -1, 0, 8 (B1 for at least two of 8, -1, 0, 8)
	(b)		Correct curve	2	M1 (ft) for at least 5 points plotted correctly A1 for a fully correct curve
	(c)	$x^2 - 2x - 3 = 0$ OR $(x - 3)(x + 1) = 0$	3 and −1	2	M1 for the straight line $y = 3$ drawn to intersect the "graph" from (a) A1 for both solutions **OR** M1 for identifying $y = 3$ from the table A1 for both solutions **OR** M1 for $(x \pm 3)(x \pm 1)$ A1 for both solutions

PAPER: 1MA0_2H					
Question		**Working**	**Answer**	**Mark**	**Notes**
*16		Angle *POT* = 180 – 90 – 32 = 58 (angle between radius and tangent = 90° and sum of angles in a triangle = 180°) Angle *OST* =angle *OTS* = 58÷2 (ext angle of a triangle equal to sum of int opp angles and base angles of an isos triangle are equal) or (angle at centre = 2x angle at circumference) OR Angle *SOT* = 90 + 32 = 122 (ext angle of a triangle equal to sum of int opp angles) (180 – 122) ÷ 2 (base angles of an isos triangle are equal)	29	5	B1 for angle OTP = 90°, quoted or shown on the diagram M1 for a method that leads to 180 – (90 + 32) or 58 shown at *TOP* M1 for completing the method leading to "58"÷2 or 29 shown at *TSP* A1 cao C1 for "angle between radius and tangent = 90°" **and** one other correct reason given from theory used NB: C0 if inappropriate rules listed OR B1 for angle OTP = 90°, quoted or shown on the diagram M1 for a method that leads to 122 shown at *SOT* M1 for (180 – "122") ÷ 2 or 29 shown at *TSP* A1 cao C1 for "angle between radius and tangent = 90°" **and** one other correct reason given from theory used NB: C0 if inappropriate rules listed

PAPER: 1MA0_2H					
Question		**Working**	**Answer**	**Mark**	**Notes**
17	(a)		Box plot overlay	2	M1 for a box drawn with at least 2 correct points from LQ, Med and UQ A1 for a fully correct box plot
	(b)		Comparison of a measure of spread plus a comparison of medians (in context)	2	B1 for a correct comparison of a measure of spread (using either range or iqr) B1 for a correct comparison of medians For the award of both marks at least one of the comparisons made must be in the context of the question.
18		$3p^2 = y + 4$ $p^2 = \frac{y+4}{3}$	$p = \sqrt{\frac{y+4}{3}}$	3	M1 for clear intention to add 4 to both sides or divide all terms by 3(with at least 3 terms) M1 for clear intention to find the square root from p^2 = (expression in y) A1 for $p = \sqrt{\frac{y+4}{3}}$ oe (accept ± a correct root)
19	(a)		$3(2 + 3x)$	1	B1 for $3(2 + 3x)$
	(b)		$(y + 4)(y – 4)$	1	B1 for $(y + 4)(y – 4)$
	(c)		$(2p − 5)(p + 2)$	2	M1 for $(2p \pm 5)(p \pm 2)$ A1 for $(2p − 5)(p + 2)$

PAPER: 1MA0_2H					
Question		**Working**	**Answer**	**Mark**	**Notes**
20		cos y = 2.25 ÷ 6 y = cos^{-1} (2.25 ÷ 6) OR 6cos 75 = 1.55…	The ladder is not safe because y is not near to 75	3	M1 for cos y = 2.25 ÷ 6 oe M1 for cos^{-1} (2.25 ÷ 6) C1 for sight of 67-68 and a statement eg this angle is NOT (near to) 75° and so the ladder is not steep enough and so not safe. OR M1 for cos 75 = x ÷ 6 M1 for 6cos 75 C1 for sight of 1.55(29…) and a statement eg that 2.25 NOT (near to) 1.55 and so the ladder is not steep enough and so not safe.
21			48 or 49	2	M1 for $\frac{460}{460+320+165}\times 100$ (=48.67 ….) or $\frac{460}{9.5}$ or $\frac{460}{9.45}$ A1 for 48 or 49
22			1.33	3	M1 for $3.4 = \frac{k}{5^2}$ oe or 3.4×5^2 (=85) M1 for '3.4×5^2' ÷ 8^2 A1 for answer in range 1.32 to 1.33 or $\frac{85}{64}$
23		d: UB = 54.5 (or 54.499), LB = 53.5 C: UB = 170.5 (or 170.499), LB = 169.5 170.5 ÷ 53.5 169.5 ÷ 54.5	3.19 3.11..	4	B1 for any one correct bound quoted M1 for 170.5 ÷ 53.5 or 169.5 ÷ 54.5 A1 for UB = answer in range 3.18 to 3.19 from correct working A1 for LB = 3.11.. from correct working

PMT

PAPER: 1MA0_2H					
Question		**Working**	**Answer**	**Mark**	**Notes**
24	(a)		18.2	2	M1 for ½ × 6 × 7 × sin60 A1 for answer in range 18.1 to 18.2
	(b)		6.56	3	M1 for $6^2 + 7^2 - 2 \times 6 \times 7 \times \cos 60$ M1 for correct order of operation eg 36 + 49 – 42 (=43) A1 for answer in range 6.55 to 6.56
25			$x = 2.87$, $y = -0.87$ and $x = -0.87$, $y = 2.87$	6	M1 for $x^2 + (2 - x)^2 = 9$ M1 for $4 - 4x + x^2$ A1 for $2x^2 - 4x - 5 = 0$ oe 3 term simplified quadratic M1 for a correct method to solve their quadratic Eg $x = \frac{4 \pm \sqrt{(16 - 4 \times 2 \times -5)}}{4}$ A1 for $x = 2.87$, $y = -0.87$ or better A1 for $x = -0.87$, $y = 2.87$ or better Award marks for equivalent algebraic expressions. Apply the same scheme as above for y first.

PMT

Modifications to the mark scheme for Modified Large Print (MLP) papers.

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 1MA0_2H			
Question		**Modification**	**Notes**
1		Model as well as diagram provided	Standard mark scheme
3	(a)	*y*-axis 2 cm for 1000. *x*-axis 2 cm for ½ Table: 8200 changed to 8000	Points plotted at (1,8000) and (3.5,5000
8		Braille diagram labelled A B E D C	Standard mark scheme
12	(a)	MLP only: *x* changed to *y*.	Standard mark scheme
13		Crosses at A and B changed to a filled-in circle. Braille: AB is joined with a line.	Standard mark scheme

PAPER: 1MA0_2H			
Question		**Modification**	**Notes**
15		Table: wording added "There are four spaces to fill." *x*-axis 2 cm for ½. *y*-axis 2 cm for 1. Labelling remains same as on standard paper.	Standard mark scheme
17		Both box plots put on one page and labelled 'Girls' and 'Boys' Girls: LQ changed to 15, UQ changed to 40 Boys: figures changed to 5 20 35 40 50 2 cm grids.	Standard mark scheme
20		'horizontal ground' and 'vertical wall' labelled.	Standard mark scheme
24		Braille: Information about measurements given	Standard mark scheme

PMT

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481
Email publication.orders@edexcel.com

Order Code UG037224 Summer 2013

For more information on Edexcel qualifications, please visit our website
www.edexcel.com

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual
Llywodraeth Cynulliad Cymru
Welsh Assembly Government
Rewarding Learning

PMT

Mark Scheme (Results)

Summer 2016

Pearson Edexcel GCSE
In Mathematics B (2MB01)
Higher (Calculator) Unit 3

PMT

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2016
Publications Code 5MB3H_01_1606_MS
All the material in this publication is copyright
© Pearson Education Ltd 2016

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Note that in some cases a correct answer alone will not score marks unless supported by working; these situations are made clear in the mark scheme. Examiners should be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will award marks for the quality of written communication (QWC).
The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labelling conventions.

ii) *select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Partial answers shown (usually indicated in the ms by brackets) can be awarded the method mark associated with it (implied).
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks; transcription errors may also gain some credit. Send any such responses to review for the Team Leader to consider.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded (embedded answers).

12 Parts of questions
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers
Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

Guidance on the use of codes within this mark scheme
M1 – method mark A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

PAPER: 5MB3H/01					
Question		**Working**	**Answer**	**Mark**	**Notes**
1*			Decision (supported)	3	M1 for correct first step e.g. 15 ÷ 3 M1 for complete method C1 for correct decision with accurate figures
2	(a)		Correct diagram	2	B2 cao (B1 for correct orientation)
	(b)		Enlargement, sf 4, centre (0, 1)	3	B1 for enlargement B1 for scale factor 4 B1 for centre (0, 1)
3			$w = \frac{d+5}{2}$	2	M1 for attempt to add 5 to both sides or divide both sides by 2 as a first step, eg $d + 5 = 2w$ oe or $\frac{d}{2} = w - \frac{5}{2}$ oe A1 for $w = \frac{d+5}{2}$ oe

PMT

PAPER: 5MB3H/01					
Question		Working	Answer	Mark	Notes
4			22.5	4	M1 for $4(x - 8) = 2x + 13$ M1 for expansion of bracket or division of all terms by 4, eg $4x - 32 = 2x + 13$, or $x - 8 = \frac{2x}{4} + \frac{13}{4}$ M1 for isolating x and number terms eg $2x = 45$, $\frac{x}{2} = \frac{45}{4}$ A1 for $\frac{45}{2}$ or 22.5 OR M1 for $(180 - 64) \div 2$ (= 58) M1 for $4(x - 8)$ = "58" or $2x + 13$ = "58" or "58" – 13 (= 45) M1 for isolating x and number terms eg $4x = 90$, $2x = 45$ or "45" ÷ 2 A1 for $\frac{45}{2}$ or 22.5 OR M1 for $64 + 4(x - 8) + 2x + 13$ M1 for $64 + 4(x - 8) + 2x + 13 = 180$ M1 for isolating x and number terms eg $6x = 135$ A1 for $\frac{45}{2}$ or 22.5
5			Correct region shaded	3	M1 for a circle centre M or N (accept arc of sufficient length to define the region) M1 for circle centre M radius 5 cm **and** circle centre N radius 3 cm (accept arc of sufficient length to define the region) A1 for correct region shaded

PAPER: 5MB3H/01					
Question		Working	Answer	Mark	Notes
*6			Conclusion (supported)	4	M1 for $\pi \times 120^2$ (= 45 216 – 45 249) M1 for "$\pi \times 120^2$" ÷ 1800 A1 for 25 – 25.2 C1 ft (dep on M2) for appropriate conclusion from their figures OR M1 for $\pi \times 120^2$ (= 45 216 – 45 249) M1 for 20 × 1800 A1 for 36 000 and 45 216 – 45 249 C1 ft (dep on M2) for appropriate conclusion from their figures OR M1 for $\pi \times 120^2$ (= 45 216 – 45 249) M1 for "$\pi \times 120^2$" ÷ 20 A1 for 2260 – 2263 C1 ft (dep on M2) for appropriate conclusion from their figures OR M1 for 1800 × 20 M1 for 36 000 ÷ π (=11 457 – 11 465) A1 for 107(.0…) C1 ft (dep on M2) for appropriate conclusion from their figures
7			161.50	5	M2 for a correct method to decrease 6720 by 20%, eg 6720 × 0.8 (= 5376) or 6720 × 0.2 (= 1344 and 6720 – 1344 (= 5376)) (M1 for a correct method to find 20% of 6720 eg 6720 × 0.2 or $\frac{20}{100} \times 6720$ (= 1344)) M1 for subtracting 1500 (= 3876) after a percentage calculation M1 "3876" ÷ 24 after the subtraction of 1500 A1 for 161.5(0)

PMT

PAPER: 5MB3H/01

Question		Working		Answer	Mark	Notes
8		3	42	3.7	4	B2 for a trial between 3 and 4 exclusive (B1 for a trial at 3 or 4) B1 for a different trial of $3.7 < x \leq 3.75$ B1 (dep on at least one previous B1) for 3.7 NB. Trials should be evaluated to at least 2 sf truncated or rounded for values of x correct to 1 decimal place. Trials should be evaluated to at least 1 dp truncated or rounded for values of x correct to 2dp. **NB** No working scores 0 marks even if answer is correct
		3.1	45.(291)			
		3.2	48.(768)			
		3.3	52.(437)			
		3.4	56.(304)			
		3.5	60.(375)			
		3.6	64.(656)			
		3.7	69.(153)			
		3.8	73.(872)			
		3.9	78.(819)			
		4	84			
		3.71	69.6(14811)			
		3.72	70.(078848)			
		3.73	70.5(45117)			
		3.74	71.(013624)			
		3.75	71.4(84375)			
9				618	4	M1 for a method to find an area which is part of the cross section M1 (dep) for a complete method to find the total area of the cross section M1 (dep M1) for their cross sectional area × 12 A1 cao **OR** M1 for a method to find the volume of cuboid, 8 × 8 × 12 (= 768) M1 for a method to find the volume of the triangular prism, ½ ×5 × 5 × 12(= 150) M1 (dep M2) for a complete method for the volume of the prism A1 cao

5MB3H_01					
Question		Working	Answer	Mark	Notes
10	(a)		15.2	3	M1 for $12.7^2 + 8.3^2$ or $161.29 + 68.89$ or 230.18 M1 for √"$(12.7^2 + 8.3^2)$" or √"$161.29 + 68.89$" or $\sqrt{230.18}$ A1 for 15.2 or 15.17(168…)
	(b)		16.1	3	M1 for $\sin 40° = \frac{DE}{25}$ oe or $\frac{DE}{\sin 40} = \frac{25}{\sin 90}$ oe M1 for 25 sin 40° A1 for 16.1 or 16.06(969…)
11	(a)		4.081	2	M1 for 8.875 or $\frac{71}{8}$ or 2.979…or 4.08 A1 for 4.081 or 4.0809(5…)…
	(b)		4×10^{-3}	2	M1 for 4×10^n, $n \neq -3$ or $a \times 10^{-3}$, $a \neq 4$ or 0.004 or 0.4×10^{-2} or 1/250 A1 cao
12			$x < 2$	2	M1 for an attempt to isolate x and number terms or multiply all terms by 2, or for $\frac{3}{2}x < 3$ or $-\frac{3}{2}x < -3$ or $x = 2$ A1 cao
13	(a)		0, 6, 4, –6	2	B2 for all values correct (B1 for any one value correct)
	(b)		Graph drawn	2	M1 ft (provided B1 in (a) for at least 6 points plotted correctly from their table A1 cao for correct curve drawn from (–4, –6) to (3, –6)
	(c)		1.5 to 1.6 and –2.5 to –2.6	2	M1 for correct use $y = 2$ (may be implied by one correct estimate) or correct use of formula. A1 for 1.5 to 1.6 and –2.5 to –2.6

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
14*		Sher Yas **Old 19300 12560** New 20265 13502 **Inc 965 942**	Correct comparison (supported)	3	M1 for a correct method to calculate the population at the beginning of the 10 year period for at least one place. eg $\frac{20265}{1.05}$ (= 19 300) oe or $\frac{13502}{1.075}$ (= 12 560) oe A1 for 965 (Sherbury) **and** 942(Yaston) C1 (dep on M1) ft for statement comparing increases leading to conclusion based on two comparable amounts eg increase in Sherbury is greater than increase in Yaston
15		$20x + 10y = 35$ $6x - 10y = -48$ $x = -\frac{1}{2}$ $y = \frac{9}{2}$ OR $12x + 6y = 21$ $12x - 20y = -96$ $26y = 117$ $y = \frac{9}{2}$ $x = -\frac{1}{2}$	$-\frac{1}{2}, \frac{9}{2}$	4	M1 for a correct process to eliminate either variable (condone one arithmetic error) A1 cao for either x or y M1 (dep on M1) for correct substitution of found value into one of the equations or appropriate method after starting again (condone one arithmetic error) A1 cao OR M1 for full method to rearrange and substitute to eliminate either variable (condone one arithmetic error) A1 cao for either x or y M1 (dep on M1) for correct substitution of found value into one of the equations or appropriate method after starting again (condone one arithmetic error) A1 cao **NB** Trial and improvement methods score 0 marks unless both x and y are correct
16			226	2	M1 for angle $ABC = 180 - 67$ (= 113) A1 cao OR M1 for obtuse angle $AOC = 2 \times 67$ (= 134) A1 cao

5MB3H_01					
Question		Working	Answer	Mark	Notes
17			– 2.82, 19.82	3	M1 for $\frac{--17 \pm \sqrt{(-17)^2 - 4\times1\times-56}}{2\times1}$ (condone one sign error) or $\left(x - \frac{17}{2}\right)^2$ oe M1 for $\frac{17 \pm \sqrt{513}}{2}$ or for method leading to $\frac{17}{2} \pm \sqrt{\frac{513}{4}}$ oe A1 for answers in the range 19.82 to 19.825; – 2.82 to –2.825
18			21.2	4	M1 for correct use of sine rule to find *AB* or *BC*, eg $\frac{6}{\sin 25} = \frac{AB}{\sin 120}$ or $\frac{6}{\sin 25} = \frac{BC}{\sin(180-120-25)}$ M1 for (*AB* =) $\frac{6}{\sin 25} \times \sin 120$ (= 12.2(951…)) or (*BC* =) $\frac{6}{\sin 25} \times \sin 35$ (= 8.1(431…) M1(dep M2) for method to find area, eg 0.5 × 6 × "*AB*" sin35 or 0.5 × 6 × "*BC*" sin 120 A1 for answer in the range 21.1 – 21.2 OR (with perpendicular from *A* meeting *BC* extended at a point *X*) M1 for method to find *AX* M1 for method to find *BX* M1(dep M2) for 0.5 × "*AX*" × "*BC*" A1 for answer in the range 21.1 – 21.2
*19			0.38 because the LB and UB agree to that number of figures	5	B1 for 10.75 or 10.85 or 10.84999… B1 for 75.055 or 75.065 or 75.064999… M1 for $\sqrt{\frac{10.85}{75.055}}$ as UB **OR** $\sqrt{\frac{10.75}{75.065}}$ as LB A1 (dep on all previous marks) for both 0.3802… and 0.3784 must clearly come from working with correct values C1 for 0.38 from 0.3784… and 0.3802 … **and** "both LB and UB round to 0.38" **NB** An answer of 038(932….) without working scores no marks

PMT

5MB3H_01					
Question		Working	**Answer**	**Mark**	**Notes**
20			7.35	3	M1 for $y \propto \frac{1}{\sqrt{x}}$ or $y = \frac{k}{\sqrt{x}}$ or $9 = \frac{k}{\sqrt{4}}$ M1 for $k = 9\sqrt{4}$ (=18) A1 for 7.35 or 7.348…
*21			$\frac{2}{3}$ (7**a** + 3**b**) and correct conclusion	4	M1 for correct vector for $\overrightarrow{OY}$, or $\overrightarrow{AP}$ eg $(\overrightarrow{OY}) = \overrightarrow{OA} + \overrightarrow{AY}$, or $\overrightarrow{OY} = \overrightarrow{OA} + \frac{2}{3}\overrightarrow{AP}$ may include terms in **a** and **b,** eg. 6**a** + $\overrightarrow{AY}$, 4**b** + 4**a** – **b** + $\overrightarrow{PY}$ or $\overrightarrow{AP} = \overrightarrow{AO} + \overrightarrow{OB} + \overrightarrow{BP}$ or –6**a** + 4**b** + 4**a** – **b** or – 2**a** + 3**b** M1 for $(\overrightarrow{AY}) = \frac{2}{3}\overrightarrow{AP}$ or ⅔(–6**a** + 4**b** + 4**a** – **b**) or ⅔(–2**a** + 3**b**) or $(\overrightarrow{PY} = \frac{1}{3}\overrightarrow{PA}$ or ⅓(– 4**a** + **b** – 4**b** + 6**a**) or ⅓(2**a** – 3**b**) M1 for correct expression for $\overrightarrow{OY}$ in terms of **a** and **b,** eg $\frac{14}{3}$ **a** + 2**b** C1 for $\frac{2}{3}$ (7**a** + 3**b**) and "$\overrightarrow{OY}$ is parallel to the vector 7**a** +3**b**" **oe**

Modifications to the mark scheme for Modified Large Print (MLP) papers.

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q2	(a)	Grid enlarged. Shading changed to dotty shading. Labels added to shapes. Image labelled shape A. Rotation given and labelled shape B. Question changed to 'Describe fully the single transformation which maps shape A to shape B. One row top and bottom, one column left and right has been removed. Three answer lines given.	2 cao (B1 for correct orientation in third quadrant or correct quarter turn)
Q2	(b)	Grid enlarged. Shading changed to dotty shading. Labels added next to the triangles 'triangle P' and 'triangle Q'. One column removed from the left.	B1 for enlargement B1 for scale factor 4 B1 for centre (0, 1)

PMT

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q4		Diagram enlarged. MLP only – x changed to y. Braille only – measurements given in the text.	M1 for $4(x - 8) = 2x + 13$ M1 for expansion of bracket or division of all terms by 4, eg $4x - 32 = 2x + 13$, $x - 8 = 2x/4 + 13/4$ M1 for isolating x and number terms eg $2x = 45$, $x/2 = 45/4$ A1 for 45/2 or 22.5 OR M1 for $(180 - 64) \div 2$ (= 58) M1 for $4(x - 8) = 58$ or $2x + 13 = 58$ M1 for isolating x and number terms eg $4x = 90$, $2x = 45$ A1 for 45/2 or 22.5 OR M1 for $64 + 4(x - 8) + 2x + 13$ M1 for $64 + 4(x - 8) + 2x + 13 = 180$ M1 for isolating x and number terms eg $6x = 135$ A1 for 45/2 or 22.5
Q5		Diagram kept the same size as original. Scale moved above the diagram and also put in the question paper.	B1 for circle centre M radius 5 cm B1 for circle centre N radius 3 cm B1 for correct region shaded

PMT

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q6		Diagram enlarged. Cross changed to a filled in circle. Arrow heads removed.	M1 for $\pi \times 120^2$ (= 45238….) M1 for "$\pi \times 120^2$" ÷ 1800 A1 for 25.0 – 25.2 C1 for "surface is large enough as 25 > 20" oe OR M1 for $\pi \times 120^2$ (= 45238….) M1 for 20 × 1800 A1 for 36000 C1 for "surface is large enough as 36000 < 45238..." oe OR M1 for $\pi \times r^2 = 1800 \times 20$ M1 for 36000 ÷ π (=11459..) A1 for 107.0… C1 for "surface is large enough as 107…< 120" oe
Q9		Model provided for all candidates. Diagram enlarged and provided for MLP.	M1 for a method to find an area which is part of the cross section M1 (dep) for a complete method to find the total area of the cross section M1 (dep M2) for a complete method for the volume of the prism A1 cao OR M1 for a method to find the volume of cuboid, 8 × 8 × 12 M1 for a method to find the volume of the triangular prism M1 (depM2) for a complete method for the volume of the prism A1 cao

PMT

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q10	(a)	Diagram enlarged. Braille only: measurements given in the text.	M1 for $12.7^2 + 8.3^2$ or 161.29 + 68.89 or 230.18 M1 for $\sqrt{161.29 + 68.89}$ or $\sqrt{230.18}$ A1 for 15.2 or 15.17(168…)
Q10	(b)	Diagram enlarged. Braille only – measurements given in the text.	M1 for $\sin 40° = \frac{DE}{25}$ oe M1 for 25 sin 40° A1 for 16.1 or 16.06(969…)
Q13	(a)	Wording added 'There are four spaces to fill.'	B2 for all values correct (B1 for any one value correct)
Q13	(b)	Grid enlarged.	M1 ft for at least 6 points plotted correctly from their table A1 cao for correct curve drawn from (–4, –6) to (3, –6)
Q16		Diagram enlarged. Dot added at centre.	M1 for angle ABC = 180 – 67 (= 113) A1 cao OR M1 for obtuse angle AOC = 2 × 67 (= 134) A1 cao

PMT

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q18		Diagram enlarged.	M1 for correct use of sine rule to find *AB* or *BC*, eg $\frac{6}{\sin 25} = \frac{AB}{\sin 120}$ M1 for (*AB* =) $\frac{6}{\sin 25} \times \sin 120$ or (*BC* =) $\frac{6}{\sin 25} \times \sin 35$ M1(dep M2) for method to find area, eg 0.5×6×*AB*sin35 or 0.5×6×*BC* sin 120 A1 for answer in the range 21.1 – 21.2 OR (with perpendicular from *A* meeting *BC* extended at a point *X*) M1 for method to find *AX* M1 for method to find *BX* M1(dep M2) for 0.5 × "*AX*" × "*BC*" A1 for answer in the range 21.1 – 21.2
Q21		Diagram enlarged.	M1 for correct vector for $\overrightarrow{OY}$, eg $(\overrightarrow{OY}) = \overrightarrow{OA} + \overrightarrow{AY}$, may include terms in **a** and **b,** eg. 6**a** + $\overrightarrow{AY}$, 4**b** + 4**a** – **b** + $\overrightarrow{PY}$ M1 for $(\overrightarrow{AY} =) \frac{2}{3}\vec{A}P$ or $(\overrightarrow{PY} =) \frac{1}{3}\overrightarrow{PA}$ M1 for correct expression for $\overrightarrow{OY}$ in terms of **a** and **b,** eg $\frac{14}{3}$**a** + 2**b** C1 for $\frac{2}{3}$(7**a** + 3**b)** and "$\overrightarrow{OY}$ is parallel to the vector 7**a** +3**b" oe**

PMT

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London WC2R 0RL

Question 16

The majority of students did not know how to work with general iterative processes such as this. It was common to see students substituting in a succession of integers starting with 0 or 1. This typically brought up one correct value (1.817…) but didn't gain credit as it was not a correct process to find x_1. Of those who did correctly find the first value, most were able to go on to gain all three values. Some however went too far, and then put the wrong values as their answers (such as x_2, x_3 and x_4) and therefore lost the accuracy mark. Very few students had any idea how to complete part (b) with many either leaving it blank or just guessing random values.

Question 17

Clear algebraic working was lacking for a good number of candidates in part (a) and was the cause of them not gaining the mark, particularly not showing $x^4 = x^2 + 5$. Those that clearly showed each step in the rearrangement were rewarded.
In part (b) many were able to gain some credit for use of the iteration formula, but many went beyond the requested value of x_3 and lost the accuracy mark. Some candidates also found the values of x_1, x_2, and x_3, and then found the mean of these 3 values and gave this as their answer,

losing the accuracy mark. In terms of substitution, it is worth noting here, like with use of the quadratic formula, that ensuring the lid of the square root sign extends fully is important to show correct substitution.

Question 17

Another familiar style question, but performance suggests it is not one that many students had spent much time preparing for. Part (a) saw some awarded 1 mark, but very few were able to gain 2 marks as they didn't complete their explanation, normally missing out a statement about a change of signs. The rearrangement in part (b) only carried one mark, but again it seems many have little experience of this when factorisation is required, and poor incorrect algebra was commonplace. Part (c) only required substitution and evaluation, and it was not surprising to see that more students were able to gain marks in this part compared to the others. Many scoring 1, 2 or 3 marks. Those who gained 2 normally carried out extra iterations and therefore should be encouraged to read the question carefully to ensure they complete the correct number.