Chhetri Academy

chhetriacademy.com

GCSE MATHS

Geometry and measures

Edexcel 1MA1 | Subtopic collection Topic 39

Right-angled trigonometry

Higher tier

Answers and examiner comments

58	2003-2025	Higher
question groups	available papers	tier

Using this booklet

1. Answer numbers match the companion question booklet.
2. Follow the original question number on marking tables.
3. Examiner comments are included where available.

Name _____ **Date** _____

Some entries use full source marking material in an appendix. Any unavailable mark schemes are identified beside the relevant answer.

Original question 10

The full source mark scheme begins on page 94. Use the original paper code, tier and question number shown above.

Original question 14

The full source mark scheme begins on page 94. Use the original paper code, tier and question number shown above.

Original question 21

No matching mark scheme was available in the supplied archive for this question.

Original question 24

No matching mark scheme was available in the supplied archive for this question.

Question		Working	Answer	Mark	Notes
20	(a)	$\frac{90}{600}$	$\frac{3}{20}$	2	M1 $\frac{90}{600}$ A1 $\frac{3}{20}$ cao [SC: B1 for 0.15 or 15% if M0 scored]
	(b)	$\frac{180}{600}\times 100$ **OR** $\frac{180}{600}=\frac{30}{100}$	30	2	M1 $\frac{180}{600}\times 100$ A1 cao **OR** M1 $\frac{180}{600}=\frac{30}{100}$ or attempt to cancel to 100 A1 cao
	(c)	$600-(90+180)$ = 330 blue or green 330÷3	110	2	M1 [“$600-(90+180)$”]÷3 A1 cao [SC: B1 for an answer of 140 or 170 if M0 scored]

1380/1F

Question		Working	Answer	Mark	Notes
23		$\cos x = \frac{8.2}{10.6} = 0.77358\ldots$ $x = \cos^{-1}\frac{8.2}{10.6} = 39.323\ldots$	39.3	3	M1 for $\cos x = \frac{8.2}{10.6}$ or $\cos \frac{8.2}{10.6}$ M1 for $\cos^{-1}\frac{8.2}{10.6}$ A1 for 39.3 – 39.33 SC: M2A0 for 0.686 or 43.69 or 39.2 or 39.37… or 39.4

20	$\cos 58^\circ = \frac{AB}{16}$ $AB = 16 \times \cos 58^\circ = 8.4787$	8.48	3	M1 for $\cos 58^\circ = \frac{AB}{16}$ M1 (dep) for $16 \times \cos 58^\circ$ A1 for 8.47- 8.48 [SC:M2 A0 for 1.9-1.91 [RAD] or 9.8-9.81 [GRAD]]

1380_4H

Question	Working	Answer	Mark	Notes
21	AB = 8 cos 37° = 8 (0.7986...) = 6.389...	6.39	3	M1 for cos37 = $\frac{AB}{8}$ M1 for AB = 8 cos 37° or 6.4 seen (dep on 1st M1) A1 for 6.38 - 6.39 OR M1 for $\frac{AB}{Sin53} = \frac{8}{Sin90}$ M1 for $AB = \frac{8Sin53}{Sin90} AB$ or 6.4 seen (dep on 1st M1) A1 for for 6.38 - 6.39 SC M2A0 for 6.12 (radians) or 6.69 (grad)

13	(a)	$\sqrt{\frac{920-364.5661765}{0.046}}$ $= \sqrt{12074.64834...}$	109.88(47047)	2	B2 for 109.88(….. (B1 for 12074 seen …. or 555.43seen …or 109 to 110)
	(b)		110	1	B1 f.t. provided answer to (a) is more than 3 s.f.

1380_4H

Question		Working	Answer	Mark	Notes
25	(a)	$BC = \sqrt{8^2 - 3^2} = \sqrt{55} =$ 7.416198... CD = 7.416… ÷ sin 50° = 9.6811….	9.68	4	M1 for $8^2 - 3^2$ oe M1(dep) for $\sqrt{8^2 - 3^2}$ or 7.41… or 7.42 seen Or M1 for $A = \cos^{-1}\left(\frac{3}{8}\right)$ (=67.98°) M1(dep) for $3 \times \tan'67.98'$ or 7.41… or 7.42 seen M1 for '7.4……' ÷ sin 50 A1 for 9.67 – 9.69 SC B3 for -28.2 to -28.3 using rad or 10.4 – 10.5 using grad
	(b)	CE^2 =19^2+9.68^2 –2(19)(9.68) cos 40 = 361+93.7024–367.84(0.766) = 172.920612 CE = 13.1499…	13.1	3	M1 for (CE^2 =) 19^2 + '$(9.68)^2$' – 2 (19) '(9.68)' cos 40 M1(dep) for correct order of evaluation to reach $\sqrt{'172.920612'}$ A1 for 13.1– 13.15 SC B2 26.4(5805…) or 26.5 used radians or 12.5(3449…) used gradians

Question		Working	Answer	Mark	Notes
20	(a)	$\frac{8}{\sin 62}$	9.06	3	M1 for $\sin 62 = \frac{8}{PR}$ **or** $\cos(90\text{-}62) = \frac{8}{PR}$ **or** $\frac{\sin 90}{PR} = \frac{\sin 62}{8}$ oe M1 for (PR=) $\frac{8}{\sin 62}$ **or** $\frac{8}{\cos(90-62)}$ **or** $\sin 90 \times \frac{8}{\sin 62}$ A1 for 9.06 – 9.061 SC: B2 for -10.82 to -10.83 using rad **or** 9.672 to 9.674 using grad **or** For methods involving trig or Pythagoras and then trig or Pythag No marks until a correct trig or pythag statement linking SR = 4.25(36…) and PR For example M1 for $(PR^2=)\ 8^2 + 4.25(36\ldots)^2$ or $\cos 62 = \frac{4.25(36...)}{PR}$ M1 for $\sqrt{64 + 18.0(9...)}$ or $\frac{4.25(36...)}{\cos 62}$ A1 9.06 – 9.061

1380_4H					
Question		**Working**	**Answer**	**Mark**	**Notes**
20	(b)	$QR^2 = 14^2 + 9.06^2 - 2.\ 14.\ 9.06\ \cos 62$ $= 196 + 82.08 - 253.68\cos 62$ $= 158.98\ldots.$	12.6	4	B1 for angle $QPR = 62^o$ M1 for $QR^2 = 14^2 + \text{`}9.06\text{'}^2 - 2\times 14\times \text{`}9.06\text{'}\times \cos 62$ M1 for correct order of evaluation **or** 158.9… A1 (ft *PR*) for 12.6 – 12.62 **or** For methods using trigonometry and Pythagoras No marks until a correct Pythag statement with *QR* as only unknown (Let M be on *PQ* such that angle *RMQ* is 90°) For example B1 for angle $QPR = 62^o$ M1 for $(QR^2 =)\ 8^2 + (14 - PR\cos 62)^2$ M1 for $\sqrt{64+\text{'}94.995\ldots\text{'}}$ **or** 158.9… A1 (ft *PR*) for 12.6 – 12.62 SC: B3 for 10.3(5511) or 10.4 using rad **or** 11.6(402014) using grad

Question		Working	Answer	Mark	Notes
16		$\cos x = \frac{6.4}{9.6}$ $x = \cos^{-1}\frac{6.4}{9.6} =$	48.2	3	M1 for $\cos x = \frac{6.4}{9.6}$ or $\cos x = 0.66(6...)$ or $\cos x = 0.67$ M1 for $\cos^{-1}\frac{6.4}{9.6}$ or $\cos^{-1} 0.66(6...)$ or $\cos^{-1} 0.67$ A1 for 48.1 – 48.2 **OR** Correct use of Pythagoras and then trigonometry, no marks until M1 for $\sin x = \frac{'7.155'}{9.6}$ or $\tan x = \frac{'7.155'}{6.4}$ or $\sin x = \frac{'7.155'}{9.6} \times \sin 90$ or $\cos x = \frac{6.4^2 + 9.6^2 - '7.155'^2}{2 \times 6.4 \times 9.6}$ M1 for $\sin^{-1}\frac{'7.155'}{9.6}$ or $\tan^{-1}\frac{'7.155'}{6.4}$ or $\sin^{-1}\left(\frac{'7.155'}{9.6} \times \sin 90\right)$ or $\cos^{-1}\left(\frac{6.4^2 + 9.6^2 - '7.155'^2}{2 \times 6.4 \times 9.6}\right)$ A1 for 48.1 – 48.2 SC B2 for 0.841... (using rad) or 53.5... (using grad)

1MA0_2H

Question		Working	Answer	Mark	Notes
18		$BD^2 + 12^2 = 16^2$ oe BD= $\sqrt{256-144}$ (=10.58...) sin 40 = $\frac{'10.58'}{CD}$ $CD = \frac{'10.58'}{\sin 40}$	16.5	5	M1 for $BD^2 + 12^2 = 16^2$ oe or $16^2 - 12^2$ or 112 seen M1 for $\sqrt{256-144}$ or $\sqrt{112}$ (=10.58...) M1 for sin 40 = $\frac{'10.58'}{CD}$ or cos 50 = $\frac{'10.58'}{CD}$ M1 for (CD =) $\frac{'10.58'}{\sin 40}$ or $\frac{'10.58'}{\cos 50}$ A1 for 16.4 – 16.5 **OR** M1 for $BD^2 + 12^2 = 16^2$ oe or $16^2 - 12^2$ or 112 seen M1 for $\sqrt{256-144}$ or $\sqrt{112}$ (=10.58..) M1 for (BC =) '10.58'× tan 50 or $\frac{'10.58'}{\tan 40}$ (=12.6…) M1 for $\sqrt{'12.6'^2+'10.58...'^2}$ A1 for 16.4 – 16.5

1MA0_2H

Question		Working	Answer	Mark	Notes
14		$AB = 5 \quad \sin 36 = \frac{5}{AD}$ $AD = \frac{5}{\sin 36}$ Or $\sin 36 = \frac{5}{BC}$ $BC = \frac{5}{\sin 36}$ $AD = BC$ OR $\cos 54 = \frac{5}{BC}$ $BC = \frac{5}{\cos 54}$	8.51	4	B1 $AB = 5$ M1 $\sin 36 = \frac{5}{AD}$ or $\frac{\sin 36}{5} = \frac{\sin 90}{AD}$ M1 $AD = \frac{5}{\sin 36}$ or $AD = \frac{5 \sin 90}{\sin 36}$ A1 8.5 – 8.51 OR M1 $\sin 36 = \frac{5}{BC}$ or $\frac{\sin 36}{5} = \frac{\sin 90}{BC}$ M1 $BC = \frac{5}{\sin 36}$ or $BC = \frac{5 \sin 90}{\sin 36}$ B1 $AD = 'BC'$ A1 8.5 – 8.51 OR B1 angle $DCB = 54$ or angle $DBC = 36$ M1 $\cos 54 = \frac{5}{BC}$ M1 $BC = \frac{5}{\cos 54}$ A1 8.5 – 8.51 NB Other methods such as tan + Pythagoras must be complete methods and will earn M2

5MB3H_01

13	(a)		11.5	3	M1 for $13^2 - 6^2$ or $169 - 36$ or 133 M1 (dep on M1) for $\sqrt{"13^2 - 6^2"}$ **or** $\sqrt{133}$ A1 for answer in the range 11.5 − 11.6
	(b)		47.2	3	M1 for cos (*RPQ*)= $\frac{17}{25}$ oe **OR** sin *PQR* = $\frac{17}{25}$with *PQR* clearly identified M1 for (*RPQ* =+) $\cos^{-1}\frac{17}{25}$oe **OR** *PQR* = $\sin^{-1}\frac{17}{25}$with *PQR* clearly identified A1 for answer in the range 47.1 − 47.2 SC : B2 for an answer of 0.823(033...) **or** 52.3(95...) **or** 52.4

1MA0_2H

Question		Working	Answer	Mark	Notes
17		$\sqrt{\frac{2.73\ldots}{0.732\ldots}}$	1.931851…	2	M1 for 2.73… or 0.732…or 3.73…or 1.931 or 1.932 or 1.93 or $(1+\sqrt{3})$ or $(\sqrt{3}-1)$ or $(2+\sqrt{3})$ **or** 1.65... **or** 0.855... A1 for 1.9318(5…) SC: B1 for 2.5127(17...)

Question		Working	Answer	Mark	Notes
20			15.0	3	M1 for $8^2 + 8^2 - 2 \times 8 \times 8 \times \cos 140$ M1 (dep) for correct order of evaluation **or** 226.(05...) A1 for answer in range 15.0 − 15.04 **OR** M1 for $\frac{PR}{\text{SIN } 140} = \frac{8}{\sin\left(\left(\frac{180-140}{2}\right)\right)}$ M1 for $PR = \frac{8}{\sin\left(\left(\frac{180-140}{2}\right)\right)} \times \sin 140$ A1 for answer in range 15.0 − 15.04 **OR** M1 for $8 \times \sin 70$ or $8 \times \cos 20$ M1 for $2 \times 8 \times \sin 70$ **or** $2 \times 8 \times \cos 20$ A1 for answer in range 15.0 − 15.04

1MA0_2H

Question		Working	Answer	Mark	Notes
20		cos y = 2.25 ÷ 6 y = $\cos^{-1}$ (2.25 ÷ 6) OR 6cos 75 = 1.55…	The ladder is not safe because y is not near to 75	3	M1 for cos y = 2.25 ÷ 6 oe M1 for $\cos^{-1}$ (2.25 ÷ 6) C1 for sight of 67-68 and a statement eg this angle is NOT (near to) 75° and so the ladder is not steep enough and so not safe. OR M1 for cos 75 = x ÷ 6 M1 for 6cos 75 C1 for sight of 1.55(29…) and a statement eg that 2.25 NOT (near to) 1.55 and so the ladder is not steep enough and so not safe.

Question		Working	Answer	Mark	Notes
10	(a)	$24.5^2 + 10.6^2$ (= 712.61) $\sqrt{712.61}$	26.7	3	M1 for (GJ^2 =) $24.5^2 + 10.6^2$ or 600.25 + 112.36 or 712.61 M1 for $\sqrt{24.5^2 + 10.6^2}$ or $\sqrt{712.61}$ A1 for answer in the range 26.69 – 26.7
	(b)	$\cos x = \frac{7}{18}$ $x = \cos^{-1}\left(\frac{7}{18}\right)$	67.1	3	M1 for $\cos(x) = \frac{7}{18}$ oe M1 for $(x =) \cos^{-1}\left(\frac{7}{18}\right)$ or $\cos^{-1}(0.388...)$ or $\cos^{-1}(0.38)$ A1 for answer in the range 67.1 – 67.17 SC: B2 for an answer of 1.1(713...) or 1.2 or 74.5(717...) or 74.6

PAPER: 5MB3H_01

PAPER: 1MA0_2H

Question	Working	Answer	Mark	Notes
18		49.5	4	M1 for $\tan 54 = \frac{\text{height}}{6}$ M1 for (height =) 6 × tan54 (=8.2-8.3) M1 for $\frac{1}{2}$ × '8.258..' × 12 A1 for 49.2 - 50 **OR** M1 for $\cos 54 = \frac{6}{AC}$ M1 for (AC =) $\frac{6}{\cos 54}$ (=10.2(07...)) M1 for $\frac{1}{2}$ × 12 × '10.207' × sin54 A1 for 49.2 - 50 **OR** M1 for $\frac{AC}{\sin 54} = \frac{12}{\sin 72}$ M1 for (AC =) $\frac{12}{\sin 72}$ × sin54 (=10.2(07...)) M1 for $\frac{1}{2}$ × 12 × '10.207' × sin54 A1 for 49.2 – 50

PAPER: 1MA0_2H

11		66.7	3	M1 for tan (y =) $\frac{86}{37}$ (= 2.3243..) M1 (dep) for $\tan^{-1}$ "2.32(43..)" = or $\tan^{-1}$ $(\frac{86}{37})$ (accept 'shift tan' or 'inv tan' for $\tan^{-1}$) A1 for answer in the range 66.6° to 66.8° [SC: B1 for an answer in the range 23.2 to 23.3 if M0 scored]

11		66.7	3	M1 for tan (y =) $\frac{86}{37}$ (= 2.3243..) M1 (dep) for $\tan^{-1}$ "2.32(43..)" = or $\tan^{-1}$ $(\frac{86}{37})$ (accept 'shift tan' or 'inv tan' for $\tan^{-1}$) A1 for answer in the range 66.6° to 66.8° [SC: B1 for an answer in the range 23.2 to 23.3 if M0 scored]

11			9.54	4	M1 for $10^2 - 5^2$ (= 75) or (*BD* =) 10 × cos 30 (= 8.66...) M1 for "75" + 4^2 (= 91) or "8.66..."^{2} + 4^2 (= 91) M1 for $\sqrt{(10^2 - 5^2 + 4^2)}$ or $\sqrt{(\text{"8.66..."}^2 + 4^2)}$ A1 for 9.53 − 9.54

PAPER: 5MB3H_01

19		BC = $\frac{12}{\tan 60}$ = 6.92(8…) DE = 6.92(…) × tan 30 = 4 CE = 12 + 4 AC = $\frac{12}{sin60}$ = 13.8(5…) CE = $\frac{13.8(5\ldots)}{cos30}$	16 with supporting working	4	M1 for a method to find BC or AC or AD B1 for angle EAD = 30˚ or AED = 60° or ACD = 30˚ or CAD = 60° M1 for a method to find CE A1 for 15.9-16.1 with supporting working

PAPER: 1MA0_2H

21		Eg $BQ=\sqrt{(5^2+6^2)}=\sqrt{61}$ Angle $ABQ=\tan^{-1}(6\div5)=50.194^o$ $BP=\sqrt{61}\div\cos30=9.018$ cm Angle $PBC=90–30–50.194^o=9.806^o$ $BC=9.018\times\cos9.806=8.886...$ **OR** $BQ=\sqrt{(5^2+6^2)}=\sqrt{61}$ $PQ=\sqrt{61}\tan30=4.509$ Angle $BQA=\tan^{-1}(5\div6)=39.806$ Angle $PQD=180–90–39.806=50.194^o$ $QD=4.509\times\cos50.194=2.886...$	8.89	5	M1 for correct use of Pythagoras M1 for correct use of appropriate trig ratio M1 for correct use of appropriate inverse trig ratio M1 for correct method to calculate angle *CBP* or *PQD* A1 for 8.88 – 8.89

PAPER: 1MA0_2H

Question		Working	Answer	Mark	Notes
17			30.1	4	M1 for a correct trigonometric statement to find an unknown angle, eg. $\sin(30+x)$ or $\cos A = \frac{10.4+5.2}{18}$ or $\frac{\sin ADC}{18} = \frac{\sin 30}{10.4}$ M1 for a complete method to find the angle, eg. $\sin^{-1}\left(\frac{10.4+5.2}{18}\right)$ (= 60.07...) or $\cos^{-1}\left(\frac{10.4+5.2}{18}\right)$ (= 29.92..) or $\sin^{-1}\left(\frac{18 \times \sin 30}{10.4}\right)$ (= 59.92.. or 180 – 59.92.. = 120.07..) M1 (dep on M2) for a fully complete method to find angle x, eg. "60.07.". – 30 or 60 – "29.92.." or 90 – "59.92.." A1 for answer in the range 30.07 to 30.1 OR M1 for $(BC^2=)\ 18^2 - (10.4 + 5.2)^2$ or $BC^2 + (10.4 + 5.2)^2 = 18^2$ M1 for $(BC =)\ \sqrt{18^2 - (10.4 + 5.2)^2}$ (= 8.97...) M1 (dep on M2) for a fully complete method to find angle x, eg. $\tan^{-1}\left(\frac{5.2}{"8.97..."}\right)$ A1 for answer in the range 30.07 to 30.1

PAPER: 1MA0_2H

Original question 10

The full source mark scheme begins on page 148. Use the original paper code, tier and question number shown above.

Question	Working	Answer	Mark	Notes
15		28.9	5	M1 for $\sin 62 = \frac{BD}{15}$ **or** $\frac{BD}{\sin 62} = \frac{15}{\sin 90}$ oe M1 for (BD =) $15 \times \sin 62$ **or** $\frac{15}{\sin 90} \times \sin 62$ oe (= 13.24...) M1 for $\tan BCD = \frac{"13.24"}{24}$ oe **or** $\tan BDC = \frac{24}{"13.24"}$ with BDC clearly identified M1 for $BCD = \tan^{-1}\frac{"13.24"}{24}$ oe **or** $BDC = \tan^{-1}\frac{24}{"13.24"}$ with BDC clearly identified A1 for 28.8 – 28.9 **OR** M1 for $\cos(90 - 62) = \frac{BD}{15}$ M1 for (BD =) $15 \times \cos(90 - 62)$ (= 13.24...) M1 for $\tan BCD = \frac{"13.24"}{24}$ oe **or** $\tan BDC = \frac{24}{"13.24"}$ with BDC clearly identified M1 for $BCD = \tan^{-1}\frac{"13.24"}{24}$ oe **or** $BDC = \tan^{-1}\frac{24}{"13.24"}$ with BDC clearly identified A1 for 28.8 – 28.9

PAPER: 1MA0/2H

22	cos PBQ= $\frac{10^2+10^2-x^2(2-\sqrt{3})}{200}$ $= \frac{200-x^2(2-\sqrt{3})}{200}$	Proof	B1 M1 M1 M1 A1	(indep) for stating $\cos 30 = \frac{\sqrt{3}}{2}$ for $PQ^2 = 10^2 + 10^2 - 2 \times 10 \times 10 \times \cos PBQ$ **or** $AC^2 = x^2 + x^2 - 2 \times x \times x \times \cos 30$ $(=x^2(2-\sqrt{3}))$ oe for $\cos PBQ = \frac{10^2+10^2-PQ^2}{2\times10\times10}$ (implies previous M1) for $\cos PBQ = \frac{10^2+10^2-(x^2+x^2-2\times x\times x\times\cos 30)}{2\times10\times10}$ conclusion of proof with all working seen

Modifications to the mark scheme for Modified Large Print (MLP) papers

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 1MA1_1H			
Question		**Modification**	**Mark scheme notes**
1		Diagram enlarged. Right axis has been labelled. Crosses have been changed to solid circles.	Standard mark scheme but in (c) accept 12 to 14
4		Diagram enlarged. Arrows removed. Dashed lines at the top and left of the square extended. Dashes made longer and thicker.	Standard mark scheme
5		Diagram enlarged. Left hand side and top of shape labelled as well. Wording added 'The marked angles are right angles.' Braille only: will add labels *A B C D* etc. and information about the diagrams.	Standard mark scheme
10		*x* has been changed to *y*	Standard mark scheme with *x* replaced by *y*
11		Diagram enlarged.	Standard mark scheme if on 2mm grid, otherwise apply greater tolerance.
13		Table turned to vertical format.	Standard mark scheme
15		Only changes are to the formula box for a cone: diagram enlarged, dashes made longer and thicker, arrow heads changed to open headed arrows.	Standard mark scheme
18		Diagram enlarged. Dashed line added from *B* to *D*	Standard mark scheme
19		Diagram enlarged. Diagonal line added from *A* to *C* and *X* has been marked on the line as the midpoint.	Standard mark scheme
21		Diagram enlarged. Angle arcs made smaller.	Standard mark scheme

Question		Modification	Mark scheme notes
22		Diagram enlarged. Labels '*x* cm' added to the left and right side of the diagram. Dotted lines added between *B P* and *B Q*.  	Standard mark scheme

7		5.86	M1 A1	for $\sin 23 = \frac{AB}{15}$ NB Allow any alternative equivalent method to form an equation in AB 5.8 to 5.9

Question	Working	Answer	Mark	Notes
18	Note *DOC*=*DOA*, *ADO*=*CDO*	21.6	P1 P1 P1 P1 A1	Recognises that *OAD* or *OCD* is 90° or right angle for using trigonometry to set up an equation in *DOA* or *ADO* eg Cos *DOA* = $\frac{5}{9}$ for using inverse trigonometry to find *DOA* or *ADO* eg *DOA* = $\text{Cos}^{-1}\frac{5}{9}$ (= 56.25…) for a complete process to find arc length *ABC* or *AC* eg $\frac{360-2\times"56.25.."}{360}\times2\times\pi\times5$ (=21.598..) or $\frac{2\times"56.25.."}{360}\times2\times\pi\times5$ (=9.8174..) for answer in the range 21.5 to 21.65

Question	Working	Answer	Mark	Notes
7		32.3	P1	for using Pythagoras to find length of third side of triangle, eg $7.5^2 - 6^2$ or $6^2 + x^2 = 7.5^2$ or uses trigonometry to find angle in triangle, eg $\sin A = \frac{6}{7.5}$ or $\cos B = \frac{6}{7.5}$
			P1	(dep P1) for complete process to find length of third side of triangle eg $\sqrt{7.5^2 - 6^2}$ or $\sqrt{56.25 - 36}$ or $\sqrt{20.25}$ (= 4.5) or uses trigonometry to find base length of triangle, eg 7.5 × cos "A" or 7.5 × sin "B" or $\frac{6}{\tan "A"}$
			P1	(dep P2) for 24 – 10 – "4.5" (= 9.5)
			P1	(indep) for process to find angle CDA, eg $\tan CDA = \frac{6}{\text{base}}$ from right- angled triangle
			A1	for answer in the range 32.2 to 32.3

8 (a)		2.7560…	M1	for 1.0654(059…), 0.1402(633…), 7.5957(541…), 2.756 truncated or rounded to no less than 2dp
			A1	for 2.7560(….)
(b)		2.76	B1	for 2.76 ft from (a)

Paper: 1MA1/2H

Question	Working	Answer	Mark	Notes
20		68.5	B1	for angle $OAB = 90°$ or angle $OCB = 90°$, may be seen on diagram
			P1	for a process to find the length of AB or the length of CB ($= 10\sqrt{3}$ oe) eg $10 \times \tan 60°$ ($= 17.3...$) or the length of OB ($= 20$), eg $10 \div \cos 60°$
			P1	for a process (dep previous P1) to find the area of the triangle OAB ($= 50\sqrt{3}$ oe) or area of triangle OCB ($= 50\sqrt{3}$ oe) or area of kite $OABC$ ($= 100\sqrt{3}$ oe)
			P1	for a process to find the area of the sector OAC e.g. $\frac{1}{3} \times \pi \times 10^2$ ($= 104.7...$),accept rounded or truncated to 3 significant figures or more
			A1	for 68.4 − 68.6

Paper: 1MA1/2H

18	39.5	P1	for a start to a process eg, for a correct trigonometric statement, eg $\sin 48 = \frac{7.3}{AC}$ **or** $\cos 42 = \frac{7.3}{AC}$ **or** $\frac{AC}{\sin 90} = \frac{7.3}{\sin 48}$ **OR** angle *CAH* unambiguously identified on a diagram	Must include correct values
		P1	for a complete correct process to find *AC*, eg $(AC =) \frac{7.3}{\sin(48)}$ (=9.8..) **or** $(AC =) \frac{7.3}{\cos(42)}$ (=9.8..) **or** $(AC =) \sin 90 \times \frac{7.3}{\sin 48}$ (=9.8..)	
		P1	for a correct statement using angle *CAH*, eg $\tan(CAH) = \frac{8.1}{"9.8..."}$ **OR** $\sqrt{8.1^2 + "9.8"^2}$ (=12.7…) **and** $\frac{\sin CAH}{8.1} = \frac{\sin 90}{"12.7"}$	
		A1	for answer in the range 39.5 – 39.51	If an answer is given in the range but then incorrectly rounded award full marks.

Paper: 1MA1/2H

5	(a)	50.5	M1	for cos $ABC = \frac{7}{11}$ (0.63…) oe	Must be a complete statement for cos, sin or tan with all three elements present.
			A1	for answer in the range 50.4 to 50.51	If an answer is in the range 50.4 to 50.51 is given in the working space then incorrectly rounded, award full marks.
	(b)	Increase (supported)	C1	States increase with supporting reason eg " $\frac{7}{10}$ is greater than $\frac{7}{11}$" " 0.636 is less than 0.7" …."cos increases as angle decreases" "decreasing the denominator increases the value of the fraction" "angle is now 45.6" (accept 45.5 – 45.6)	If figures are given they must be correct (truncated or rounded).

Paper: 1MA1/3H

Question	Answer	Mark	Mark scheme	Additional guidance
19	$\frac{5}{3}$	P1	for process to derive an equation in x, eg $\frac{x}{4x-1}=\frac{6x+5}{12x+31}$	
		P1	for complete process to remove fractions, eg $x(12x + 31) = (6x + 5)(4x – 1)$	Must be correct use of brackets
		P1	for process to reduce to a quadratic equation, eg $12x^2 – 17x – 5 = 0$	Award for correct LHS only.
		P1	for process to solve the quadratic equation by factorisation or use of quadratic formula, eg $(4x + 1)(3x – 5) = 0$	Award for correct LHS only. Accept substitution into the formula; $\frac{--17\pm\sqrt{(-17)^2-4\times12\times-5}}{2\times12}$
		A1	for $\frac{5}{3}$ oe	Accept answers in the range 1.66 to 1.67 as equivalent

Paper: 1MA1/3H

8	(a)	1	B1	cao	
	(b)	8	M1	starts process, eg $\cos(60) = \frac{4}{x}$ or $0.5 = \frac{4}{x}$ oe or $\sin 30 = \frac{4}{x}$ or $\frac{\sin 30}{4} = \frac{\sin 90}{x}$ oe	All three elements of cos, 4, x must be present in an equation. eg cos = 4/x is acceptable but cos(4/x) is insufficient
			A1	cao	

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
13	84.9	P1	shows a process to find the radius or diameter eg $44 = 2 \times \pi \times r$ **or** $r = \frac{22}{\pi}$ **or** $d = \frac{44}{\pi}$ **or** $r = 7.0028$ **or** $d = 14.0056..$	Allow r in the range 7 to 7.1 and d in the range 14 to 14.1 Could be shown on the diagram.
		P1	(dep on P1) complete method to find the area eg ½ × "d" ^{2} × Sin60 oe, ½ × 14 × Tan60 oe, ½ × 14 × $\sqrt{14^2 - 7^2}$ oe	
		A1	for answer in the range 84.8 to 85	If the correct answer in the range is given in working and then rounded incorrectly award full marks.

Question	Answer	Mark	Mark scheme	Additional guidance
6	17.3	P1	for full process to find either angle eg (180 − 90) ÷ (2+3) ×2 (=36) **or** for 36 or 54 seen as an angle	May be seen on diagram Condone correct values if incorrectly placed.
		P1	for a correct equation using trigonometry eg cos [*A*] = 14÷ *AB*	This must be shown as an equation with all four elements (eg cos, [*A*], 14, *AB*) present. [*A*] could be 36 or any angle clearly and unambiguously identified as *A*. This also applies to [*B*] with Sine.
		P1	(dep previous P mark) for rearranging their trigonometry equation to make *AB* the subject eg (*AB* =) "14 ÷ cos 36"	
		A1	for an answer in the range 17.3 to 17.4	If an answer is shown in the range in working and then incorrectly rounded award full marks.

Paper: 1MA1/3H

Question	Answer	Mark	Mark scheme	Additional guidance
12	73.6	P1	for correct initial use of Pythagoras eg $5^2 + 5^2$ (=50) **or** a trigonometric ratio in the form $\frac{5 \div 2}{0.5AC}$ = sin 45 oe	
		P1	for finding the length of half of the diagonal eg $\sqrt{"50"} \div 2$ (= 3.5...) **or** $0.5AC = \frac{5 \div 2}{\sin 45}$ (=3.5...) oe	do not accept $\sqrt{20} \div 2$
		P1	for process to use tan eg tan *TAC* = (12 ÷ "3.5..") (=3.3..) **or** complete alternative method arriving at an equation with the subject as sin *TAC* or cos *TAC*	
		A1	for an answer in the range 73.58 to 74.1	

Question	Answer	Mark	Mark scheme	Additional guidance
14	$\frac{1}{2}$	M1	for $\frac{1}{\sqrt{3}} \times \frac{\sqrt{3}}{2}$ **or** $\frac{\sqrt{3}}{3} \times \frac{\sqrt{3}}{2}$ **or** $(\frac{1}{2} \div \frac{\sqrt{3}}{2}) \times \frac{\sqrt{3}}{2}$ **OR** $\tan 30 = \frac{1}{\sqrt{3}}$ oe **or** $\sin 60 = \frac{\sqrt{3}}{2}$	
		A1	for $\frac{1}{2}$ or 0.5	

Paper: 1MA1/2H

Question	Answer	Mark	Mark scheme	Additional guidance
5	9.85	M1	for $\sin(38) = \frac{AB}{16}$ oe **or** alternative method to find *AB*	
		A1	for an answer in the range 9.76 to 9.92	

Question	Answer	Mark	Mark scheme	Additional guidance
7	0.319	M1 A1	for partial method eg 1.70(499…) or 16.74 or $\frac{837}{50}$ or 0.101(8516...) or 0.102 or 0.32 for 0.319(1419…)	Accept 0.319 or better. Condone incorrect digits after the 0.319; isw incorrect rounding if 0.319(1419…) is shown in working.

14	36	P1 P1 P1 A1	for process to find an expression for the area of triangle eg $\frac{1}{2} \times 24 \times AE \times \sin 30$ (= $6AE$) (dep P1) for process to link the area of rectangle with the area of the triangle eg $2 \times \frac{1}{2} \times 24 \times AE \times \sin 30$ (= $12AE$) or for $AB = 12$ (indep) for use of given ratio eg $AE = 3AB$ oe, eg area of rectangle = $AE \times AB = 3x \times x$ cao	Accept any correct expression, eg $\frac{1}{2} \times 24 \times y \times \sin 30$ May be shown on the diagram by labelling AE and AB with, for example, $3x$, x or x, $\frac{1}{3}x$ or $\frac{3}{4}x$, $\frac{1}{4}x$ Do not accept 3, 1 or 1, $\frac{1}{3}$ or $\frac{3}{4}$, $\frac{1}{4}$ for this mark.

Paper: 1MA1/3H

12	32.1	P1	starts process, eg $\sin 40 = \frac{DB}{8.6}$ oe or for 8.6 × sin 40 (=5.52797…)	Accept values rounded or truncated to 2 dp.
		P1	complete process to find *ED*, eg (8.6 × sin40) ÷ 2 (=2.76...)	
		P1	process to find angle *EAD*, eg $\tan^{-1}\left(\frac{"2.76...."}{4.4}\right)$ or $\tan^{-1}$ (“0.628…”)	
		A1	answer in range 32.09 to 32.2	If an answer in the range is seen in working and then incorrectly rounded award full marks

Paper: 1MA1/2H

18	17.6	P1 P1 P1 P1 A1	for correct trig statement, eg $\sin 30 = \frac{h}{6}$ for complete process to find h, eg $6 \times \frac{1}{2}$ (= 3) for correct substitution into the area of a trapezium formula, eg $\frac{1}{2}(a+b) \times "3" = 66$ **or** $a + b = 44$ **or** $\frac{1}{2}(2x+3x) \times h = 66$ for complete correct process to find the length of AB, eg $\left[\frac{66 \times 2}{3} \div (2 + "3")\right] \times 2$ cao	An answer of $\frac{88}{5}$ gets P4 A0

Paper: 1MA1/1H

6	(a)	17.8	M1	for $\tan 56 = \frac{x}{12}$ or $(BC) = 12 \times \tan 56$ oe **or** alternative method to find *BC*	For any alternative method candidates must arrive at an equation with BC as the only unknown
			A1	for an answer in the range 17.7 to 17.8	If an answer in the range 17.7 to 17.8 is given in the working space then incorrectly rounded, award full marks.
	(b)	33.6	M1	for $\cos x = \frac{15}{18}$ **or** $\cos x = 0.83..$ **or** $x = \cos^{-1}\frac{15}{18}$ **or** alternative method to find *x*	For any alternative method candidates must arrive at an equation with *x* as the only unknown
			A1	for an answer in the range 33.5 to 33.91	If an answer in the range 33.5 to 33.91 is given in the working space then incorrectly rounded, award full marks.

Paper: 1MA1/2H

Paper: 1MA1/2H

Question	Answer	Mark	Mark scheme	Additional guidance
8	0.739	M1	for partial evaluation, eg 6.07(5732...) **or** $\sqrt{7.2}$ (= 2.68(328...)) **or** 7.80(778…) **or** $\sqrt[3]{7.80(7782818...)}$ (= 1.98(3851871…))	
		A1	for an answer in the range 0.739 to 0.747	If an answer is given in the range in working and then rounded incorrectly award full marks.

12	23.4	M1	for stating that $AC = 8$ or for a relationship that may be used to find AC eg $(AC =)\ 8 \times \tan 45$ or $\tan 45 = \frac{AC}{8}$	May be seen on diagram May use the sine rule
		M1	for relationship that may be used to find AB, eg $\sin(20) =$ "8" $\div AB$ or $(AB =) \frac{"8"}{\sin 20}$	
		A1	for answer in the range 23.3 to 23.4	If an answer is given in the range in working and then rounded incorrectly award full marks.
			Alternative	
		M1	for a relationship that may be used to find AD eg $\cos(45) = 8 \div AD$ oe or $(AD =)$ 11.3(13...)	May be seen on diagram
		M1	for a relationship that may be used to find AB, eg $\frac{AB}{\sin 45} = \frac{"11.3"}{\sin 20}$	
		A1	for answer in the range 23.3 to 23.4	If an answer is given in the range in working and then rounded incorrectly award full marks.

Paper: 1MA1/3H

18	30.6	P1	for process to find *TC*, eg. (*TC* =) $14 \times \frac{3}{3+4}$ (= 6)	Lengths of *TC*, *TD*, *SD* may be seen on the diagram
		P1	for process to find *TD*, eg. (*TD* =) $\sqrt{14^2 + "6"^2}$ or $\sqrt{232}$ or $2\sqrt{58}$ (= 15.2…)	
		P1	for process to find *SD,* using area of a trapezium, 147 = 0.5 × (*SD* + 12) × 14, or *SD*= 9	
		P1	for $\tan^{-1}\left(\frac{"9"}{"15.2..."}\right)$	A complete set of processes to find the angle is needed where an alternative route is involved with more than one stage in the working
		A1	for answer in the range 30.4 to 30.7	If an answer is given in the range in working and then rounded incorrectly award full marks.

Paper: 1MA1/3H

22	30	P1 A1	for a correct trig statement using angle ACF, eg $\sin x = \frac{6.8}{13.6}$ **or** $x = \sin^{-1}\left(\frac{6.8}{13.6}\right)$ **or** $\sin x = 0.5$ **or** $\frac{6.8}{\sin x} = \frac{13.6}{\sin 90}$ oe cao	Check diagram for working

Paper: 1MA1/1H

19	46.989...	B1	for stating at least one bound, 9.25 or 9.35 or 12.55 or 12.65	Accept $9.34\dot{9}$ or 9.3499(…) or $12.64\dot{9}$ or 12.6499(…)
		P1	for correct use of LB and UB, eg $\sin x = \frac{[\text{LB of } AB]}{[\text{UB of } AC]}$	$9.25 \leq \text{LB} < 9.3$ $12.6 < \text{UB} \leq 12.65$ $\sin(x) = 0.731(\ldots)$
		A1	for answer in the range 46.98 to 47 from correct working	Answer only award no marks.

Paper: 1MA1/2H

Question	Answer	Mark	Mark scheme	Additional guidance
10	1.25	B1	for $(\sin a =) \frac{7-2x}{8}$ or $(\tan b =) \frac{1+x}{4}$	
		P1	for start of a process to solve $\frac{7-2x}{8} = \frac{1+x}{4}$, eg $4(7-2x) = 8(1+x)$ or $28-8x = 8+8x$ or $7-2x = 2(1+x)$ or $7-2x = 2+2x$ or $\frac{7-2x}{8} = \frac{2(1+x)}{8}$ or $\frac{7-2x}{8} = \frac{2x+2}{8}$	Must come from a correct equation
		A1	for 1.25 oe	

Paper: 1MA1/2H

Question	Answer	Mark	Mark scheme	Additional guidance
20	10.8	P1	for process to find FC, eg $\tan 52 = \frac{8}{FC}$ $(FC = 6.25(028..))$	
		P1	for process that will lead to side length of ABC, eg $\sin 60 = \frac{"6.25" \times 1.5}{BC}$ **or** $\cos 30 = \frac{"6.25" \times 1.5}{BC}$ **or** $("6.25" \times 1.5)^2 + (0.5x)^2 = x^2$ oe	
		A1	for answer in range 10.8 to 10.83	

Paper: 1MA1/2H

8	14.3	M1	for a correct statement for AB using trigonometry, eg $\tan 62 = \frac{AB}{7.6}$ **or** (AB =) $7.6 \times \tan 62$	
		A1	answer in the range 14.28 to 14.3	If an answer is given in the range in working and then rounded incorrectly award full marks

Question	Answer	Mark	Mark scheme	Additional guidance
7	60.3	M1	for a correct trig statement, eg $\tan x = \frac{7}{4}$	Other methods are possible but only award this mark at the point of an equation with x as the only unknown
		A1	60.2 to 60.42	

10	12.6	M1	for partial evaluation, eg $(4.777(294722))^2$ **or** $\sqrt[3]{5.969(47697)}$ **or** 22.822(54486) **or** 1.814(081699)	
		A1	for answer in the range 12.58 to 12.6	12.58077014 If an answer is shown in the range in working and then incorrectly rounded award full marks

Paper: 1MA1/3H

GCSE

Mark Scheme

Mathematics A (1387)

June 2003

Edexcel
Success through qualifications

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information please call our Customer Response Centre on 0870 240 9800, or visit our website at www.edexcel.org.uk.

June 2003

Publications Code UG014129

All the material in this publication is copyright
© London Qualifications Ltd 2003

CONTENTS

Notes on Marking Principles 2
Paper 5501 4
Paper 5502 11
Paper 5503 17
Paper 5504 24
Paper 5505 32
Paper 5506 40

NOTES ON MARKING PRINCIPLES

NOTES ON MARKING PRINCIPLES

1 Three types of mark are available

M marks - awarded for correct working seen or implied.
A marks - conditional accuracy marks which are awarded for accurate working following the award of M marks.
B marks - unconditional accuracy marks (independent of M).

2 Abbreviations

cao - correct answer only.
ft - follow through.
$\surd$ - Denotes a "follow through" answer.

SC - special case.
isw - ignore subsequent working.
oe - or equivalent (and appropriate).
NB: a candidate cannot benefit from both isw and ft.

3 If no working shown, then

correct answers normally score full marks
incorrect (even though nearly correct) answers score no marks.

4 Marking instructions

Misread loses A marks (and sometimes B marks) on that part but ft can be allowed on subsequent parts. M marks can still be earned. If in doubt contact your team leader.
If there is a wrong answer in the answer space DO CHECK the working in the body of the script.

- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.
- If there is a wrong answer in the answer space DO CHECK the working in the body of the script.
- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.

5 Style of marking

Answer correct: tick and write part mark in margin NEXT TO BRACKETED MARK.
Answer incorrect: cross, but show M, A or B marks if any earned in body of script and transfer the total of these to the margin next to the bracket mark.
Total for each double page at bottom right page (except for back if used), FINAL TOTAL IN RELEVANT BOX ON FRONT COVER.
Nought in margin for fully incorrect question or page AND FOR NO ATTEMPT.
Where no attempt has been made a line should be put in the answer space and zero in the margin next to the bracketed mark
CHOICE OF METHOD – No marks unless one answer is in answer space – then mark that.
CROSSED OUT WORK - if not replaced this should be marked (if legible).
There must always be a mark next to bracketed mark in the margin/

6 Follow Through Marks

Follow throughs are guided by two principles:

(a) Follow throughs which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous, do not award.
(b) Follow throughs which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

7 Probability

Probability answers must be given as fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least two decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.

8 Linear Equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

9 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

Paper 5501

No	Working	Answer	Mark	Notes
1 (a)		46	1	B1 cao
(b)		3.4	1	B1 oe
(c)		Arrow at 430	1	B1 allow ± half graduation
(d)		Arrow at 3.7	1	B1 allow ± half graduation Accept indications other than arrows as long as they are clear
2		1.60 2.05	2	B1) Condone B1) reversal
3 (a)	$\frac{18}{24}$ or $\frac{9}{12}$ or $\frac{6}{8}$	$\frac{3}{4}$	2	B2 for $\frac{3}{4}$ cao (B1 for $\frac{18}{24}$ or $\frac{9}{12}$ or $\frac{6}{8}$) SC B1 for $\frac{1}{4}$ only
(b)		16 squares shaded	1	B1 cao
4 (a)		line	1	B1 within overlay tolerance
(b)		midpoint	1	B1 within overlay tolerance ft from (a) ± 0.2cm
(c)		rectangle	1	B1 for rectangle 6 cm ± 0.2cm by 4 cm ± 0.2cm
5		kilograms, kg litre, *l* or cubic metres, m^3 inches, in	3	B1 B1 B1
6 (a)		parallel lines marked	1	B1
(b)		right angle marked	1	B1
(c)(i)		acute	1	B1
(ii)		reflex	1	B1

Paper 5501

No	Working	Answer	Mark	Notes
7 (i)		sphere	1	B1
(ii)		cylinder	1	B1 Accept circular prism
(iii)		pyramid	1	B1 Condone omission of "triangular" Accept tetrahedron
8 (a)(i)		40	2	B1 cao
(ii)		50		B1 cao
(b)		5 complete symbols	1	B1 cao
(c)		 and oe inc	1	B1
9 (i)		9, 37, 56, 59, 75	5	B1 cao
(ii)		0.067, 0.56, 0.6, 0.605, 0.65		B1 cao Ignore trailing zeros
(iii)		–10, –6, –4, 2, 5		B1 cao
(iv)		$\frac{2}{5}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}$		B2 for all 4 correct (B1 for any 3 in correct order) SC B1 for all 4 in reverse order (applies to(iv) only)
10 (a)			1	B1
(b)		Plotting (4, 24)	1	B1 ft from their matchsticks
(c)		60	1	B1 cao
(d)		$m = 6n$	2	B2 for $m = 6n$ oe (B1 for $6n$ oe or m = multiple of n except $m = n$)
11 (i)		6, 12	4	B1 cao
(ii)		4, 16		B1 cao
(iii)		3, 4, 6 or 3, 4, 6, 12		B1 Condone omission of 12
(iv)		8, 27		B1 cao

Paper 5501

No	Working	Answer	Mark	Notes
12		2.43	4	B1 for 36 or 0.36 B1 for 96 or 0.96 B1 for 125 or 1.25 If none of first 3 B1s awarded then SC B1 for four 24s and five 25s seen OR 4 × 24 and 5 × 25 seen B1 for 2.43 cao
13 (a)(i)		Edinburgh and Plymouth	3	B1 for Edinburgh or −7 B1 for Plymouth or 5
(ii)		12		B1ft from (i) if one positive and one negative
(b)		Cardiff and Belfast London and Plymouth	2	B1 for Cardiff and Belfast OR −6 and −4 B1 for London and Plymouth OR 3 and 5
14		5 9 8	3	B1 cao B1 cao B1 cao
15 (a)			2	✓ / × / 2 / 0 / B2 1 / 0 / B1 1 / 1 / B1 2 / 1 / B1
(b)	$2\times2\times2$	8	2	M1 for $2\times2\times2$ A1 for 8 cao
16 (a)		30	1	B1 cao
(b)		3	1	B1 cao
(c)	$2 \times 29 = 58$ $5 \times 30 = 150$ $2 \times 31 = 62$ $1\times32 = 32$ $\frac{302}{10} = 30.2$	30.2	3	M1 for freq × no pins M1 (dep on 1st M1) for totalling and ÷ 10 A1 for 30.2 cao

Paper 5501

No	Working	Answer	Mark	Notes
17		7 5 5 13 33 52 23	3	B3 all correct (B2 for 4, 5 or 6 correct B1 for 2 or 3 correct)
18 (a)(i) (ii) (iii) (iv) (b)		$4c$ p^4 $8g$ $10pr$ OR $10rp$ $10y - 15$	4 1	B1 oe B1 cao B1 oe B1 B1 cao Accept $10y + -15$
19		$\frac{2}{3}$	3	M1 for 3 rows (9 squares) shaded M1 for 2 columns (10 squares) shaded A1 for $\frac{2}{3}$
	$\frac{3}{5} = \frac{9}{15}$ $\frac{2}{3} = \frac{10}{15}$	$\frac{2}{3}$	3	M1 for $\frac{3}{5} = \frac{9}{15}$ M1 for $\frac{2}{3} = \frac{10}{15}$ A1 for $\frac{2}{3}$
	$\frac{3}{5} = 0.6$ $\frac{2}{3} = 0.66$ or 0.67 or better	$\frac{2}{3}$	3	M1 for $\frac{3}{5} = 0.6$) Accept M1 for $\frac{2}{3} = 0.66$ or 0.67 or better) percentages A1 for $\frac{2}{3}$

Paper 5501

No	Working	Answer	Mark	Notes
20 (a)	955 × 48 7640 38200 45840 OR 48 × 955 240 2400 43200 45840	458.40	3	M1 for complete correct method (condone one computational error) A2 for 458.40 cao (A1 for digits 4584 OR ft if M1 awarded)
(b)	14. 5 48 | 696. 0 48 216 192 24 0 24 0	14.50	3	M1 for 1 as first digit in answer and remainder 21 M1 (dep) 4 as second digit in answer A1 for 14.50 (Accept 14.5)
21 (a)		$12x$	1	B1 oe
(b)		$12x + 10y$	2	B2 oe ft from (a) (B1 $12x$ + multiple of y or $10y$ seen) SC B1 for $x = 12x + 10y$ OR $y = 12x + 10y$

Paper 5501

No	Working	Answer	Mark	Notes
22	$\frac{1}{3}+\frac{1}{4}=\frac{4}{12}+\frac{3}{12}=\frac{7}{12}$ $1-\frac{7}{12}=\frac{5}{12}$	$\frac{5}{12}$	3	M1 for $\frac{4}{12}$ **and** $\frac{3}{12}$ oe A2 for $\frac{5}{12}$ oe (A1 for $\frac{7}{12}$) B1 for 1–"$\left(\frac{1}{3}+\frac{1}{4}\right)$" correctly evaluated
	$1-\frac{1}{3}=\frac{2}{3}$, $1-\frac{1}{4}=\frac{3}{4}$ $\frac{8}{12}-\frac{3}{12}=\frac{5}{12}$ or $\frac{9}{12}-\frac{4}{12}=\frac{5}{12}$	$\frac{5}{12}$	3	B1 for $\frac{2}{3}$ or $\frac{3}{4}$ seen M1 for $\frac{8}{12}-\frac{3}{12}=\frac{5}{12}$ or $\frac{9}{12}-\frac{4}{12}=\frac{5}{12}$ A1 for $\frac{5}{12}$ oe
		$\frac{5}{12}$	3	M1 for 0.25 and 0.33 or better A1 for 0.58 or better A1 for $0.41\dot{6}$ or recurring
23 (a) (b)(i) (ii)	 180 – (54 + 54)	54 72 Reason	1 3	B1 cao M1 for 180 – (54 + 54) A1 ft from (a) if $x < 90$ B1 for mentioning isosceles and equal or base angles or equal sides and equal or base angles

Paper 5501

No	Working	Answer	Mark	Notes
24 (a)		Bryani	2	M1 for 4×9 or $4 \times 3 \times 3$ or $4 \times x \times x$ or square x first or square 3 first A1 SC 4×3^2 with Bryani scores B2
(b)		64	1	B1 cao
25 (a)		 dotted line may be solid	3	B2 for rectangle base 3 squares and height 4 squares (B1 for rectangle with one correct dimension) B1 for horizontal line 1 cm from top) SC B2 for completely correct elevation on its side
(b)			2	B2 for perspective drawing showing slant face and cutout B1 for perspective drawing with either slant face cutout omitted
26 (a)	20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$	40	2	M1 for 20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$ A1 cao
(b)	$\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes	line from (45, 20) to (65, 0)	2	M1 for $\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes seen A1 for correct line SC If M0, B1 for line from (45, 20) to (t, 0) where $t > 45$ or B1 for a line of the correct gradient.

Paper 5502

No	Working	Answer	Mark	Notes
1 (a)		$\frac{7}{100}$	1	B1 cao accept 0.07
(b)		0.18	1	B1 cao
(c)	20 in 100 oe	40	2	M1 for sight of 20 in 100 or 20 × 2 A1 cao
2 (a)		6cm	2	B1 for 6 ± 0.2 or 60 ± 2 B1 indep for cm or mm consistent with 1st B1
(b)		At centre	1	B1 within overlay
(c)		Circle drawn	1	B1 all within overlay
3		See diagram	3	B3 all correct – see separate sheet (B2 for 3 correct B1 for 2 correct)
4 (a)		$\frac{1}{4}$ oe	1	B1 cao
(b)		0.75	1	B1 cao
(c)		75%	1	B1 cao
(d)(i)		9	2	B1 accept answer in range 9 - 9.2
(ii)		15 - 16		B1 accept answers in range15 - 16
5 (a)		9:30	1	B1 cao
(b)		2hrs 45 min	2	B2 for 2hr 45 min or 2 $\frac{3}{4}$ hr or 165 minutes B1 2:45 or 2.45 or 165 or 45min + 1hr + 1hr oe
(c)		17	1	B1 cao
6 (a)	96 × 4	3.84	2	M1 for 96 × 4 or digits 384 A1 cao
(b)	3 × 96 + 40 = 328	(0).56	2	M1 for 3 × 96 + 40 or digits 328 or digits 56 A1 cao accept 56p

Paper 5502

No	Working	Answer	Mark	Notes
7 (a)		54 000	1	B1 cao accept 54 thousand
(b)		50 000	1	B1 (accept ten thousand or 10 000) oe
8 (a)		14	1	B1 cao
(b)		6	1	B1 cao
(c)		Correct reflection	2	B2 fully correct (B1 correct reflection in a line parallel to the mirror line or condoning 1 block error in shape or position of shape)
9 (a)		Missing horiz label 1 (and 6) missing on vertical scale	2	B1 B1
(b)		Correct graph	2	B1 for bar up to 4 for yellow B1 for bar up to 2 for green
(c)		Blue	1	B1 cao
(d)	3 + 5 + 4 + 2	14	1	B1 ft from (b)
(e)		$\frac{3}{14}$	1	B1 ft on '14'
10	Barry (8) because you double Kath (7) because you add, 1,2,3		2	B1 oe B1 oe SC: B1 for correct rules only
11 (a)		$2n$	1	B1 for $2n$ or $n + n$ OR $2 \times n$ OR $n \times 2$ OR $n2$
(b)		$2n + 15$	1	B1 for "$2n$" + 15 oe
(c)		20q	1	B1 cao
12 (a)	1 + 3 + 5 + 8 + 5	22	2	M1 add frequencies A1 cao
(b)		No, is>no of cups of coffee in the table	1	B1 'average cannot be bigger than 6' oe OR 'Average must be less than 6 oe'
13 (a)		Trapezium	1	B1 cao ignore spelling
(b)		(2, 3)	1	B1 cao
(c)		Isosceles	1	B1 cao ignore spelling
(d)		Q correct	1	B1 cao

Paper 5502

No	Working	Answer	Mark	Notes
14 (a)		250 000	1	B1 cao
(b)	$\frac{28}{4}$	7	2	M1 for $\frac{28}{4}$ oe or "250000" × 28 A1 cao **SC** B1 for 7 000 000
15 (a)		10	1	B1 cao
(b)		5.5	1	B1 ±0.3 pounds
(c)	$\frac{110}{22}$	50	3	M1 for use of graph at 11 or $\frac{110}{22}$ A1 for 5 A1 cao **SC** B2 for 49.5 – 50.6

Paper 5502

No	Working	Answer	Mark	Notes
16 (a)	$269.30 - 56.80 = 212.50$ $\frac{212.50}{42.50}$	6	2	M1 for $\frac{269.30 - 56.80}{42.50}$ or 5 seen A1 cao
(b)	5% of £269.30 £269.30 – "£13.465" OR $\frac{95}{100} \times 269.30$	255.83 or 255.84	3	M1 for $(5 \div 100) \times 269.30$ M1 for $269.30 - $ "13.465" A1 cao OR M2 for $\frac{95}{100} \times 269.30$ A1 cao **Alternative Method:** M1 for $\frac{5}{100} \times 56.80$ $(= 2.84)$ and $\frac{5}{100} \times 42.50$ (=2.12(5)) (OR 53.96 AND 40.38 (40.375) seen M1 for $56.80 -$ "2.84" (= 53.96) $42.50 -$ "2.12(5)" (= 40.375 or 40.38) "5" × "40.375" + "53.96" A1 cao
17	$4.1^2 \times 1.07 = 16.81 \times 1.07$	17.9867	2	M1 for ("4.1") followed by squaring, or sight of 16.81 A1 cao SC: *B1 for 18 or better with no working*
18 (a)		60	1	B1 cao
(b)	$360 - 60 - 90 - 90$	120	2	M1 for $360 -$ " 60 " $- 90 - 90$ or $180 -$ "60" A1 cao
(c)	6×2	12	2	M1 for 6×2 A1 cao for 12

Paper 5502

No	Working	Answer	Mark	Notes
19 (a)(i)	$240 \times 5 = 1200$	1250	3	B1 cao 1250
(ii)	$\frac{50}{'1250'}$	$\frac{1}{25}$		M1 cao $\frac{50}{'1250'}$ A1 for $\frac{1}{25}$ in its simplest form
(b)	$\frac{60}{100} \times 1000 = 600$	12:5	3	M1 for $\frac{60}{100} \times 1000$ oe A1 for 600 A1 cao
20 (a)		$x + 2$	1	B1 accept $2 + x$ but not $x = x + 2$
(b)	$x + 5 + x + 5 + x + 2 + x + 2$	$4x + 14$	2	M1 adding 4 sides, two of which are '$x + 2$' (all sides to be linear expressions in x) **SC** $x + 5 + x + 2 \times 2$ gets M1 A1 for correct simplified answer or $(20 - 14) \div 4$ oe gets M1
(c)	'$4x + 14$' = 20	1.5 oe	2	M1 for equation A1 cao
21	$\sum f = 90$	Angles drawn, labelled	3	M1 for 1 person = 4°or one angle correct in table or pie chart A1 any 2 correctly drawn angles in pie chart A1 fully correct chart labelled
22 (a)		$2p - q$	2	B1 cao for $2p$ B1 cao for $-q$ accept $(-q + 2p)$, $2p - 1q$ and $2p + -q$
(b)	$5x = 3 + 4$	1.4	2	M1 for either (+3 or sight of 7) or (÷ 5 or sight of 0.8 and 0.6) A1 cao accept $\frac{7}{5}$ or $1\frac{2}{5}$

Paper 5502

No	Working	Answer	Mark	Notes
23 (a)		$\frac{4\times5}{2}$	1	B1 cao
(b)	$1+2+3+4+5+6+7+8$	$\frac{8\times9}{2}$	1	B1 cao
(c)	$\frac{100\times101}{2}$	5050	1	B1 cao
24	0 | 5 7 8 8 1 | 0 0 0 0 2 5 5 5 6 2 | 0 0 0 4 5 3 | 3 5 Key 1 | 2 = 12 (min)	See working	3	B1 for stem 0, 1, 2, 3 or 0, 10, 20, 30 B1 for accurate unordered leaves condone 1 error or omission B1 for key and ordered leaves all correct
25	$3.2\times2.8=8.96$ $2\times4.5\times2.8=25.2$ $2\times4.5\times28.8=28.8$ $\frac{62.96}{2.5}\times2.99$	£75.30	5	M1 for area of any face found correctly M1 for 2 areas found correctly A1 for 62.96 or 54 M1 for $\frac{'62.96'}{2.5}\times2.99$ Al cao Alternate method for candidates who round up $\frac{"62.92"}{2.5}$ M1 for "26" × 2.99 A1 for £77.74 cao **SC**: *for top included B2 for 71.92 m^2 seen or* B3 for £86.02 seen **SC** B4 for £64.58 or £65.78 seen
26	$2.5\times10\,000$	25 000	2	M1 for $2.5\times100\times100$ A1 cao

Paper 5503

No	Working	Answer	Mark	Notes
1 (a)(i)		$8g$	2	B1 oe
(ii)		$10rp$		B1 for $10pr$ or $10\ rp$
(b)		$10y - 15$	1	B1 cao accept $10y$ - + 15
(c)	$6x + 8 - 12x + 15$	$-6x + 23$	2	M1 for 3 correct terms out of 4 A1 cao
2		$m = 6n$	2	B2 for $m = 6n$ oe accept $6 \times n$, n 6 (B1 for $6n$ alone, or $6n$ +1 oe OR m = multiple of n except $m = n$)
3 (i)		0.067, 0.56, 0.6, 0.605, 0.65	1	B1 cao Ignore trailing zeros
(ii)		$-10, -6, -4, 2, 5$	1	B1 cao
(iii)		$\frac{2}{5}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}$	2	B2 all four correct (B1 any three in correct order) SC: B1 all 4 in reverse order
4		7 5 5 13 33 52 23	3	B3 all correct (B2 for 4, 5 or 6 correct entries) (B1 for 2, 3 correct entries)

Paper 5503

No	Working	Answer	Mark	Notes
5		$\frac{2}{3}$	3	M1 for 3 rows (9 squares) shaded M1 for 2 columns (10 squares) shaded A1
	OR $\frac{3}{5}=\frac{9}{15}$ $\frac{2}{3}=\frac{10}{15}$ Therefore $\frac{2}{3}>\frac{3}{5}$	$\frac{2}{3}$		M1 for $\frac{3}{5}=\frac{9}{15}$ M1 for $\frac{2}{3}=\frac{10}{15}$ A1
	OR $\frac{3}{5}=0.6$ or percent $\frac{2}{3}=0.66$ or 0.67 or better Therefore $\frac{2}{3}>\frac{3}{5}$	$\frac{2}{3}$		M1 for $\frac{3}{5}=0.6$ or percent M1 for $\frac{2}{3}=0.66$ or 0.67 or better A1
6	$2\times 29=58$ $5\times 30=150$ $2\times 31=62$ $1\times 32=32$ $\frac{302}{10}=30.2$	30.2	3	M1 for freq × no. pins (at least 3) M1 for totalling and for ÷ 10 (dep on 1st M1) A1 cao
7 (a)			2	B1 for 180° rotation (wrong centre) B1 cao
(b)			3	B1 for any enlargement sf other than 1 B1 for all sides halved B1 for position

Paper 5503

No	Working	Answer	Mark	Notes
8 (a)		$12x$	1	B1 oe
(b)		$12x + 10y$	2	B2 oe or ft from (a) (B1 for $12x$ + a multiple of y or $10y$ seen) **SC** B1 for $x = 12x + 10y$ or $y = 12x + 10y$
9	$\frac{1}{3}+\frac{1}{4}=\frac{4}{12}+\frac{3}{12}=\frac{7}{12}$ $1-\frac{7}{12}=\frac{5}{12}$ OR $1-\frac{1}{3}=\frac{2}{3}, 1-\frac{1}{4}=\frac{3}{4}$ $\frac{9}{12}-\frac{4}{12}=\frac{5}{12}, \frac{8}{12}-\frac{3}{12}=\frac{5}{12}$	$\frac{5}{12}$	3	M1 for $\frac{4}{12}$ and $\frac{3}{12}$ oe A2 for $\frac{5}{12}$ oe (A1 for getting $\frac{7}{12}$) (B1 for $1 - \frac{"1}{3} + \frac{1"}{4}$ correctly evaluated) OR B1 for $\frac{2}{3}$ or $\frac{3}{4}$ seen M1 for $\frac{3}{4}-\frac{1}{3}=\frac{9}{12}-\frac{4}{12}$ or $\frac{2}{3}-\frac{1}{4}=\frac{8}{12}-\frac{3}{12}$ A1 for $\frac{5}{12}$ oe OR M1 0.25 and 0.33 or better A1 for 0.58 or better A1 for 0.416 *recurring*
10 (a)		54	1	B1 cao
(b)(i)	180 – (54 +54)	72	3	M1 for 180 – (“54” + “54”) A1 for 72° ft from (a) if $x < 90°$
(ii)				B1 for mentioning isosceles and equal or base angles or equal sides and equal or base angles

Paper 5503

No	Working	Answer	Mark	Notes
11 (a)	Bryani was correct $4 \times 3^2 = 4 \times 9 = 36$	Bryani	2	M1 for 4×9 or $4 \times 3 \times 3$ or "square the three/x then multiply by four" A1 Bryani **SC** 4×3^2 with Bryani gets B2
(b)			1	B1 cao
12 (a)			3	B2 for rectangle height 4 squares, base 3 squares (B1 for rectangle with one correct dimension) B1for line 1 square from the top **SC** B2 for completely correct elevation on its side
(b)			2	B2 for perspective drawing showing slant face and cut out (B1 for perspective drawing with either slant face or cut out omitted or one aspect incorrect)
13 (a)	20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$	40	2	M1 for 20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$ A1 cao
(b)	$\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes seen	Line from (45, 20) to (65, 0)	2	M1 for $\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes seen A1 for correct line **SC** If M0, B1 for line from (45, 20) to (t, 0) where $t > 45$ or a line of the correct gradient
14 (i)		119.31	3	B1 cao
(ii)		119 310		B1 cao
(iii)		1.23		B1 cao

Paper 5503

No	Working	Answer	Mark	Notes
15	$\frac{10}{100}\times 12000$ 12 000 – 1200 = 10 800 $\frac{10800}{10}=1080$ 10 800 – 1080 = £9720	£9720	3	M1 for $\frac{10}{100}\times 12000$ or sight of 1200 or 2400 or 10 800 or 9600 M1 (dep) for $\frac{10}{100}\times(12\,000-\frac{10}{100}\times 12000)$ or sight of 1080 A1 cao Alternative mark scheme M2 for $12000\times(1-\frac{10}{100})^2$ (M1 for $12000\times(1-\frac{10}{100})$ A1 cao
16 (a)	$2p = 6$	$p = 3$	2	M1 for $7p - 5p = 8 - 2$ or $2p$ or 6 A1 cao
(b)	$7r - 5r = -20 - 2$	-11	2	M1 for $7r + 2 = 5r - 20$ or $\frac{7r}{5} + \frac{2}{5} = r - 4$ or $7r - 5r = 20 - 2$ or $\frac{7r}{5} - r = -4 - \frac{2}{5}$ A1 cao
17		$5n + 1$	2	B2 oe (B1 for $5n$ seen) NB: n = gets B1 max

Paper 5503

No	Working	Answer	Mark	Notes
18 (a)	–1, 0, 1		2	B2 for –1, 0, 1 (B1 for –1, 0 or 0,1 or –1,1 or –2, -1, 0, 1 only)
(b)	(–1, –1), (0, –1), (1, –1), (0, 0), (1, 0), (1, 1)		3	B3 for 6 points correct B2 for 3 points correct B1 for 1 point correct **NB** –B1 each additional point over six
19 (a)	Triangle with vertices at (0,0) (0,–2) and (3,0)		2	M1 for correct orientation A1 cao
(b)	Rotation, 180°, centre (0,1) Enlargement sf - 1 centre (0,1)		2	B2 for 180° 'rotation' centre (0, 1) B2 for Enlargement sf - 1 centre (0,1) (B1 for any two of the three parts) NB: B0 if additional transformation is included
20	Bisector of $\angle BAC$ Arc around A Region		3	B3 cao (B2 for either two correct boundaries, no shading/ wrong shading or one correct boundary, one incorrect boundary with valid shading) (B1 for either two incorrect boundaries but one drawn from A and one intersection, with valid shading or one correct boundary) Ignore shading outside the triangle
21		Length Volume Area	3	B1 for Length B1 for Volume B1 for Area
22 (a)	Unbiased question with choices		2	B1 for unbiased question B1 for at least 2 choices
(b)			2	Classification 1: A biased question Classification 2: A restricted sample of people Classification 3: Not specifying a range of foods Classification 4: Nothing to do with eating habits B2 reasons which satisfy 2 different classifications (B1 a reason which satisfies one classification)



Paper 5503

No	Working	Answer	Mark	Notes
23 (a)	$6 \times 10^2 \times 8 \times 10^4$ 48×10^6	4.8×10^7	3	M1 for $6 \times 10^a \times 8 \times 10^b$ oe, a and b integers including 0 A1 for 48×10^6 oe A1 cao
(b)	200 000 + 30 000	230000	2	B2 cao (B1 for sight of 200 000 or 30 000 or 2.3×10^5 or 23×10^4)
24 (i)		64	4	B1 cao
(ii)		3		B1 cao
(iii)	$\sqrt{16 \times 9} = \sqrt{144}$	12		B2 cao (B1 for sight of $\sqrt{2^4} \times \sqrt{9}$ or better, or 144 seen)
25 (i)	Tangent 90° to diameter /radius/ line from (through) centre	27°	4	B1 for 27° cao B1 for reason
(ii)	180 –(90+ "27") angle in semicircle (is 90°)/Alternate segments /angle at centre twice at circumference	63°		B1 ft for 90–"27" if not 63° B1 for reason
26 (a)(i)		152	2	B1 cao
(ii)		177		B1 cao
(b)			3	B1 for median marked at 167 B1 ft for postion of box with its ends at "152" and "177" B1 for position of whiskers with ends at 132 and 182 NB: For any points plotted between 141 and 149 give a tolerance of an extra ± 1 square
27	$x^2 + xy + xy + y^2$	$x^2 + 2xy + y^2$	2	M1 for at least 3 of the 4 terms correct A1 cao
		25	2	M1 for recognising 3.47 + 1.53 (= 5) A1 cao

Paper 5504

No	Working	Answer	Mark	Notes
1 (a)	$4.1^2 \times 1.07 = 16.81 \times 1.07$	17.9867	2	M1 for (4.1) followed by squaring, or sight of 16.81 A1 cao SC: B1 for 18 or better with no working
(b)		$(1.6 + 3.8 \times 2.4) \times 4.2$	1	B1 cao Allow additional brackets if they give an expression with value 45.024
2 (a)	$269.30 - 56.80 = 212.50$	6	2	M1 for $\frac{269.30 - 56.80}{42.50}$ or 5 seen A1 cao
(b)	5% of £269.30 £269.30 – "£13.465" OR $\frac{95}{100} \times 269.30$	255.83 or 255.84	3	M1 for $(5 \div 100) \times 269.30$ M1 (dep) for 269.30 – "13.465" A1 cao OR M2 for $\frac{95}{100} \times 269.30$ A1 cao **Alternative Method:** M1 for $\frac{5}{100} \times 56.80$ (= 2.84) and $\frac{5}{100} \times 42.50$ (=2.12(5)) (OR 53.96 AND 40.38 (40.375) seen M1 for 56.80 – "2.84" (= 53.96) 42.50 – "2.12(5)" (= 40.375 or 40.38) "5" × "40.375" + "53.96" A1 cao

Paper 5504

No	Working	Answer	Mark	Notes
3 (a)		60	1	B1 cao
(b)	360 – 60 – 90 – 90	120	2	M1 for 360 – "60" – 90 – 90 or 180 – "60" A1 cao
(c)	6 × 2	12	2	M1 for 6 × 2 A1 cao 12
(d)		Correct drawing	2	B2 for triangle and construction lines (see overlay) (B1 for 1 line of length 4cm and correct arcs crossing OR for correct triangle with either no arcs or incorrect arcs) SC: B1 similar triangle drawn with construction lines
4 a(i)	240 × 5 = 1200	1250	3	B1 cao 1250
(ii)	$\frac{50}{1250}$	$\frac{1}{25}$		M1 $\frac{50}{"1250"}$ A1 oe in its simplest form
(b)	$\frac{60}{100} \times 1000 = 600$ 600: 250	12:5	3	M1 for $\frac{60}{100} \times 1000$ oe A1 for 600 A1 cao
5 (a)		$x + 2$	1	B1
(b)	$x + 5 + x + 5 + x + 2 + x + 2$	$4x + 14$	2	M1 adding 4 sides, two of which must be '$x + 2$' (all sides to be linear expressions in x) A1 for correct simplified answer
(c)	"$4x + 14$" = 20	1.5	2	M1 for equation "$4x + 14$" = 20 OR $\frac{20-14}{4}$ oe A1 cao

Paper 5504

No	Working	Answer	Mark	Notes
6 (a)		$2p - q$	2	B1 cao for $2p$ B1 cao for $-q$ accept $-q + 2p$ and $2p - 1q$
(b)	$5x = 3 + 4$	1.4	2	M1 for either (+3 or sight of 7) or (÷ 5 or sight of 0.8 and 0.6) A1 cao accept $\frac{7}{5}$ or $1\frac{2}{5}$
7 (a)		$\frac{4\times5}{2}$	1	B1 cao
(b)	$1 + 2 + 3 + 4 + 5 + 6 + 7 + 8$	$\frac{8\times9}{2}$	1	B1 cao
(c)	$\frac{100\times101}{2}$	5050	1	B1 cao
(d)		$\frac{n(n+1)}{2}$	2	B2 cao (B1 for any quadratic in n)
8	$3.2 \times 2.8 = 8.96$ $2 \times 4.5 \times 2.8 = 25.2$ $2 \times 4.5 \times 3.2 = 28.8$ $\frac{62.96}{2.5} \times 2.99$	75.30	5	M1 for area of any face found correctly M1 for 2 areas seen A1 for 62.96 or 54 M1 for $\frac{'62.96'}{2.5} \times 2.99$ A1 cao Alternate method for candidates who round up $\frac{"62.96"}{2.5}$ M1 for "26" x 2.99 A1 for £77.74 cao SC: for top included B2 for 71.92 m^2 seen or B3 for £86.02 or £86.71 SC: B4 for £64.58 or £65.78 seen

Paper 5504

No	Working	Answer	Mark	Notes
9	2.5×10000	25000	2	M1 for 2.5×10000 or $2.5 \times 100 \times 100$ A1 cao
10 (a)	$\sum f = 90$ (88), 144, 32, 96	Angles drawn, labelled	3	M1 for 1 person = 4° or one angle correct in table or pie chart A1 any 2 angles correctly drawn in pie chart A1 fully correct chart labelled
(b)	$0.38 + 0.27 + 0.15$	0.20	2	M1 1 – sum A1 cao
11	0 | 5 7 8 8 1 | 0 0 0 0 2 5 5 5 6 2 | 0 0 0 4 5 3 | 3 5 Key 1 | 3 = 13 (min)	See working column	3	B1 for stem 0, 1, 2, 3 or 0, 10, 20, 30 B1 for accurate unordered leaves condone 1 error or omission B1 for key and ordered leaves all correct
12 (a)	$V = \pi \times 4^2 \times 10$	502 – 503	2	M1 for $\pi \times 4^2 \times 10$ A1 502 – 503
(b)	$P^2 = 10^2 + 8^2$ $P = \sqrt{164}$	$\sqrt{164} < 13$	3	M1 for sight of a correct right-angled triangle M1 for $10^2 + 8^2$ A1 for conclusion based on a correct calculation or 12.8 seen
13 (a)	2×30	$2 \times 2 \times 3 \times 5$	4	M1 for systematic method, eg division, factor trees (at least one prime) A1 cao
	2×48	$2^5 \times 3$		M1 for systematic method, eg division, factor trees (at least one prime) A1 cao
(b)		12	1	B1 cao
(c)	$2^5 \times 3 \times 5$	480	2	B2 cao B1 for $2^5 \times 3 \times 5$ or any correct common multiple

Paper 5504

No	Working	Answer	Mark	Notes
14 (a)		$150 < C \leq 200$	2	M1 use of cum freq to find the cost of the 20th or 20.5^{th} car OR $\frac{1}{2}\Sigma f$ or $\frac{1}{2}(\Sigma f + 1)$ A1 eg 150 to 200
(b)		No, because the 21st value is in the same interval	1	B1 20.5^{th} or 21st in same interval or an alternative correct explanation
(c)	80% = 5200 $\frac{5200}{80} \times 100$	6500	3	M1 for (100 – 20) % = 5200 M1 for $\frac{5200}{"80"} \times 100$ A1 cao
15 (a)	$x^2(x + 1) = 230$	AG	2	M1 for $x \times x \times (x+1)$ or $x \times x \times x + 1$ oe A1 cao from $x \times x \times (x+1)$
(b)	‘5 – 150 5.1 – 158.7 5.2 – 167.6 5.3 – 177.0 5.4 – 186.6 5.5 – 196.6 5.6 – 207.0 5.7 – 217.7 5.8 – 228.8 5.9 –240.2 5.85 – 234.4	5.8	4	B 2 for trial between 5.8 and 5.9 inclusive (B1 for different trial between 5 and 6 inclusive) B1 for different trial between 5.8 and 5.85 (not including 5.8) B1 (dep on at least one previous B1) cao for 5.8, 5.81, 5.811

Paper 5504

No	Working	Answer	Mark	Notes
16	$\pi \times \left(\frac{15}{2}\right)^2 = 176.715$	88.4cm^2	3	M1 for $\pi \times \left(\frac{15}{2}\right)^2$ A1 88.3 – 88.4 B1(ind) for cm²
17 (a)	$5 = 0.5x + 1$	8	2	M1 for $5 = 0.5x + 1$ A1 cao
(b)		$y = \frac{1}{2}x + c$	1	B1 $y = \frac{1}{2}x + c$, c ≠ 1, oe
(c)		$x = 2y - 2$ OR $x = 2(y - 1)$	2	M1 for correctly multiplying both sides by 2 or correctly isolating $\frac{x}{2}$ A1 for $x = 2(y - 1)$, $x = \frac{y-1}{0.5}$, $x = \frac{y-1}{\frac{1}{2}}$ oe SC: B1 for $x = 2y - 1$
18	$4x - 6y = 22$ $15x + 6y = 54$ $19x = 76$	$x = 4,\ y = -1$	4	M1 for coefficients of x or y the same followed by correct operation , one arithmetical error A1 cao M1(dep on previous M mark) for sub for other variable A1 cao Trial and improvement –0 unless both x and y correct values found

Paper 5504

No	Working	Answer	Mark	Notes
19 (a)	$SF = \frac{10}{6}$ $\frac{10}{6} \times 4.8 = 8$	8	2	M1 for sight of $\frac{10}{6}$ or $\frac{6}{10}$ or 1.67 or better or $\frac{CD}{10} = \frac{4.8}{6}$ A1 cao
(b)	$\frac{10}{6} \times 4.5 - 4.5 = 3$	19.8	2	M1 for use of SF from (a) to find *BC* or *AC* and adding 4 sides A1 cao
20	$\frac{6 \times 10^{15}}{3.2 \times 10^{8}}$ 1.875×10^{7}	4.3×10^{3}	3	B3 for 4.3×10^{3} to 4.34×10^{3} (B2 for 1.875×10^{7} oe or 4300 to 4340 or final answer of 1.9×10^{7}) (B1 for sight of 6×10^{15} oe or 3.2×10^{8} oe)
21	$8.5 \times \tan 38 = 8.5 \times 0.7813$	6.64	3	M1 for correct use of trig, eg $\tan 38 = \frac{opp}{8.5}$ M1 for 8.5 × tan38 A1 6.64 – 6.641

Paper 5504				
No	**Working**	**Answer**	**Mark**	**Notes**
22 (a)		No, as you would expect about 100. Yes, as it is possible to get 200 sixes with a fair dice	1	B1 for a consistent answer
(b)	$\frac{1}{6}, \frac{5}{6}$ + labels		3	B1 for $\frac{5}{6}$ on the red dice, *not six* branch B1 for a fully complete tree diagram with all branches labelled B1 for $\frac{1}{6}$, $\frac{5}{6}$ on all remaining branches as appropriate

Paper 5505

No	Working	Answer	Mark	Notes
1 (i) (ii) (iii)		119.31 119310 1.23	3	B1 cao B1 cao B1 cao
2	$\frac{10}{100}\times 12000 = 1200$ 12 000 – 1200 = 10 800 10 800 ÷ 10 =1080 10 800 – 1080 = £9720	9720	3	M1 for $\frac{10}{100}\times 12\,000$ or sight of 1200 or 2400 or 10 800 or 9600 M1 (dep) for $\frac{10}{100}\times(12\,000 - \frac{10}{100}\times 12\,000)$ or sight of 1080 A1 cao Alternative markscheme M2 for $12000\times\left(1-\frac{10}{100}\right)^2$ (M1 for $12000\times\left(1-\frac{10}{100}\right)$) A1 cao
3	$7r - 5r = -20 - 2$	–11	2	M1 for $7r + 2 = 5r - 20$ or $\frac{7r}{5} + \frac{2}{5} = r - 4$ or $7r - 5r = -20 - 2$ or $\frac{7r}{5} - r = -4 - \frac{2}{5}$ A1 cao

Paper 5505

No	Working	Answer	Mark	Notes
4 (a)		–1, 0, 1	2	B2 for –1, 0, 1 only (B1 for –1, 0 or 0,1 or –1, 1 or –2, -1, 0, 1 only)
(b)		(–1,–1)(0,–1) (1,–1) (0, 0) (1, 0) (1, 1)	3	B3 for 6 points correct (B2 for 3 points correct) (B1 for 1 point correct) *NB: –B1 for each additional point over six*
5		$5n + 1$	2	B2 oe (B1 for $5n$ seen) NB: n = gets B1 max
6 (a)	Triangle with vertices at (0,0) (0,–2) and (3,0)		2	M1 for correct orientation A1 cao
(b)	Rotation, 180°, centre (0,1) Enlargement sf –1 centre (0,1)		2	B2 for 180° 'rotation' centre (0, 1) or for Enlargement sf - 1 centre (0,1) (B1 for any two of the three parts) NB: B0 if additional transformation is included
7	Bisector of $\angle BAC$ Arc around A Region	See overlay	3	B3 cao (B2 for either two correct boundaries, no shading/ wrong shading or one correct boundary, one incorrect boundary with valid shading) (B1 for either two incorrect boundaries but one drawn from A and one intersection, with valid shading or one correct boundary) Ignore shading outside the triangle
8		Length Volume Area	3	B1 for Length B1 for Volume B1 for Area



Paper 5505

No	Working	Answer	Mark	Notes
9 (a)	Unbiased question with choices		2	B1 for unbiased question B1 for at least 2 choices
(b)	Leading question and a restricted sample		2	Classification 1: A biased question Classification 2: A restricted sample of people Classification 3: Not specifying a range of foods Classification 4: Nothing to do with eating habits B2 reasons which satisfy 2 different classifications (B1 a reason which satisfies one classification)
10 (a)	$6 \times 10^2 \times 8 \times 10^4$ $48 \times 10^6 = 4.8 \times 10^7$	4.8×10^7	3	M1 for $6 \times 10^a \times 8 \times 10^b$ oe, a and b integers including 0 A1 for 48×10^6 oe A1 cao
(b)	200 000 + 30 000 = 230 000	230 000	2	B2 cao (B1 for sight of 200 000 or 30 000 or 2.3×10^5 or 23×10^4)
11 (a)	$x^2 + xy + xy + y^2$	$x^2 + 2xy + y^2$	2	M1 for at least 3 of the 4 terms correct A1 cao
(b)		25	2	M1 for recognising 3.47 + 1.53 (= 5) A1 cao
12 (i)	Tangent 90° to diameter /radius/ line from (through) centre	27°	4	B1 for 27° cao B1 for reason
(ii)	180 –(90+ "27") angle in semicircle (is 90°)/Alternate segments /angle at centre twice at circumference	63°		B1 ft for 90–"27" if not 63° B1 for reason
13 (i)		p^9	1	B1 cao
(ii)		$6q^6$	2	B2 for $6q^6$ (B1 for sight of $\frac{6q^9}{q^3}$ or $3q \times 2q^5$ or $3q^4 \times 2q^2$ or $6 \times q \times q \times q \times q \times q \times q$ or final answer of the form kq^6, $k > 0$)

Paper 5505

No	Working	Answer	Mark	Notes
14 (a)(i)		152	2	B1 cao
(ii)		177		B1 cao
(b)			3	B1 for median marked at 167 B1 ft for postion of box with its ends at "152" and "177" B1 for position of whiskers with ends at 132 and 182 NB: For any points plotted between 141 and 149 give a tolerance of an extra ± 1 square
15	(arc =) $\frac{40}{360} \times 2\pi \times 9$ $= 2\pi$	$2\pi + 18$	4	M1 for $\frac{40}{360} \times$ M1 for $2\pi \times 9$ M1 (dep) for $\frac{40}{360} \times 2\pi \times 9$ oe A1 for $\frac{18 \times \pi}{9} + 18$ oe exact form
16 (i)		1	1	B1 cao
(ii)		$\frac{1}{16}$	1	B1 cao accept 0.0625
(iii)		64	1	B1 cao condone ± 64

Paper 5505

No	Working	Answer	Mark	Notes
17 (a)	$F \propto 1/x^2$ $F = k/x^2$ $4 = k / 3^2$ $F = 36/x^2$	$F = \frac{36}{x^2}$	3	M1 for $F = k/x^2$ seen or implied. ($k \neq 1$) M1 (dep) for subst. or sight of $k = 36$ A1 for $F = 36/x^2$
(b)		9	1	B1 ft for 9 (ft on $F = kx^n$, $n \neq 0$)
(c)	$64 = \frac{36}{x^2}$ $x^2 = \frac{36}{64}$ $x = \pm\frac{3}{4}$	$\frac{6}{8}$	2	M1 for $x^2 = \frac{"36"}{64}$ A1 for $\frac{6}{8}$ oe (condone $\pm$) *SC: Use of $F = kx^2$ max M1 M1 A0 B1 ft M0 A0* *SC: Use of* $F = \frac{k}{\sqrt{x}}$ *max M1 M1 A0 B0 M1* ($\sqrt{x} = \frac{4\sqrt{3}}{64}$) *A0*
18	$(5+\sqrt{3})(5-\sqrt{3}) = 5\times5 - 5\sqrt{3} + 5\sqrt{3} - \sqrt{3}\sqrt{3}$ $= 5\times5 - 3$ $\frac{22}{\sqrt{22}} = \frac{22\sqrt{22}}{22}$	$\sqrt{22}$	3	B1 for correct expansion 25 - $5\sqrt{3}$ + $5\sqrt{3}$ - $\sqrt{3}$ $\sqrt{3}$ with 1st three terms reducing to 25 **without** any errors seen B1 (indep) for $\sqrt{3}$ $\sqrt{3}$ = 3 B1 for $\sqrt{22}$ coming from $\frac{22}{\sqrt{22}}$ (S.C $\frac{(5+\sqrt{3})(5-\sqrt{3})\sqrt{22}}{22}$ gets B1)

Paper 5505

No	Working	Answer	Mark	Notes
19 (a)		60 40	2	B1 cao B1 cao
(b)		correct bars	2	B1 for $30 < x \leq 40$ with an area of 2½ squares B1 for $40 < x \leq 70$ with an area of 3 squares *SC:* $\frac{0}{4}$ *give M1 if clearly using area or frequency density*
20 (a)	$6x + 8 - 12x + 15$	$-6x + 23$	2	M1 for 3 of the 4 terms $6x$, $+8$, $-12x$, $+15$ correct A1 cao
(b)		$32x^5y^{15}$	2	B2 cao (B1 for two of 32, x^5, y^{15})
(c)	$\frac{(n+1)(n-1)}{n+1} \times \frac{2}{n-2}$ $\frac{2(n-1)}{n-2}$	$\frac{2(n-1)}{n-2}$	3	M1 for $k(n + 1)(n - 1)$ M1 dep for $\frac{(n+1)(n-1)}{(n+1)} = n - 1$ A1 for $\frac{2(n-1)}{n-2}$
21	Vertices at $(-3, -1\frac{1}{2}), (-4\frac{1}{2}, -1\frac{1}{2}), (-3, -4\frac{1}{2})$		3	B1 for all sides $\times 1\frac{1}{2}$ B1 for correct orientation with 2 vertices almost correct B1 cao

Paper 5505

No	Working	Answer	Mark	Notes
22	Total = 3 + 5 + 2 (=10) $\frac{3}{10}\times\frac{3}{10}\times\frac{5}{10}\left(=\frac{45}{1000}\right), \frac{3}{10}\times\frac{3}{10}\times\frac{2}{10}\left(=\frac{18}{1000}\right)$ $\frac{5}{10}\times\frac{5}{10}\times\frac{3}{10}\left(=\frac{75}{1000}\right), \frac{5}{10}\times\frac{5}{10}\times\frac{2}{10}\left(=\frac{50}{1000}\right)$ $\frac{2}{10}\times\frac{2}{10}\times\frac{3}{10}\left(=\frac{12}{1000}\right), \frac{2}{10}\times\frac{2}{10}\times\frac{5}{10}\left(=\frac{20}{1000}\right)$ $3\times\left(\frac{"45"}{1000}+\frac{"18"}{1000}+\frac{"75"}{1000}+\frac{"50"}{1000}+\frac{"12"}{1000}+\frac{"20"}{1000}\right)$ $\frac{660}{1000}$	$\frac{660}{1000}$ oe	5	M3 for all six expressions seen OR their combined equivalents (M2 for four expressions seen OR their combined equivalents) (M1 for two expressions seen OR their combined equivalents) M1 sum of 18 relevant products condone 1 slip A1 for $\frac{660}{1000}$ oe *SC: without replacement maximum M4 A0* SC: Just 2 beads: Answer either $\frac{38}{100}$ oe OR $\frac{28}{90}$ oe B1

Paper 5505

No	Working	Answer	Mark	Notes
23 (a)(i)		$6\mathbf{b} - 6\mathbf{a}$	2	B1 for $6\mathbf{b} - 6\mathbf{a}$ oe
(ii)		$6\mathbf{a}$		B1 for $6\mathbf{a}$ oe
(b)	$\overrightarrow{EX} = \overrightarrow{EB} + \overrightarrow{BX}$ $= 12\mathbf{b} + ½\ \overrightarrow{BC}$	$12\mathbf{b} - 3\mathbf{a}$	2	M1 for $\overrightarrow{EX} = \overrightarrow{EB} + \overrightarrow{BX}$ oe vector journey in a form ready for straightforward substitution A1 for $12\mathbf{b} - 3\mathbf{a}$ oe
(c)	$\overrightarrow{AY} = \frac{5}{3}\overrightarrow{AB}$ or $\overrightarrow{BY} = \frac{2}{3}\overrightarrow{AB}$ $\overrightarrow{EY} = 16\mathbf{b} - 4\mathbf{a}$ or $\overrightarrow{XY} = 4\mathbf{b} - \mathbf{a}$ $\overrightarrow{EY} = 4\overrightarrow{XY}$ or $\overrightarrow{EX} = 3\overrightarrow{XY}$ or $\overrightarrow{EY} = \frac{4}{3}\overrightarrow{EX}$	Printer Answer	3	B1 for either $\overrightarrow{AY} = \frac{5}{3}\overrightarrow{AB}$ or $\overrightarrow{BY} = \frac{2}{3}\overrightarrow{AB}$ oe B1 ft for either $\overrightarrow{EY} = 16\mathbf{b} - 4\mathbf{a}$ or $\overrightarrow{XY} = 4\mathbf{b} - \mathbf{a}$ ft only on parts (*a*) and (*b*) B1 for either $\overrightarrow{EY} = 4\overrightarrow{XY}$ or $\overrightarrow{EX} = 3\overrightarrow{XY}$ or $\overrightarrow{EY} = \frac{4}{3}\overrightarrow{EX}$ oe **plus** conclusion of *E*, *X*, *Y* on the same straight line
24 (a)(i)		$(5, -4)$	4	B1 cao
(ii)		$(2, -9)$		B1 cao
(iii)		$(2, 4)$		B1 cao
(iv)		$(1, -4)$		B1 cao
(b)		$(x-2)^2 - 4$	4	B4 for $(x-2)^2 - 4$ oe eg. $x^2 - 4x$ (B3 for $(x+2)^2 - 4$ or $(x-2)^2 + 4$) (B2 for $x^2 - 4$ or $(x-2)^2$ OR $x^2 + bx$, $b \neq 0$ OR $(x+2)^2 + 4$ OR $f(x-2) - 4$) (B1 for $x^2 + 4$ or $(x+2)^2$ or $ax^2 + bx$ or $x^2 + bx + c$ OR $x - 2 - 4$ or $x^2 - 2 - 4$, $a, b, c \neq 0$)

Paper 5506

No	Working	Answer	Mark	Notes
1 (a)	$V = \pi \times 4^2 \times 10$	502 – 503 cm^3	2	M1 for $\pi \times 4^2 \times 10$ A1 502 – 503
(b)	$P^2 = 10^2 + 8^2$ $P = \sqrt{164}$	$\sqrt{164} < 13$	3	M1 for sight of correct right angled triangle M1 for $10^2 + 8^2$ A1 for conclusion based on a correct calculation Or 12.8 seen
2 (a)(i)	2×30	$2 \times 2 \times 3 \times 5$	4	M1 for systematic method, eg division, factor trees (at least one prime) A1 cao
(ii)	2×48	$2^5 \times 3$		M1 for systematic method, division, factor trees (at least one prime) A1 cao
(b)		12	1	B1 cao
(c)	$2^5 \times 3 \times 5$	480	2	B2 cao B1 for $2^5 \times 3 \times 5$ or any correct common multiple

Paper 5506

No	Working	Answer	Mark	Notes
3 (a)		$150 < C \leq 200$	2	M1 use of cum freq to find the cost of the 20th or 20.5^{th} car OR $\frac{1}{2}\Sigma f$ th or $\frac{1}{2}(\Sigma f + 1)$ th car. A1 eg 150 to 200, 150 – 200
(b)		No, because the 21st value is in the same interval	1	B1 for 20.5^{th} or 21st value in the same internal consistent with 'a' OR Refers to the median value being low in the interval (statement to be mathematically correct) See additional sheet
(c)	80% = 5200 $\frac{5200}{80} \times 100$	6500	3	M1 for (100 – 20)% = 5200 M1 for $\frac{5200}{"80"} \times 100$ A1 cao

Paper 5506

No	Working	Answer	Mark	Notes
4 (a)	$x^2(x + 1) = 230$	AG	2	M1 for $x \times x \times (x+1)$ or $x \times x \times x + 1$ oe, $x^2(x+1)$, $x^2 \times x + 1$ A1 cao from $x \times x \times (x+1)$, no need to see 230
(b)	5 – 150 6 – 252 5.1 – 158.7 5.2 – 167.6 5.3 – 177.0 5.4 – 186.6 5.5 – 196.6 5.6 – 207.0 5.7 – 217.7 5.8 – 228.8 5.9 –240.2 5.85 – 234.4	5.8	4	B2 for trial between 5.8 and 5.9 inclusive evaluated (B1 for trial between 5 and 6 inclusive evaluated) B1 for different trial between 5.8 and 5.85 (not including 5.8) B1 dep on at least are previous B1 5.8, 5.81, 5.811
5	$\pi \times \left(\frac{15}{2}\right)^2 = 176.715$	88.4 cm^2	3	M1 for $\pi \times \left(\frac{15}{2}\right)^2$ seen A1 88.3 – 88.4 B1 for cm² (independent)

Paper 5506

No	Working	Answer	Mark	Notes
6 (a)	$5 = 0.5x + 1$	8	2	M1 for $5 = 0.5x + 1$ A1 cao
(b)		$y = \frac{1}{2}x + c$	1	B1for $y = \frac{1}{2}x + c$, $c \neq 1$, oe
(c)		$x = 2y - 2$ OR $x = 2(y - 1)$	2	M1for correctly multiplying both sides by 2 or correctly isolating $\frac{x}{2}$ A1 for $x = 2(y - 1)$, $x = \frac{y-1}{0.5}$, $\frac{y-1}{\frac{1}{2}}$ oe SC B1 for x= 2y -1
7	$4x - 6y = 22$ $15x + 6y = 54$ $19x = 76$	$x = 4$, $y = -1$	4	M1 for coefficients of x or y the same followed by correct operation, allow one arithmetical error A1 cao M1 (dep) for correct sub for other variable A1 cao Trial and improvement 0 marks unless both correct values of x and y found
8 (a)	SF = $\frac{10}{6}$ $\frac{10}{6} \times 4.8 = 8$	8	2	M1 for sight of $\frac{10}{6}$ or $\frac{6}{10}$ or 1.67 or better or $\frac{CD}{10} = \frac{4.8}{6}$ A1 cao
(b)	$\frac{10}{6} \times 4.5 - 4.5 = 3$	19.8	2	M1 for use of SF from "a" to find AC or BC or $\frac{BC}{4.5} = \frac{4}{6}$ and adding 4 sides A1 cao

Paper 5506

No	Working	Answer	Mark	Notes
9	$\frac{6\times10^{15}}{3.2\times10^{8}}$ 1.875×10^{7}	4.3×10^{3}	3	B3 for 4.3×10^{3} to 4.34×10^{3} (B2 for 1.875×10^{7} oe or 4300 to 4340, final answer of 1.9×10^{7} B1 for sight of 6×10^{15} oe or 3.2×10^{8} oe)
10	$8.5\times\tan38$ $=8.5\times0.7813$ $\frac{8.5}{\sin(90-38)}=\frac{AB}{\sin38}$ $AB=\frac{8.5\times\sin38}{\sin(90-38)}$ $=\frac{5.2331}{0.788}=6.64$	6.64	3	M1 for correct use of trig, eg $\tan 38=\frac{opp}{8.5}$ M1 for $8.5\times\tan38$ A1 6.64 – 6.641 OR M1 for correct substitution into the sine rule M1 (dep) for correct rearrangement for AB= A1 6.64 – 6.641
11 (a)		No, as you would expect about 100. Yes, as it is possible to get 200 sixes with a fair dice	1	B1 for a consistent answer See additional sheet

Paper 5506

No	Working	Answer	Mark	Notes
(b)	$\frac{1}{6}, \frac{5}{6}$ + labels		3	B1 for $\frac{5}{6}$ on the red dice, *not six* branch B1 for a fully complete tree diagram with all branches labelled B1 for $\frac{1}{6}$ and $\frac{5}{6}$ on all remaining branches as appropriate
(c)(i)	$\left(\frac{1}{6}\right)^2$	$\frac{1}{36}$	2	M1 $\left(\frac{1}{6}\right)^2$ or $\frac{1}{6}\times\frac{1}{6}$ only or 0.28 A1 $\frac{1}{36}$ or 0.03 or better
(ii)	$1-\left(\frac{5}{6}\right)^2$ OR $\frac{1}{6}\times\frac{5}{6}+\frac{5}{6}\times\frac{1}{6}+\frac{1}{6}\times\frac{1}{6}$	$\frac{11}{36}$	3	M2 for $1-\left(\frac{5}{6}\right)^2$ or $1-\frac{5}{6}\times\frac{5}{6}$ A1 cao OR M1 for $\frac{1}{6}\times\frac{5}{6}$ oe M1 for 2 or 3 only of $\frac{1}{6}\times\frac{5}{6}, \frac{5}{6}\times\frac{1}{6}$, "a" A1 for $\frac{11}{36}$ or 0.31 or better

Paper 5506

No	Working	Answer	Mark	Notes
12 (a)	$\pi \times 30 \times \frac{7.5^2}{3} - \pi \times 10 \times \frac{2.5^2}{3} = 1767 - 65$	1700	3	M1 for either $\pi \times 30 \times \frac{7.5^2}{3}$ or $\pi \times 10 \times \frac{2.5^2}{3}$ M1 (dep) for difference A1 1700 – 1702 **SC** B1 Using d instead of r, 6800 – 6808
(b)	$\frac{S}{2\pi d} = \sqrt{h^2 + d^2}$ $\left(\frac{S}{2\pi d}\right)^2 = h^2 + d^2$	$h = \sqrt{\frac{S^2 - 4\pi^2 d}{4\pi^2 d^2}}$	3	M1 for correctly isolating $\sqrt{h^2 + d^2}$ or $h^2 + d^2$ or $h + d$ or kh^2 or kh M1(indep) squaring both sides A1 $h = \sqrt{\frac{S^2 - 4\pi^2 d^4}{4\pi^2 d^2}}$, $h = \frac{\sqrt{S^2 - 4\pi^2 d^4}}{2\pi d}$ $h = \sqrt{\left(\frac{S}{2\pi d}\right)^2 - d^2}$
(c)	$\left(\frac{30}{20}\right)^2 \times 450$ or $450 \div \left(\frac{20}{30}\right)^2$	1012.5	2	M1 for sight of correct SF^2 including 4:9 A1 1010 to 1013

Paper 5506

No	Working	Answer	Mark	Notes
13 (a)	$\frac{2x(x+20)}{2} = 400$	As given	2	M1 $\frac{2x(x+20)}{2}$ or $\frac{2x \times x + 20}{2}$ or $2x(x + 20) = 800$ A1 cao following correct working, no need for = 400 *SC B1* $2x \times x + \frac{1}{2} \times 2x(10 - \frac{x}{2}) \times 2$
(b)	$\frac{-20 \pm \sqrt{20^2 - 4 \times 1 \times (-400)}}{2}$ $= \frac{-20 \pm 44.721}{2}$	12.361	3	M1 for correct sub, up to signs, in the quad formula A1 for 44.7 or $\sqrt{2000}$ A1 for 12.3606 – 12.361, ignore negative solution T.I B3 for 12.361 OR Completing the square M1 for $(x+10)^2$ seen A1 for $-10 \pm \sqrt{500}$ A1 for 12.3606 – 12.361 ignore negative solution

Paper 5506

No	Working	Answer	Mark	Notes
14 (a)	$0.5\times 8\times 15\times \sin 70^0$	56.4	2	M1 for correct sub into area formula A1 56.38 – 56.4
(b)	$AB^2 = 8^2 + 15^2 - 2\times 8\times 15\times \cos 70^0 = 206.9$ EITHER $0.5\times AB\times CX$ ='56.38' OR $\frac{\sin B}{8} = \frac{\sin 70}{\sqrt{'206.9'}}$ $B = 31.5$ 15 sin '31.5'	7.84	4	M1 for correct sub into cos rule A1 for 206.9 - 207 or 14.38 – 14.4 EITHER M1 for use of area rule to find CX A1 7.83 – 7.84 OR M1 for correct use of sine rule to find sin B or sin A and then sight of 15sin B *or* 8 sin A A1 7.83 – 7.84

Paper 5506

No	Working	Answer	Mark	Notes
15 (a)	$4a^2 - 4a + 1 - (4b^2 - 4b + 1) =$ $4(a^2 - b^2) - 4(a - b)$ $4(a - b)(a + b - 1)$ OR $((2a - 1) - (2b - 1))((2a - 1) + (2b - 1))$ $(2a - 2b)(2a + 2b - 2)$	AG	3	Expansion Method M1 for a correct expansion of any one of the three terms M1(dep) on an attempt to expand all 3 terms and show LHS = RHS A1 fully correct algebra RHS exp is $4(a^2 + ab - a - ba - b^2 + b)$ OR Factorisation Method M1 for attempt to use difference of 2 squares on LHS M1 for one bracket correctly simplified A1 fully correct
(b)	Any 2 odd square numbers have the above form If a and b are both even or odd then $a - b$ is even, so $4(a - b)$ is a multiple of 8 If one of a,b is odd, then $a + b - 1$ is even, so $4(a + b - 1)$ is a multiple of 8		3	B1 'any 2 square nos have the above form' (may be implied by sight of $(2a - 1)^2 - (2b - 1)^2$ in part (b)) B1 first reason B1 second reason SC B1 for $(2r + 1)^2 - (2r - 1)^2$ B1 for 8r
16 (a)	$g_L = \frac{2 \times 4.495}{1.35^2 \times \sin 30.5}$ $g_u = \frac{2 \times 4.505}{1.25^2 \times \sin 29.5}$	9.719 11.710	4	B2 for any 4 of 4.505, 1.25, 29.5, 4.495, 1.35, 30.5 seen (B1 for any two or three seen) B1 for 11.710 – 11.7103 B1 cao 9.719 – 9.71904
(b)	Round, until lower and upper bounds agree	10	1	B1 for 10 + reason " they agree to this level of accuracy"

Paper 5506

No	Working	Answer	Mark	Notes
17 (a)(i)		xy	3	B1 cao
(ii)		y^2		B1 for y^2 or $y \times y$
(iii)		$\frac{x}{2}$		B1 for $\frac{x}{2}$ or $0.5x$ or $2^{-1}x$
(b)	Divide to get $2y = 1$	$q = -1$	2	M1 for $2y = 1$ or $\frac{x}{2} = 32$ or $p + q = 5$ or $1 + p + 2q = 5$
		$p = 6$		A1 cao
18 (a)	$x^2 - 2mx + m^2 - k$	$k = m^2$	2	M1 for correct exp of $(x - m)^2$ or correct completion of the square eg $\left(x - \frac{2m}{2}\right)^2 - \left(\frac{2m}{2}\right)^2$
				A1 cao
				SC B1 for $k = -m^2$
(b)(i)	Min value is $-m^2$	$-m^2$	3	M1 for recognition that min value occurs when $(x - m)^2 = 0$ (either (b)(i) or (b)(ii) correct implies this M1)
(ii)	$x = m$	m		A1 ft on 'k', " $-k$" gets M1 A0
				A1 cao
19	$0.06 \times 0.05 = 0.003$	No	2	M1 for 0.06×0.05
				A1 correct conclusion based on 0.003 or 0.06 x 0.05 stated as $\neq 0.0011$
				OR M1 for statement that for the two events to be independent P (BL and CL) = P(BL) × P(CL)

Paper 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
20		50 50 4	3	B1 50 or $\frac{100}{2}$ B1 for 50 or "a" B1 4 or $\frac{360}{90}$ oe

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4LN

Telephone 01623 467467
Fax 01623 450481

Order Code UG014129 June 2003

For more information on Edexcel qualifications please contact our
Customer Response Centre on 0870 240 9800
or email: enquiries@edexcel.org.uk
or visit our website: www.edexcel.org.uk

London Qualifications Limited. Registered in England and Wales no.4496750
Registered Office: Stewart House, 32 Russell Square, London WC1B 5DN



PMT

Mark Scheme (Results)

Summer 2016

Pearson Edexcel GCSE
In Mathematics B (2MB01)
Higher (Calculator) Unit 3

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2016
Publications Code 5MB3H_01_1606_MS
All the material in this publication is copyright
© Pearson Education Ltd 2016

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Note that in some cases a correct answer alone will not score marks unless supported by working; these situations are made clear in the mark scheme. Examiners should be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will award marks for the quality of written communication (QWC).
The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labelling conventions.

ii) *select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 **With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Partial answers shown (usually indicated in the ms by brackets) can be awarded the method mark associated with it (implied).
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks; transcription errors may also gain some credit. Send any such responses to review for the Team Leader to consider.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 **Follow through marks**

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 **Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.

10 **Probability**

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded (embedded answers).

12 Parts of questions
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers
Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

Guidance on the use of codes within this mark scheme
M1 – method mark A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

PMT

PAPER: 5MB3H/01					
Question		**Working**	**Answer**	**Mark**	**Notes**
1*			Decision (supported)	3	M1 for correct first step e.g. 15 ÷ 3 M1 for complete method C1 for correct decision with accurate figures
2	(a)		Correct diagram	2	B2 cao (B1 for correct orientation)
	(b)		Enlargement, sf 4, centre (0, 1)	3	B1 for enlargement B1 for scale factor 4 B1 for centre (0, 1)
3			$w = \frac{d+5}{2}$	2	M1 for attempt to add 5 to both sides or divide both sides by 2 as a first step, eg $d + 5 = 2w$ oe or $\frac{d}{2} = w - \frac{5}{2}$ oe A1 for $w = \frac{d+5}{2}$ oe

PMT

PAPER: 5MB3H/01					
Question		Working	Answer	Mark	Notes
4			22.5	4	M1 for $4(x - 8) = 2x + 13$ M1 for expansion of bracket or division of all terms by 4, eg $4x - 32 = 2x + 13$, or $x - 8 = \frac{2x}{4} + \frac{13}{4}$ M1 for isolating x and number terms eg $2x = 45$, $\frac{x}{2} = \frac{45}{4}$ A1 for $\frac{45}{2}$ or 22.5 OR M1 for $(180 - 64) \div 2$ (= 58) M1 for $4(x - 8)$ = "58" or $2x + 13$ = "58" or "58" – 13 (= 45) M1 for isolating x and number terms eg $4x = 90$, $2x = 45$ or "45" ÷ 2 A1 for $\frac{45}{2}$ or 22.5 OR M1 for $64 + 4(x - 8) + 2x + 13$ M1 for $64 + 4(x - 8) + 2x + 13 = 180$ M1 for isolating x and number terms eg $6x = 135$ A1 for $\frac{45}{2}$ or 22.5
5			Correct region shaded	3	M1 for a circle centre M or N (accept arc of sufficient length to define the region) M1 for circle centre M radius 5 cm **and** circle centre N radius 3 cm (accept arc of sufficient length to define the region) A1 for correct region shaded

PAPER: 5MB3H/01					
Question		Working	Answer	Mark	Notes
*6			Conclusion (supported)	4	M1 for $\pi \times 120^2$ (= 45 216 – 45 249) M1 for "$\pi \times 120^2$" ÷ 1800 A1 for 25 – 25.2 C1 ft (dep on M2) for appropriate conclusion from their figures OR M1 for $\pi \times 120^2$ (= 45 216 – 45 249) M1 for 20 × 1800 A1 for 36 000 and 45 216 – 45 249 C1 ft (dep on M2) for appropriate conclusion from their figures OR M1 for $\pi \times 120^2$ (= 45 216 – 45 249) M1 for "$\pi \times 120^2$" ÷ 20 A1 for 2260 – 2263 C1 ft (dep on M2) for appropriate conclusion from their figures OR M1 for 1800 × 20 M1 for 36 000 ÷ π (=11 457 – 11 465) A1 for 107(.0…) C1 ft (dep on M2) for appropriate conclusion from their figures
7			161.50	5	M2 for a correct method to decrease 6720 by 20%, eg 6720 × 0.8 (= 5376) or 6720 × 0.2 (= 1344 and 6720 – 1344 (= 5376)) (M1 for a correct method to find 20% of 6720 eg 6720 × 0.2 or $\frac{20}{100} \times 6720$ (= 1344)) M1 for subtracting 1500 (= 3876) after a percentage calculation M1 "3876" ÷ 24 after the subtraction of 1500 A1 for 161.5(0)

PMT

PAPER: 5MB3H/01					
Question		**Working**	**Answer**	**Mark**	**Notes**
8		3 42 3.1 45.(291) 3.2 48.(768) 3.3 52.(437) 3.4 56.(304) 3.5 60.(375) 3.6 64.(656) 3.7 69.(153) 3.8 73.(872) 3.9 78.(819) 4 84 3.71 69.6(14811) 3.72 70.(078848) 3.73 70.5(45117) 3.74 71.(013624) 3.75 71.4(84375)	3.7	4	B2 for a trial between 3 and 4 exclusive (B1 for a trial at 3 or 4) B1 for a different trial of $3.7 < x \leq 3.75$ B1 (dep on at least one previous B1) for 3.7 NB. Trials should be evaluated to at least 2 sf truncated or rounded for values of x correct to 1 decimal place. Trials should be evaluated to at least 1 dp truncated or rounded for values of x correct to 2dp. **NB** No working scores 0 marks even if answer is correct
9			618	4	M1 for a method to find an area which is part of the cross section M1 (dep) for a complete method to find the total area of the cross section M1 (dep M1) for their cross sectional area × 12 A1 cao **OR** M1 for a method to find the volume of cuboid, 8 × 8 × 12 (= 768) M1 for a method to find the volume of the triangular prism, ½ ×5 × 5 × 12(= 150) M1 (dep M2) for a complete method for the volume of the prism A1 cao

5MB3H_01					
Question		Working	Answer	Mark	Notes
10	(a)		15.2	3	M1 for $12.7^2 + 8.3^2$ or $161.29 + 68.89$ or 230.18 M1 for √"$(12.7^2 + 8.3^2)$" or √"$161.29 + 68.89$" or $\sqrt{230.18}$ A1 for 15.2 or 15.17(168…)
	(b)		16.1	3	M1 for $\sin 40° = \frac{DE}{25}$ oe or $\frac{DE}{\sin 40} = \frac{25}{\sin 90}$ oe M1 for 25 sin 40° A1 for 16.1 or 16.06(969…)
11	(a)		4.081	2	M1 for 8.875 or $\frac{71}{8}$ or 2.979…or 4.08 A1 for 4.081 or 4.0809(5…)…
	(b)		4×10^{-3}	2	M1 for 4×10^n, $n \neq -3$ or $a \times 10^{-3}$, $a \neq 4$ or 0.004 or 0.4×10^{-2} or 1/250 A1 cao
12			$x < 2$	2	M1 for an attempt to isolate x and number terms or multiply all terms by 2, or for $\frac{3}{2}x < 3$ or $-\frac{3}{2}x < -3$ or $x = 2$ A1 cao
13	(a)		0, 6, 4, –6	2	B2 for all values correct (B1 for any one value correct)
	(b)		Graph drawn	2	M1 ft (provided B1 in (a) for at least 6 points plotted correctly from their table A1 cao for correct curve drawn from (–4, –6) to (3, –6)
	(c)		1.5 to 1.6 and –2.5 to –2.6	2	M1 for correct use $y = 2$ (may be implied by one correct estimate) or correct use of formula. A1 for 1.5 to 1.6 and –2.5 to –2.6

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
14*		Sher Yas **Old 19300 12560** New 20265 13502 **Inc 965 942**	Correct comparison (supported)	3	M1 for a correct method to calculate the population at the beginning of the 10 year period for at least one place. eg $\frac{20265}{1.05}$ (= 19 300) oe or $\frac{13502}{1.075}$ (= 12 560) oe A1 for 965 (Sherbury) **and** 942(Yaston) C1 (dep on M1) ft for statement comparing increases leading to conclusion based on two comparable amounts eg increase in Sherbury is greater than increase in Yaston
15		$20x + 10y = 35$ $6x - 10y = -48$ $x = -\frac{1}{2}$ $y = \frac{9}{2}$ OR $12x + 6y = 21$ $12x - 20y = -96$ $26y = 117$ $y = \frac{9}{2}$ $x = -\frac{1}{2}$	$-\frac{1}{2}, \frac{9}{2}$	4	M1 for a correct process to eliminate either variable (condone one arithmetic error) A1 cao for either x or y M1 (dep on M1) for correct substitution of found value into one of the equations or appropriate method after starting again (condone one arithmetic error) A1 cao OR M1 for full method to rearrange and substitute to eliminate either variable (condone one arithmetic error) A1 cao for either x or y M1 (dep on M1) for correct substitution of found value into one of the equations or appropriate method after starting again (condone one arithmetic error) A1 cao **NB** Trial and improvement methods score 0 marks unless both x and y are correct
16			226	2	M1 for angle $ABC = 180 - 67$ (= 113) A1 cao OR M1 for obtuse angle $AOC = 2 \times 67$ (= 134) A1 cao

5MB3H_01					
Question		Working	**Answer**	**Mark**	**Notes**
17			– 2.82, 19.82	3	M1 for $\frac{--17 \pm \sqrt{(-17)^2 - 4\times1\times-56}}{2\times1}$ (condone one sign error) or $\left(x - \frac{17}{2}\right)^2$ oe M1 for $\frac{17 \pm \sqrt{513}}{2}$ or for method leading to $\frac{17}{2} \pm \sqrt{\frac{513}{4}}$ oe A1 for answers in the range 19.82 to 19.825; – 2.82 to –2.825
18			21.2	4	M1 for correct use of sine rule to find *AB* or *BC*, eg $\frac{6}{\sin 25} = \frac{AB}{\sin 120}$ or $\frac{6}{\sin 25} = \frac{BC}{\sin(180-120-25)}$ M1 for (*AB* =) $\frac{6}{\sin 25} \times \sin 120$ (= 12.2(951…)) or (*BC* =) $\frac{6}{\sin 25} \times \sin 35$ (= 8.1(431…) M1(dep M2) for method to find area, eg 0.5 × 6 × "*AB*" sin35 or 0.5 × 6 × "*BC*" sin 120 A1 for answer in the range 21.1 – 21.2 OR (with perpendicular from *A* meeting *BC* extended at a point *X*) M1 for method to find *AX* M1 for method to find *BX* M1(dep M2) for 0.5 × "*AX*" × "*BC*" A1 for answer in the range 21.1 – 21.2
*19			0.38 because the LB and UB agree to that number of figures	5	B1 for 10.75 or 10.85 or 10.84999… B1 for 75.055 or 75.065 or 75.064999… M1 for $\sqrt{\frac{10.85}{75.055}}$ as UB **OR** $\sqrt{\frac{10.75}{75.065}}$ as LB A1 (dep on all previous marks) for both 0.3802… and 0.3784 must clearly come from working with correct values C1 for 0.38 from 0.3784… and 0.3802 … **and** "both LB and UB round to 0.38" **NB** An answer of 038(932….) without working scores no marks

5MB3H_01					
Question		Working	**Answer**	**Mark**	**Notes**
20			7.35	3	M1 for $y \propto \frac{1}{\sqrt{x}}$ or $y = \frac{k}{\sqrt{x}}$ or $9 = \frac{k}{\sqrt{4}}$ M1 for $k = 9\sqrt{4}$ (=18) A1 for 7.35 or 7.348…
*21			$\frac{2}{3}$ (7**a** + 3**b**) and correct conclusion	4	M1 for correct vector for $\overrightarrow{OY}$, or $\overrightarrow{AP}$ eg $(\overrightarrow{OY}) = \overrightarrow{OA} + \overrightarrow{AY}$, or $\overrightarrow{OY} = \overrightarrow{OA} + \frac{2}{3}\overrightarrow{AP}$ may include terms in **a** and **b,** eg. 6**a** + $\overrightarrow{AY}$, 4**b** + 4**a** – **b** + $\overrightarrow{PY}$ or $\overrightarrow{AP} = \overrightarrow{AO} + \overrightarrow{OB} + \overrightarrow{BP}$ or –6**a** + 4**b** + 4**a** – **b** or – 2**a** + 3**b** M1 for $(\overrightarrow{AY}) = \frac{2}{3}\overrightarrow{AP}$ or ⅔(–6**a** + 4**b** + 4**a** – **b**) or ⅔(–2**a** + 3**b**) or $(\overrightarrow{PY} = \frac{1}{3}\overrightarrow{PA}$ or ⅓(– 4**a** + **b** – 4**b** + 6**a**) or ⅓(2**a** – 3**b**) M1 for correct expression for $\overrightarrow{OY}$ in terms of **a** and **b,** eg $\frac{14}{3}$ **a** + 2**b** C1 for $\frac{2}{3}$ (7**a** + 3**b**) and "$\overrightarrow{OY}$ is parallel to the vector 7**a** +3**b**" **oe**

Modifications to the mark scheme for Modified Large Print (MLP) papers.

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q2	(a)	Grid enlarged. Shading changed to dotty shading. Labels added to shapes. Image labelled shape A. Rotation given and labelled shape B. Question changed to 'Describe fully the single transformation which maps shape A to shape B. One row top and bottom, one column left and right has been removed. Three answer lines given.	2 cao (B1 for correct orientation in third quadrant or correct quarter turn)
Q2	(b)	Grid enlarged. Shading changed to dotty shading. Labels added next to the triangles 'triangle P' and 'triangle Q'. One column removed from the left.	B1 for enlargement B1 for scale factor 4 B1 for centre (0, 1)

PMT

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q4		Diagram enlarged. MLP only – x changed to y. Braille only – measurements given in the text.	M1 for $4(x - 8) = 2x + 13$ M1 for expansion of bracket or division of all terms by 4, eg $4x - 32 = 2x + 13$, $x - 8 = 2x/4 + 13/4$ M1 for isolating x and number terms eg $2x = 45$, $x/2 = 45/4$ A1 for 45/2 or 22.5 OR M1 for $(180 - 64) \div 2$ (= 58) M1 for $4(x - 8) = 58$ or $2x + 13 = 58$ M1 for isolating x and number terms eg $4x = 90$, $2x = 45$ A1 for 45/2 or 22.5 OR M1 for $64 + 4(x - 8) + 2x + 13$ M1 for $64 + 4(x - 8) + 2x + 13 = 180$ M1 for isolating x and number terms eg $6x = 135$ A1 for 45/2 or 22.5
Q5		Diagram kept the same size as original. Scale moved above the diagram and also put in the question paper.	B1 for circle centre M radius 5 cm B1 for circle centre N radius 3 cm B1 for correct region shaded

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q6		Diagram enlarged. Cross changed to a filled in circle. Arrow heads removed.	M1 for $\pi \times 120^2$ (= 45238….) M1 for "$\pi \times 120^2$" ÷ 1800 A1 for 25.0 – 25.2 C1 for "surface is large enough as 25 > 20" oe OR M1 for $\pi \times 120^2$ (= 45238….) M1 for 20 × 1800 A1 for 36000 C1 for "surface is large enough as 36000 < 45238..." oe OR M1 for $\pi \times r^2 = 1800 \times 20$ M1 for 36000 ÷ π (=11459..) A1 for 107.0… C1 for "surface is large enough as 107…< 120" oe
Q9		Model provided for all candidates. Diagram enlarged and provided for MLP.	M1 for a method to find an area which is part of the cross section M1 (dep) for a complete method to find the total area of the cross section M1 (dep M2) for a complete method for the volume of the prism A1 cao OR M1 for a method to find the volume of cuboid, 8 × 8 × 12 M1 for a method to find the volume of the triangular prism M1 (depM2) for a complete method for the volume of the prism A1 cao

PMT

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q10	(a)	Diagram enlarged. Braille only: measurements given in the text.	M1 for $12.7^2 + 8.3^2$ or $161.29 + 68.89$ or 230.18 M1 for $\sqrt{161.29 + 68.89}$ or $\sqrt{230.18}$ A1 for 15.2 or 15.17(168…)
Q10	(b)	Diagram enlarged. Braille only – measurements given in the text.	M1 for $\sin 40° = \frac{DE}{25}$ oe M1 for $25 \sin 40°$ A1 for 16.1 or 16.06(969…)
Q13	(a)	Wording added 'There are four spaces to fill.'	B2 for all values correct (B1 for any one value correct)
Q13	(b)	Grid enlarged.	M1 ft for at least 6 points plotted correctly from their table A1 cao for correct curve drawn from (–4, –6) to (3, –6)
Q16		Diagram enlarged. Dot added at centre.	M1 for angle ABC = 180 – 67 (= 113) A1 cao OR M1 for obtuse angle AOC = 2 × 67 (= 134) A1 cao

PMT

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q18		Diagram enlarged.	M1 for correct use of sine rule to find AB or BC, eg $\frac{6}{\sin 25} = \frac{AB}{\sin 120}$ M1 for $(AB =) \frac{6}{\sin 25} \times \sin 120$ or $(BC =) \frac{6}{\sin 25} \times \sin 35$ M1(dep M2) for method to find area, eg $0.5\times6\times AB\sin35$ or $0.5\times6\times BC \sin 120$ A1 for answer in the range 21.1 – 21.2 OR (with perpendicular from A meeting BC extended at a point X) M1 for method to find AX M1 for method to find BX M1(dep M2) for $0.5 \times$ "AX" $\times$ "BC" A1 for answer in the range 21.1 – 21.2
Q21		Diagram enlarged.	M1 for correct vector for $\overrightarrow{OY}$, eg $(\overrightarrow{OY}) = \overrightarrow{OA} + \overrightarrow{AY}$, may include terms in **a** and **b,** eg. $6\mathbf{a} + \overrightarrow{AY}$, $4\mathbf{b} + 4\mathbf{a} - \mathbf{b} + \overrightarrow{PY}$ M1 for $(\overrightarrow{AY} =) \frac{2}{3}\vec{A}P$ or $(\overrightarrow{PY} =) \frac{1}{3}\overrightarrow{PA}$ M1 for correct expression for $\overrightarrow{OY}$ in terms of **a** and **b,** eg $\frac{14}{3}\mathbf{a} + 2\mathbf{b}$ C1 for $\frac{2}{3}(7\mathbf{a} + 3\mathbf{b})$ and "$\overrightarrow{OY}$ is parallel to the vector $7\mathbf{a} + 3\mathbf{b}$" **oe**

PMT

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London WC2R 0RL

Question 14

Students who gave the exact value of either tan30º or sin60º gained the first mark. Various strategies were used to find these values. Some students worked them out by drawing an appropriate triangle; some used patterns in a table with angles of 0º, 30º, 45º, 60º and 90º and some simply recalled them from memory. Sometimes sin30º ÷ cos30º was used to find the value of tan30º. However, the strategies used were not always successful and incorrect values were very common. When only one of the two values was correct it was more often $\sin 60^\circ = \frac{\sqrt{3}}{2}$. In some responses the values were not labelled. If only one value was correct examiners had to assume from the calculation shown that the order was tan30º×sin60º. Some of the students who

got to $\frac{1}{\sqrt{3}} \times \frac{\sqrt{3}}{2}$ did not go on to get full marks because they gave $\frac{\sqrt{3}}{2\sqrt{3}}$ as the final answer or simplified it to 2. The question required the answer to be given in its simplest form. Some students used incorrect values such as $\tan 30^\circ = 1$ and $\sin 60^\circ = 0.5$ which just happened to give an answer of 0.5. When the "correct" answer clearly came from incorrect working a mark of 0 was awarded.

Question 5

As should be expected on the higher tier, the majority of students were able to gain both marks on this simple trigonometry question for a correct answer in range. Most students completed the question using the sine ratio, however some used a combination of other trigonometric ratios and Pythagoras's Theorem. Provided they didn't prematurely round their interim values, they ended with an answer in range and still gained both marks. As ever some students used the wrong ratio, and this typically meant zero marks. There were a small number of students who had their calculator set in radians; in this instance if the method was shown then the method mark could be awarded.

Question 7

This question was done well by the great majority of students who were awarded both marks. Some students gave an answer of 0.32 believing this was correct to 3 significant figures. These students were given full marks provided they had shown a more accurate version of the answer in the working space. The request to give the answer correct to 3 significant figures served as a guide to students and correct answers given to more than 3 significant figures gained full credit. Students are therefore advised to write down their full calculator display before rounding.

Question 14

A small minority of students scored full marks for their responses to this question. Weaker students often showed little or no working and if they did show working, it was often characterised by assuming that the triangles were right angled and using trigonometry to fit such a situation. More able students often did realise that references to area in the question together with the diagram showing non right angled triangles suggested use of the formula $\frac{1}{2}ab\sin C$. However, some of these students tried to find the length of AE instead of using the ratio $AB : AE$ to express AE as an unknown variable, for example $3x$. Students who used $AB = x$ and $AE = 3x$ scored at least one mark and often went on to write down and solve an equation using the given fact that the sum of the areas of the triangles was equal to the area of the rectangle. Examiners were surprised at the number of students who equated the area of one triangle to the area of the rectangle which led to the incorrect final answer of 18.

Question15

Some students appeared to have little knowledge about the content covered in this question and either did not attempt the question or revealed little understanding through their responses to the question. In general, little working was seen and students may have benefited from writing down the transformations represented by the equations $y = f(-x)$ and $y = f(x) - 3$.
Very few students did this and so some denied themselves the award of a mark for showing an understanding of the types of transformation involved in the question. However, examiners were able to award a good number of students at least one mark for getting an x coordinate as -7 or a y coordinate as -1. Students who gave a final answer of $(-10, -5)$ following on from an intermediate stage of $(-7, -2)$ could not be awarded any marks because they had not discriminated between which coordinates were changed by which transformation. Some students attempted to substitute values for x and y, often 7 and 2 respectively. The best students did show a good understanding of the notation and transformations involved and gave a fully correct answer.

Question 12

This was not well done with many not attempting any trigonometry. Those that did often got the first mark for the process of making a correct start to trigonometry, but then spoilt their working. Others, having calculated 5.52… went on to use Pythagoras's Theorem but then lost their way. It was not uncommon to see students using 5.52.. as the length of *ED*. Quite a few students over complicated the process of solution by using the sine rule or even cosine rule in an attempt to solve, with few of these showing any success due to error in manipulation. Some students, who showed a completely correct process, rounded or truncated their interim values to the extent that their final answer was outside the range required for the final mark.

Question 18

This proved to be a challenging question and finding the length of *AB* proved to be beyond most students. Some of those who attempted the question identified that they need to find the perpendicular height of the trapezium and gained the first mark for writing a correct trigonometric statement such as sin30 = *h*/6. Some students could not recall the value of sin30 and did not complete the process to find the height. After finding the height many students struggled to make any further progress, failing to realise that they should use the formula for the area of a trapezium. It was common to see the trapezium split into two triangles and a rectangle and attempts at using Pythagoras. A good proportion of those that did substitute correctly into the formula went on to show a complete process to find the length of *AB*. A small number of students gave $\frac{88}{5}$ as the final answer and lost the accuracy mark.

Question 6

It was evident that students had spent some time studying trigonometry, but again the learning did not seem to be secure. Many students were not able to identify the correct ratio for either part. Of those who did, many either substituted incorrectly, or they gained the first mark but were unable to gain the second as they struggled with the rearrangement. This was most evident in part (b) but was still seen in part (a).
Of those who attempted an alternative method, a good number were again able to evidence understanding of Pythagoras' Theorem, but were again unable to choose the correct ratio to gain the first mark.

Question 8

Another familiar question, and another answered well by many. A significant number of students did struggle, and it seemed it was the denominator of the fraction including the

fractional power that caused the issues. They could gain a single mark for a correct partial evaluation, but it was common to either see no marks or both marks.

Students need to make sure they are able use the power button correctly on their calculators so they are able to process complex calculations. There were many instances where just the answer was written down which lost possible marks for a partial evaluation. 1.43 was often seen as the cube root was taken of everything. Also 7.5 was multiplied by $\frac{1}{2}$ rather than to the power. Another common incorrect answer was 1.427... which came from taking the cube root of the whole fraction rather than just the denominator.

Question 12

In answering this question, most students showed a good understanding of how to apply trigonometrical ratios to this 2 stage problem. However, a significant proportion of them made errors in manipulating the ratio used in the second stage of the problem.

It was envisaged that students would use the trigonometry of right angled triangles to solve this problem and most did. However, a large number of students used the sine rule to get their answer. Examiners were surprised how few students realised that triangle *ACD* was isosceles and so they were not able to write down the length of AC without further calculation. However, they usually used tan 45 = $\frac{AC}{8}$ correctly to get 8 cm for the length of *AC*. Most students were then able to write down a relationship which would enable them to find the length of *AB*, and this was awarded some credit. However, too many of these students did not rearrange their equation correctly and instead of getting $AB = \frac{8}{\sin 20}$, wrote down $AB = 8 \times \sin 20$, leading to an incorrect answer.

Question 18

It is encouraging to report that the majority of students gained some credit for their attempt at this question. The length of *TC* was often found successfully and a good proportion of students were also able to find the length of at least one of *SD* and *TD*. Only the higher attaining students were able to identify the angle which was required and put everything together in order to find the size of this angle, that is the size of angle *DTS*. Errors commonly seen included, substituting 12 and 14 incorrectly into the formula for the area of the cross section and making an error in the use of Pythagoras' rule to find the length of *DT*.

Question 22

Many students drew the lines *FC* and *AC* on the diagram and identified *ACF* as the angle they were attempting to find. Those who drew only the line *FC* often failed to identify angle *ACF*. After identifying the correct angle some students made no further progress and attempts at using Pythagoras were quite common. About half of those who got the first mark, often for a statement such as $\sin x = \frac{6.8}{13.6}$ or $x = \sin^{-1}\left(\frac{6.8}{13.6}\right)$, were then able to give a correct answer. Some students evaluated 6.8 ÷ 13.6 incorrectly, obtaining 2 for example, but many did not attempt to simplify $\frac{6.8}{13.6}$. Had students recognised that 6.8 is half of 13.6 more might have gained the second mark. Some that did get to $\sin x = \frac{1}{2}$ then gave an answer such as 45 or 60. Students who gained the first mark for a correct sine rule statement were often unable to rearrange this correctly to find the value of *x*. Some students presented the exact trig values in tabular form but were unable to apply them to the question.

Question 19

Many candidates, including some of the weaker ones, were able to gain a mark here for a suitable bound, it was good to see that they were applying their knowledge of error intervals into a more complex context. However, many then struggled to select the correct bound in their expression for $\sin(x)$ and so gained no further credit. Candidates who did select the correct bounds and achieved 2 marks, typically then did gain the final mark, although a number forgot to find the inverse sine and were left with an answer of 0.731... rather than 46.989… Some candidates attempted the question in the incorrect order and worked with $\sin(x)$ and then attempted to find bounds of this answer. This method gained no marks.

Question 10

This question started with an independent B mark for a correct expression for either sin a or tan b. It was pleasing to see so many students gaining this credit and then setting the values equal to each other to form an equation. Many did then struggle to solve this question and the process mark was a correct start to do so. To gain this, students had to have a correct process to clear both fractions or to write them with a common denominator with a correct matching numerator. A significant number formed the correct equation but didn't gain the P mark as they only multiplied the equations by 4, thus only clearing one denominator.
Those who were able to clear the denominators, often then went on to solve correctly and gave an answer of 1.25 or $\frac{5}{4}$.

Question 20

Although there were not as many steps as we often see on a 3D trigonometry problem here, many candidates struggled to make a start to the question. The first step to find FC proved conceptually too difficult and many either didn't attempt, or used a method that would lead to VC not FC. This could then lead to FC with a second step using Pythagoras, but this was normally missing.
If learners did find FC, they then had to use the ratio to find MC along with Pythagoras to form an equation for the side length. Alternatively, they could use MC along with either the $60°$ or $30°$ angle and trigonometry. Neither were seen often, but those who got this far generally gained the accuracy mark.

Question 8

Of those who scored 0, they usually chose the cosine ratio instead of the tan ratio or used the tan ratio incorrectly, writing $\tan 62 = \frac{7.6}{AB}$. A number of learners opted for the sine rule to successfully get their answer.

Question 7

A straightforward trigonometry question to find an unknown angle, and one where many gained both marks. The efficient method was to use tan, but that wasn't the only method seen. There were several learners that used Pythagoras' Theorem as first step and then either sin or cos. Both methods were acceptable, but the method mark was only awarded when they got as far as an equation where the missing angle was the only unknown.
Whilst correct, and perfectly acceptable, these alternatives, are far less efficient, and often led to arithmetic errors, or accuracy errors due to premature rounding. Learners should be encouraged to use the most efficient method where possible and reminded not to round until a final answer is obtained. There were some who identified tan as the correct ratio to use but made the mistake of writing the numbers the wrong way round.

Question 10

Many responses correctly gave the final answer of 12.6, which was often the only value written in the answer space. In some cases, the full calculator value was shown before rounding. This was helpful as, where 12.58 was incorrectly rounded to 12.5, both marks could still be awarded if the full value was shown. However, responses that only stated 12.5 without any supporting working or intermediate value were not credited.

Some responses showed calculations in radians rather than degrees. This resulted in no marks being awarded, highlighting the importance of checking calculator settings before the exam.

Other frequent errors included –2.582… from cube rooting only the 17.4 rather than the full denominator, 2.157… from not squaring the 1.83, and 2.63… from not squaring the numerator. In these cases, partial credit was often awarded for correct intermediate steps, such as a correct numerator of 22.822… or a correct denominator of 1.814…