Chhetri Academy

chhetriacademy.com

GCSE MATHS

Geometry and measures

Edexcel 1MA1 | Subtopic collection Topic 46

Similar shapes (length, area, volume)

Higher tier

Answers and examiner comments

59	2003-2025	Higher
question groups	available papers	tier

Using this booklet

1. Answer numbers match the companion question booklet.
2. Follow the original question number on marking tables.
3. Examiner comments are included where available.

Name _____ **Date** _____

Some entries use full source marking material in an appendix. Any unavailable mark schemes are identified beside the relevant answer.

Original question 8

The full source mark scheme begins on page 84. Use the original paper code, tier and question number shown above.

Original question 12

The full source mark scheme begins on page 84. Use the original paper code, tier and question number shown above.

Original question 16

No matching mark scheme was available in the supplied archive for this question.

Original question 10

No matching mark scheme was available in the supplied archive for this question.

Original question 11

The full source mark scheme begins on page 138. Use the original paper code, tier and question number shown above.

Original question 20

The full source mark scheme begins on page 199. Use the original paper code, tier and question number shown above.

Original question 12

No matching mark scheme was available in the supplied archive for this question.

Original question 21

No matching mark scheme was available in the supplied archive for this question.

22	(a)	$6 \times \frac{15}{10}$	9	2	M1 for sight of $\frac{15}{10}$ or $\frac{10}{15}$ or $\frac{10}{6}$ or $\frac{6}{10}$ oe seen A1 cao **NB** ratios get M0 unless of the form 1:*n*
	(b)	$12 \times \frac{10}{15}$ oe	8	2	M1 for correct use of $\frac{15}{10}$ or $\frac{10}{15}$ or $\frac{15}{12}$ or $\frac{12}{15}$ or $\frac{"9"}{6}$ or $\frac{6}{"9"}$ oe A1 for 8 or ft from $12 \times 6 \div$ '9'

1380 4H

1380/4H

Question		Working	Answer	Mark	Notes
18	(a)	$12\times\frac{6}{4}$	18	2	M1 for sight of $\frac{6}{4}$ oe or $\frac{4}{6}$ oe or $\frac{12}{4}$ oe or $\frac{4}{12}$ oe or a ratio eg. 6:4 oe or decimal eg. 1.5 oe A1 cao
	(b)	$15\times\frac{4}{6}$	10	2	M1 for $15\times\frac{4}{6}$ oe or $\frac{15}{"18"}\times 12$ oe A1 cao

Question	Working	Answer	Mark	Notes
23	50 : 18 = 25 : 9 Ratio of length = 5 : 3 Ratio of volume = 125 : 27 $500\times\frac{27}{125}$	108	4	M1 for ratio of length =√50:√18 or √25 : √9 (or 5 : 3 or 3 : 5 seen) or 1.66-1.67 or 0.6 M1 for ratio of volume (mass) = "5"^{3} : "3"^{3} or $25^{\frac{3}{2}}$: $9^{\frac{3}{2}}$ oe or 4.62-4.63 or 0.216 M1 (dep on at least M1) for 500 × "$\frac{9^{\frac{3}{2}}}{25^{\frac{3}{2}}}$" or 500 ÷"4.62" or 500÷"4.63" or 500×"0.216" A1 cao [NB: ratios do not need to be presented as ratios as long as the relationship is clear]

Question	Working	Answer	Mark	Notes
19 (a)	15 ÷ 10 8 × 1.5	12	2	M1 for 15 ÷ 10 or 1.5 or $\frac{3}{2}$ or $\frac{2}{3}$ A1 cao
(b)	½ × (8 +"a") × 5	50	2	NB : ft from (a) provided '*DC*' > 8 M1 for $\frac{(8+"a")\times 5}{2}$ A1 ft **or** M1 for (8× 5) + ½("*DC*" − 8) × 5 A1 ft **or** M1 for ½ × "*DC*" × 15 − ½ × 8 × 10 A1 ft **or** M1 for ½ × 8 × 10 × "1.5²" − ½ × 8 × 10 A1 ft

1380_3H

1380/3H

17		$PBC = 90 - PAC$ $BCP = 90 - (90 - PAC)$	Proof	3	M1 for $PBC = 90 - PAC$ or $PAC = 90 - PBC$ or $ACP = 90 - PCB$ M1 for $BCP = 90 - (90 - PAC)$ or $PAC = 90 - (90 - BCP)$ oe A1 for $PAC = PCB$ and $PCA = PBC$ and $APC = CPB$ B1 SC if M0 awarded for $APC = BPC = 90°$ or statement matching the 3 equal sets of angles $PAC = PCB$ and $PCA = PBC$ and $APC = CPB$

1380_3H

17	(a)	$BC \div 12 = 10 \div 6$ $BC = 10 \times 12 \div 6$	20	2	M1 for $12 \div 6$ or $6 \div 12$ or $10 \div 6$ or $6 \div 10$ oe or a decimal equivalent including 1.6, 1.66...., 1.67 or 1.7 A1 19.9 – 20.4
	(b)	$PR \div 18 = 6 \div 10$ $PR = 6 \times 18 \div 10$	10.8	2	M1 for $6 \times 18 \div 10$ oe or $18 \div (1.6, 1.66..., 1.67, 1.7)$ oe or a complete method ft '20' eg $12 \div$'20'$\times 18$ A1 for 10.8

1380_4H

Question		Working	Answer	Mark	Notes
17		$\frac{15+6}{15}\times 12.5$	17.5	3	M1 for $\frac{DE}{12.5}=\frac{15+6}{15}$ oe **or** $\frac{15}{15+6}$ **or** $\frac{15+6}{15}$ **or** $\frac{7}{5}$ **or** $\frac{5}{7}$ **or** $\frac{2}{5}$ **or** $\frac{5}{2}$ (1.4 or 0.4 or 2.5 or 0.714…) M1 for $\frac{15+6}{15}\times 12.5$ **or** $\frac{7}{5}\times 12.5$ oe **or** $12.5+\frac{2}{5}\times 12.5$ oe A1 cao

Original question 18

The full source mark scheme begins on page 251. Use the original paper code, tier and question number shown above.

Question		Working	Answer	Mark	Notes
1		15 ÷ 3 × 4 =	20 cm	3	M1 for 4 × 5 or 3 × 5 or $\frac{3}{15}$, $\frac{15}{3}$, $\frac{3}{4}$, $\frac{4}{3}$, or equivalent values, or 4:20 oe, or identification of 5 as the scale factor of enlargement. A1 cao C1 (indep) for units: cm stated on answer line or with "20" in the working space if not given on answer line.

Question		Working	Answer	Mark	Notes
17	(a)	$1:2^2$ or $2^2:1$ $80 \times 2^2 = 80 \times 4 =$	320	2	M1 for sight of $1:2^2$ or $2^2:1$ or 2^2 or $\frac{1}{4}$ for ratio of area or 80×4 or identification of 4 as the scale factor A1 cao
	(b)	$1:2^3$ or $2^3:1$ 171700×2^3 $= 171700 \times 8 =$ OR $h_a=\frac{171700\times 3}{\pi\times40^2}$ $=102.47589$ $h_b = h_a\times2=204.95..$ $vol_b= \frac{1}{3}\pi\times80^2\times204.95..$	1 373 600	3	M1for sight of $1:2^3$ or $2^3:1$ or 2^3 or $\frac{1}{8}$ for ratio of volumes or identification of 8 as the scale factor M1 for $2^3 \times 171700$ A1 cao OR M1 for complete calculation to find the height of A (=102.47589..) M1 (dep) for $h_a\times2$ and used to find vol_b A1 cao

Question		Working	Answer	Mark	Notes
25	(a)		640	2	M1 for $80\times\left(\frac{8}{4}\right)^3$ or $80\div\left(\frac{4}{8}\right)^3$ A1 cao
	(b)		40	2	M1 for $160\div\left(\frac{8}{4}\right)^2$ or $160\times\left(\frac{4}{8}\right)^2$ or ft their scale factor from (a) A1 cao

1MA0_1H

Question		Working	Answer	Mark	Notes
20		SF = $(x^2 - 1) \div 2\,(x - 1)$ = $(x - 1) \times (x + 1) \div 2\,(x - 1)$ = ½ $(x + 1)$ Area *DEF* = $4 \times \left[\frac{1}{2}(x+1)\right]^2$ = $(x + 1)^2$ = $x^2 + 2x + 1$		4	M1 $(x^2 - 1) \div 2\,(x - 1)$ or SF $\times 2\,(x - 1) = (x^2 - 1)$ M1 ½ $(x + 1)$ or $(x - 1) \times (x + 1) \div 2\,(x - 1)$ M1 $4 \times \left(\frac{x+1}{2}\right)^2$ or $4 \times \left(\frac{x^2-1}{2(x-1)}\right)^2$ C1 fully correct convincing process OR M1 $(x^2 + 2x + 1) \div 4$ M1 $\sqrt{(x^2 + 2x + 1) \div 4}$ or $\sqrt{(x+1)(x+1) \div 4}$ or $(x + 1) \div 2$ M1 $2\,(x - 1) \times (x + 1) \div 2$ C1 fully correct convincing process

15	(a)	$\frac{1}{2} \times (4 + 12) \times 10$	80	2	M1 for a fully correct method for area of *QRST* A1 cao
	(b)	**For example** $\frac{PT+10}{PT} = \frac{12}{4} = 3$ $PT + 10 = 3PT$ $2PT = 10$	5	3	M1 for a correct scale factor or ratio using two corresponding sides from two similar triangles or two sides within the same triangle (may be seen within an equation) eg. $\frac{12}{4}$oe **or** 4 : 12 oe **or** $\frac{PT}{4}$**or** $\frac{PS}{12}$ **or** $\frac{12}{12-4}$ etc. M1 for a correct equation with *PT* or *PS* as the only variable **or** complete correct method using scale factor A1 cao

1MA0_2H

Question		Working	Answer	Mark	Notes
22		$\frac{1}{3} \times \pi \times 15^2 \times 40$ $-\frac{1}{3} \times \pi \times 7.5^2 \times 20$	8250	4	B1 for 15cm as diameter or 7.5 cm as radius of smaller cone (may be marked on diagram or used in a formula) M1 for a numerical expression for the volume of one cone eg. $\frac{1}{3} \times \pi \times 15^2 \times 40$ (=9424...) **or** $\frac{1}{3} \times \pi \times 7.5^2 \times 20$ (=1178...) M1 for $\frac{1}{3} \times \pi \times 15^2 \times 40$ oe $-\frac{1}{3} \times \pi \times 7.5^2 \times 20$ oe A1 for answer in the range 8240 − 8250 **OR** B1 for 2^3 M1 for a numerical expression for the volume of the large cone eg. $\frac{1}{3} \times \pi \times 15^2 \times 40$ (=9424...) M1 volume of frustrum = $\frac{7}{8} \times \frac{1}{3} \times \pi \times 15^2 \times 40$ oe A1 for answer in the range 8240 − 8250

Question		Working	Answer	Mark	Notes
22			1200 cm^3	4	M1 for $10 \times 2 \times 2$ and 15×2 M1 for "40" × "30" A1 for 1200 B1 (indep) for cm^3 OR M1 for 10×15 or 2^3 or 8 indicated as scale factor M1 for $10 \times 15 \times 2 \times 2 \times 2$ A1 for 1200 B1 (indep) for cm^3 SC B2 for 600 cm^3 (B1 for 600)

16		$120 \div 20 = 6$ $6^2 = 36$ $36 \times 300 = 10\ 800$	10 800	3	M1 $120 \div 20$ (= 6) oe, can be implied by $120^2 \div 20^2$ M1 '6'$^2 \times 300$ A1 cao

Question	Working	Answer	Mark	Notes
13		32.2	4	M1 for $BC = \frac{10}{4} \times 5$ (= 12.5) M1 for $EC = \frac{10}{4} \times 5.8 - 5.8$ (=8.7) M1(dep on at least M1) for '12.5' + '8.7' + 5 + 6 A1 cao OR M1 for $BC = \frac{5}{4} \times 10$ (= 12.5) M1 for $EC = \frac{5.8}{5} \times$ '12.5' – 5.8 (= 8.7) M1 (dep on at least M1) for '12.5' + '8.7' + 5 + 6 A1 cao OR M1 for $BC = \frac{5}{4} \times 10$ (= 12.5) M1 for $EC = \frac{6}{4} \times 5.8$ (= 8.7) M1 (dep on at least M1) for '12.5' + '8.7' + 5 + 6 A1 cao OR M1 for 4 + 5 + 5.8 (= 14.8) M1 for $\frac{10}{4} \times$ '14.8' (= 37) M1 (dep on at least M1) for '37' – 4 – 5.8 + 5 A1 cao

PAPER: 5MB3H_01

Question	Working	Answer	Mark	Notes
13		32.2	4	M1 for $BC = \frac{10}{4} \times 5$ (= 12.5) M1 for $EC = \frac{10}{4} \times 5.8 - 5.8$ (=8.7) M1(dep on at least M1) for '12.5' + '8.7' + 5 + 6 A1 cao OR M1 for $BC = \frac{5}{4} \times 10$ (= 12.5) M1 for $EC = \frac{5.8}{5} \times$ '12.5' – 5.8 (= 8.7) M1 (dep on at least M1) for '12.5' + '8.7' + 5 + 6 A1 cao OR M1 for $BC = \frac{5}{4} \times 10$ (= 12.5) M1 for $EC = \frac{6}{4} \times 5.8$ (= 8.7) M1 (dep on at least M1) for '12.5' + '8.7' + 5 + 6 A1 cao OR M1 for 4 + 5 + 5.8 (= 14.8) M1 for $\frac{10}{4} \times$ '14.8' (= 37) M1 (dep on at least M1) for '37' – 4 – 5.8 + 5 A1 cao

PAPER: 5MB3H_01

PAPER: 1MA0_1H

Question		Working	Answer	Mark	Notes
20			6	3	M1 for $\frac{15}{10}$ (=1.5) or $\frac{10}{15}$ (=0.66..) or $\frac{16}{10}$(=1.6) or $\frac{10}{16}$(=0.625) M1 for $\frac{15}{10}$ × 16 (=24) oe A1 cao OR M1 for $\frac{15}{16}$(=0.9375) or $\frac{16}{15}$ (=1.066…) or $\frac{16}{10}$(=1.6) or $\frac{10}{16}$(=0.625) M1 for $\frac{15}{16}$× 10 (=9.375) oe A1 20.625 oe

17	(a)		7.5	2	M1 for sight of $\frac{9}{6}$ (=1.5) oe or $\frac{6}{9}$ (=0.66..) oe or $\frac{5}{6}$ (=0.83..) oe or $\frac{6}{5}$ (=1.2) oe or a ratio, eg 6:9 oe or decimal, eg 1.5 oe A1 cao
	(b)		8	2	M1 for $12 \times \frac{6}{9}$ oe or $12 \div \frac{9}{6}$ oe or $\frac{12}{"7.5"} \times 5$ oe A1 cao

PAPER: 1MA0_2H

PAPER: 1MA0_1H

Question		Working	Answer	Mark	Notes
18	(a)		7.5	2	M1 for $\frac{12}{18}$ oe **or** $\frac{18}{12}$ oe **or** $\frac{12}{5}$ oe **or** $\frac{5}{12}$ oe A1 cao
	(b)		45	3	M1 for $\left("\frac{3}{2}"\right)^2$ oe or $\left("\frac{2}{3}"\right)^2$ oe M1 for complete method to find area of shaded region, eg $36 \times "1.5"^2 - 36$ A1 cao (SC B2 for 81)

3			21	2	M1 for 120 ÷ 30 (= 4) or 30 ÷ 120 (= 0.25) or w/30 = 84/120 oe A1 cao

21		$3^2 \times 180$	1620	2	M1 for using a scale factor of 3^2 (= 9) A1 cao

Original question 9

The full source mark scheme begins on page 271. Use the original paper code, tier and question number shown above.

Original question 22

The full source mark scheme begins on page 290. Use the original paper code, tier and question number shown above.

21		3.75 oe	3	M1 for a correct scale factor or ratio using two corresponding sides from similar triangles or two sides from the same triangle (may be seen in an equation) e.g $\frac{6}{1.5}$ oe or $\frac{1.5}{6}$ oe or $\frac{5}{6}$ or $\frac{6}{5}$ etc. (accept these written as ratios) M1 for a complete method to find ED A1

PAPER: 1MA0/1H

17		16	2	M1 for 18 ÷ 24 (= 0.75) or 24 ÷ 18 (= 1.333) A1 cao

Question	Working	Answer	Mark	Notes
15		18	4	M1 for correct initial use of Pythagoras eg $(AB^2 =)\ 10^2 - 6^2$ (= 64) or $AB = 8$ M1 (dep M1) for "$\sqrt{64}$" ÷ 2 (= 4) M1 for method to find area of trapezium eg $\frac{1}{2} \times$ "4" $\times (6 \div 2 + 6)$ A1 cao OR M1 for correct initial use of Pythagoras eg $(AB^2 =)\ 10^2 - 6^2$ (= 64) or $AB = 8$ M1 (dep M1) for method to find area of ΔABC eg $\frac{1}{2} \times$ "$\sqrt{64}$" $\times 6$ (= 24) or area of ΔAED $\frac{1}{2} \times 6 \div 2 \times$ "4" (= 6) or $24 \times (\frac{1}{2})^2$ (= 6) M1 for a complete method to find area of $EDBC$ e.g $\frac{3}{4} \times$ "24" eg "24" − "6" A1 cao

5 (a)		3.9	M1	for a ratio of $\frac{8.1}{5.4}$ (=1.5) oe or $\frac{5.4}{8.1}$ (=0.66..) oe or $\frac{2.6}{5.4}$ (= 0.48..) oe or $\frac{5.4}{2.6}$ (= 2.07..) oe
			A1	cao
(b)		2.05	M1	for $\frac{5.4}{8.1} \times$ 6.15 oe (= 4.1) or $\frac{2.7}{8.1} \times$ 6.15 oe or ft "scale factor" from (a)
			A1	cao

Paper: 1MA1/2H

Question	Working	Answer	Mark	Notes
15		Proof	C1	for identifying one pair of equal angles with a correct reason, e.g. (angle) *BAE* = (angle) *CDE*; angles in the same segment are equal or angles at the circumference subtended on the same arc are equal or for identifying two pairs of equal angles with no correct reasons given (angles must be within the appropriate triangles)
			C1	for identifying a second pair of equal angles with a correct reason, e.g. (angle) *AEB* = (angle) *DEC*; opposite angles or vertically opposite angles are equal or for identifying the three pairs of equal angles with no correct reasons given
			C1	for stating the three pairs of equal angles of the two triangles e.g. *ABE* = *DCE*, *BEA* = *CED*, *EAB* = *EDC* with fully correct reasons

22		2, 14.5	P1 A1 P1 A1 C1	for scale factor of $\frac{12}{3}$ **or** $\frac{3}{12}$ **or** $\frac{15}{12}$ **or** $\frac{12}{15}$ **or** $\frac{8}{12}$ **or** $\frac{12}{8}$ **or** $\frac{15}{8}$ oe **or** correctly identifies 2 pairs of corresponding sides for x=2 for complete method to find other value for x eg $\frac{15}{8}\times 12 - 8$ for x = 14.5 Describes both assumptions for similarity

14		Shown	M1 M1 A1	for $\sqrt[3]{\frac{8}{27}}$ (= $\frac{2}{3}$) or $\sqrt[3]{\frac{27}{8}}$ (= $\frac{3}{2}$) or 2 : 3 or 3 : 2 for $\left(\sqrt[3]{\frac{8}{27}}\right)^2$ (= $\frac{4}{9}$) or $\left(\sqrt[3]{\frac{27}{8}}\right)^2$ (= $\frac{9}{4}$) or 4 : 9 or 9 : 4 132 from correct arithmetic

Paper: 1MA1/3H

13	6.50	M1 M1 A1	for method to find ratio or scale factor of lengths or volumes eg $\sqrt{3} : 2$ or $1 : 1.15(47....)$ or $0.86(60\ldots) : 1$ or $\sqrt{27} : 8$ oe for complete method to find ratio of volumes and use to find required volume eg $10 \div (\text{"}1.15...\text{"})^3$ or $10 \times (\text{“}0.86\ldots\text{”})^3$ for answer in the range 6.49 to 6.53	Scale factors may just be seen as 1.15..., 0.86...etc If an answer is given within the range then incorrectly rounded to 3 sig figs, award full marks. Accept 6.5

Paper: 1MA1/3H

15	3 : 10	P1 P1 P1 A1	process to find ratio of lengths **A**:**B** = $\sqrt{4}$:$\sqrt{25}$ (= 2:5 or $\frac{2}{5}$ or 2, 5) for process to find ratio of lengths **B**:**C** = $\sqrt[3]{27}$: $\sqrt[3]{64}$ (= 3:4 or $\frac{3}{4}$ or 3, 4) for process to write as one ratio eg. finding a common multiple of 3 and 5 **or** 6 : 15 : 20 oe cao	Accept working in fractions for the award of process marks but the final answer must be in correct simplified ratio notation

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
17	4	P1	for process to find ratio of corresponding lengths, eg. $\sqrt{4}:\sqrt{9}$ (= 2 : 3)	
		P1	for process to find ratio of volumes, eg "2"^{3} : "3"^{3} (= 8 : 27)	
		P1	for "27" ÷ "8" (= 3.375)	This may be seen by checking their volume, eg. "8" × 4 (= 32) and "8" × 3 (= 24)
		A1	for rounding to give an answer of 4 from correct working	An answer of 4 with no supportive working gets no marks

Paper: 1MA1/1H

18	1.95	P1 P1 P1 P1 A1	for correct substitution into the cosine rule, eg $3.4^2 = 6.1^2 + 6.2^2 - 2 \times 6.1 \times 6.2 \times \cos BCA$ for a full process to find *BCA* eg (cos *BCA* =) $\frac{6.1^2 + 6.2^2 - 3.4^2}{2 \times 6.1 \times 6.2}$ **or** (*BCA* =) 32(.08046913…) correct substitution into the sine rule, eg $\frac{DC}{\sin("32.08..." \times \frac{2}{5})} = \frac{6.2}{\sin(180 - "32.08..." - ("32.08..." \times \frac{2}{5})}$ for complete process to find *DC* eg (*DC* =) $\frac{6.2 \times \sin "12.832"}{\sin "135.088"}$ Answer in the range 1.94 to 1.951	Can be any angle within triangle *ABC* P2 can be awarded for *BCA* = 32(.08046913…) Must not come from incorrect processing

Paper: 1MA1/3H

Paper: 1MA1/3H

Question	Answer	Mark	Mark scheme	Additional guidance
19	Proof	P1	for start to process to find area of *ABCDEF*, eg area of equilateral triangle $= \frac{1}{2} \times x \times x \times \sin 60$ $\left(= \frac{\sqrt{3}}{4}x^2\right)$ **OR** for start to process to find area of *FGHIJK*, eg area of equilateral triangle $= \frac{1}{2} \times px \times px \times \sin 60$ $\left(= \frac{\sqrt{3}}{4}p^2x^2\right)$	Any correct process to find the area of part of the hexagon is acceptable for this mark, eg $\frac{1}{2} \times x \times x \times \sin 120$ or $\frac{1}{2} \times (x + 2x) \times \frac{\sqrt{3}}{2}x$ Allow sin 60 left in expressions for the first 3 marks.
		P1	for complete process of finding area of *ABCDEF*, eg $6 \times \frac{1}{2} \times x \times x \times \sin 60$ or $6 \times \frac{1}{2} \times x \times x \times \frac{\sqrt{3}}{2}$ $\left(= \frac{3\sqrt{3}}{2}x^2\right)$ oe **OR** for complete process of finding area of *FGHIJK*, eg $6 \times \frac{1}{2} \times px \times px \times \frac{\sqrt{3}}{2}$ $\left(= \frac{3\sqrt{3}}{2}p^2x^2\right)$ oe	
		P1	for process of finding area of *ABCDEF* eg $\frac{3\sqrt{3}}{2}x^2$ oe **AND** for process of finding area of *FGHIJK*, eg $p^2 \times \frac{3\sqrt{3}}{2}x^2$ oe	
		C1	correct algebra leading to given result, $\frac{3\sqrt{3}}{2}(p^2 - 1)x^2$	Accept $\frac{3\sqrt{3}}{2}x^2(p^2 - 1)$ as final result.

Paper: 1MA1/3H

10	14.14	P1	works out scale factor eg (9 + 6) ÷ 6 (= 2.5) **OR** for start of process to find angle *DBE* eg sin $B = \frac{2}{6}$ oe	Note method can be carried out in either order
		P1	uses Pythagoras eg $6^2 - 2^2$ (= 32) or $\sqrt{32}$ (= 5.6...) **OR** calculates *AC* eg 2 × "2.5" (= 5) **OR** for complete process to find angle *DBE* eg $\sin^{-1}\left(\frac{2}{6}\right)$ (= 19.4...)	May be seen on diagram
		P1	complete process to find *CB* eg "2.5" × "√32" (= 10√2) **or** $\sqrt{(9+6)^2 - "5"^2}$ (= 10√2) **OR** uses trigonometry, eg 15 × cos "19.4..."	
		A1	14.1 to 14.15	If the answer is given within the range but then rounded incorrectly award full marks.

Paper: 1MA1/3H	

Paper: 1MA1/3H

Question	Answer	Mark	Mark scheme	Additional guidance
15	Proof	C1	for angle PQX = angle SRX as angles in the same segment are equal (or angles at the circumference subtended from the same arc/chord of a circle are equal)	Underlined words need to be shown; reasons need to be linked to their method.
			or angle QPX = angle RSX as angles in the same segment are equal (or angles at the circumference subtended from the same arc/chord of a circle are equal)	
			or angle PXQ = angle SXR as vertically opposite angles/ vertically opposite angles are equal	Could be shown on the diagram
			or for identifying two pairs of corresponding equal angles with no reason given	
		C1	for identifying two pairs of corresponding equal angles with correct reasons given	
		C1	for stating that the triangles are similar because all three pairs of corresponding angles are equal with complete reasons given.	Note that the students third/final reason may be: Angles in a triangle add up to 180

Paper: 1MA1/2H

Question	Answer	Mark	Mark scheme	Additional guidance
17	3.6	P1	process to find the volume scale factor, eg 1587.762 ÷ 58.806 (= 27) **or** 58.806 ÷ 1587.762 (= 0.037…)	
		P1	process to find the height of **B**, eg 2 × 43.74 ÷ 8.1 (= 10.8) **or** process to find the area of **A,** eg 43.74 ÷ $(\sqrt[3]{"27"})^2$ (= 4.86) **or** 43.74 × $(\sqrt[3]{"0.037..."})^2$ (= 4.86)	
		P1	complete process to find height of **A**, eg "10.8" ÷ $\sqrt[3]{"27"}$ **or** "4.86" × 2 ÷ (8.1 ÷ $\sqrt[3]{"27"}$)	
		A1	cao	

5	(a)	16	M1	for a ratio of $\frac{20}{5}$ or $\frac{5}{20}$ or 4 or 0.25 or $\frac{5}{4}$ or $\frac{4}{5}$ or 1.25 or 0.8 oe	
			A1	cao	
	(b)	5.5	M1	for 22 × "0.25" or 22 ÷ "4" oe	
			A1	oe	

Paper: 1MA1/3H

23	Shown	M1 M1 C1	for start to process using ratio, eg 1 : 1 + 2 : 1 + 2 + 3 (= 1 : 3 : 6) for process to write down ratio of areas of triangles eg $1^2 : 3^2 : 6^2$ (= 1 : 9 : 36) for correct working leading to answer, eg 1 : 9 – 1 : 36 – 9 = 1 : 8 : 27	Values may not be seen in a ratio, can still award M marks May be seen combined with algebra

Paper: 1MA1/2H

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
13	187.5	M1	for $\left(\frac{5}{2}\right)^3$ or $\left(\frac{2}{5}\right)^3$ or 2.5^3 or $\left(\frac{1}{2.5}\right)^3$ or $2^3 : 5^3$	
		M1	for a complete method, eg $12 \div 8 \times 125$ or 12×2.5^3	
		A1	oe	Accept $\frac{375}{2}$

Question	Answer	Mark	Mark scheme	Additional guidance
11	12	P1	for process of working with area scale factor, eg area SF = 50^2 (= 2500) or 0.5 × 0.5 (= 0.25) or 48 × 50 × 50 (= 120 000) **or** for process which uses 1 (m^2) = 10 000 (cm^2), eg $\frac{48}{10\,000}$	
		P1	for complete process, eg $\frac{"120\,000"}{100\times100}$ or “0.25” × 48	
		A1	cao	

Question	Answer	Mark	Mark scheme	Additional guidance
15	Shown	M1	for method to find comparable ratios eg (21.6 + 58.4) ÷ 32 (= 2.5) **and** (32 + 22) ÷ 21.6 (= 2.5) **or** 32 ÷ (21.6 + 58.4) (= 0.4) **and** 21.6 ÷ (32 + 22) (= 0.4) **or** 21.6 ÷ 32 (= 0.675) **and** (32 + 22) ÷ (21.6 + 58.4) (= 0.675) **or** 32 ÷ 21.6 (= 1.4̇8̇) **and** (21.6 + 58.4) ÷ (32 + 22) (= 1.4̇8̇)	Fractions may be used. Accept 3 sf or more.
		C1	for correct conclusion supported by the correct evaluation of 2 ratios to show they are equal	We do not require reference to the common angle.

Question	Answer	Mark	Mark scheme	Additional guidance
19	2 : 5	P1	for using similar triangles to form an equation eg $\frac{AB}{BC}=\frac{AD}{AB}$ oe or $\frac{AB}{4}=\frac{25}{AB}$ oe or $\frac{AB}{4k}=\frac{25k}{AB}$ oe or $AB:4=25:AB$ oe or $BC\times\text{sf}=BD\div\text{sf}$ oe or $4\times\text{sf}=25\div\text{sf}$ oe **or** for working with the perpendicular height of triangle ABC eg $(h^2=)\ 25^2-23^2\ (=96)$ or $(h=)\ \sqrt{25^2-23^2}\ (=\sqrt{96})$	May use x or any other letter for AB Accept $AB = 4 \times$ sf, $AB = 25 \div$ sf
		P1	for process to find AB eg $(AB=)\ \sqrt{4\times25}\ (=10)$ oe or $\sqrt{"96"+2^2}\ (=10)$ **or** for process to find the scale factor eg $\sqrt{\frac{25}{4}}\ (=\frac{5}{2}$ oe)	
		A1	oe	

Question	Answer	Mark	Mark scheme	Additional guidance
17	12 : 15 : 5	P1	for process to find ratio of heights of **L** and **M** eg $\sqrt[3]{64} : \sqrt[3]{125}$ (= 4 : 5) oe or $\sqrt[3]{\frac{64}{64}} : \sqrt[3]{\frac{125}{64}}$ $\left(=1:1.25\right)$ oe or $\sqrt[3]{\frac{64}{125}} : \sqrt[3]{\frac{125}{125}}$ $\left(=0.8:1\right)$ oe	Condone not written as a ratio as long as clear $\frac{125}{64}=1.953...$ $\frac{64}{125}=0.512$
		P1	for process to find ratio of heights of **M** and **P** eg $\sqrt{144}$: $\sqrt{16}$ (= 12 : 4 = 3 : 1) oe or $\sqrt{\frac{144}{16}} : \sqrt{\frac{16}{16}}$ $\left(=3:1\right)$ oe or $\sqrt{\frac{144}{144}} : \sqrt{\frac{16}{144}}$ $\left(=1:0.\dot{3}\right)$ oe	Condone not written as a ratio as long as clear $\frac{144}{16}=9$ $\frac{16}{144}=0.\dot{1}$
		P1	(dep on P2) for process to find ratio of heights of all 3, eg "(4 : 5)" × 3 and "(3 : 1)" × 5 or $\left(1:1.25\right)\times 12$ and $\left(3:1\right)\times 5$ or $\left(0.8:1\right)$ and $\left(1:0.\dot{3}\right)$	
		A1	for 12 : 15 : 5 oe	Can ISW incorrect simplification of a correct ratio

Paper: 1MA1/2H

12	64 : 25	P1	for start of process to find ratio of lengths of **A** to **B**, eg $\sqrt[3]{64}$ (= 4) **or** $\sqrt[3]{125}$ (= 5) **or** 4 : 5	
		P1	for $\sqrt[3]{125} \div 2$ (= 2.5) oe **or** $\left("\sqrt[3]{64}"\right)^2$ (= 16)	
		P1	for process to find ratio of areas of **A** to **C**, eg "4"^{2} : "2.5"^{2} (= 16 : 6.25)	
		A1	for 64 : 25 oe in form a : b where a and b are integers	

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
13	18.9	P1	for using length scale factor to find *AB*, eg (*AB* =) $10.8 \div \left(\frac{3}{2}\right)$ or (*AB* =) $10.8 \times \left(\frac{2}{3}\right)$ (= 7.2) for using length scale factor to find *BC*, eg (*BC* =) $10.8 \div \frac{6.3}{6.3-4.2}$ or (*BC* =) $10.8 \times \frac{6.3-4.2}{6.3}$ (= 3.6) **or** finds area scale factor, eg $\left(\frac{3}{2}\right)^2$ or $\left(\frac{2}{3}\right)^2$	Can use a combination of skills but must have a complete process to find *AB* or *BC* to score this mark
		P1	for a complete process to find the area of trapezium, eg $\frac{6.3+4.2}{2} \times (10.8 - "7.2")$ or $\frac{(6.3-4.2) \times "3.6"}{2} + "3.6" \times 4.2$ or $\frac{10.8 \times 6.3}{2} - \frac{7.2 \times 4.2}{2}$ or $\frac{10.8 \times 6.3}{2} - \frac{10.8 \times 6.3}{2} \div \left("\frac{3}{2}"\right)^2$ or $\frac{10.8 \times 6.3}{2} - \frac{10.8 \times 6.3}{2} \times \left("\frac{2}{3}"\right)^2$	
		A1	accept trailing zeros eg 18.90	

Paper: 1MA1/3H

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
8 (a)	Proof	M1	for identifying angle *A* as common **or** for angle *AED* = angle *ACB* or angle *ADE* = angle *ABC* with appropriate reason(s) eg corresponding angles are equal or co-interior angles add up to 180 and angles on a straight line add up to 180	
		C1	for completing the proof by identifying a second pair of equal angles with appropriate reason(s) **and** a statement that the angles in each triangle are the same	Statement can be implied by identifying a third pair of equal angles (no reason needed)
(b)	6	P1	for a scale factor of 1.5 or $\frac{2}{3}$ oe **or** for $\frac{AE}{20}=\frac{18}{30}$ oe	
		P1	for a process to find the length of *AE*, eg $18 \div 1.5\ (=12)$ oe **or** $(AE =)\frac{20\times18}{30}$ oe **or** for a complete process to find the length of *EC*, eg $18\times(1-\frac{2}{3})$ oe	
		A1	cao	

Question	Answer	Mark	Mark scheme	Additional guidance
16 (a)	4.9	P1	for beginning to work with scale factors eg $\left(\frac{h}{12}\right)^3$ **or** $\left(\frac{12}{h}\right)^3$ **or** $\frac{90}{1350}\left(=\frac{1}{15}\right)$ **or** $\frac{1350}{90}\left(=15\right)$ **or** 90 : 1350 oe	
		P1	for equating scale factors eg $\left(\frac{h}{12}\right)^3=\frac{90}{1350}$ **or** $\left(\frac{12}{h}\right)^3="15"$ **or** for finding the linear scale factor, eg $\sqrt[3]{\frac{90}{1350}}$ **or** $\sqrt[3]{"15"}\left(=2.46(6...)\right)$	
		P1	for a complete method eg $\sqrt[3]{12^3\times\frac{90}{1350}}$ **or** $12\times\sqrt[3]{\frac{90}{1350}}$ **or** $\frac{12}{"\left(\sqrt[3]{"15"}\right)"}$	
		A1	for answers in the range 4.8 to 4.9	
(b)	$(15)^{\frac{2}{3}}$	B1	for $15^{\frac{2}{3}}$ **or** $\left(\sqrt[3]{15}\right)^2$ **or** $\sqrt[3]{15^2}$ **or** $225^{\frac{1}{3}}$	If given in an acceptable form, then given as a decimal, isw and award

GCSE

Mark Scheme

Mathematics A (1387)

June 2003

Edexcel
Success through qualifications

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information please call our Customer Response Centre on 0870 240 9800, or visit our website at www.edexcel.org.uk.

June 2003

Publications Code UG014129

All the material in this publication is copyright
© London Qualifications Ltd 2003

CONTENTS

Notes on Marking Principles 2
Paper 5501 4
Paper 5502 11
Paper 5503 17
Paper 5504 24
Paper 5505 32
Paper 5506 40

NOTES ON MARKING PRINCIPLES

NOTES ON MARKING PRINCIPLES

1 Three types of mark are available

M marks - awarded for correct working seen or implied.
A marks - conditional accuracy marks which are awarded for accurate working following the award of M marks.
B marks - unconditional accuracy marks (independent of M).

2 Abbreviations

cao - correct answer only.
ft - follow through.
√ - Denotes a "follow through" answer.

SC - special case.
isw - ignore subsequent working.
oe - or equivalent (and appropriate).
NB: a candidate cannot benefit from both isw and ft.

3 If no working shown, then

correct answers normally score full marks
incorrect (even though nearly correct) answers score no marks.

4 Marking instructions

Misread loses A marks (and sometimes B marks) on that part but ft can be allowed on subsequent parts. M marks can still be earned. If in doubt contact your team leader.
If there is a wrong answer in the answer space DO CHECK the working in the body of the script.

- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.
- If there is a wrong answer in the answer space DO CHECK the working in the body of the script.
- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.

5 Style of marking

Answer correct: tick and write part mark in margin NEXT TO BRACKETED MARK.
Answer incorrect: cross, but show M, A or B marks if any earned in body of script and transfer the total of these to the margin next to the bracket mark.
Total for each double page at bottom right page (except for back if used), FINAL TOTAL IN RELEVANT BOX ON FRONT COVER.
Nought in margin for fully incorrect question or page AND FOR NO ATTEMPT.
Where no attempt has been made a line should be put in the answer space and zero in the margin next to the bracketed mark
CHOICE OF METHOD – No marks unless one answer is in answer space – then mark that.
CROSSED OUT WORK - if not replaced this should be marked (if legible).
There must always be a mark next to bracketed mark in the margin/

6 Follow Through Marks

Follow throughs are guided by two principles:

(a) Follow throughs which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous, do not award.

(b) Follow throughs which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

7 Probability

Probability answers must be given as fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least two decimal places (unless tenths).

Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.

8 Linear Equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

9 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

Paper 5501

No	Working	Answer	Mark	Notes
1 (a)		46	1	B1 cao
(b)		3.4	1	B1 oe
(c)		Arrow at 430	1	B1 allow ± half graduation
(d)		Arrow at 3.7	1	B1 allow ± half graduation Accept indications other than arrows as long as they are clear
2		1.60 2.05	2	B1) Condone B1) reversal
3 (a)	$\frac{18}{24}$ or $\frac{9}{12}$ or $\frac{6}{8}$	$\frac{3}{4}$	2	B2 for $\frac{3}{4}$ cao (B1 for $\frac{18}{24}$ or $\frac{9}{12}$ or $\frac{6}{8}$) SC B1 for $\frac{1}{4}$ only
(b)		16 squares shaded	1	B1 cao
4 (a)		line	1	B1 within overlay tolerance
(b)		midpoint	1	B1 within overlay tolerance ft from (a) ± 0.2cm
(c)		rectangle	1	B1 for rectangle 6 cm ± 0.2cm by 4 cm ± 0.2cm
5		kilograms, kg litre, *l* or cubic metres, m^3 inches, in	3	B1 B1 B1
6 (a)		parallel lines marked	1	B1
(b)		right angle marked	1	B1
(c)(i)		acute	1	B1
(ii)		reflex	1	B1

Paper 5501

No	Working	Answer	Mark	Notes
7 (i)		sphere	1	B1
(ii)		cylinder	1	B1 Accept circular prism
(iii)		pyramid	1	B1 Condone omission of "triangular" Accept tetrahedron
8 (a)(i)		40	2	B1 cao
(ii)		50		B1 cao
(b)		5 complete symbols	1	B1 cao
(c)		 and oe inc	1	B1
9 (i)		9, 37, 56, 59, 75	5	B1 cao
(ii)		0.067, 0.56, 0.6, 0.605, 0.65		B1 cao Ignore trailing zeros
(iii)		–10, –6, –4, 2, 5		B1 cao
(iv)		$\frac{2}{5}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}$		B2 for all 4 correct (B1 for any 3 in correct order) SC B1 for all 4 in reverse order (applies to(iv) only)
10 (a)			1	B1
(b)		Plotting (4, 24)	1	B1 ft from their matchsticks
(c)		60	1	B1 cao
(d)		$m = 6n$	2	B2 for $m = 6n$ oe (B1 for $6n$ oe or m = multiple of n except $m = n$)
11 (i)		6, 12	4	B1 cao
(ii)		4, 16		B1 cao
(iii)		3, 4, 6 or 3, 4, 6, 12		B1 Condone omission of 12
(iv)		8, 27		B1 cao

Paper 5501

No	Working	Answer	Mark	Notes
12		2.43	4	B1 for 36 or 0.36 B1 for 96 or 0.96 B1 for 125 or 1.25 If none of first 3 B1s awarded then SC B1 for four 24s and five 25s seen OR 4 × 24 and 5 × 25 seen B1 for 2.43 cao
13 (a)(i)		Edinburgh and Plymouth	3	B1 for Edinburgh or −7 B1 for Plymouth or 5
(ii)		12		B1ft from (i) if one positive and one negative
(b)		Cardiff and Belfast London and Plymouth	2	B1 for Cardiff and Belfast OR −6 and −4 B1 for London and Plymouth OR 3 and 5
14		5 9 8	3	B1 cao B1 cao B1 cao
15 (a)			2	✓ × : 2 0 B2; 1 0 B1; 1 1 B1; 2 1 B1
(b)	$2 \times 2 \times 2$	8	2	M1 for $2 \times 2 \times 2$ A1 for 8 cao
16 (a)		30	1	B1 cao
(b)		3	1	B1 cao
(c)	$2 \times 29 = 58$ $5 \times 30 = 150$ $2 \times 31 = 62$ $1 \times 32 = 32$ $\frac{302}{10} = 30.2$	30.2	3	M1 for freq × no pins M1 (dep on 1st M1) for totalling and ÷ 10 A1 for 30.2 cao

Paper 5501

No	Working	Answer	Mark	Notes
17		7 5 5 13 33 52 23	3	B3 all correct (B2 for 4, 5 or 6 correct B1 for 2 or 3 correct)
18 (a)(i) (ii) (iii) (iv) (b)		$4c$ p^4 $8g$ $10pr$ OR $10rp$ $10y - 15$	4 1	B1 oe B1 cao B1 oe B1 B1 cao Accept $10y + - 15$
19		$\frac{2}{3}$	3	M1 for 3 rows (9 squares) shaded M1 for 2 columns (10 squares) shaded A1 for $\frac{2}{3}$
	$\frac{3}{5} = \frac{9}{15}$ $\frac{2}{3} = \frac{10}{15}$	$\frac{2}{3}$	3	M1 for $\frac{3}{5} = \frac{9}{15}$ M1 for $\frac{2}{3} = \frac{10}{15}$ A1 for $\frac{2}{3}$
	$\frac{3}{5} = 0.6$ $\frac{2}{3} = 0.66$ or 0.67 or better	$\frac{2}{3}$	3	M1 for $\frac{3}{5} = 0.6$) Accept M1 for $\frac{2}{3} = 0.66$ or 0.67 or better) percentages A1 for $\frac{2}{3}$

Paper 5501

No	Working	Answer	Mark	Notes
20 (a)	955 × 48 7640 38200 45840 OR 48 × 955 240 2400 43200 45840	458.40	3	M1 for complete correct method (condone one computational error) A2 for 458.40 cao (A1 for digits 4584 OR ft if M1 awarded)
(b)	14. 5 48 | 696. 0 48 216 192 24 0 24 0	14.50	3	M1 for 1 as first digit in answer and remainder 21 M1 (dep) 4 as second digit in answer A1 for 14.50 (Accept 14.5)
21 (a)		$12x$	1	B1 oe
(b)		$12x + 10y$	2	B2 oe ft from (a) (B1 $12x$ + multiple of y or $10y$ seen) SC B1 for $x = 12x + 10y$ OR $y = 12x + 10y$

Paper 5501

No	Working	Answer	Mark	Notes
22	$\frac{1}{3}+\frac{1}{4}=\frac{4}{12}+\frac{3}{12}=\frac{7}{12}$ $1-\frac{7}{12}=\frac{5}{12}$	$\frac{5}{12}$	3	M1 for $\frac{4}{12}$ **and** $\frac{3}{12}$ oe A2 for $\frac{5}{12}$ oe (A1 for $\frac{7}{12}$) B1 for 1–"$\left(\frac{1}{3}+\frac{1}{4}\right)$" correctly evaluated
	$1-\frac{1}{3}=\frac{2}{3}$, $1-\frac{1}{4}=\frac{3}{4}$ $\frac{8}{12}-\frac{3}{12}=\frac{5}{12}$ or $\frac{9}{12}-\frac{4}{12}=\frac{5}{12}$	$\frac{5}{12}$	3	B1 for $\frac{2}{3}$ or $\frac{3}{4}$ seen M1 for $\frac{8}{12}-\frac{3}{12}=\frac{5}{12}$ or $\frac{9}{12}-\frac{4}{12}=\frac{5}{12}$ A1 for $\frac{5}{12}$ oe
		$\frac{5}{12}$	3	M1 for 0.25 and 0.33 or better A1 for 0.58 or better A1 for $0.41\dot{6}$ or recurring
23 (a)		54	1	B1 cao
(b)(i)	180 – (54 + 54)	72	3	M1 for 180 – (54 + 54) A1 ft from (a) if $x < 90$
(ii)		Reason		B1 for mentioning isosceles and equal or base angles or equal sides and equal or base angles

Paper 5501

No	Working	Answer	Mark	Notes
24 (a)		Bryani	2	M1 for 4×9 or $4 \times 3 \times 3$ or $4 \times x \times x$ or square x first or square 3 first A1 SC 4×3^2 with Bryani scores B2
(b)		64	1	B1 cao
25 (a)		 dotted line may be solid	3	B2 for rectangle base 3 squares and height 4 squares (B1 for rectangle with one correct dimension) B1 for horizontal line 1 cm from top) SC B2 for completely correct elevation on its side
(b)			2	B2 for perspective drawing showing slant face and cutout B1 for perspective drawing with either slant face cutout omitted
26 (a)	20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$	40	2	M1 for 20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$ A1 cao
(b)	$\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes	line from (45, 20) to (65, 0)	2	M1 for $\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes seen A1 for correct line SC If M0, B1 for line from (45, 20) to (t, 0) where $t > 45$ or B1 for a line of the correct gradient.

Paper 5502

No	Working	Answer	Mark	Notes
1 (a)		$\frac{7}{100}$	1	B1 cao accept 0.07
(b)		0.18	1	B1 cao
(c)	20 in 100 oe	40	2	M1 for sight of 20 in 100 or 20 × 2 A1 cao
2 (a)		6cm	2	B1 for 6 ± 0.2 or 60 ± 2 B1 indep for cm or mm consistent with 1st B1
(b)		At centre	1	B1 within overlay
(c)		Circle drawn	1	B1 all within overlay
3		See diagram	3	B3 all correct – see separate sheet (B2 for 3 correct B1 for 2 correct)
4 (a)		$\frac{1}{4}$ oe	1	B1 cao
(b)		0.75	1	B1 cao
(c)		75%	1	B1 cao
(d)(i)		9	2	B1 accept answer in range 9 - 9.2
(ii)		15 - 16		B1 accept answers in range15 - 16
5 (a)		9:30	1	B1 cao
(b)		2hrs 45 min	2	B2 for 2hr 45 min or 2 $\frac{3}{4}$ hr or 165 minutes B1 2:45 or 2.45 or 165 or 45min + 1hr + 1hr oe
(c)		17	1	B1 cao
6 (a)	96 × 4	3.84	2	M1 for 96 × 4 or digits 384 A1 cao
(b)	3 × 96 + 40 = 328	(0).56	2	M1 for 3 × 96 + 40 or digits 328 or digits 56 A1 cao accept 56p

Paper 5502

No	Working	Answer	Mark	Notes
7 (a)		54 000	1	B1 cao accept 54 thousand
(b)		50 000	1	B1 (accept ten thousand or 10 000) oe
8 (a)		14	1	B1 cao
(b)		6	1	B1 cao
(c)		Correct reflection	2	B2 fully correct (B1 correct reflection in a line parallel to the mirror line or condoning 1 block error in shape or position of shape)
9 (a)		Missing horiz label 1 (and 6) missing on vertical scale	2	B1 B1
(b)		Correct graph	2	B1 for bar up to 4 for yellow B1 for bar up to 2 for green
(c)		Blue	1	B1 cao
(d)	3 + 5 + 4 + 2	14	1	B1 ft from (b)
(e)		$\frac{3}{14}$	1	B1 ft on '14'
10	Barry (8) because you double Kath (7) because you add, 1,2,3		2	B1 oe B1 oe SC: B1 for correct rules only
11 (a)		$2n$	1	B1 for $2n$ or $n + n$ OR $2 \times n$ OR $n \times 2$ OR $n2$
(b)		$2n + 15$	1	B1 for "$2n$" + 15 oe
(c)		20q	1	B1 cao
12 (a)	1 + 3 + 5 + 8 + 5	22	2	M1 add frequencies A1 cao
(b)		No, is>no of cups of coffee in the table	1	B1 'average cannot be bigger than 6' oe OR 'Average must be less than 6 oe'
13 (a)		Trapezium	1	B1 cao ignore spelling
(b)		(2, 3)	1	B1 cao
(c)		Isosceles	1	B1 cao ignore spelling
(d)		Q correct	1	B1 cao

Paper 5502

No	Working	Answer	Mark	Notes
14 (a)		250 000	1	B1 cao
(b)	$\frac{28}{4}$	7	2	M1 for $\frac{28}{4}$ oe or "250000" × 28 A1 cao **SC** B1 for 7 000 000
15 (a)		10	1	B1 cao
(b)		5.5	1	B1 ±0.3 pounds
(c)	$\frac{110}{22}$	50	3	M1 for use of graph at 11 or $\frac{110}{22}$ A1 for 5 A1 cao **SC** B2 for 49.5 – 50.6

Paper 5502

No	Working	Answer	Mark	Notes
16 (a)	$269.30 - 56.80 = 212.50$ $\frac{212.50}{42.50}$	6	2	M1 for $\frac{269.30 - 56.80}{42.50}$ or 5 seen A1 cao
(b)	5% of £269.30 £269.30 – "£13.465" OR $\frac{95}{100} \times 269.30$	255.83 or 255.84	3	M1 for $(5 \div 100) \times 269.30$ M1 for $269.30 -$ "13.465" A1 cao OR M2 for $\frac{95}{100} \times 269.30$ A1 cao **Alternative Method:** M1 for $\frac{5}{100} \times 56.80$ $(= 2.84)$ and $\frac{5}{100} \times 42.50$ (=2.12(5)) (OR 53.96 AND 40.38 (40.375) seen M1 for $56.80 -$ "2.84" (= 53.96) $42.50 -$ "2.12(5)" (= 40.375 or 40.38) "5" × "40.375" + "53.96" A1 cao
17	$4.1^2 \times 1.07 = 16.81 \times 1.07$	17.9867	2	M1 for ("4.1") followed by squaring, or sight of 16.81 A1 cao SC: *B1 for 18 or better with no working*
18 (a)		60	1	B1 cao
(b)	$360 - 60 - 90 - 90$	120	2	M1 for $360 -$ " 60 " $- 90 - 90$ or $180 -$ "60" A1 cao
(c)	6×2	12	2	M1 for 6×2 A1 cao for 12

Paper 5502

No	Working	Answer	Mark	Notes
19 (a)(i)	$240 \times 5 = 1200$	1250	3	B1 cao 1250
(ii)	$\frac{50}{'1250'}$	$\frac{1}{25}$		M1 cao $\frac{50}{'1250'}$ A1 for $\frac{1}{25}$ in its simplest form
(b)	$\frac{60}{100} \times 1000 = 600$	12:5	3	M1 for $\frac{60}{100} \times 1000$ oe A1 for 600 A1 cao
20 (a)		$x + 2$	1	B1 accept $2 + x$ but not $x = x + 2$
(b)	$x + 5 + x + 5 + x + 2 + x + 2$	$4x + 14$	2	M1 adding 4 sides, two of which are '$x + 2$' (all sides to be linear expressions in x) **SC** $x + 5 + x + 2 \times 2$ gets M1 A1 for correct simplified answer or $(20 - 14) \div 4$ oe gets M1
(c)	'$4x + 14$' = 20	1.5 oe	2	M1 for equation A1 cao
21	$\sum f = 90$	Angles drawn, labelled	3	M1 for 1 person = 4°or one angle correct in table or pie chart A1 any 2 correctly drawn angles in pie chart A1 fully correct chart labelled
22 (a)		$2p - q$	2	B1 cao for $2p$ B1 cao for $-q$ accept $(-q + 2p)$, $2p - 1q$ and $2p + -q$
(b)	$5x = 3 + 4$	1.4	2	M1 for either (+3 or sight of 7) or (÷ 5 or sight of 0.8 and 0.6) A1 cao accept $\frac{7}{5}$ or $1\frac{2}{5}$

Paper 5502

No	Working	Answer	Mark	Notes
23 (a)		$\frac{4\times5}{2}$	1	B1 cao
(b)	$1+2+3+4+5+6+7+8$	$\frac{8\times9}{2}$	1	B1 cao
(c)	$\frac{100\times101}{2}$	5050	1	B1 cao
24	0 | 5 7 8 8 1 | 0 0 0 0 2 5 5 5 6 2 | 0 0 0 4 5 3 | 3 5 Key 1 | 2 = 12 (min)	See working	3	B1 for stem 0, 1, 2, 3 or 0, 10, 20, 30 B1 for accurate unordered leaves condone 1 error or omission B1 for key and ordered leaves all correct
25	$3.2\times2.8=8.96$ $2\times4.5\times2.8=25.2$ $2\times4.5\times28.8=28.8$ $\frac{62.96}{2.5}\times2.99$	£75.30	5	M1 for area of any face found correctly M1 for 2 areas found correctly A1 for 62.96 or 54 M1 for $\frac{'62.96'}{2.5}\times2.99$ Al cao Alternate method for candidates who round up $\frac{"62.92"}{2.5}$ M1 for "26" × 2.99 A1 for £77.74 cao **SC**: *for top included B2 for 71.92 m^2 seen or* B3 for £86.02 seen **SC** B4 for £64.58 or £65.78 seen
26	$2.5\times10\,000$	25 000	2	M1 for $2.5\times100\times100$ A1 cao

Paper 5503

No	Working	Answer	Mark	Notes
1 (a)(i)		$8g$	2	B1 oe
(ii)		$10rp$		B1 for $10pr$ or $10\ rp$
(b)		$10y - 15$	1	B1 cao accept $10y$ - + 15
(c)	$6x + 8 - 12x + 15$	$-6x + 23$	2	M1 for 3 correct terms out of 4 A1 cao
2		$m = 6n$	2	B2 for $m = 6n$ oe accept $6 \times n$, n 6 (B1 for $6n$ alone, or $6n$ +1 oe OR m = multiple of n except $m = n$)
3 (i)		0.067, 0.56, 0.6, 0.605, 0.65	1	B1 cao Ignore trailing zeros
(ii)		$-10, -6, -4, 2, 5$	1	B1 cao
(iii)		$\frac{2}{5}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}$	2	B2 all four correct (B1 any three in correct order) SC: B1 all 4 in reverse order
4		7 5 5 13 33 52 23	3	B3 all correct (B2 for 4, 5 or 6 correct entries) (B1 for 2, 3 correct entries)

Paper 5503

No	Working	Answer	Mark	Notes
5		$\frac{2}{3}$	3	M1 for 3 rows (9 squares) shaded M1 for 2 columns (10 squares) shaded A1
	OR $\frac{3}{5}=\frac{9}{15}$ $\frac{2}{3}=\frac{10}{15}$ Therefore $\frac{2}{3}>\frac{3}{5}$	$\frac{2}{3}$		M1 for $\frac{3}{5}=\frac{9}{15}$ M1 for $\frac{2}{3}=\frac{10}{15}$ A1
	OR $\frac{3}{5}=0.6$ or percent $\frac{2}{3}=0.66$ or 0.67 or better Therefore $\frac{2}{3}>\frac{3}{5}$	$\frac{2}{3}$		M1 for $\frac{3}{5}=0.6$ or percent M1 for $\frac{2}{3}=0.66$ or 0.67 or better A1
6	$2\times 29=58$ $5\times 30=150$ $2\times 31=62$ $1\times 32=32$ $\frac{302}{10}=30.2$	30.2	3	M1 for freq × no. pins (at least 3) M1 for totalling and for ÷ 10 (dep on 1st M1) A1 cao
7 (a)			2	B1 for 180° rotation (wrong centre) B1 cao
(b)			3	B1 for any enlargement sf other than 1 B1 for all sides halved B1 for position

Paper 5503

No	Working	Answer	Mark	Notes
8 (a)		$12x$	1	B1 oe
(b)		$12x + 10y$	2	B2 oe or ft from (a) (B1 for $12x$ + a multiple of y or $10y$ seen) **SC** B1 for $x = 12x + 10y$ or $y = 12x + 10y$
9	$\frac{1}{3}+\frac{1}{4}=\frac{4}{12}+\frac{3}{12}=\frac{7}{12}$ $1-\frac{7}{12}=\frac{5}{12}$ OR $1-\frac{1}{3}=\frac{2}{3}, 1-\frac{1}{4}=\frac{3}{4}$ $\frac{9}{12}-\frac{4}{12}=\frac{5}{12}, \frac{8}{12}-\frac{3}{12}=\frac{5}{12}$	$\frac{5}{12}$	3	M1 for $\frac{4}{12}$ and $\frac{3}{12}$ oe A2 for $\frac{5}{12}$ oe (A1 for getting $\frac{7}{12}$) (B1 for $1 - \frac{"1"}{3} + \frac{1"}{4}$ correctly evaluated) OR B1 for $\frac{2}{3}$ or $\frac{3}{4}$ seen M1 for $\frac{3}{4}-\frac{1}{3}=\frac{9}{12}-\frac{4}{12}$ or $\frac{2}{3}-\frac{1}{4}=\frac{8}{12}-\frac{3}{12}$ A1 for $\frac{5}{12}$ oe OR M1 0.25 and 0.33 or better A1 for 0.58 or better A1 for 0.416 *recurring*
10 (a)		54	1	B1 cao
(b)(i)	180 – (54 +54)	72	3	M1 for 180 – (“54” + “54”) A1 for 72° ft from (a) if $x < 90°$
(ii)				B1 for mentioning isosceles and equal or base angles or equal sides and equal or base angles

Paper 5503

No	Working	Answer	Mark	Notes
11 (a)	Bryani was correct $4 \times 3^2 = 4 \times 9 = 36$	Bryani	2	M1 for 4×9 or $4 \times 3 \times 3$ or "square the three/x then multiply by four" A1 Bryani **SC** 4×3^2 with Bryani gets B2
(b)			1	B1 cao
12 (a)			3	B2 for rectangle height 4 squares, base 3 squares (B1 for rectangle with one correct dimension) B1for line 1 square from the top **SC** B2 for completely correct elevation on its side
(b)			2	B2 for perspective drawing showing slant face and cut out (B1 for perspective drawing with either slant face or cut out omitted or one aspect incorrect)
13 (a)	20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$	40	2	M1 for 20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$ A1 cao
(b)	$\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes seen	Line from (45, 20) to (65, 0)	2	M1 for $\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes seen A1 for correct line **SC** If M0, B1 for line from (45, 20) to (t, 0) where $t > 45$ or a line of the correct gradient
14 (i)		119.31	3	B1 cao
(ii)		119 310		B1 cao
(iii)		1.23		B1 cao

Paper 5503

No	Working	Answer	Mark	Notes
15	$\frac{10}{100}\times 12000$ 12 000 – 1200 = 10 800 $\frac{10800}{10}=1080$ 10 800 – 1080 = £9720	£9720	3	M1 for $\frac{10}{100}\times 12000$ or sight of 1200 or 2400 or 10 800 or 9600 M1 (dep) for $\frac{10}{100}\times(12\,000-\frac{10}{100}\times 12000)$ or sight of 1080 A1 cao Alternative mark scheme M2 for $12000\times(1-\frac{10}{100})^2$ (M1 for $12000\times(1-\frac{10}{100})$ A1 cao
16 (a)	$2p = 6$	$p = 3$	2	M1 for $7p - 5p = 8 - 2$ or $2p$ or 6 A1 cao
(b)	$7r - 5r = -20 - 2$	-11	2	M1 for $7r + 2 = 5r - 20$ or $\frac{7r}{5} + \frac{2}{5} = r - 4$ or $7r - 5r = 20 - 2$ or $\frac{7r}{5} - r = -4 - \frac{2}{5}$ A1 cao
17		$5n + 1$	2	B2 oe (B1 for $5n$ seen) NB: n = gets B1 max

Paper 5503

No	Working	Answer	Mark	Notes
18 (a)	–1, 0, 1		2	B2 for –1, 0, 1 (B1 for –1, 0 or 0,1 or –1,1 or –2, -1, 0, 1 only)
(b)	(–1, –1), (0, –1), (1, –1), (0, 0), (1, 0), (1, 1)		3	B3 for 6 points correct B2 for 3 points correct B1 for 1 point correct **NB** –B1 each additional point over six
19 (a)	Triangle with vertices at (0,0) (0,–2) and (3,0)		2	M1 for correct orientation A1 cao
(b)	Rotation, 180°, centre (0,1) Enlargement sf - 1 centre (0,1)		2	B2 for 180° 'rotation' centre (0, 1) B2 for Enlargement sf - 1 centre (0,1) (B1 for any two of the three parts) NB: B0 if additional transformation is included
20	Bisector of $\angle BAC$ Arc around A Region		3	B3 cao (B2 for either two correct boundaries, no shading/ wrong shading or one correct boundary, one incorrect boundary with valid shading) (B1 for either two incorrect boundaries but one drawn from A and one intersection, with valid shading or one correct boundary) Ignore shading outside the triangle
21		Length Volume Area	3	B1 for Length B1 for Volume B1 for Area
22 (a)	Unbiased question with choices		2	B1 for unbiased question B1 for at least 2 choices
(b)			2	Classification 1: A biased question Classification 2: A restricted sample of people Classification 3: Not specifying a range of foods Classification 4: Nothing to do with eating habits B2 reasons which satisfy 2 different classifications (B1 a reason which satisfies one classification)



Paper 5503

No	Working	Answer	Mark	Notes
23 (a)	$6 \times 10^2 \times 8 \times 10^4$ 48×10^6	4.8×10^7	3	M1 for $6 \times 10^a \times 8 \times 10^b$ oe, a and b integers including 0 A1 for 48×10^6 oe A1 cao
(b)	200 000 + 30 000	230000	2	B2 cao (B1 for sight of 200 000 or 30 000 or 2.3×10^5 or 23×10^4)
24 (i)		64	4	B1 cao
(ii)		3		B1 cao
(iii)	$\sqrt{16 \times 9} = \sqrt{144}$	12		B2 cao (B1 for sight of $\sqrt{2^4} \times \sqrt{9}$ or better, or 144 seen)
25 (i)	Tangent 90° to diameter /radius/ line from (through) centre	27°	4	B1 for 27° cao B1 for reason
(ii)	180 –(90+ "27") angle in semicircle (is 90°)/Alternate segments /angle at centre twice at circumference	63°		B1 ft for 90–"27" if not 63° B1 for reason
26 (a)(i)		152	2	B1 cao
(ii)		177		B1 cao
(b)			3	B1 for median marked at 167 B1 ft for postion of box with its ends at "152" and "177" B1 for position of whiskers with ends at 132 and 182 NB: For any points plotted between 141 and 149 give a tolerance of an extra ± 1 square
27	$x^2 + xy + xy + y^2$	$x^2 + 2xy + y^2$	2	M1 for at least 3 of the 4 terms correct A1 cao
		25	2	M1 for recognising 3.47 + 1.53 (= 5) A1 cao

Paper 5504

No	Working	Answer	Mark	Notes
1 (a)	$4.1^2 \times 1.07 = 16.81 \times 1.07$	17.9867	2	M1 for (4.1) followed by squaring, or sight of 16.81 A1 cao SC: B1 for 18 or better with no working
(b)		$(1.6 + 3.8 \times 2.4) \times 4.2$	1	B1 cao Allow additional brackets if they give an expression with value 45.024
2 (a)	269.30 – 56.80 = 212.50	6	2	M1 for $\frac{269.30 - 56.80}{42.50}$ or 5 seen A1 cao
(b)	5% of £269.30 £269.30 – "£13.465" OR $\frac{95}{100} \times 269.30$	255.83 or 255.84	3	M1 for (5 ÷ 100) × 269.30 M1 (dep) for 269.30 – "13.465" A1 cao OR M2 for $\frac{95}{100} \times 269.30$ A1 cao **Alternative Method:** M1 for $\frac{5}{100} \times 56.80$ (= 2.84) and $\frac{5}{100} \times 42.50$ (=2.12(5)) (OR 53.96 AND 40.38 (40.375) seen M1 for 56.80 – "2.84" (= 53.96) 42.50 – "2.12(5)" (= 40.375 or 40.38) "5" × "40.375" + "53.96" A1 cao

Paper 5504

No	Working	Answer	Mark	Notes
3 (a)		60	1	B1 cao
(b)	360 – 60 – 90 – 90	120	2	M1 for 360 – "60" – 90 – 90 or 180 – "60" A1 cao
(c)	6 × 2	12	2	M1 for 6 × 2 A1 cao 12
(d)		Correct drawing	2	B2 for triangle and construction lines (see overlay) (B1 for 1 line of length 4cm and correct arcs crossing OR for correct triangle with either no arcs or incorrect arcs) SC: B1 similar triangle drawn with construction lines
4 a(i)	240 × 5 = 1200	1250	3	B1 cao 1250
(ii)	$\frac{50}{1250}$	$\frac{1}{25}$		M1 $\frac{50}{"1250"}$ A1 oe in its simplest form
(b)	$\frac{60}{100} \times 1000 = 600$ 600: 250	12:5	3	M1 for $\frac{60}{100} \times 1000$ oe A1 for 600 A1 cao
5 (a)		$x + 2$	1	B1
(b)	$x + 5 + x + 5 + x + 2 + x + 2$	$4x + 14$	2	M1 adding 4 sides, two of which must be '$x + 2$' (all sides to be linear expressions in x) A1 for correct simplified answer
(c)	"$4x + 14$" = 20	1.5	2	M1 for equation "$4x + 14$" = 20 OR $\frac{20-14}{4}$ oe A1 cao

Paper 5504

No	Working	Answer	Mark	Notes
6 (a)		$2p - q$	2	B1 cao for $2p$ B1 cao for $-q$ accept $-q + 2p$ and $2p - 1q$
(b)	$5x = 3 + 4$	1.4	2	M1 for either (+3 or sight of 7) or (÷ 5 or sight of 0.8 and 0.6) A1 cao accept $\frac{7}{5}$ or $1\frac{2}{5}$
7 (a)		$\frac{4\times5}{2}$	1	B1 cao
(b)	$1 + 2 + 3 + 4 + 5 + 6 + 7 + 8$	$\frac{8\times9}{2}$	1	B1 cao
(c)	$\frac{100\times101}{2}$	5050	1	B1 cao
(d)		$\frac{n(n+1)}{2}$	2	B2 cao (B1 for any quadratic in n)
8	$3.2 \times 2.8 = 8.96$ $2 \times 4.5 \times 2.8 = 25.2$ $2 \times 4.5 \times 3.2 = 28.8$ $\frac{62.96}{2.5} \times 2.99$	75.30	5	M1 for area of any face found correctly M1 for 2 areas seen A1 for 62.96 or 54 M1 for $\frac{'62.96'}{2.5} \times 2.99$ A1 cao Alternate method for candidates who round up $\frac{"62.96"}{2.5}$ M1 for "26" x 2.99 A1 for £77.74 cao SC: for top included B2 for 71.92 m^2 seen or B3 for £86.02 or £86.71 SC: B4 for £64.58 or £65.78 seen

Paper 5504

No	Working	Answer	Mark	Notes
9	2.5×10000	25000	2	M1 for 2.5×10000 or $2.5 \times 100 \times 100$ A1 cao
10 (a)	$\sum f = 90$ (88), 144, 32, 96	Angles drawn, labelled	3	M1 for 1 person = 4° or one angle correct in table or pie chart A1 any 2 angles correctly drawn in pie chart A1 fully correct chart labelled
(b)	$0.38 + 0.27 + 0.15$	0.20	2	M1 1 – sum A1 cao
11	0 | 5 7 8 8 1 | 0 0 0 0 2 5 5 5 6 2 | 0 0 0 4 5 3 | 3 5 Key 1 | 3 = 13 (min)	See working column	3	B1 for stem 0, 1, 2, 3 or 0, 10, 20, 30 B1 for accurate unordered leaves condone 1 error or omission B1 for key and ordered leaves all correct
12 (a)	$V = \pi \times 4^2 \times 10$	502 – 503	2	M1 for $\pi \times 4^2 \times 10$ A1 502 – 503
(b)	$P^2 = 10^2 + 8^2$ $P = \sqrt{164}$	$\sqrt{164} < 13$	3	M1 for sight of a correct right-angled triangle M1 for $10^2 + 8^2$ A1 for conclusion based on a correct calculation or 12.8 seen
13 (a)	2×30	$2 \times 2 \times 3 \times 5$	4	M1 for systematic method, eg division, factor trees (at least one prime) A1 cao
	2×48	$2^5 \times 3$		M1 for systematic method, eg division, factor trees (at least one prime) A1 cao
(b)		12	1	B1 cao
(c)	$2^5 \times 3 \times 5$	480	2	B2 cao B1 for $2^5 \times 3 \times 5$ or any correct common multiple

Paper 5504

No	Working	Answer	Mark	Notes
14 (a)		$150 < C \leq 200$	2	M1 use of cum freq to find the cost of the 20th or 20.5^{th} car OR $\frac{1}{2}\Sigma f$ or $\frac{1}{2}(\Sigma f + 1)$ A1 eg 150 to 200
(b)		No, because the 21st value is in the same interval	1	B1 20.5^{th} or 21st in same interval or an alternative correct explanation
(c)	80% = 5200 $\frac{5200}{80} \times 100$	6500	3	M1 for (100 – 20) % = 5200 M1 for $\frac{5200}{"80"} \times 100$ A1 cao
15 (a)	$x^2(x + 1) = 230$	AG	2	M1 for $x \times x \times (x+1)$ or $x \times x \times x + 1$ oe A1 cao from $x \times x \times (x+1)$
(b)	‘5 – 150 5.1 – 158.7 5.2 – 167.6 5.3 – 177.0 5.4 – 186.6 5.5 – 196.6 5.6 – 207.0 5.7 – 217.7 5.8 – 228.8 5.9 –240.2 5.85 – 234.4	5.8	4	B 2 for trial between 5.8 and 5.9 inclusive (B1 for different trial between 5 and 6 inclusive) B1 for different trial between 5.8 and 5.85 (not including 5.8) B1 (dep on at least one previous B1) cao for 5.8, 5.81, 5.811

Paper 5504

No	Working	Answer	Mark	Notes
16	$\pi \times \left(\frac{15}{2}\right)^2 = 176.715$	88.4cm^2	3	M1 for $\pi \times \left(\frac{15}{2}\right)^2$ A1 88.3 – 88.4 B1(ind) for cm²
17 (a)	$5 = 0.5x + 1$	8	2	M1 for $5 = 0.5x + 1$ A1 cao
(b)		$y = \frac{1}{2}x + c$	1	B1 $y = \frac{1}{2}x + c$, c ≠ 1, oe
(c)		$x = 2y - 2$ OR $x = 2(y - 1)$	2	M1 for correctly multiplying both sides by 2 or correctly isolating $\frac{x}{2}$ A1 for $x = 2(y - 1)$, $x = \frac{y-1}{0.5}$, $x = \frac{y-1}{\frac{1}{2}}$ oe SC: B1 for $x = 2y - 1$
18	$4x - 6y = 22$ $15x + 6y = 54$ $19x = 76$	$x = 4,\ y = -1$	4	M1 for coefficients of x or y the same followed by correct operation , one arithmetical error A1 cao M1(dep on previous M mark) for sub for other variable A1 cao Trial and improvement –0 unless both x and y correct values found

Paper 5504

No	Working	Answer	Mark	Notes
19 (a)	SF $= \frac{10}{6}$ $\frac{10}{6} \times 4.8 = 8$	8	2	M1 for sight of $\frac{10}{6}$ or $\frac{6}{10}$ or 1.67 or better or $\frac{CD}{10} = \frac{4.8}{6}$ A1 cao
(b)	$\frac{10}{6} \times 4.5 - 4.5 = 3$	19.8	2	M1 for use of SF from (a) to find *BC* or *AC* and adding 4 sides A1 cao
20	$\frac{6 \times 10^{15}}{3.2 \times 10^{8}}$ 1.875×10^{7}	4.3×10^{3}	3	B3 for 4.3×10^{3} to 4.34×10^{3} (B2 for 1.875×10^{7} oe or 4300 to 4340 or final answer of 1.9×10^{7}) (B1 for sight of 6×10^{15} oe or 3.2×10^{8} oe)
21	$8.5 \times \tan 38 = 8.5 \times 0.7813$	6.64	3	M1 for correct use of trig, eg $\tan 38 = \frac{opp}{8.5}$ M1 for 8.5 × tan38 A1 6.64 – 6.641

Paper 5504

No	Working	Answer	Mark	Notes
22 (a)		No, as you would expect about 100. Yes, as it is possible to get 200 sixes with a fair dice	1	B1 for a consistent answer
(b)	$\frac{1}{6}, \frac{5}{6}$ + labels		3	B1 for $\frac{5}{6}$ on the red dice, *not six* branch B1 for a fully complete tree diagram with all branches labelled B1 for $\frac{1}{6}$, $\frac{5}{6}$ on all remaining branches as appropriate

Paper 5505

No	Working	Answer	Mark	Notes
1 (i) (ii) (iii)		119.31 119310 1.23	3	B1 cao B1 cao B1 cao
2	$\frac{10}{100}\times 12000 = 1200$ 12 000 – 1200 = 10 800 10 800 ÷ 10 =1080 10 800 – 1080 = £9720	9720	3	M1 for $\frac{10}{100}\times 12\,000$ or sight of 1200 or 2400 or 10 800 or 9600 M1 (dep) for $\frac{10}{100}\times(12\,000 - \frac{10}{100}\times 12\,000)$ or sight of 1080 A1 cao Alternative markscheme M2 for $12000\times\left(1-\frac{10}{100}\right)^2$ (M1 for $12000\times\left(1-\frac{10}{100}\right)$) A1 cao
3	$7r - 5r = -20 - 2$	–11	2	M1 for $7r + 2 = 5r - 20$ or $\frac{7r}{5} + \frac{2}{5} = r - 4$ or $7r - 5r = -20 - 2$ or $\frac{7r}{5} - r = -4 - \frac{2}{5}$ A1 cao

Paper 5505

No	Working	Answer	Mark	Notes
4 (a)		–1, 0, 1	2	B2 for –1, 0, 1 only (B1 for –1, 0 or 0,1 or –1, 1 or –2, -1, 0, 1 only)
(b)		(–1,–1)(0,–1) (1,–1) (0, 0) (1, 0) (1, 1)	3	B3 for 6 points correct (B2 for 3 points correct) (B1 for 1 point correct) *NB: –B1 for each additional point over six*
5		$5n + 1$	2	B2 oe (B1 for $5n$ seen) NB: n = gets B1 max
6 (a)	Triangle with vertices at (0,0) (0,–2) and (3,0)		2	M1 for correct orientation A1 cao
(b)	Rotation, 180°, centre (0,1) Enlargement sf –1 centre (0,1)		2	B2 for 180° 'rotation' centre (0, 1) or for Enlargement sf - 1 centre (0,1) (B1 for any two of the three parts) NB: B0 if additional transformation is included
7	Bisector of $\angle BAC$ Arc around A Region	See overlay	3	B3 cao (B2 for either two correct boundaries, no shading/ wrong shading or one correct boundary, one incorrect boundary with valid shading) (B1 for either two incorrect boundaries but one drawn from A and one intersection, with valid shading or one correct boundary) Ignore shading outside the triangle
8		Length Volume Area	3	B1 for Length B1 for Volume B1 for Area



Paper 5505

No	Working	Answer	Mark	Notes
9 (a)	Unbiased question with choices		2	B1 for unbiased question B1 for at least 2 choices
(b)	Leading question and a restricted sample		2	Classification 1: A biased question Classification 2: A restricted sample of people Classification 3: Not specifying a range of foods Classification 4: Nothing to do with eating habits B2 reasons which satisfy 2 different classifications (B1 a reason which satisfies one classification)
10 (a)	$6 \times 10^2 \times 8 \times 10^4$ $48 \times 10^6 = 4.8 \times 10^7$	4.8×10^7	3	M1 for $6 \times 10^a \times 8 \times 10^b$ oe, a and b integers including 0 A1 for 48×10^6 oe A1 cao
(b)	200 000 + 30 000 = 230 000	230 000	2	B2 cao (B1 for sight of 200 000 or 30 000 or 2.3×10^5 or 23×10^4)
11 (a)	$x^2 + xy + xy + y^2$	$x^2 + 2xy + y^2$	2	M1 for at least 3 of the 4 terms correct A1 cao
(b)		25	2	M1 for recognising 3.47 + 1.53 (= 5) A1 cao
12 (i)	Tangent 90° to diameter /radius/ line from (through) centre	27°	4	B1 for 27° cao B1 for reason
(ii)	180 –(90+ "27") angle in semicircle (is 90°)/Alternate segments /angle at centre twice at circumference	63°		B1 ft for 90–"27" if not 63° B1 for reason
13 (i)		p^9	1	B1 cao
(ii)		$6q^6$	2	B2 for $6q^6$ (B1 for sight of $\frac{6q^9}{q^3}$ or $3q \times 2q^5$ or $3q^4 \times 2q^2$ or $6 \times q \times q \times q \times q \times q \times q$ or final answer of the form kq^6, $k > 0$)

Paper 5505

No	Working	Answer	Mark	Notes
14 (a)(i)		152	2	B1 cao
(ii)		177		B1 cao
(b)			3	B1 for median marked at 167 B1 ft for postion of box with its ends at "152" and "177" B1 for position of whiskers with ends at 132 and 182 NB: For any points plotted between 141 and 149 give a tolerance of an extra ± 1 square
15	(arc =) $\frac{40}{360} \times 2\pi \times 9$ $= 2\pi$	$2\pi + 18$	4	M1 for $\frac{40}{360} \times$ M1 for $2\pi \times 9$ M1 (dep) for $\frac{40}{360} \times 2\pi \times 9$ oe A1 for $\frac{18 \times \pi}{9} + 18$ oe exact form
16 (i)		1	1	B1 cao
(ii)		$\frac{1}{16}$	1	B1 cao accept 0.0625
(iii)		64	1	B1 cao condone ± 64

Paper 5505

No	Working	Answer	Mark	Notes
17 (a)	$F \propto 1/x^2$ $F = k/x^2$ $4 = k / 3^2$ $F = 36/x^2$	$F = \frac{36}{x^2}$	3	M1 for $F = k/x^2$ seen or implied. ($k \neq 1$) M1 (dep) for subst. or sight of $k = 36$ A1 for $F = 36/x^2$
(b)		9	1	B1 ft for 9 (ft on $F = kx^n$, $n \neq 0$)
(c)	$64 = \frac{36}{x^2}$ $x^2 = \frac{36}{64}$ $x = \pm\frac{3}{4}$	$\frac{6}{8}$	2	M1 for $x^2 = \frac{"36"}{64}$ A1 for $\frac{6}{8}$ oe (condone $\pm$) *SC: Use of $F = kx^2$ max M1 M1 A0 B1 ft M0 A0* *SC: Use of* $F = \frac{k}{\sqrt{x}}$ *max M1 M1 A0 B0 M1* ($\sqrt{x} = \frac{4\sqrt{3}}{64}$) *A0*
18	$(5+\sqrt{3})(5-\sqrt{3}) = 5\times5 - 5\sqrt{3} + 5\sqrt{3} - \sqrt{3}\sqrt{3}$ $= 5\times5 - 3$ $\frac{22}{\sqrt{22}} = \frac{22\sqrt{22}}{22}$	$\sqrt{22}$	3	B1 for correct expansion 25 - $5\sqrt{3}$ + $5\sqrt{3}$ - $\sqrt{3}$ $\sqrt{3}$ with 1st three terms reducing to 25 **without** any errors seen B1 (indep) for $\sqrt{3}$ $\sqrt{3}$ = 3 B1 for $\sqrt{22}$ coming from $\frac{22}{\sqrt{22}}$ (S.C $\frac{(5+\sqrt{3})(5-\sqrt{3})\sqrt{22}}{22}$ gets B1)

Paper 5505

No	Working	Answer	Mark	Notes
19 (a)		60 40	2	B1 cao B1 cao
(b)		correct bars	2	B1 for $30 < x \leq 40$ with an area of 2½ squares B1 for $40 < x \leq 70$ with an area of 3 squares *SC:* $\frac{0}{4}$ *give M1 if clearly using area or frequency density*
20 (a)	$6x + 8 - 12x + 15$	$-6x + 23$	2	M1 for 3 of the 4 terms $6x$, $+8$, $-12x$, $+15$ correct A1 cao
(b)		$32x^5y^{15}$	2	B2 cao (B1 for two of 32, x^5, y^{15})
(c)	$\frac{(n+1)(n-1)}{n+1} \times \frac{2}{n-2}$ $\frac{2(n-1)}{n-2}$	$\frac{2(n-1)}{n-2}$	3	M1 for $k(n+1)(n-1)$ M1 dep for $\frac{(n+1)(n-1)}{(n+1)} = n-1$ A1 for $\frac{2(n-1)}{n-2}$
21	Vertices at $(-3, -1\frac{1}{2}), (-4\frac{1}{2}, -1\frac{1}{2}), (-3, -4\frac{1}{2})$		3	B1 for all sides $\times 1\frac{1}{2}$ B1 for correct orientation with 2 vertices almost correct B1 cao

Paper 5505

No	Working	Answer	Mark	Notes
22	Total = 3 + 5 + 2 (=10) $\frac{3}{10}\times\frac{3}{10}\times\frac{5}{10}\left(=\frac{45}{1000}\right), \frac{3}{10}\times\frac{3}{10}\times\frac{2}{10}\left(=\frac{18}{1000}\right)$ $\frac{5}{10}\times\frac{5}{10}\times\frac{3}{10}\left(=\frac{75}{1000}\right), \frac{5}{10}\times\frac{5}{10}\times\frac{2}{10}\left(=\frac{50}{1000}\right)$ $\frac{2}{10}\times\frac{2}{10}\times\frac{3}{10}\left(=\frac{12}{1000}\right), \frac{2}{10}\times\frac{2}{10}\times\frac{5}{10}\left(=\frac{20}{1000}\right)$ $3\times\left(\frac{"45"}{1000}+\frac{"18"}{1000}+\frac{"75"}{1000}+\frac{"50"}{1000}+\frac{"12"}{1000}+\frac{"20"}{1000}\right)$ $\frac{660}{1000}$	$\frac{660}{1000}$ oe	5	M3 for all six expressions seen OR their combined equivalents (M2 for four expressions seen OR their combined equivalents) (M1 for two expressions seen OR their combined equivalents) M1 sum of 18 relevant products condone 1 slip A1 for $\frac{660}{1000}$ oe *SC: without replacement maximum M4 A0* SC: Just 2 beads: Answer either $\frac{38}{100}$ oe OR $\frac{28}{90}$ oe B1

Paper 5505

No	Working	Answer	Mark	Notes
23 (a)(i)		$6\mathbf{b} - 6\mathbf{a}$	2	B1 for $6\mathbf{b} - 6\mathbf{a}$ oe
(ii)		$6\mathbf{a}$		B1 for $6\mathbf{a}$ oe
(b)	$\overrightarrow{EX} = \overrightarrow{EB} + \overrightarrow{BX}$ $= 12\mathbf{b} + \frac{1}{2}\overrightarrow{BC}$	$12\mathbf{b} - 3\mathbf{a}$	2	M1 for $\overrightarrow{EX} = \overrightarrow{EB} + \overrightarrow{BX}$ oe vector journey in a form ready for straightforward substitution A1 for $12\mathbf{b} - 3\mathbf{a}$ oe
(c)	$\vec{AY} = \frac{5}{3}\vec{AB}$ or $\vec{BY} = \frac{2}{3}\vec{AB}$ $\vec{EY} = 16\mathbf{b} - 4\mathbf{a}$ or $\vec{XY} = 4\mathbf{b} - \mathbf{a}$ $\vec{EY} = 4\vec{XY}$ or $\vec{EX} = 3\vec{XY}$ or $\vec{EY} = \frac{4}{3}\vec{EX}$	Printer Answer	3	B1 for either $\vec{AY} = \frac{5}{3}\vec{AB}$ or $\vec{BY} = \frac{2}{3}\vec{AB}$ oe B1 ft for either $\vec{EY} = 16\mathbf{b} - 4\mathbf{a}$ or $\vec{XY} = 4\mathbf{b} - \mathbf{a}$ ft only on parts (*a*) and (*b*) B1 for either $\vec{EY} = 4\vec{XY}$ or $\vec{EX} = 3\vec{XY}$ or $\vec{EY} = \frac{4}{3}\vec{EX}$ oe **plus** conclusion of *E*, *X*, *Y* on the same straight line
24 (a)(i)		$(5, -4)$	4	B1 cao
(ii)		$(2, -9)$		B1 cao
(iii)		$(2, 4)$		B1 cao
(iv)		$(1, -4)$		B1 cao
(b)		$(x-2)^2 - 4$	4	B4 for $(x-2)^2 - 4$ oe eg. $x^2 - 4x$ (B3 for $(x+2)^2 - 4$ or $(x-2)^2 + 4$) (B2 for $x^2 - 4$ or $(x-2)^2$ OR $x^2 + bx$, $b \neq 0$ OR $(x+2)^2 + 4$ OR $f(x-2) - 4$) (B1 for $x^2 + 4$ or $(x+2)^2$ or $ax^2 + bx$ or $x^2 + bx + c$ OR $x - 2 - 4$ or $x^2 - 2 - 4$, $a, b, c \neq 0$)

Paper 5506

No	Working	Answer	Mark	Notes
1 (a)	$V = \pi \times 4^2 \times 10$	502 – 503 cm^3	2	M1 for $\pi \times 4^2 \times 10$ A1 502 – 503
(b)	$P^2 = 10^2 + 8^2$ $P = \sqrt{164}$	$\sqrt{164} < 13$	3	M1 for sight of correct right angled triangle M1 for $10^2 + 8^2$ A1 for conclusion based on a correct calculation Or 12.8 seen
2 (a)(i)	2×30	$2 \times 2 \times 3 \times 5$	4	M1 for systematic method, eg division, factor trees (at least one prime) A1 cao
(ii)	2×48	$2^5 \times 3$		M1 for systematic method, division, factor trees (at least one prime) A1 cao
(b)		12	1	B1 cao
(c)	$2^5 \times 3 \times 5$	480	2	B2 cao B1 for $2^5 \times 3 \times 5$ or any correct common multiple

Paper 5506

No	Working	Answer	Mark	Notes
3 (a)		$150 < C \leq 200$	2	M1 use of cum freq to find the cost of the 20th or 20.5^{th} car OR $\frac{1}{2}\Sigma f$ th or $\frac{1}{2}(\Sigma f + 1)$ th car. A1 eg 150 to 200, 150 – 200
(b)		No, because the 21st value is in the same interval	1	B1 for 20.5^{th} or 21st value in the same internal consistent with 'a' OR Refers to the median value being low in the interval (statement to be mathematically correct) See additional sheet
(c)	80% = 5200 $\frac{5200}{80} \times 100$	6500	3	M1 for (100 – 20)% = 5200 M1 for $\frac{5200}{"80"} \times 100$ A1 cao

Paper 5506

No	Working	Answer	Mark	Notes
4 (a)	$x^2(x + 1) = 230$	AG	2	M1 for $x \times x \times (x+1)$ or $x \times x \times x + 1$ oe, $x^2(x+1)$, $x^2 \times x + 1$ A1 cao from $x \times x \times (x+1)$, no need to see 230
(b)	5 – 150 6 – 252 5.1 – 158.7 5.2 – 167.6 5.3 – 177.0 5.4 – 186.6 5.5 – 196.6 5.6 – 207.0 5.7 – 217.7 5.8 – 228.8 5.9 –240.2 5.85 – 234.4	5.8	4	B2 for trial between 5.8 and 5.9 inclusive evaluated (B1 for trial between 5 and 6 inclusive evaluated) B1 for different trial between 5.8 and 5.85 (not including 5.8) B1 dep on at least are previous B1 5.8, 5.81, 5.811
5	$\pi \times \left(\frac{15}{2}\right)^2 = 176.715$	88.4 cm^2	3	M1 for $\pi \times \left(\frac{15}{2}\right)^2$ seen A1 88.3 – 88.4 B1 for cm² (independent)

Paper 5506

No	Working	Answer	Mark	Notes
6 (a)	$5 = 0.5x + 1$	8	2	M1 for $5 = 0.5x + 1$ A1 cao
(b)		$y = \frac{1}{2}x + c$	1	B1for $y = \frac{1}{2}x + c$, $c \neq 1$, oe
(c)		$x = 2y - 2$ OR $x = 2(y - 1)$	2	M1for correctly multiplying both sides by 2 or correctly isolating $\frac{x}{2}$ A1 for $x = 2(y - 1)$, $x = \frac{y-1}{0.5}$, $\frac{y-1}{\frac{1}{2}}$ oe SC B1 for x= 2y -1
7	$4x - 6y = 22$ $15x + 6y = 54$ $19x = 76$	$x = 4$, $y = -1$	4	M1 for coefficients of x or y the same followed by correct operation, allow one arithmetical error A1 cao M1 (dep) for correct sub for other variable A1 cao Trial and improvement 0 marks unless both correct values of x and y found
8 (a)	SF = $\frac{10}{6}$ $\frac{10}{6} \times 4.8 = 8$	8	2	M1 for sight of $\frac{10}{6}$ or $\frac{6}{10}$ or 1.67 or better or $\frac{CD}{10} = \frac{4.8}{6}$ A1 cao
(b)	$\frac{10}{6} \times 4.5 - 4.5 = 3$	19.8	2	M1 for use of SF from "a" to find AC or BC or $\frac{BC}{4.5} = \frac{4}{6}$ and adding 4 sides A1 cao

Paper 5506

No	Working	Answer	Mark	Notes
9	$\frac{6\times10^{15}}{3.2\times10^{8}}$ 1.875×10^{7}	4.3×10^{3}	3	B3 for 4.3×10^{3} to 4.34×10^{3} (B2 for 1.875×10^{7} oe or 4300 to 4340, final answer of 1.9×10^{7} B1 for sight of 6×10^{15} oe or 3.2×10^{8} oe)
10	$8.5\times\tan38$ $=8.5\times0.7813$ $\frac{8.5}{\sin(90-38)}=\frac{AB}{\sin38}$ $AB=\frac{8.5\times\sin38}{\sin(90-38)}$ $=\frac{5.2331}{0.788}=6.64$	6.64	3	M1 for correct use of trig, eg $\tan 38=\frac{opp}{8.5}$ M1 for $8.5\times\tan38$ A1 6.64 – 6.641 OR M1 for correct substitution into the sine rule M1 (dep) for correct rearrangement for AB= A1 6.64 – 6.641
11 (a)		No, as you would expect about 100. Yes, as it is possible to get 200 sixes with a fair dice	1	B1 for a consistent answer See additional sheet

Paper 5506

No	Working	Answer	Mark	Notes
(b)	$\frac{1}{6}, \frac{5}{6}$ + labels		3	B1 for $\frac{5}{6}$ on the red dice, *not six* branch B1 for a fully complete tree diagram with all branches labelled B1 for $\frac{1}{6}$ and $\frac{5}{6}$ on all remaining branches as appropriate
(c)(i)	$\left(\frac{1}{6}\right)^2$	$\frac{1}{36}$	2	M1 $\left(\frac{1}{6}\right)^2$ or $\frac{1}{6}\times\frac{1}{6}$ only or 0.28 A1 $\frac{1}{36}$ or 0.03 or better
(ii)	$1-\left(\frac{5}{6}\right)^2$ OR $\frac{1}{6}\times\frac{5}{6}+\frac{5}{6}\times\frac{1}{6}+\frac{1}{6}\times\frac{1}{6}$	$\frac{11}{36}$	3	M2 for $1-\left(\frac{5}{6}\right)^2$ or $1-\frac{5}{6}\times\frac{5}{6}$ A1 cao OR M1 for $\frac{1}{6}\times\frac{5}{6}$ oe M1 for 2 or 3 only of $\frac{1}{6}\times\frac{5}{6}, \frac{5}{6}\times\frac{1}{6}$, "a" A1 for $\frac{11}{36}$ or 0.31 or better

Paper 5506

No	Working	Answer	Mark	Notes
12 (a)	$\pi \times 30 \times \frac{7.5^2}{3} - \pi \times 10 \times \frac{2.5^2}{3} = 1767 - 65$	1700	3	M1 for either $\pi \times 30 \times \frac{7.5^2}{3}$ or $\pi \times 10 \times \frac{2.5^2}{3}$ M1 (dep) for difference A1 1700 – 1702 **SC** B1 Using d instead of r, 6800 – 6808
(b)	$\frac{S}{2\pi d} = \sqrt{h^2 + d^2}$ $\left(\frac{S}{2\pi d}\right)^2 = h^2 + d^2$	$h = \sqrt{\frac{S^2 - 4\pi^2 d}{4\pi^2 d^2}}$	3	M1 for correctly isolating $\sqrt{h^2 + d^2}$ or $h^2 + d^2$ or $h + d$ or kh^2 or kh M1(indep) squaring both sides A1 $h = \sqrt{\frac{S^2 - 4\pi^2 d^4}{4\pi^2 d^2}}$, $h = \frac{\sqrt{S^2 - 4\pi^2 d^4}}{2\pi d}$ $h = \sqrt{\left(\frac{S}{2\pi d}\right)^2 - d^2}$
(c)	$\left(\frac{30}{20}\right)^2 \times 450$ or $450 \div \left(\frac{20}{30}\right)^2$	1012.5	2	M1 for sight of correct SF^2 including 4:9 A1 1010 to 1013

Paper 5506

No	Working	Answer	Mark	Notes
13 (a)	$\frac{2x(x+20)}{2} = 400$	As given	2	M1 $\frac{2x(x+20)}{2}$ or $\frac{2x \times x + 20}{2}$ or $2x(x + 20) = 800$ A1 cao following correct working, no need for = 400 *SC B1* $2x \times x + \frac{1}{2} \times 2x(10 - \frac{x}{2}) \times 2$
(b)	$\frac{-20 \pm \sqrt{20^2 - 4 \times 1 \times (-400)}}{2}$ $= \frac{-20 \pm 44.721}{2}$	12.361	3	M1 for correct sub, up to signs, in the quad formula A1 for 44.7 or $\sqrt{2000}$ A1 for 12.3606 – 12.361, ignore negative solution T.I B3 for 12.361 OR Completing the square M1 for $(x+10)^2$ seen A1 for $-10 \pm \sqrt{500}$ A1 for 12.3606 – 12.361 ignore negative solution

Paper 5506

No	Working	Answer	Mark	Notes
14 (a)	$0.5\times 8\times 15\times \sin 70^0$	56.4	2	M1 for correct sub into area formula A1 56.38 – 56.4
(b)	$AB^2 = 8^2 + 15^2 - 2\times 8\times 15\times \cos 70^0 = 206.9$ EITHER $0.5\times AB\times CX = '56.38'$ OR $\frac{\sin B}{8} = \frac{\sin 70}{\sqrt{'206.9'}}$ $B = 31.5$ 15 sin '31.5'	7.84	4	M1 for correct sub into cos rule A1 for 206.9 - 207 or 14.38 – 14.4 EITHER M1 for use of area rule to find CX A1 7.83 – 7.84 OR M1 for correct use of sine rule to find sin B or sin A and then sight of 15sin B *or* 8 sin A A1 7.83 – 7.84

Paper 5506

No	Working	Answer	Mark	Notes
15 (a)	$4a^2 - 4a + 1 - (4b^2 - 4b + 1) =$ $4(a^2 - b^2) - 4(a - b)$ $4(a - b)(a + b - 1)$ OR $((2a - 1) - (2b - 1))((2a - 1) + (2b - 1))$ $(2a - 2b)(2a + 2b - 2)$	AG	3	Expansion Method M1 for a correct expansion of any one of the three terms M1(dep) on an attempt to expand all 3 terms and show LHS = RHS A1 fully correct algebra RHS exp is $4(a^2 + ab - a - ba - b^2 + b)$ OR Factorisation Method M1 for attempt to use difference of 2 squares on LHS M1 for one bracket correctly simplified A1 fully correct
(b)	Any 2 odd square numbers have the above form If a and b are both even or odd then $a - b$ is even, so $4(a - b)$ is a multiple of 8 If one of a,b is odd, then $a + b - 1$ is even, so $4(a + b - 1)$ is a multiple of 8		3	B1 'any 2 square nos have the above form' (may be implied by sight of $(2a - 1)^2 - (2b - 1)^2$ in part (b)) B1 first reason B1 second reason SC B1 for $(2r + 1)^2 - (2r - 1)^2$ B1 for 8r
16 (a)	$g_L = \frac{2 \times 4.495}{1.35^2 \times \sin 30.5}$ $g_u = \frac{2 \times 4.505}{1.25^2 \times \sin 29.5}$	9.719 11.710	4	B2 for any 4 of 4.505, 1.25, 29.5, 4.495, 1.35, 30.5 seen (B1 for any two or three seen) B1 for 11.710 – 11.7103 B1 cao 9.719 – 9.71904
(b)	Round, until lower and upper bounds agree	10	1	B1 for 10 + reason " they agree to this level of accuracy"

Paper 5506

No	Working	Answer	Mark	Notes
17 (a)(i)		xy	3	B1 cao
(ii)		y^2		B1 for y^2 or $y \times y$
(iii)		$\frac{x}{2}$		B1 for $\frac{x}{2}$ or $0.5x$ or $2^{-1}x$
(b)	Divide to get $2y = 1$	$q = -1$	2	M1 for $2y = 1$ or $\frac{x}{2} = 32$ or $p + q = 5$ or $1 + p + 2q = 5$
		$p = 6$		A1 cao
18 (a)	$x^2 - 2mx + m^2 - k$	$k = m^2$	2	M1 for correct exp of $(x - m)^2$ or correct completion of the square eg $\left(x - \frac{2m}{2}\right)^2 - \left(\frac{2m}{2}\right)^2$
				A1 cao
				SC B1 for $k = -m^2$
(b)(i)	Min value is $-m^2$	$-m^2$	3	M1 for recognition that min value occurs when $(x - m)^2 = 0$ (either (b)(i) or (b)(ii) correct implies this M1)
(ii)	$x = m$	m		A1 ft on 'k', " $-k$" gets M1 A0
				A1 cao
19	$0.06 \times 0.05 = 0.003$	No	2	M1 for 0.06×0.05
				A1 correct conclusion based on 0.003 or 0.06 x 0.05 stated as $\neq 0.0011$
				OR M1 for statement that for the two events to be independent P (BL and CL) = P(BL) × P(CL)

Paper 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
20		50 50 4	3	B1 50 or $\frac{100}{2}$ B1 for 50 or "a" B1 4 or $\frac{360}{90}$ oe

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4LN

Telephone 01623 467467
Fax 01623 450481

Order Code UG014129 June 2003

For more information on Edexcel qualifications please contact our
Customer Response Centre on 0870 240 9800
or email: enquiries@edexcel.org.uk
or visit our website: www.edexcel.org.uk

London Qualifications Limited. Registered in England and Wales no.4496750
Registered Office: Stewart House, 32 Russell Square, London WC1B 5DN



GCSE

Edexcel GCSE

Mathematics A 1387

Summer 2005
Mark scheme

Edexcel GCSE
Mathematics A 1387

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information please call our Customer Services on 0870 240 9800, or visit our website at www.edexcel.org.uk.

Summer 2005

Publications Code UG016720

All the material in this publication is copyright
© Edexcel Ltd 2005

CONTENTS

Notes on Marking Principles 1

Paper 5521/01 3

Paper 5521/02 12

Paper 5523/03 18

Paper 5523/04 27

Paper 5525/05 37

Paper 5525/06 45

NOTES ON MARKING PRINCIPLES

1 Types of mark

- M marks: method marks
- A marks: accuracy marks
- B marks: unconditional accuracy marks (independent of M marks)

2 Abbreviations

cao – correct answer only
ft – follow through
isw – ignore subsequent working
SC: special case
oe – or equivalent (and appropriate)
dep – dependent
indep - independent

3 No working

If no working is shown then correct answers normally score full marks
If no working is shown then incorrect (even though nearly correct) answers score no marks.

4 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.
If there is no answer on the answer line then check the working for an obvious answer.

5 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

6 **Ignoring subsequent work**
It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

7 **Probability**
Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

8 **Linear equations**
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

9 **Parts of questions**
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

Paper 5521/01					
No		Working	Answer	Mark	Notes
1	(a)		17252	1	B1 cao
	(b)		5400	1	B1 cao
	(c)		thousands, 1000, 4000	1	B1
2	(a)		grams, g	3	B1 oe spelling
			centimetres, cm		B1 oe spelling
			millilitres, m*l*, cm^3		B1 oe spelling
	(b)		5	1	B1 cao
3	(a)		106, 102	1	B1 cao ignore extras
	(b)		eg take away 4	1	B1 could be indicated on the diagram
	(c)		46	1	B1 cao

Paper 5521/01

No	Working	Answer	Mark	Notes
4	286 × 43 = 11440 + 12298 = 12298 43 × 286 = 258 + 3440 + 8600 = 12298 286 × 40 = 1140 286 × 3 = 858 1140+858=12298 x | 40 | 3 200 | 8000 | 600 | 8600 80 | 3200 | 240 | 3440 6 | 240 | 18 | 258 12298 2 8 6 1 | 0 8 | 3 2 | 2 4 | 4 2 | 0 6 | 2 4 | 1 8 | 3 2 9 8	12298	3	M2 for complete correct method (condone one computational error) (M1 for complete correct method with two computational errors) A1 cao OR B2 inside of grid completed (condone missing zeros and one error) (B1 2 or 3 errors) B1 cao
5 (a) (b)(i) (ii) (c) (d)(i) (ii) (iii)		18, 69 18 or 36 16 or 36 factor 18 11 or 88 69	1 1 1 1 3	B1 B1 B1 B1 B1 cao B1 B1 cao

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
6	(a)		April & May	1	B1 for both
	(b)		Daffodil	1	B1
	(c)		Feb	1	B1
	(d)		Crocus	1	B1
	(e)(i)		$\frac{1}{5}$	2	B1 for $\frac{1}{5}$ oe
	(ii)		× from 56 mm to 64 mm from 0		B1 A single mark on the line, between 56 mm and 64 mm measured from end 0
7	(a)	$\frac{40}{100}$	$\frac{2}{5}$	2	B2 for $\frac{2}{5}$ B1 for $\frac{40}{100}$ or $\frac{4}{10}$ or $\frac{20}{50}$ or $\frac{8}{20}$
	(b)		0.98	1	B1 cao
	(c)		7 500 000	1	B1 cao
	(d)		25	1	B1 cao
	(e)		60	1	B1 cao

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
8	(a)(i)		(0, 2)	2	B1 cao
	(ii)		(4, 1)		B1 cao
	(b)		$(2, 1\frac{1}{2})$ marked	1	B1 Allow 2mm tolerance from $(2, 1\frac{1}{2})$
9	(a)		2.40	1	B1 cao could be indicated on the diagram
	(b)	1.60 + 0.40	2.00	2	M1 for appropriate sum or product in £ or p or 200 seen eg 1.60 + 0.40, 160 + 40, 0.80 + 0.80 + 0.40, 80 + 80 + 40, 0.08×25, 0.80×2.5, 200 A1 cao
	(c)	$1\div0.8$ or $2.50\div2$	1.25	2	M1 for $1.00\div0.8$ or $2.50\div2$ or 125 or appropriate combination eg $1+\frac{1}{2}\times0.50$ A1 cao
10	(a)		hexagon	1	B1 Condone spelling error
	(b)		Sum of angles at a point is 360°	2	B1 for 360 seen B1 for "point", "complete turn" or "a circle" or similar unless accompanied by an incorrect angle SC If neither B1 scored, award B1 for a clear indication that the size of an angle, other than *x*, is 90° or a right angle (may be on diagram)
	(c)	$30\times4+8\times2$	136	2	M1 $30\times4+8\times2$ or attempt to sum 5 or 6 lengths A1 cao

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
11	(a)		13, 67, 76, 103, 130	5	B1 cao
	(b)		$-7, -3, -1, 0, 5$		B1 cao
	(c)		0.07, 0.072,0.7, 0.702, 0.72		B1 cao
	(d)		0.6, $\frac{2}{3}$, 70%, $\frac{3}{4}$		B2 (B1 for any 3 in correct order)
12	(a)		16 30	1	B1 Accept 4 30 pm Do not accept 4 30
	(b)(i)	$33.56 \div 4$ oe	8.39	3	M1 for $33.56 \div 4$ oe eg $3356 \div 4$, division by 2 twice A1 cao
	(ii)		9		B1 ft from "8.39" unless whole number of pounds
13	(a)		6 cm^2	3	B2 for 6 cao for numerical answer (B1 for $5.5 < \text{Area} \leq 7$) then B1 (indep) for cm^2 with or without numerical answer
	(b)	See diagram	Correct shape	2	B2 (B1 for any 2 sides correct or a correct enlargement scale factor $\neq 1$ or 2)

Paper 5521/01				
No	**Working**	**Answer**	**Mark**	**Notes**
14 (a)	$(4+3)\times10$	70	2	M1 for (4 + 3) × 10 A1 cao
(b)	$120\div10-3$	9	2	M1 for $\frac{120}{10}$ or 12 seen eg $12\times10=120$ A1 cao
(c)		$C=10(n+3)$	3	B3 for $C=10(n+3)$ oe such as $C=(n+3)\times10$ (B2 for correct RHS or $C=n+3\times10$, $C=10n+3$ oe B1 for C = some other linear expression in n or for $n+3\times10$, $10n+3$ etc) Note: $C = n$ scores no marks
15		11 13 16 8 21	2	B2 all correct (B1 for 2 correct)

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
16	(a)		$2p + 4q$	2	B2 for $2p + 4q$ (accept $2 \times p$ etc) (B1 for $2p$ or $4q$)
	(b)		$2y^2$	1	B1 accept $2 \times y^2$ oe inc $2 \times y \times y$
	(c)		$3c + 4d$	2	B2 for $3c + 4d$ (accept $3 \times c$ etc) (B1 for $3c$ or $4d$)
	(d)		$8pq$	1	B1 accept in any order but must not include $\times$ sign
17	(a)(i)		60	2	B1 cao
	(ii)		eg top triangle is equilateral		B1 for reason
	(b)		150	2	M1 for $\frac{180 - "60"}{2} + 90$ A1 ft from (a)(i) if $x < 90$ SC B1 for "60" + 90 if $x < 90$
18			40	2	M1 for 60×2 or 120 or $60 \div 3$ or 20 or $\frac{120}{180}$ A1 cao
19			correct drawing	2	B2 Condone hidden detail shown with solid lines and missing lines on front face (B1 for a correct sketch with other incorrect sketch(es) or for prism with correct cross section >1 cube wide or for attempt to draw prism with correct cross section or prism with correct plan and side elevation)

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
20		$\frac{600}{3\times10}$ or $\frac{640}{3.2\times10}$	$20\text{-}21\frac{1}{3}$	2	M1 for rounding at least two of the numbers to 1 sf or for sight of 640, 3.2 or 640, 32 or 600, 32 or 30 seen A1 for $20\text{-}21\frac{1}{3}$ Note: 20.3125 scores M0 A0
21	(a)		Points plotted	1	B1 $\pm$ 1 full (2 mm) square
	(b)		positive	1	B1 cao
	(c)		Line of best fit	1	B1 Must pass through (42.5, 1.45), (42.5, 1.55) AND (67.5, 1.75), (67.5, 1.85)
	(d)		~ 1.65	1	B1 ft from single line segment with positive gradient $\pm$ 1 full (2 mm) square
22	(a)	eg $50\times\frac{2000}{500}$	200	2	M1 for $\frac{2000}{500}$ or 4 seen A1 cao
	(b)	eg $400\times\frac{750}{500}$	600	2	M1 for $\frac{750}{500}$ or 1.5 seen or 400 + 200 A1 cao

Paper 5521/01					
No		**Working**	**Answer**	**Mark**	**Notes**
23	(a)	$4\times3-2\times1$ $12-2$	10	3	M1 for 3×4 (=12) or 1×2 or attempt to divide diagram up into rectangles M1 "12"– "2" or sum of areas of rectangles A1 cao
	(b)(i)	$\frac{10}{100}\times680$ or $680\div10$ 680 + 68	748	5	M1 $\frac{10}{100}\times680$ or 680÷10 or 68 seen M1 (dep) 680 + "68" or M2 for 680×1.10 A1 cao
	(ii)	"748"÷50 or 14.96	15		M1 For "748"÷50 or 14.96 Accept "748" rounded up or down to next 50 followed by $\div50$ A1 ft from (b)(i) rounded up SC B1 for 680 (seen) leading to 14

Paper 5521/02					
No		**Working**	**Answer**	**Mark**	**Notes**
1	(a)	Plain llll lll 8 Chicken lll 3 Bovril llll 5 S & Vin llll 4		3	M1 for attempt to tally A1 for 1 frequency correct or all tallies correct A1 for all frequencies correct (accept for /20)
	(b)		4	1	B1 ft
	(c)		Plain or 8	1	B1 ft
2	(a)(i)		11	2	B1 cao
	(ii)		16		B1 cao
	(b)	See diagram	Correct lines	2	B2 cao for both lines correct (B1 for one line correct)
	(c)		12	2	B2 cao (B1 for 11 or 13)
3	(a)		580	1	B1 for 580 (± 2) could be written on line
	(b)		7.2	1	B1 for 7.2 ± 0.02 could be written on line
	(c)		Arrow at 48	1	B1 allow ± half graduation
	(d)		Arrow at 6.7	1	B1 allow ± half graduation
4	(i)		Cylinder	2	B1 ignore spelling
	(ii)		Cuboid		B1 ignore spelling
5	(a)	£10 – (£2.15 + £2.30)	5.55	4	M1 £2.15 + £2.30 A1 for 4.45 M1 £10 – "4.45" A1 cao
	(b)	£60 ÷ £2.80 = 21.42857	21	2	M1 for £60 ÷ 2.80 or sight of digits 214… A1 for 21
	(c)	$120 \times 25 \div 100$	30	2	M1 ¼ of £120 (oe) A1 cao SC B2 for £90

Paper 5521/02					
No		**Working**	**Answer**	**Mark**	**Notes**
6	(a)(i)		143^0	2	B1 for 143 ($\pm 2^0$)
	(ii)		Obtuse		B1 for obtuse (ignore spelling)
	(b)	See diagram	Accurate drawing	1	B1 for accurate drawing ± 2mm
7	(a)(i)		5	2	B1 cao
	(ii)		23		B1 cao
	(b)	× 2 – 1		1	B1 for explaining a suitable method
	(c)	See their diagram		1	B1 for a correct diagram
	(d)		14, 17	2	B2 cao for both (B1 for one only ft from their 14)
8	(a)		90	1	B1 accept –90
	(b)		540	1	B1 accept –540
	(c)		Jupiter	1	B1 accept –150
	(d)		- 230	1	B1 cao

Paper 5521/02					
No		Working	Answer	Mark	Notes
9	(a)	2658 – 2430 = 228 "228" × 32	72.96	4	M1 2658 – 2430 A1 228 M1 "228" × 32 or "228" × 0.32 or digits 7296 seen A1 cao **Or** M1 for 2430 × 32 (or digits 77760 seen) or 2658 × 32 (or digits 85056 seen) A1 if 1 correct M1 for "85056" – "77760" or 7296 seen A1 cao
	(b)	$\frac{2}{5} \times 145 = 58$ 145 – "58"	87	3	M1 $\frac{2}{5} \times 145$ (or M1 $\frac{3}{5}$ seen) A1 58 (or M1 $\frac{3}{5} \times 145$) A1 for 87 ft
	(c)(i) (ii)		80 125	2	B1 for 80 (± 1) B1 125 (± 3)

Paper 5521/02					
No		**Working**	**Answer**	**Mark**	**Notes**
10	(a)		1.5 – 2.0	1	B1 for height between 1.5m – 2.0m inclusive
	(b)	Height of man × "2.5"	3 – 6	3	B3 for height between 3m – 6m inclusive (B2 for multiplying (a) by a number between 2 and 3 inclusive) (B1 for multiplying (a) by a number)
11		61 – 19 = 42 42 ÷ 3 = 14	14	2	M1 for – 19 or 42 seen A1 cao
12	(a)		5	1	B1
	(b)	4+5+5+5+4+3+2+1+4+5 = 38 mean = 38 ÷ 10 = 3.8	3.8	2	M1 for attempt to add and ÷ 10 or 3.7 or 3.9 seen A1 for 3.8 SC B1 for 33.5 seen
13	(a)		$3x$	1	B1 cao Accept $3 \times x$, $x3$, $x \times 3$, $x + x + x$
	(b)		x - 9	1	B1 for x – 9 cao
14	(a)	14.44 – 8.660254038	5.77974(…)	2	M1 for 14.44 seen or 8.66(…) or 5.7 or 5.8 or better rounded or truncated
	(b)		6	1	A1 cao B1 ft
15		15 ÷ 24	62.5	2	M1 for 15 ÷ 24 or 1500 ÷ 24 or sight of digits 625 A1 cao
16		2.10 × 450	945	2	M1 for digits 210 × 450 or sight of digits 945 A1 cao
17		See diagram	$2(y + y)$ $2y + 2y$	2	B1 for $2(y + y)$ B1 for $2y + 2y$ (Deduct B1 for each additional tick (>2) to min 0)
18		360° ÷ 18 (=20) Sector angles: G= 60; S= 80; B=220; Correct sectors labelled correctly Use angle measurer	Angles drawn, labelled	4	B4 for fully correct and labelled pie chart (B3 for all angles correct or for a labelled pie chart with 2 correct angles) (B2 for labelled pie chart with 1 correct angle drawn) (B1 for 360° ÷ 18 or 20 seen or implied)

Paper 5521/02					
No		Working	Answer	Mark	Notes
19	(a)		Correct plane	2	B2 for a correct plane defined by showing at least 2 lines. (B1 for a line of symmetry on one face)
	(b)		Correct net	2	B2 cao (B1 for 2 equilateral triangles joined appropriately to at least one rectangle or for 1 equilateral triangle joined appropriately to one of 3 rectangles)
	(c)		Correct drawing	2	B1 for two extra sides of length 6cm (± 2mm) B1 for construction arcs 6cm from each of the ends of the given line.
20	(a)		15	1	B1 for 15 (± 1)
	(b)		15	1	B1 for 15 (± 0.4)
	(c)			2	B1 horiz. line from (2, 20) to (3, 20) B1 line from (3, 20) to (5,0) or horizontal translation of it SC B1 for any journey ending at (5,0)
21	(a)	$x+4+x+x+4+x$	$4x+8$	2	M1 for attempting to add x, $x + 4$, x, $x + 4$ may be implied by $4x + a$ ($a > 0$) A1 for $4x + 8$ or $4(x + 2)$
	(b)	$4x + 8 = 54$ $4x = 46$ $x = 11.5$ Length = "11.5" + 4	15.5	3	M1 for "$4x + 8$" = 54 A1 cao for 11.5 seen B1 ft for "11.5" + 4
22		0.4 + 0.15 1 – "0.55"	0.45	2	M1 for 1 – sum A1 for 0.45 o.e. SC B1 for 0.81
23	(a)		3:1	1	B1 cao
	(b)	$\pi \times 2.45$	7.7	2	M1 for $\pi \times 2.45$ (accept π as 3.1 or better) A1 for 7.59 to 7.70

Paper 5521/02					
No		**Working**	**Answer**	**Mark**	**Notes**
24		7×10000	70 000	2	M1 for 7 X 10 000 or 7 X 100 X 100 A1 cao
25		5.40 ÷ 3 × 7	12.60	3	M1 for 5.40 ÷ 3 or sight of 1.8 M1 dep for "1.80" × 7 A1 for 12.6 or equivalent
26		7.60 x $\frac{17.5}{100}$ = 1.33 7.60 + 1.33 = 8.93 1650 x "8.93"	£14 734.50	4	M1 for 7.60 x $\frac{17.5}{100}$ or 1.33 seen or 7.60 x 1.175 (oe) (Award M1 for 10%, 5% and 2½% correctly calculated) A1 for 8.93 or 893 M1 for 1650 x "8.93" or digits 147345 seen A1 cao Accept 14734.5 **Alternative** M1 for 1650 × 7.6(0) or 12540 seen M1 for "12540" × $\frac{17.5}{100}$ or 2194.5 seen or "12540"× 1.175 (oe) (Award M1 for 10%, 5% and 2½% correctly calculated) M1 for "12540" + "2194.5" (dep on both previous method marks) or digits 147345 seen A1 cao accept 14 734.5

Paper 5523/03				
No	**Working**	**Answer**	**Mark**	**Notes**
1		Correct shape	2	B2 B1 for any 2 sides correct, or a correct enlargement scale factor ≠1 or 2.
2 (a)		11 13 16 8 21	2	B2 all correct (B1 for 2 correct) sign
(b)		$\frac{31}{80}$	1	B1 oe
3 (a)		$2p+3q$	2	B2 for $2p+3q$ (accept $2\times p$ etc) (B1 for $2p$ or $3q$ or $2p3q$)
(b)		$2y^2$	1	B1 accept $2\times y^2$ or $2\times y\times y$
(c)		$3c+4d$	2	B2 for $3c+4d$ (accept $3\times c$ etc) (B1 for $3c$ or $4d$ or $3c4d$)
(d)		$8pq$	1	B1 accept in any order but must not include ×
4 (a)(i)		60	2	B1 cao
(ii)		eg top triangle is equilateral		B1 for reason
(b)		150	2	M1 $\frac{(180-"60")}{2}+90$ A1 ft from (a)(i) if $x < 90°$ SC:B1 for answer from "60" + 90 if $x < 90°$

Paper 5523/03					
No		Working	Answer	Mark	Notes
5	(a)	$4\times3-2\times1$ $12-2$	10	3	M1 for 3×4 (=12) or 1×2 or attempt to divide diagram up into rectangles M1 "12" – "2" A1 cao
	(b)(i)	$\frac{10}{100}\times680$ or $680\div10$ 680 + 68	748	5	M1 $\frac{10}{100}\times680$ or $680\div10$ or 68 seen M1 (dep) 680 +"68" (or M2 for 680×1.10) A1 cao
	(ii)	"748" ÷ 50 or 14.96	15		M1 for "748" $\div50$ or 14.96; accept "748" rounded up or down to next 50 followed by $\div50$ A1 ft from (b)(i) rounded up SC B1 for 680 (seen) leading to 14
6	(a)	$2\times5+5\times-3=10-15=$	-5	2	M1 substitute e.g 2×5 and 5×-3 or 10 and -15 A1 cao
	(b)	$5\times4^2-7$ $5\times16-7$	73	3	M1 substitution e.g $5\times4^2-7$; do not accept 54^2-7 M1 $5\times16-7$ or $5\times4\times4-7$ or $80-7$ (NB 4^2 as 4×4) A1 cao

Paper 5523/03					
No		**Working**	**Answer**	**Mark**	**Notes**
7		679 × 28 = 5432 + 13580 = 19012 or 28 × 679 = 252 + 1960 + 16800 = 19012 or  	190.12	3	M1 for an attempt to multiply the units and tens, or correct partitioning M1 complete correct method (condone one arithmetic error) A1 for 190.12 cao OR M1 for putting the numbers in a grid M1 for multiplying out and addition (condone one error) A1 answer shown with point OR M1 for correct partitioning M1 679×20 and 679×8 calculated oe (condone one error) A1 cao
8	(a)		Points plotted	1	B1 ±1 full (2mm) square
	(b)		positive	1	B1 cao
	(c)		Line of best fit	1	B1 must pass through (42.5, 1.45), (42.5, 1.55) AND (67.5, 1.75), (67.5, 1.85)
	(d)		approx 1.65	1	B1 ft from single line segment with positive gradient ± 1 full (2mm) square

Paper 5523/03					
No		**Working**	**Answer**	**Mark**	**Notes**
9	(a)	eg $50 \times \frac{2000}{500}$	200	2	M1 $2000 \div 500$ or 4 seen A1 cao
	(b)	eg $400 \times \frac{750}{500}$	600	2	M1 $750 \div 500$ or 1.5 seen or 400 + 200 A1 cao
10			$C = 10(n+3)$	3	B3 for $C = 10(n+3)$ oe such as $C = (n+3) \times 10$ (B2 for correct RHS or $C = n + 3 \times 10$, $C = 10n + 3$ etc B1 for C = some other linear expression in n or $n + 3 \times 10$, $10n + 3$ etc) NB: C=n scores no marks
11	(a)		$p(p+6)$	2	B2 for $p(p+6)$ or $p \times (p+6)$ (B1 for $p(ap+b)$ where a,b are numbers or $p+6$ seen on it's own, or part of an expression)
	(b)	$x^2 - 4x + 7x - 28$	$x^2 + 3x - 28$	2	M1 for 4 terms correct ignoring signs (e.g $x^2, 4x, 7x, 28$) or 3 terms with correct signs (e.g $x^2, -4x, 7x, -28$) A1 cao
12			correct drawing	2	B2. Condone hidden detail shown with solid lines, or missing lines on front face. (B1 for : one sketch correct with other sketches incorrect cross-section correct with depth > 1 cube, correct plan and side elevation)

Paper 5523/03					
No		**Working**	**Answer**	**Mark**	**Notes**
13		$\frac{600}{3\times10}$ or $\frac{640}{3.2\times10}$	20 to 21 $\frac{1}{3}$	2	M1 For rounding at least two of the numbers to 1 sf, or for sight of 640, 3.2 or 640, 32 or 600, 32 or 30 seen A1 for 20 to 21 $\frac{1}{3}$ NB: 20.3125 scores M0 A0
14	(a)		correct reflection	2	B2 (B1 reflection in line other than x=3)
	(b)		reflection in $y = x$	2	B2 cao Accept the word "reflected" (B1 any statement including the word "reflection")
15	(a)(i)		5^6	1	B1 accept 15125, 5^{4+2}
	(ii)		5^3	1	B1 accept 125, 5^{9-6}
	(b)	$x + y = 10$ and $x - y = 4$	$x = 7$ $y = 3$	3	M1 for either $x + y = 10$ or $x - y = 4$ A2 for both values correct [(A1 for one value correct) If M0, award B3 for both values correct or B2 for one value correct, otherwise B0] SC B2 for $x = 3$ and $y = 7$

Paper 5523/03					
No		**Working**	**Answer**	**Mark**	**Notes**
16	(a)	$5-3x=2x+2$ $5-2=2x+3x$	$\frac{3}{5}$	3	B1 for $2x+2$ seen OR $2.5-1.5x=x+1$ M1 for correct rearrangement of 4 terms A1 for $\frac{3}{5}$ oe
	(b)		$-3,-2,-1,0,1,2$	2	B2 (B1 for 5 correct and not more than one incorrect integers)
17			question + response boxes oe	2	1st aspect: One question with time period (eg each night); ignore other questions. 2nd aspect: Response list (at least two), not overlapping.* 3rd aspect: Some mention of units (eg hours) in either question or responses Award B2 for all three aspects, or B1 for just two aspects. * 0-1, 2-3, 4-5 is OK, but 0-1, 1-2, 2-3 is not OK.

Paper 5523/03					
No		**Working**	**Answer**	**Mark**	**Notes**
18	(a)	$\frac{2}{3}\times\frac{3}{4}=\frac{6}{12}$	$\frac{1}{2}$	2	M1 for $\frac{6}{12}$ or $\frac{3}{6}$ or $\frac{2\times3}{3\times4}$ A1 accept 0.5
	(b)	$1+2+\frac{8}{12}+\frac{9}{12}$	$4\frac{5}{12}$	3	M1 for attempt to convert to fractions with common denominator e.g two fractions, denominator of 12 A1 correct conversion : $\frac{8}{12}$ and $\frac{9}{12}$, or $\frac{20}{12}$ and $\frac{33}{12}$ seen (oe) A1 cao for $4\frac{5}{12}$ OR attempts to convert to decimals: must use at least 2dp M1 0.66+0.75 (or 1.66+2.75) or 0.67+0.75 etc A1 4.41, 4.417, 4.416 A1 4.416 (*recurring*)
19		$2\times\frac{1}{2}\times6\times8$ or 48 $8\times9+6\times9+10\times9$ or 72 +54 + 90	264 cm^2	4	M1 attempt to find the area of one face; $\frac{1}{2}\times6\times8$ or (8×9) or (6×9) or (10×9) or 72 or 54 or 90 or 24 M1 all five faces with an intention to add A1 cao numerical answer of 264 B1 (indep) cm^2 with or without numerical answer

Paper 5523/03					
No		**Working**	**Answer**	**Mark**	**Notes**
20			$\frac{\pi ab^3}{3d}$ $3(c+d)^3$ $3\pi bc^2$	3	B3 (B1 for each one correct) Nb –B1 for each of the 4th,5th,6th tick
21	(a)	$x+0.3+0.2+x=1$	0.25	2	M1 for $x+0.3+0.2+x=1$ oe, or $0.5\div 2$ A1 oe
	(b)	0.3×200	60	2	M1 0.3×200 A1 cao Accept 60 out of 200 (in words) SC B1 for $\frac{60}{200}$
22	(a)		$(-12)-4\quad -2\quad (0)8$	3	B3 for all correct [(B1 for each one correct)
	(b)		points plotted accurately points joined with smooth curve	2	B1 ± 1 full (2mm) square ft table if at least B1 awarded (all 5 points plotted) B1 ft for any smooth curve if previous B1 gained NB: curve must pass within 1 full square of the points
23			$\frac{1}{4}$ on LH branch $\frac{2}{3}$ & $\frac{1}{3}$ & $\frac{2}{3}$ on RH branches	2	B1 B1
24			m=3 n=5	2	B1 for 3 B1 for 5 (B2 for $2^3\times 5$ or $2\times 2\times 2\times 5$) SC: award B1 only if m=5, n=3 or for 8×5 seen

Paper 5523/03					
No		Working	Answer	Mark	Notes
25		$\frac{5--1}{-1-2} = -2$	$y = -2x + 5$	4	M1 for clear attempt to find gradient eg fraction with -1,5 in numerator, 2,-1 in denominator A1 for -2 B2 ft for $y =$"-2"$x + 5$ oe (eg $y = \frac{-6}{3}x + 5$) (B1 for $y = mx + 5$ or , $-2x+5$ or $y =$"-2"$x + c$)
26	(a)(i)		150	2	B1 accept 150 or 210
	(ii)				B1 for angle at the centre is twice the angle at the circumference
	(b)	$360 - 90 - 90 -$"150"	30	3	B1 identifies angle between radius and tangent as $90°$ (may be in working or on diagram) M1 $360° - 90 - 90 -$"150" A1 ft from (a)(i) excluding a negative answer OR B1 for 90 M1 for $2 \times (180 - 90 -$"$\frac{150}{2}$"$)$ A1 ft from (a)(i) excluding a negative answer OR B3 for $180 -$(a)

Paper 5523/04					
No		**Working**	**Answer**	**Mark**	**Notes**
1	(a)	14.44 – 8.660254038	5.77974(….)	2	M1 for 14.44 seen or 8.66(……) seen or 5.7 or 5.8 or better, rounded or truncated A1 cao
	(b)		6	1	B1 ft
2		15 ÷ 24	62.5	2	M1 for 15 ÷ 24 or 1500 ÷ 24 or sight of digits 625 A1 cao
3	(a)	2.10 × 450	945	2	M1 for digits 210 × 450 or sight of digits 945 A1 cao
	(b)	63÷2.10	30	2	M1 for 63 ÷ digits 210 A1 cao
4		See diagram	2(y + y) 2y + 2y	2	B1 for 2(y + y) B1 for 2y + 2y (Deduct B1 for each additional tick (>2) to min 0)
5		360° ÷ 18 (=20) Sector angles: G= 60; S= 80; B=220; Correct sectors labelled correctly Use angle measurer	Angles drawn, labelled	4	B4 for fully correct and labelled pie chart (B3 for all angles correct or for a labelled pie chart with two angles correct) (B2 for labelled pie chart with one correct angle drawn) (B1 for 360÷18 or 20 seen or implied)
6	(a)		Correct plane	2	B2 for a correct plane defined by showing at least 2 lines. (B1 for a line of symmetry on one face)
	(b)		Correct net	2	B2 cao (B1 for 2 equilateral triangles joined appropriately to at least one rectangle or for 1 equilateral triangle joined appropriately to one of the three rectangles)
	(c)		Correct drawing	2	B1 for two extra sides of length 6 cm (± 2mm) B1 for construction arcs 6cm from each of the ends of the given line
7		61 – 19 = 42 42 ÷ 3 = 14	14	2	M1 for –19 or 42 seen or 3x + 19 A1 cao

Paper 5523/04					
No		Working	Answer	Mark	Notes
8	(a) (b) (c)		15 15	1 1 2	B1 cao for 15(±1) B1 cao for 15(±0.4) B1 horiz. line from (2,20) to (3,20) B1 line from (3,20) to (5,0) or horiz. translation of it SC: B1 for any journey ending at (5,0)
9	(a)	x+4+x+x+4+x	4x+8	2	M1 for attempting to add $x+4, x, x+4, x$ may be implied by 4x+a, a>0 A1 for 4x+8 or 4(x + 2)
	(b)	4 x + 8 = 54 4 x = 46 x = 11.5 Length = "11.5" + 4	15.5	3	M1 for "4x + 8" = 54 A1 cao for 11.5 seen B1 ft for "11.5"+ 4
10		0.4 + 0.15 1 – 0.55	0.45	2	M1 for 1 – sum A1 for 0.45 oe SC: B1 for 0.81
11	(a) (b)	 π × 2.45	3:1 7.7	1 2	B1 cao M1 for π × 2.45 (accept π as 3.1 or better) A1 for 7.59 to 7.70
12		7 × 10000	70000	2	M1 for 7 × 10000 or 7 × 100 × 100 A1 cao
13		5.40 ÷ 3 × 7	12.60	3	M1 for 5.40 ÷ 3 or sight of 1.8 M1 (dep) for "1.80" × 7 A1 for 12.6 or equivalent

Paper 5523/04				
No	**Working**	**Answer**	**Mark**	**Notes**
14	$7.60 \times \frac{17.5}{100} = 1.33$ 7.60 + 1.33 = 8.93 1650 × "8.93"	£14734.50	4	M1 for $7.60 \times \frac{17.5}{100}$ or 1.33 seen or 7.60 ×1.175 (oe) (Award M1 for 10%, 5% and 2½% correctly calculated) A1 for 8.93 or 893 M1 for 1650 × "8.93" or digits 147345 seen A1 cao Accept 14734.5 OR M1 for 1650×7.6 or 12540 seen M1 for "12540" $\times \frac{17.5}{100}$ or 2194.5 seen or "12540"×1.175 (oe) (Award M1 for 10%, 5%, and 2½% correctly calculated) M1 for "12540" + "2194.5" (dep on both previous M marks) or digits 147345 seen A1 cao accept 14734.5

Paper 5523/04					
No		**Working**	**Answer**	**Mark**	**Notes**
15		$285 \times 1000/(60 \times 60) = 79.1\dot{6}$		3	M2 for $285 \times 1000 \div 60 \div 60$ or $80 \times 60 \times 60 \div 1000$ or for a correct method to obtain two comparable values e.g $80 \times 60 \times 60$ and 285×1000 (M1 for $285 \div 60 \div 60$ or 0.079(….) seen or $80 \times 60 \times 60$ or 288000 seen or for 285×1000 or 285000 seen or $80 \div 1000$ or 0.08 seen) A1 for 288 or 79.(….) or for two correctly calculated comparable values e.g 288000 and 285000
16	(a)	$4x + 12 = 6$ $4x = -6$	-1.5	3	B1 for $4x + 12$ or $x + 3 = \frac{6}{4}$ M1 for a correct re-arrangement of their 3 terms to isolate $4x$ or x A1 for -1.5 oe
	(b)	$v - u = 5t$	$\frac{v-u}{5}$	2	M1 for isolating $\pm 5t$ or $\pm t$ or for dividing through by 5 A1 oe

Paper 5523/04					
No		**Working**	**Answer**	**Mark**	**Notes**
17		3→15 4→48 3.1→17.3(91) 3.2→19.9(68) 3.3→22.7(37) 3.4→25.7(04) 3.5→28.8(75) 3.4→25.7(04) 3.3→22.7(37) 3.35→24.1(95375)	3.3	4	B2 for trial between 3.3 and 3.4 inclusive (B1 for trial between 3 and 4 inclusive) B1 for different trial between 3.3 and 3.4 exclusive B1 (dep on at least one previous B1) for 3.3 NB trials should be evaluated to at least 1 dp truncated or rounded
18	(a)	36 ÷ (7+3+2) "3" × 7	21	3	M1 for 36 ÷ (7+3+2) M1 (dep) for "3" × 7 or 3 or 2 A1 cao
	(b)	$51.5 \times \frac{8.5}{100} = 4.3775$ 51.5 – 4.3775 = 47.1225	47 or 47.1 or 47.12	4	M1 for 51.5 × $\frac{8.5}{100}$ or 4.37(75) seen M1 (dep) for 51.5 – "4.37(75)" A1 for 47 or better B1 (indep) for rounding their answer correctly to the nearest whole number or 1 or 2 d.p OR M1 for $51.5 \times \frac{100-8.5}{100}$ M1 for $51.5 \times$"0.915" or 0.515 × "91.5" A1 for 47 or better B1 (indep) for rounding their answer correct to the nearest whole number or 1 or 2 d.p

Paper 5523/04					
No		**Working**	**Answer**	**Mark**	**Notes**
19	(a)		Angle in a semicircle	1	B1 oe
	(b)	$12^2 + 16^2 = 400$ $\sqrt{400} = 20$	20	3	M1 for $12^2 + 16^2$ M1 for $\sqrt{144+256}$ A1 cao
	(c)	$\pi \times 10^2$	314	3	M1 for $\pi \times \left(\frac{"20"}{2}\right)^2$ M1 (indep) for correct order of evaluation of $\pi \times r^2$ for any r A1 for 314 – 315 inclusive

Paper 5523/04					
No		Working	Answer	Mark	Notes
20	(a)	$(1 \times 10)+(3 \times 15)+(5 \times 30)+(7 \times 35)+$ $(9 \times 25)+(11 \times 5) = 730$ "730" $\div$ 120 = 6.08333	6.08	4	M1 for use of *fx* with *x* consistent within intervals (including end points) M1 (dep) for use of midpoints M1 (dep on 1st M1) for use of $\frac{\sum fx}{\sum f}$ A1 6.08 to 6.085
	(b)		(10),25,55,90,115, 120	1	B1 for all correct
	(c)		graph	2	B1 ft for 5 or 6 points plotted correctly $\pm$ 1 full (2mm) square at the end of interval dep on sensible table (condone 1 addition error) B1(dep) for points joined by curve or line segments provided no gradient is negative – ignore any part of graph outside range of their points. (SC: B1 if 5 or 6 points plotted not at end but consistent within each interval and joined)

Paper 5523/04				
No	**Working**	**Answer**	**Mark**	**Notes**
(d)		72–74	2	M1 (ft dep on graph being cf) for reading from graph at 7 A1 ft ±1 full (2 mm) square Or B2 for $72-74$
21 (a) (b) (c) (d)		a^7 $15x^3y^4$ $x-1$ $(x+3)(x-3)$	1 2 1 1	B1 accept a^{4+3} B2 cao (B1 for two of 15, x^3, y^4 in a product) B1 cao B1 cao
22	80% = 220 $220 \div 80 \times 100$	275	3	M1 for recognising that 80% is equivalent to 220 M1 for $220 \div 80 \times 100$ oe A1 cao

Paper 5523/04					
No		**Working**	**Answer**	**Mark**	**Notes**
23			$x = 3$ $y = 0.5$	3	M1 for coefficients of x or y the same followed by correct operation, condone one arithmetical error M1 (dep) for substituting found value in one equation A1 cao SC: B1 for one correct answer only if M's not awarded
24			1.4×10^{10}	2	B2 for 1.4×10^{10} or 1.44×10^{10} (B1 for 14.4×10^{9} or 14400,000,000 or 14000,000,000 or 14×10^{9})
25	(a)	$\tan x = \frac{1.9}{3.2}$ $x = \tan^{-1}\left(\frac{1.9}{3.2}\right) = 30.7$	30.7	3	M1 for $\tan x = \frac{1.9}{3.2}$ or $\tan \frac{1.9}{3.2}$ M1 for $\tan^{-1}\left(\frac{1.9}{3.2}\right)$ A1 for 30.6 – 30.7
	(b)	90 + "30.7"	121	1	B1 (indep) ft for 90 + "30.7" rounded to 3 or 4 s.f

Paper 5523/04				
No	**Working**	**Answer**	**Mark**	**Notes**
26	SF$=\frac{12}{9}$ $\frac{12}{9}\times 6=8$	2	2	M1 for $\frac{12}{9}$ or $\frac{9}{12}$ or 1.33… seen or 0.75 seen or 8 seen or $\frac{6}{9}$ or $\frac{9}{6}$ or 0.66… or 1.5 or $\frac{1}{3}$ or 3 oe seen A1 cao

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
1	(a)	$x^2-4x+7x-28$	$x^2+3x-28$	2	M1 for 4 terms correct ignoring signs (e.g $x^2,4x,7x,28$) or 3 terms with correct signs (e.g $x^2,-4x,7x,-28$) A1 cao
	(b)		y^4+2y^2	2	B2 cao B1 for y^4 or $2y^2$
	(c)		$p(p+6)$	2	B2 for $p(p+6)$ or $p\times(p+6)$ (B1 for $p(ap+b)$ where a,b are numbers or $p+6$ seen on it's own, or part of an expression)
	(d)		$3x(2x-3y)$	2	B2 (B1 for $3(2x^2-3xy)$ or $x(6x-9y)$ or $3x(...)$)
2			question + response boxes oe	2	1st aspect: One question with time period (eg each night); ignore other questions. 2nd aspect: Response list (at least two), not overlapping.* 3rd aspect: Some mention of units (eg hours) in either question or responses Award B2 for all three aspects, or B1 for just two aspects. * 0-1, 2-3, 4-5 is OK, but 0-1, 1-2, 2-3 is not OK.
3	(b)		reflection in $y=x$	2	B2 cao accept the word "reflected" (B1 any statement including the word "reflection")
4	(a)	$5-3x=2x+2$ $5-2=2x+3x$	$\frac{3}{5}$	3	B1 for $2x+2$ seen OR $2.5-1.5x=x+1$ M1 for correct rearrangement of 4 terms A1 for $\frac{3}{5}$ oe
	(b)		$-3,-2,-1,0,1,2$	2	B2 (B1 for 5 correct and not more than one incorrect integers)

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
5	(a)	$\frac{2}{3}\times\frac{3}{4}=\frac{6}{12}$	$\frac{1}{2}$	2	M1 for $\frac{6}{12}$ or $\frac{3}{6}$ or $\frac{2\times3}{3\times4}$ A1 accept 0.5
	(b)	$1+2+\frac{8}{12}+\frac{9}{12}$	$4\frac{5}{12}$	2	M1 for attempt to convert to fractions with common denominator e.g two fractions, denominator of 12 A1 correct conversion : $\frac{8}{12}$ and $\frac{9}{12}$, or $\frac{20}{12}$ and $\frac{33}{12}$ seen (oe) A1 cao for $4\frac{5}{12}$ OR attempts to convert to decimals: must use at least 2dp M1 0.66+0.75 (or 1.66+2.75) or 0.67+0.75 etc A1 4.41, 4.417, 4.416 or 0.41, 0.417, 0.416 or 0.42, 4.42 A1 4.416 (*recurring*)
6	(a)(i)		5^6	1	B1 accept 15625, 5^{4+2}
	(ii)		5^3	1	B1 accept 125, 5^{9-6}
	(b)	$x+y=10$ and $x-y=4$	$x = 7$ $y = 3$	3	M1 for either $x + y = 10$ or $x-y=4$ A2 for both values correct [(A1 for one value correct) If M0, award B3 for both values correct or B2 for one value correct, otherwise B0] SC B2 for $x = 3$ or $y = 7$

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
7		$2\times\frac{1}{2}\times6\times8$ or 48 $8\times9+6\times9+10\times9$ or 72 +54 + 90	264 cm^2	4	M1 attempt to find the area of one face; $\frac{1}{2}\times6\times8$ or (8×9) or (6×9) or (10×9) or 72 or 54 or 90 or 24 or 48 M1 all five faces with an intention to add A1 cao numerical answer of 264 B1 (indep) cm^2 with or without numerical answer
8			$\frac{\pi ab^3}{3d}$ $3(c+d)^3$ $3\pi bc^2$	3	B3 (B1 for each one correct) Nb –B1 for each of the 4th,5th,6th tick
9	(a) (b)	$x+0.3+0.2+x=1$ 0.3×200	0.25 60	2 2	M1 for $x+0.3+0.2+x=1$ oe, or $0.5\div2$ A1 oe M1 0.3×200 A1 cao Accept 60 out of 200 (in words) SC B1 for $\frac{60}{200}$
10	(a) (b)		$(-12)-4\quad-2\quad(0)8$ 5 points plotted accurately points joined with smooth curve	3 2	B3 for all correct [(B1 for each one correct) B1 ±1 full (2mm) square ft table if at least B1 awarded (all 5 points plotted) B1 ft for any smooth curve if previous B1 gained NB: curve must pass within 1 full square of the points
11			m=3 n=5	2	B1 for 3 B1 for 5 (B2 for $2^3\times5$ or $2\times2\times2\times5$) SC: award B1 only if m=3, n=3, for 8×5 seen

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
12		$\frac{5--1}{-1-2} = -2$	$y = -2x + 5$	4	M1 for clear attempt to find gradient eg fraction with -1,5 in numerator, 2,-1 in denominator A1 for -2 cao B2 ft for $y =$"-2"$x + 5$ oe (eg $y = \frac{-6}{3}x + 5$) (B1 for $y = mx + 5$ or , $-2x+5$ or $y =$"-2"$x + c$)
13	(a)		$\frac{1}{4}$ on LH branch $\frac{2}{3}$ & $\frac{1}{3}$ & $\frac{2}{3}$ on RH branches	2	B1 cao B1
	(b)	$\frac{3}{4} \times \frac{2}{3} + \frac{1}{4} \times \frac{1}{3} = \frac{6}{12} + \frac{1}{12}$	$\frac{7}{12}$	3	M1 for $\frac{3}{4} \times \frac{2}{3}$ or $\frac{1}{4} \times \frac{1}{3}$ from their tree diagram M1 for sum of two products A1 for $\frac{7}{12}$ oe
	(c)	$n = 21 \times 4$ or $\frac{1}{6} : \frac{1}{4}$ oe $\frac{1}{6} \times 84$ or $21 \times \frac{2}{3}$	14	3	M1 for either $\frac{1}{3} \times \frac{3}{4} \left(= \frac{1}{4}\right)$ or $\frac{2}{3} \times \frac{1}{4} \left(= \frac{1}{6}\right)$ from their tree diagram M1 for 21×4 (=84) or $\frac{21}{3} \times 2$ A1 for 14 cao SC: B2 for 63 seen in fraction or ratio

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
14	(a)(i)		150	2	B1 accept 150 or 210
	(ii)				B1 for angle at the centre is twice the angle at the circumference
	(b)		30	3	B1 identifies angle between radius and tangent as $90°$ (may be in working or on diagram) M1 $360° - 90 - 90 - "150"$ A1 ft from (a)(i) excluding a negative answer Or B1 for 90 M1 for $2 \times (180 - 90 - "\frac{150}{2}")$ A1 ft from (a)(i) excluding a negative answer Or B3 for $180 - (a)$ SC: 180 – "210" can get B1 for $90°$ and/or B1 for "cyclic quadrilateral"
		$360 - 90 - 90 - "150"$ or $180 - "150"$			
15	(a)		0.2727…	1	B1 for 2.27 recurring or 0.2727…. oe or 0.273
	(b)	eg $x = 0.3939…$ so $100x = 39.3939…$ $99x = 39$ so $x = \frac{39}{99} = \frac{13}{33}$		3	M1 for $100x = 39.39…$ M1 dep for subtraction of both sides A1 for $\frac{13}{33}$ from correct proof **Alternative method** M1 for $13.000 \div 33$ M1 for remainders 31 and 13 A1 for 0.39 recurring SC:B1 for $\frac{39}{99}$

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
16	(a)	$d = kt^2$ $80 = k \times 4^2$	$d = 5t^2$	3	M1 for $d = kt^2$ or $d \propto t^2$ M1 sub d=80 and t=4 into their equation A1 for $d = 5t^2$ oe (cao)
	(b)		245	1	B1 ft from (a) using "k"
	(c)	$45 = 5t^2$	3	2	M1 ft from (a) for substituting d=45 into their equation A1 for 3 cao (condone inclusion of -3)
17	(a)(i) (ii)		(0, 9) (8, 25)	3	B1 cao B1 for $x = 8$ cao B1 for $y = 25$ cao SC: B1 for (25, 8)
	(b)	LHS = $\left(\frac{100-(x^2-16x+64)}{4}\right)$ = $\left(\frac{36+16x-x^2}{4}\right)$ RHS = $\left(\frac{36-2x+18x-x^2}{4}\right)$ = LHS		3	M1 for expansion of either set of brackets with at least 3 of 4 terms correct M1 for common denominator of 4 or multiplying through by 4 or reducing each numerator to a single term A1 for fully correct solution **Alternative method** M1 for $(5-\frac{(x-8)}{2})(5+\frac{(x-8)}{2})$ M1 for $(\frac{2\times5-(x-8)}{2})(\frac{2\times5+(x-8)}{2})$ A1 for $\frac{(18-x)(x+2)}{4}$

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
18	(a)	$\frac{810\pi}{90\pi}$ or 9 $\sqrt{9}$ or 3	12	3	M1 for $\frac{810\pi}{90\pi}$ or 9 or $\frac{1}{9}$ or 1:9 oe M1 for $\sqrt{\frac{810\pi}{90\pi}}$ or $\sqrt{9}$ or 3 or $\frac{1}{3}$ or $\sqrt{9}:\sqrt{1}$ oe A1 cao
	(b)	3^3 or 27 or 2700	2700π	2	SC:B1 for answer of 36 M1 for "3"^{3} or 27 or $\left(\sqrt{9}\right)^3 : (\sqrt{81})^3$ oe or 9^3 or 2700– A1 cao
19	(a)(i)		1	1	B1 cao
	(ii)		8	1	B1 cao
	(iii)	$64^{-\frac{2}{3}} = \frac{1}{64^{\frac{2}{3}}}$ or $64^{-\frac{2}{3}} = (4^2)^{-1}$	$\frac{1}{16}$	2	M1 for knowing negative power is a reciprocal or power of $\frac{1}{3}$ root is a cube root A1 cao for $\frac{1}{16}$
	(b)	$\sqrt{27} = \sqrt{9\times3}$ or $\sqrt{27} = 3\sqrt{3}$ or $\sqrt{27} = 3^{3/2}$	$\frac{5}{2}$ oe	2	M1 for $\sqrt{27} = \sqrt{9\times3}$ or $\sqrt{27} = 3^{3/2}$ A1 for $\frac{5}{2}$ oe (cao) **Alternative method** M1 for $9\times27 = 3^{2n}$ A1 for $\frac{5}{2}$ oe (cao)

Paper 5525/05					
No		**Working**	**Answer**	**Mark**	**Notes**
20	(a)(i) (ii) (b)(i) (ii)		(90, 1) (180, 0) (45, 0) (90, -3)	2 2 	B1 cao could be indicated on diagram B1 cao could be indicated on diagram B1 cao could be indicated on diagram B1 cao could be indicated on diagram
21		$\frac{1}{3}\pi x^2 h = \frac{4}{3}\pi(2x)^3$ $x^2 h = 4\times 8x^3$	$32x$	3	M1 for substitution in correct formulae M1 (dep.) for correct unsimplified expression eg $h = \frac{\frac{4}{3}\pi(2x)^3}{\frac{1}{3}\pi x^2}$ oe or $h = 8x$ oe A1 for $32x$ cao
22	(a)	$\left(\overline{OM} =\right)\ \boldsymbol{a} + 2\boldsymbol{b}$ $\left(\overline{ON} =\right) 3\boldsymbol{a}$ or $\frac{6}{2}\boldsymbol{a}$ $\left(\overline{MN} =\right)\ -\boldsymbol{a} - 2\boldsymbol{b} + 3\boldsymbol{a}$	$2\boldsymbol{a} - 2\boldsymbol{b}$	2	B2 (B1 for either $\overline{OM}$ or $\overline{ON}$ or $-\boldsymbol{a} - 2\boldsymbol{b} + 3\boldsymbol{a}$ SC: B1 for $2\boldsymbol{b} - 2\boldsymbol{a}$
	(b)	$\left(\overline{OX} =\right)\ 2\boldsymbol{a} + \boldsymbol{b}$ $\left(\overline{OY} =\right) \boldsymbol{b} + 4\boldsymbol{a}$ (½ $\overline{QR}$ =) $2\mathbf{a} - \boldsymbol{b}$ or (½ $\overline{RQ}$ =) $\boldsymbol{b} - 2\boldsymbol{a}$	$\overline{XY} = 2\boldsymbol{a}$ (hence parallel)	2	B1 for either $\overline{OX}$ or $\overline{OY}$ or (½ $\overline{QR}$) B1 for $\overline{XY} = 2\boldsymbol{a}$ or $\overline{YX} = -2\boldsymbol{a}$

Paper 5525/06					
No		Working	Answer	Mark	Notes
1	(a)	$v - u = 5t$	$\frac{v-u}{5}$	2	M1 for isolating $\pm 5t$ or $\pm t$ or for dividing through by 5 A1 oe
	(b)	$x - 3 = 5x - 25$ $22 = 4x$	$5\frac{1}{2}$	3	M1 for $x - 3 = 5(x-5)$ or $\frac{x}{5} - \frac{3}{5} = x - 5$ M1 for isolating terms in x correctly from $ax + b = cx + d$ A1 cao accept $5\frac{1}{2}$, $\frac{11}{2}$, 5.5
2	(a)	$36 \div (7+3+2)$ "3" $\times 7$	21	3	M1 for $36 \div (7+3+2)$ M1 (dep) for "3" $\times 7$ or 3 or 2 A1 cao
	(b)	$51.5 \times \frac{8.5}{100} = 4.3775$ $51.5 - 4.3775 = 47.1225$	47 or 47.1 or 47.12	4	M1 for $51.5 \times \frac{8.5}{100}$ or 4.37(75) seen M1 (dep) for 51.5 – "4.37(75)" A1 for 47 or better B1 (indep) for rounding their answer correctly to the nearest whole number or 1 or 2 d.p OR M1 for $51.5 \times \frac{100 - 8.5}{100}$ M1 for $51.5 \times$"0.915" or 0.515 ×"91.5" A1 for 47 or better B1 (indep) for rounding their answer correct to the nearest whole number or 1 or 2 d.p

Paper 5525/06					
No		**Working**	**Answer**	**Mark**	**Notes**
3		3→15 4→48 3.1→17.3(91) 3.2→19.9(68) 3.3→22.7(37) 3.4→25.7(04) 3.5→28.8(75) 3.4→25.7(04) 3.3→22.7(37) 3.35→24.1(95375)	3.3	4	B2 for trial between 3.3 and 3.4 inclusive (B1 for trial between 3 and 4 inclusive) B1 for different trial between 3.3 and 3.4 exclusive B1 (dep on at least one previous B1) for 3.3 NB trials should be evaluated to at least 1 dp truncated or rounded
4	(a)		Angle in a semicircle	1	B1 oe
	(b)	$12^2 + 16^2 = 400$ $\sqrt{400} = 20$	20	3	M1 for $12^2 + 16^2$ M1 for $\sqrt{144+256}$ A1 cao
	(c)	$\pi \times 10^2$	314	3	M1 for $\pi \times \left(\frac{"20"}{2}\right)^2$ M1 (indep) for correct order of evaluation of $\pi \times r^2$ for any r A1 for 314 – 315 inclusive

Paper 5525/06					
No		Working	Answer	Mark	Notes
5	(a)	$(1 \times 10)+(3 \times 15)+(5 \times 30)+(7 \times 35)+$ $(9 \times 25)+(11 \times 5) = 730$ "730" ÷ 120 = 6.08333	6.08	4	M1 for use of *fx* with *x* consistent within intervals (including end points) M1 (dep) for use of midpoints M1 (dep on 1st M1) for use of $\frac{\sum fx}{\sum f}$ A1 6.08 to 6.085
	(b)		(10),25,55,90, 115,120	1	B1 for all correct
	(c)		graph	2	B1 ft for 5 or 6 points plotted correctly ± 1 full (2mm) square at the end of interval dep on sensible table (condone 1 addition error) B1(dep) for points joined by curve or line segments provided no gradient is negative – ignore any part of graph outside range of their points. (SC: B1 if 5 or 6 points plotted not at end but consistent within each interval and joined)
	(d)		72 – 74	2	M1 (ft dep on graph being cf) for reading from graph at 7 A1 ft ±1 full (2 mm) square Or B2 for 72 – 74

Paper 5525/06					
No		Working	Answer	Mark	Notes
6	(a)		a^7	1	B1 accept a^{4+3}
	(b)		$15x^3y^4$	2	B2 cao (B1 for two of 15, x^3, y^4 in a product)
	(c)		$x-1$	1	B1 cao
	(d)		$(a + 3b)(a - 3b)$	2	B2 for $(a + 3b)(a - 3b)$ (B1 for $(a \pm 3b)(a \pm 3b)$
7		80% = 220 $220 \div 80 \times 100$	275	3	M1 for recognising that 80% is equivalent to 220 M1 for $220 \div 80 \times 100$ oe A1 cao
8			$x = 3$ $y = 0.5$	3	M1 for coefficients of x or y the same followed by correct operation, condone one arithmetical error M1 (dep) for substituting found value in one equation A1 cao SC: B1 for one correct answer only if Ms not awarded
9			1.4×10^{10}	2	B2 for 1.4×10^{10} or 1.44×10^{10} (B1 for 14.4×10^9 or 14400,000,000 or 14000,000,000 or 14×10^9)

Paper 5525/06					
No		Working	Answer	Mark	Notes
10	(a)	$\tan x = \frac{1.9}{3.2}$ $x = \tan^{-1}\left(\frac{1.9}{3.2}\right) = 30.7$	30.7	3	M1 for $\tan x = \frac{1.9}{3.2}$ or $\tan \frac{1.9}{3.2}$ M1 for $\tan^{-1}\left(\frac{1.9}{3.2}\right)$ A1 for 30.6 – 30.7
	(b)	90 + "30.7"	121	1	B1 (indep) ft for 90 + "30.7" rounded to 3 or 4 s.f
11	(a)	$SF = \frac{12}{9}$ $\frac{12}{9} \times 6 = 8$	2	2	M1 for $\frac{12}{9}$ or $\frac{9}{12}$ or 1.33… seen or 0.75 seen or 8 seen or $\frac{6}{9}$ or $\frac{9}{6}$ or 0.66… or 1.5 or $\frac{1}{3}$ or 3 oe seen A1 cao
	(b)	$SF = \frac{9}{12}, \frac{9}{12} \times 7 = 5.25$	5.25	2	M1 for $\frac{BE}{7} = \frac{9}{12}$ oe A1 cao

Paper 5525/06					
No		Working	Answer	Mark	Notes
12	(a)	$84 = 6.7\pi + 2 \times 6.7 + 2a$ $2a + 13.4 = 62.95\ldots$ or $2a + 34.44 = 84$	24.8	3	M1 for substituting correctly, π may be left M1 for correct rearrangement as far as $\pm 2a$ A1 for $24.7 - 24.8$
	(b)	$P = \pi r + 2r + 2a$ $P - 2a = \pi r + 2r$ $P - 2a = (\pi + 2)r$	$\frac{P-2a}{\pi+2}$	3	M1 subtracting $2a$ from each side M1 for factorising to get $(\pi+2)r$ A1 for $\frac{P-2a}{\pi+2}$ oe S.C $\frac{p-2a}{5.14}$ oe is M1 M1 A0
13		Area $\Delta ABC = \frac{1}{2} \times 14 \times 8 \times \sin 106$ (= 53.8)	53.8	3	M1 for $\frac{1}{2} \times 14 \times 8 \times \sin 106$ M1 (dep) for $56 \times 0.961(26..)$ or $107.6\ldots$ A1 $53.8 - 53.9$ SC 107.6 is B2
14	(a)		4.5	1	B1 cao
	(b)	$500 \times 1.045^{20} = 1205.857\ldots\ldots$	1205.86	2	M1 for 500×1.045^{20} A1 for $1205.85 - 1206$ (SC:B1 for $705.85 - 706$ no working)

Paper 5525/06				
No	**Working**	**Answer**	**Mark**	**Notes**
15 (a)	$6x^2 + 11x - 10 + 6x - 4 = 25$ $6x^2 + 17x - 39 = 0$		3	M1 for an expression for the area involving either $(3x-2)(2x+5)+2(3x-2)$ or $3x(3x-2)+(3x-2)(7-x)$ or $3x(2x+5)-2(7-x)$ or $(3x-2)^2 + 2(3x-2) + (3x-2)(7-x)$ where in each case at least one of 2 or 3 product terms must be correct M1 (indep) for one correct expansion involving x^2 A1 for simplification to final answer

Paper 5525/06					
No		Working	Answer	Mark	Notes
15	(b)(i)	$x = \frac{-17 \pm \sqrt{17^2 - 4\times 6\times(-39)}}{2\times 6}$ $= \frac{-17 \pm \sqrt{289+936}}{12}$ $x = +\frac{18}{12}$ or -4.33	1.5, $-\frac{13}{3}$	4	M1 for $x = \frac{-17 \pm \sqrt{17^2 - 4\times 6\times(-39)}}{2\times 6}$ up to signs in b & c M1 for $x = \frac{-17 \pm \sqrt{1225}}{12}$ A1 $x = 1.5$ or -4.33, or better OR M2 for $(3x+13)(2x-3)$ (M1 for $(3x \pm a)(2x \pm b)$ with ab = ±39 A1 x = 1.5 or – 4.33, or better
		$x^2 + \frac{17}{6}x - \frac{39}{6} = 0$ $\left(x+\frac{17}{12}\right)^2 - \left(\frac{17}{12}\right)^2 - \frac{39}{6} = 0$ $\left(x+\frac{17}{12}\right)^2 = \left(\frac{17}{12}\right)^2 + \frac{39}{6}$			OR M1 for $\left(x+\frac{17}{12}\right)^2$ *seen* M1 $\left(x+\frac{17}{12}\right)^2 = \left(\frac{17}{12}\right)^2 + \frac{39}{6}$ A1 x = 1.5 or – 4.33, or better SC:M1 for answer "1.5" with no working or T & I
	(ii)		8		B1 cao length = 8

Paper 5525/06					
No		**Working**	**Answer**	**Mark**	**Notes**
16	(a)	P(win) = $\frac{2}{5}\times\frac{3}{5}+\frac{2}{5}\times\frac{1}{5}(=\frac{8}{25})$	$\frac{8}{25}$	3	M1 for $\frac{2}{5}\times\frac{3}{5}$ or $\frac{2}{5}\times\frac{1}{5}$ or for clearly identifying in P(R) $\times$P(R) + P(B) $\times$ P(B) M1 for P(win) = " $\frac{2}{5}$ " $\times$ " $\frac{3}{5}$ " + " $\frac{2}{5}$ " $\times$ " $\frac{1}{5}$ " A1 for $\frac{8}{25}$, oe
	(b)	$\frac{8}{25}$ $\times$ 100 (=32) 100 $\times$ 20 – "32" $\times$ 50	£4	2	M1 for (" $\frac{8}{25}$ " $\times$100)$\times$50 or $\times$ 0.5 A1 cao

Paper 5525/06					
No		**Working**	**Answer**	**Mark**	**Notes**
17		Lower bound of 1200 is 1150 Upper bound of 60 is 65 $1150 \div 65$	17	4	B1 for 1150 or 1250 seen B1 for 65 or 55 seen M1 (Lower bound of load) ÷ (Upper bound of weight) Where 1150 ≤ LB load < 1200 and 60 < UB Weight ≤ 65 A1 for 17 requires fully correct working OR B1 for 1150 or 1250 seen B1 for 65 or 55 seen M1 (upper bound of load) ÷(lower bound of weight) Where 1200 ‹ UB load ≤ 1250 and 55 ≤ LB weight < 60 A1 for 22 requires fully correct working OR M2 $1200 \div 55$ A1 21.8 A1 21 requires fully correct working OR M2 $1200 \div 65$ A1 18.4(6) A1 18 requires fully correct working
18	(a)	$3^4 x^4 y^8$	$81x^4y^8$	2	B2 for $81x^4y^8$ (B1 for 2 of 81, x^4, y^8)
	(b)	$\frac{x(x-3)}{(x-5)(x-3)}$	$\frac{x}{x-5}$	3	B1 for $x(x-3)$ B1 for $(x-5)(x-3)$ B1 cao

Paper 5525/06					
No		Working	Answer	Mark	Notes
19	(a)	$6^2 - 2^2 = 32$	5.66	2	M1 for $6^2 - 2^2$ (= 32) A1 5.65 – 5.66
	(b)	V  $DVA = 2 \times \sin^{-1}(\frac{2}{6})$ OR $\cos DVA = \frac{6^2 + 6^2 - 16}{2 \times 6 \times 6}$ $= \frac{56}{72}$ $DVA = \cos^{-1}\left(\frac{56}{72}\right) = 38.94$	38.9	3	M1 $\sin x = \frac{2}{6}$ oe M1 for $DVA = 2 \times \sin^{-1}(\frac{2}{6})$ A1 38.9 – 38.95 OR M1 for $(\cos DVA =) \frac{6^2 + 6^2 - 4^2}{2 \times 6 \times 6}$ M1 for $DVA = \cos^{-1}\left(\frac{56}{72}\right)$ A1 38.9 – 38.95

Paper 5525/06				
No	**Working**	**Answer**	**Mark**	**Notes**
19 (c)	$AC^2 = 2^2 + 2^2 - 2 \times 2 \times 2 \times \cos 120^o$ $AC = \sqrt{12}$ **OR** $AN = 2 \times \sin 60 = \sqrt{3}$ **OR** $VN = \sqrt{"32"+1} = \sqrt{33}$ $\cos AVC = \frac{6^2+6^2-12}{2\times 6\times 6}$ $\cos AVC = \frac{60}{72}$ **OR** $AVC = 2 \times \sin^{-1}\frac{\sqrt{3}}{6}$, using AN **OR** $AVC = 2\times\cos^{-1}\frac{\sqrt{33}}{6}$, using VN	33.6	4	M1 for any valid method for AC or AN or VN where N is the midpoint of AC A1 for $AC^2 = 12$ or $AC = \sqrt{12}$ (= 3.46…) or $AN = \sqrt{3}$ (=1.73…)or $VN = \sqrt{33}$ (=5.74…) M1 (indep) for correct method to find angle AVC A1 33.55 – 33.6

Paper 5525/06					
No		**Working**	**Answer**	**Mark**	**Notes**
20	(a)	Graph translated 1 unit to the right passing through the points (-1,0), (1,2) and (2,0)		2	M1 for translation of $\begin{pmatrix}1\\0\end{pmatrix}$ or $\begin{pmatrix}-1\\0\end{pmatrix}$ A1 for right through the 3 points, $\pm\frac{1}{4}$ sq
	(b)	Graph stretched 2 units parallel to y-axis; passing through the points (-2,0), (0,4) and (1,0)		2	M1 for graph stretched parallel to the y-axis by scale factor 2 A1 through all 3 points; $\pm\frac{1}{4}$ sq not on grid at $x = 2$

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481

Email publications@linneydirect.com

Order Code UG016720 Summer 2005

For more information on Edexcel qualifications, please visit www.edexcel.org.uk/qualifications
Alternatively, you can contact Customer Services at www.edexcel.org.uk/ask or on 0870 240 9800

Edexcel Limited. Registered in England and Wales no.4496750
Registered Office: One90 High Holborn, London, WC1V 7BH

edexcel

Edexcel GCSE

Mathematics A 1387

June 2007

advancing learning, changing lives

Mark Scheme

Edexcel GCSE
Mathematics A 1387

Table Of Contents

Notes on Marking Principles	3 - 4
5521 / 01 Mark Scheme	5 - 11
5521 / 02 Mark Scheme	12 – 17
5523 / 03 Mark Scheme	18 – 27
5523 / 04 Mark Scheme	28 – 35
5525 / 05 Mark Scheme	36 – 43
5525 / 06 Mark Scheme	44 - 52

NOTES ON MARKING PRINCIPLES

1 **Types of mark**

M marks: method marks
A marks: accuracy marks
B marks: unconditional accuracy marks (independent of M marks)

2 Abbreviations

cao – correct answer only
ft – follow through
isw – ignore subsequent working
SC: special case
oe – or equivalent (and appropriate)
dep – dependent
indep - independent

3 No working

If no working is shown then correct answers normally score full marks
If no working is shown then incorrect (even though nearly correct) answers score no marks.

4 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.
If there is no answer on the answer line then check the working for an obvious answer.

5 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

6 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

7 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

8 Linear equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

9 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

Remember: if you are having difficulty making a decision on how you should mark a candidate response contact your Team Leader for advice, or send the item to review.

Paper 5521_01					
No		Working	Answer	Mark	Notes
1		Cat |||| /||| 8 Dog |||| | 6 Fish || 2 Hamster |||| 4	8, 6, 2, 4	3	M1 for attempt to tally or one frequency correct in either column A1 for 1 frequency correct or all tallies correct in correct column A1 for all frequencies correct (accept if /20)
2			27, 35, 42, 67, 118	1	B1 cao
3	(a)		Diameter drawn	1	B1 for a diameter
	(b)		Right angle marked	1	B1 *R* marked correctly
	(c)		Rectangle drawn	1	B1 for a rectangle
4	(a)		40	1	B1 cao
	(b)		50	1	B1 cao
	(c)		3 full loaves 4 full loaves + 1 half loaf	2	B1 for 3 full loaves B1 for 4 full loaves + 1 half loaf
5	(a)		7252	1	B1 cao
	(b)		Three thousand and eighty six	1	B1 accept 3 thousand and eighty six (condone 0 hundred)
	(c)		4600	1	B1 accept 4600
	(d)		200	1	B1 for 200 or 2 hundred or 100 or hundred
6	(i)		Cube	2	B1 for 'cube' (accept 'cuboid') ignore spelling
	(ii)		Cylinder		B1 for 'cylinder' ignore spelling
7		5×100	500	2	B2 for 490 or 500 or 510 (B1 for either 5 or 5.0 or 100 seen)

Paper 5521_01					
No		Working	Answer	Mark	Notes
8	(a)		8.4 cm or 84 mm	2	B2 allow ±2mm (B1 for 8.4 or 84, B1 for appropriate unit)
	(b)		37°	1	B1 allow ±2°
9	(a)		Carbon black	1	B1 accept 'black carbon' accept 26%
	(b)		0.1(0)	1	B1 cao
	(c)		0.04	1	B1 cao
	(d)	$\frac{26}{100}$	$\frac{13}{50}$	2	M1 for $\frac{26}{100}$ A1 cao
10	(a)		97	1	B1 cao
	(b)		London Reading	1	B1 cao
	(c)	41 + 57 + 58	156	3	M1 for two of 41, 57, 58 M1(dep) for '41' + '57' + '58' A1 cao
11	(a)		3	1	B1 cao allow ± 0.2
	(b)		–5	1	B1 cao allow ±0.2

Paper 5521_01					
No		**Working**	**Answer**	**Mark**	**Notes**
12	(a)		Centimetres (cm) miles litres (*l*)	3	B3 (B1 for each correct answer) accept abbreviations
	(b)		300	1	B1 cao
	(c)	'1500>1400' or '1.5>1.4'	Reason	1	B1 for No and '1500>1400' or '1.5>1.4'
13	(a)	Height of bars 12, 8, 6 lines drawn between points	Bars drawn	2	B2 for 3 bars correctly drawn (B1 for 2 bars correct)
	(b)		July and August	1	B1 oe
	(c)	24 – 4	20	1	B1 (Accept '4 to 24' oe)
	(d)	The temperatures are rising	Temp's rising oe	1	B1 for reason
14	(a) (i)		$(-4, 3)$	2	B1 cao
	(ii)		$(2, -1)$		B1 cao
	(b) (i)	*D* marked at $(-4, -1)$	Point marked on grid	2	B1 for point marked at $(-4, -1)$ cao
	(ii)		$(-4, -1)$		B1 ft
15	(a)			1	B1 cao
	(b)			1	B1 cao

Paper 5521_01				
No	**Working**	**Answer**	**Mark**	**Notes**
16 (a)		(Pat +) reason	1	B1 correct comment (Pat may be implied)
(b)	21 ÷ 3	7	1	B1 cao
17 (i)		S marked at 1	3	B1 for S within ½ cm of 1
(ii)		P marked at 0		B1 for P marked at 0 cao
(iii)		Q marked at "1/3"		B1 for Q marked at 1/3 ± 1 cm use overlay
18	$6 + 6 + 3$ or $2\frac{1}{2} \times 6$	15	3	M1 for realizing 6 glasses in one bottle M1 for realizing 3 glasses in ½ a bottle A1 cao (M2 for attempt to find $2\frac{1}{2} \times 6$) oe
19	15 and 16 parts shaded Alternative 1 $\frac{3}{4} = 0.75$, $\frac{4}{5} = 0.8$ Alternative 2 $\frac{3}{4} = \frac{15}{20}$, $\frac{4}{5} = \frac{16}{20}$	$\frac{4}{5}$ + reason	3	M1 for correctly shading 15 parts for 3/4 M1 for correctly shading 16 parts for 4/5 A1 (dependent on M2) for selection of 4/5 Alternative 1 M1 for $\frac{3}{4} = 0.75$ M1 for $\frac{4}{5} = 0.8$ A1 (dep on M2) for selection of 0.8 or $\frac{4}{5}$ or $\frac{16}{20}$ Alternative 2 M1 for $\frac{3}{4} = \frac{15}{20}$ M1 for $\frac{4}{5} = \frac{16}{20}$ A1(dep on M2) for selection of $\frac{4}{5}$ or $\frac{16}{20}$

Paper 5521_01					
No		**Working**	**Answer**	**Mark**	**Notes**
20	(a)		−5 (−3) (−1) 1 3 5	2	B2 cao (B1 for any 2 or 3 correct)
	(b)	Points plotted	Line	2	B2 for line from (−1, −5) to (4, 5) (B1 ft for plotting at least 5 "points")
21		Shapes shaded on grid	6 tessellating shapes	2	B2 for fully correct with 5 or more additional shapes, no gaps (B1 for 4 shapes tessellating with at least one shape inverted, with or without the given shape ignore extras)
22	(a)		$\frac{7}{20}$	1	B1 for $\frac{7}{20}$ oe
	(b)	7 + 2 (or 20 – 11) are not lime flavour	$\frac{9}{20}$	1	B1 for $\frac{9}{20}$
	(c)		0	1	B1 for 0, zero or nought ($\frac{0}{20}$ gets B0)
23	(a)		$4a$	1	B1 accept $4\times a$, $a \times 4$, $a4$
	(b)		$12b$	1	B1 accept $12\times b$, $b \times 12$, $b12$
	(c)		$2a + 6b$	2	B2 cao (B1 for $2a$ or $6b$ seen)
	(d)		$x(x-6)$	2	B2 cao (B1 for $x(ax+b)$ where a, b are numbers not equal to 0 or $x - 6$ seen on its own, or as part of an expression)
24	(a)		40	1	B1 cao
	(b)		45	1	B1 for 42 – 48 accept ¾ hour
	(c)	40×2 or $\frac{40}{30}\times 60$ or $40\div\frac{1}{2}$	80	2	M1 for 40×2 or $\frac{40}{30}$ or $40\div\frac{1}{2}$ A1 cao NB $\frac{40}{45}\times 60$ gets M0 A0

Paper 5521_01					
No		**Working**	**Answer**	**Mark**	**Notes**
25	(a)		4560	1	B1 cao
	(b)		45.6	1	B1 cao
	(c)		2.4	1	B1 cao
26		$-7 - 3 = -10$ $2 \times -10 = -20$ $-20 \div 4$	-5	3	M1 for substitution of 2 and -7 into $p(q - 3)$ or sight of -20 or $-14 - 6$ M1dep for '-20' $\div 4$ A1 cao B1 SC for sight of -10 if M0 awarded
27	(a)		Reflection in y-axis	1	B1 cao
	(b)		Rotation by half turn about (0,0)	2	B2 cao (B1 for half turn not about (0,0).)
	(c)		Enlargement Scale factor 3 Centre (0,0)	3	B1 for 'enlargement' B1 for "scale factor 3" or 3 seen B1 for 'centre (0,0)
28	(a)		Reason	1	B1 for 'The frequencies are nearly equal' oe
	(b)	$1\times26+2\times26+3\times23+4\times25 = 247$ $247/100$	2.47	3	M1 for fx (attempting at least 2 relevant products) M1 for $\sum fx \div 100$ A1 2.47 cao

Paper 5521_01				
No	**Working**	**Answer**	**Mark**	**Notes**
29	5 × 5 × 6	150	4	M1 for attempt at 1 division (e.g. 40 ÷ 8), may be implied by marks or number on one edge of diagram or by 5 or 6 seen M1 for attempt at 3 divisions (40÷8, 40÷8, 60÷10), may be implied by marks or numbers on diagram or by 5,5 and 6 seen. M1 (dep on 1st M1) for "5" × "5" × "6" A1 cao Alternatively M1 for 40 × 40 × 60 **or** 8 × 8 × 10 **or** 96000 or 640 seen M1 for 40 × 40 × 60 **and** 8 × 8 × 10 or 96000 **and** 640 seen M1 (dep on 1st M1) for "(40 × 40 × 60)" ÷ "(8 × 8 × 10)" A1 cao SC:B1 for dividing area of one carton face by area of corresponding box face if M0

Paper 5521_02					
No		Working	Answer	Mark	Notes
1	(a)	Draw diagram.	Diagram	1	B1 cao
	(b)		13,16	1	B1 cao
	(c)		31	1	B1 cao
2	(a)		8	1	B1 cao
	(b)		14	2	B2 for 14 (B1 for 13 or 15)
	(c)		16	2	B2 for 16 (B1 for 15, 17 or 8)
3	(a)		4,7 drawn	2	B2 for car height 4 and bus height 7, (B1 for one correct)
	(b)		6	1	B1 cao
	(c)		Walk	1	B1
	(d)		27	1	B1 cao
4	(a)		40	1	B1 for 40–41 inclusive
	(b)		12	1	B1 for 11.5 – 12.5 inclusive
5	(a)		Row complete	2	B2 for 1+..+11; 36 (B1 for one of the 2 cells complete)
	(b)		Square	1	B1 "square"
6		One line of symmetry		1	B1 within 2mm of centre of base / 2mm of vertex

Paper 5521_02				
No	**Working**	**Answer**	**Mark**	**Notes**
7 (a) (i)		09 06	2	B1 (accept 9 06 oe)
(ii)		39		B1 cao
(b)		06 55	1	B1 (accept 6 55 oe)
(c)		2h 6min	1	B1 cao
(d)		15 min	2	M1 for 0906 – 0645 – "(c)" or 0906 – 0645 − 2hr 6min or 2hr 21min − "(c)" or 2hr 21min − 2hr 6 min or 141 − 126 or 20 − 5 A1 cao SC: B1 for 55 or 75 or 93 seen
8 (i)		8,10,12,20 or 30	3	B1 at least one of 8, 10, 12, 20, 30 (no extras)
(ii)		8,12 or 20		B1 at least one of 8,12, 20 (no extras)
(iii)		3 or 5		B1 3 or 5 or both (no extras)
9 (a)		C or G	1	B1 at least one of C or G (no extras)
(b)		A and F	1	B1 cao
(c)		2	1	B1 (accept –2)
10 (a)		$\frac{7}{10}$	1	B1 7/10 oe
(b)(i)		4 squares	2	B1 4 squares shaded
(ii)		80%		B1 80% or ft from unshaded part (no ft from 0% or 100%)
(c)(i)		2.5	2	B1 2.4–2.6 inclusive
(ii)		1.7		B1 1.6–1.8 inclusive
11 (a)		2	1	B1 for 2 or –2
(b)		14	1	B1 for 14 or –14

Paper 5521_02					
No		**Working**	**Answer**	**Mark**	**Notes**
12		2 × 8.50 = 17.00 3 × 4.50 = 13.50 Total = 30.50 50.00 – 30.50	19.5(0)(*p*)	3	M1 for adding 5 correct values or 2 × 8.50 + 3 × 4.50 (ignore units) or 30.5(0) or 3050 seen M1 dep for 50 − "30.50" (ignore units) (OR M1for adding at least 1 adult ticket and at least 1 child ticket **and** subtracting from 50) A1 cao SC: B1 for 24 or 37 or 2400 or 3700 seen
13	(a)		Hexagon	1	B1
	(b)(i)		120	2	B1 cao
	(ii)		Str line		B1 reference to a (straight) line and 180°
	(c)		Obtuse	1	B1 Accept "interior"
14	(a)	2, 2, 3, 3, 3, 4, 4, 4, 5, 6	3.5	2	M1 ordering the numbers (condone 1 error or omission) A1 cao
	(b)	36 ÷ 10	3.6	2	M1 sum of numbers ÷ 10 A1 cao SC B1 for 3r 6
	(c)	6 – 2	4	1	B1 cao
15	(a)		Paul	1	B1 cao
	(b)	36 ÷ 2 oe	18	1	B1 cao
	(c)	60/360 =	1/6	2	M1 60/360 oe A1 cao

Paper 5521_02					
No		**Working**	**Answer**	**Mark**	**Notes**
16		4.7 ÷ 5.9 = 0.796610169	0.7966..	2	B2 for 0.7966 or better (B1 for 0.8, 0.80, 0.79, 0.796, 0.797 or digits 59 seen
17		$6x - 7 + 7 = 38 + 7$ $6x = 45$	7.5	2	M1 $6x = 45$ or +7 both sides A1 7.5 oe; accept 45/6 oe
18		**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	3	B3 all six entries correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
19	(a)	900 × 1.70 =	1530	2	M1 900 × 1.7(0) or digits 153(0) seen A1 cao
	(b)	160 ÷ 1.70 =	£94.12 or £94.11	2	M1 160 ÷ 1.7(0) or digits 941(....) seen A1 cao
20	(a)(i)	180 – 54 (=126) "126" ÷ 2	63	3	M1 for (180–54) ÷ 2 A1 cao
	(ii)		Reason		B1 (indep) angles in triangle add to 180 OR equal angles in isosceles triangle OR equal angles and 2 sides the same (B0 if any incorrect reasoning given eg parallel, equilateral triangle)
	(b)	180 – "x"	117	1	B1 117 or ft 180 – "x" if $x < 90$
21	(a)	3 × 35 + 50	155	2	M1 for 3 × 35 + 50 or digits 155 seen A1 cao
	(b)	260 – 50 = 210 210 ÷ 35 =	6	3	M1 for 260–50 or 210 seen. M1 for "260-50" ÷ 35 or 210÷35 A1 cao SC B1 for starting at a number between 100 and 170 and adding at least two 35's and showing a total between 230 and 290 OR For adding at least three 35's, perhaps with other numbers, and showing a total between 180 and 240 (or between 230 and 290 if 50 is included in the sum)

Paper 5521_02				
No	**Working**	**Answer**	**Mark**	**Notes**
(c)		P=35h + 50	3	B3 for P=35h+50 or P=35×h+50 oe (B2 for correct RHS or P=h + 50 ×35 or P=35h+k where k is numerical oe) (B1 for P = some other linear expression in h, OR h + 50 ×35 OR 35h seen) NB: P=h scores no marks; ignore £ signs. SC B2 for $h = \frac{P-50}{35}$
22 (a)		Elevation	2	B2 for 4 vertical squares. Accept 4 by 1 rectangle. (B1 for 4 vertical squares with one square added or one parallelogram added at the top, or 3 vertical squares, or 4 horizontal squares)
(b)		Plan	2	B2 for 2 adjacent squares, vertical or horizontal. Accept 2 by 1 rectangle. (B1 for 3 adjacent horizontal or vertical squares or a rectangle with sides in the ratio 2:1)
23 (i)		5	3	B1 cao
(ii)		9		B1 cao
(iii)		6		B1 cao
24	$45.00 + 45.00 \times \frac{15}{100} =$ $45.00 + 6.75 =$	51.75	3	M2 for $45.00 + 45.00 \times \frac{15}{100}$ oe or 45.00×1.15 oe OR $45.00 + 6.75$ OR complete method or 5175 seen. (M1 for $45.00 \times \frac{15}{100}$ oe OR 6.75 seen OR 675 seen OR correct method for calculating 15% of 45) A1 cao SC Award B2 for an answer of 38.25

Paper 5521_02				
No	**Working**	**Answer**	**Mark**	**Notes**
25 (a)		Points	1	B1 all three points ±1 full square
(b)		Negative	1	B1 Negative (ignore additional descriptors unless contradictory)
(c)		lobf	1	B1 A single straight line drawn to cross between (5,30), (5,40) and (40,0), (40,15); accept freehand if considered to be straight.
(d)(i)		18–25	2	B1 18g–25g inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient.
(ii)		30–40		B1 30–40 min inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient
26		300 3 75 150	3	B3 for 4 correct answers (B2 for 2 or 3 correct answers) (B1 for 1 correct answer)
27	π × 0.65	2.04–2.05	2	M1 for π × 0.65 or 3.14 x 0.65 or 3.142 x 0.65 A1 for 2.04–2.05 SC Award B1 for 2.0 seen (not 2)
28	5 miles = 8 km 70mph÷5×8 = 112 km/h OR 120km/h ÷ 8 × 5 = 75 mph Faster than 70 mph	70mph (Great Britain) (112 km)	3	M1 5 miles = 8 km; OR 70 mph is about 100 km/h OR 1km=0.6(25) miles OR 1mile=1.6km oe M1 70 ÷ 5 × 8 (=112) or 120 ÷ 8 × 5 (=75) A1 (dep on at least M1) GB or 70 mph Refer to both answer line and working. NB GB or 70 mph without working scores 0 marks

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
1		Cuboid drawn	2	B2 for correct isometric drawing in any orientation (ignore points 'behind', mark 7 vertices only); accept lines drawn near to dots as long as there is no ambiguity. (B1 for one of the three faces drawn correctly **or** for an isometric drawing of any cuboid)
2	Different makes of car Tally Frequency	Make of car Tally Frequency	3	B1 for make of car or list of at least 3 different makes B1 for tally or tally marks B1 for frequency or totals
3		6 tessellating shapes	2	B2 for fully correct with 5 or more additional shapes, no gaps (B1 for 4 or more shapes tessellating, with at least one shape inverted, with or without the given shape, ignore extras)
4	24.90 ÷ 3 or 8.30 24.90 – '8.30' or 2 × 8.30	16.6(0)	3	M1 for 24.90 ÷ 3 or 8.30 M1 (dep) for 24.90 – “8.30” or 2 × “8.30” A1 for 16.60 or 16.6

Paper 5523_03

No	Working	Answer	Mark	Notes
5	315 × 24 24 × 315 1260 6300 7560 7200 7560 **3 1 5** 0 6 | 0 2 | 1 0 — **2** 1 2 | 0 4 | 2 0 — **4** 7 5 6 0 **300 10 5** 6000 | 200 | 100 — **20** 1200 | 40 | 20 — **4** 6000+200+100+1200+40+20 = 7560 **3 0.1 0.05** 60 | 2 | 1 — **20** 12 | 0.4 | 0.2 — **4** 60 + 2 + 1 + 12 + 0.4 + 0.2 = 75.6	75.6(0)	3	M1 for a complete method with relative place value correct. Condone 1 multiplication error, addition not necessary. **OR** M1 for a complete grid with not more than 1 multiplication error, addition not necessary. **OR** M1 for sight of a complete partitioning method, condone 1 multiplication error, final addition not necessary. A1 for 7560 or digits 756(0) A1 (dep on M1, but not previous A1) for correct placement of decimal point.

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
6	15 and 16 parts shaded Alternative 1 $\frac{3}{4} = 0.75$ or 75%, $\frac{4}{5} = 0.8$ or 80% Alternative 2 $\frac{3}{4} = \frac{15}{20}$, $\frac{4}{5} = \frac{16}{20}$	$\frac{4}{5}$ + reason	3	M1 for shading 15 parts for $\frac{3}{4}$ M1 for shading 16 parts for $\frac{4}{5}$ A1 (dep on M2) for selection of $\frac{4}{5}$ with correct shading Alternative 1 M1 for $\frac{3}{4} = 0.75$ or 75% M1 for $\frac{4}{5} = 0.8$ or 80% A1 (dep on M2) for selection of 0.8 or 80% or $\frac{4}{5}$ with correct decimals or percentages Alternative 2 M1 for $\frac{3}{4} = \frac{15}{20}$ oe M1 for $\frac{4}{5} = \frac{16}{20}$ oe A1 (dep on M2) for selection of $\frac{4}{5}$ or $\frac{16}{20}$ with equivalent fractions

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
7		$5 \times 5 \times 6$	150	4	M1 for attempt at 1 division (e.g. 40 ÷ 8), may be implied by marks or number on one edge of diagram or by 5 or 6 seen M1 for attempt at 3 divisions (40÷8, 40÷8, 60÷10), may be implied by marks or numbers on diagram or by 5,5 and 6 seen. M1 (dep on 1st M1) for "5" × "5" × "6" A1 cao Alternatively M1 for 40 × 40 × 60 **or** 8 × 8 × 10 **or** 96000 or 640 seen M1 for 40 × 40 × 60 **and** 8 × 8 × 10 or 96000 **and** 640 seen M1 (dep on 1st M1) for "(40 × 40 × 60)" ÷ "(8 × 8 × 10)" A1 cao SC:B1 for dividing area of one carton face by area of corresponding box face if M0
8	(a)		$\frac{7}{20}$	1	B1 for $\frac{7}{20}$ oe
	(b)	7 + 2 (or 20 – 11) are not lime flavour	$\frac{9}{20}$	1	B1 for $\frac{9}{20}$ oe
	(c)		0	1	B1 for 0, zero or nought ($\frac{0}{20}$ gets B0)
9	(a)		$80x$	1	B1 for $80x$ (accept $80 \times x$, $x80$, $x \times 80$) seen
	(b)		$95y$	1	B1 for $95y$ (accept $95 \times y$, $y95$, $y \times 95$) seen
	(c)		$80x + 95y$	2	M1ft for adding "$80x$" and "$95y$" (algebraic expressions only) A1 for $80x + 95y$

Paper 5523_03

No	Working	Answer	Mark	Notes
10 (a)		40	1	B1 cao
(b)		45	1	B1 for 42 to 48 (accept 3/4 hour)
(c)	40×2 or $\frac{40}{30}\times60$ or $40\div\frac{1}{2}$	80	2	M1 for 40×2 or $\frac{40}{30}$ or $40\div\frac{1}{2}$ A1 cao NB $\frac{40}{45}\times 60$ gets M0 A0
11 (a)	$3\times3-4\times2$ or $9-8$	1	2	M1 for substitution of 3 and 2 into expression or 9 and 8 seen A1 cao
(b)	$-7-3=-10$ $2\times-10=-20$ $-20\div4$	-5	3	M1 for substitution of 2 and –7 into p(q – 3) or sight of -20 or –14 – 6 M1 (dep) for "–20" ÷ 4 A1 cao SC: B1 for –10 seen if M0
12 (a)	6 | 8 9 7 7 | 8 5 9 6 3 8 | 1 3 1 7 1 9 | 0 1	6 | 7 8 9 7 | 3 5 6 8 9 8 | 1 1 1 3 7 9 | 0 1	3	M1 for unordered diagram (condone one error) A1 cao B1 for key (eg 6 | 7 = 67)
(b)(i)		Explanation	2	B1 for '(order numbers and) select middle value' oe
(ii)		79		B1 cao

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
13	(a)		Reflection in y-axis	1	B1 for triangle with vertices at (-1, 1) (-3, 1) and (-1,4)
	(b)		Rotation by half turn about (0, 0)	2	B2 for triangle with vertices (-1, -1) (-3, -1) and (-1, -4) (B1 for half turn not about (0,0))
	(c)		Enlargement Scale factor 3 Centre (0, 0)	3	B1 for 'enlargement' B1 for 'scale factor 3' or 3 seen B1 for 'centre (0,0)' B0 for any combination of transformations
14	(a)		4560	1	B1 cao
	(b)		45.6	1	B1 cao
	(c)		2.4	1	B1 cao
15	(a)	$4a-2a+5b+b$	$2a+6b$	2	B2 cao (B1 for $2a$ or $6b$ seen)
	(b)		$x(x-6)$	2	B2 cao (B1 for $x(ax+b)$ where a, b are numbers not equal to zero or $x-6$ seen on its own, or part of an expression)
	(c)		$3x-2x^3$	2	B2 cao (B1 for $3x$ or $2x^3$)
	(d)		$4x(3y+x)$	2	B2 cao (B1 for $2(6xy+2x^2)$ or $4(3xy+x^2)$ or $x(12y+4x)$ or $2x(6y+2x)$ or $4x(\quad)$)

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
16	(a)	$1 - (0.2 + 0.3 + 0.1)$	0.4	2	M1 for $1 - (0.2 + 0.3 + 0.1)$ A1 for 0.4 oe, accept $\frac{0.4}{1}$
	(b)	0.2×200	40	2	M1 for 0.2×200 A1 cao NB $\frac{40}{200}$ is M1 A0, 40 out of 200 is M1 A1
17	(a) (i)	$180 - 2 \times 25$	130	3	M1 for $180 - 2 \times 25$ A1 cao
	(ii)		Reason		B1 for mentioning isosceles and equal (or base) angles or equal sides and equal (or base) angles
	(b)	$180 - 95$	85	1	B1 cao
18	(a) (i)		7^5	3	B1 cao
	(ii)		7^4		B2 cao (B1 for sight of 7^5 or 7^{2+3} or 7×7^3 or $7^1 \times 7^3$ or $7^2 \times 7^2$ or 7^{2+3-1})
	(b)		$\frac{1}{2}$	1	B1 for $\frac{1}{2}$ or 0.5 or 2^{-1}
19	(a)		3×10^7	1	B1 cao
	(b)		0.002	1	B1 cao

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
20		Box plot	2	3 aspects: 1st aspect – vertical line for median 2nd aspect – box using correct quartiles 3rd aspect – whiskers (could be single line) drawn with correct end points B2 for fully correct box plot (B1 for 1 aspect)
21 (a)	e.g. 2 | 126 3 | 63 3 | 21 7	$2\times3\times3\times7$	2	M1 for a systematic method of at least 2 correct divisions by a prime number oe factor trees; can be implied by digits 2, 3, 3, 7 on answer line. A1 for $2\times3^2\times7$ or $2\times3\times3\times7$
(b)	$2\times3\times7$	42	2	B2 cao (B1 for 6, 14, 21 or $2\times3\times7$)
22	$\frac{8}{3}\times\frac{5}{4}=\frac{8\times5}{3\times4}=\frac{40}{12}$	$3\frac{1}{3}$	3	B1 for $\frac{8}{3}$ oe improper fraction or $\frac{5}{4}$ oe improper fraction M1 (dep on B1) for multiplying numerator and denominator of "$\frac{8}{3}$" and "$\frac{5}{4}$" A1 for $3\frac{1}{3}$ oe mixed number or $\frac{10}{3}$ OR B1 for 1.25 and 2.67 or 2.66(…) M1 (dep on B1) for correct method of multiplication A1 for $3.\dot{3}$

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
23			2	M1 for a relevant pair of intersecting arcs A1 for line drawn within guidelines, at least 3cm in length, accept broken line [SC: B1 for line drawn within guidelines if M0]
24 (a)		$-1, 0, 1, 2, 3$	2	B2 cao (-1 each error or omission)
(b)(i)		$x \geq \frac{7}{2}$	3	M1 for $2x \geq 7$, condone use of = sign or wrong equality A1 for $x \geq \frac{7}{2}$ oe as final answer
(ii)		4		SC:B1 for 3.5 or $\frac{7}{2}$ seen if M0 B1 ft from $x \geq$ "$\frac{7}{2}$"
25	$4x + 2y = 8$ $4x - 10y = 20$ $12y = -12$ $y = -1$ $4x + 2(-1) = 8$ $x = 2.5$	$x = 2.5$ $y = -1$	3	M1 for correct process to eliminate either x or y (condone one arithmetical error) M1 (dep) for substituting found value into either equation A1 for $x = 2.5$, $y = -1$ [SC: B1 for $x = 2.5$ or $y = -1$ if M0]

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
26		Interior angle of hexagon = 180 – (360 ÷ 6) = 120 360 – (90 + 120)	150	4	Alternative 1 M1 for 360 ÷ 6 A1 for 60 M1 (dep on M1) for "60" + 90 A1 cao Alternative 2 M1 for 360 ÷ 6 A1 for 60 M1 (dep on M1) for 360 – (2 × "60" + 90) A1 cao Alternative 3 M1 for (6 – 2) × 180 ÷ 6 A1 for 120 M1 (dep on M1) for 360 – (90 + "120") A1 cao
27	(a)		(16), 50, 82, 96, 100	1	B1 cao
	(b)	Cumulative freq. diag. curve/ segments	Cum. freq graph	2	B1 for 4 or 5 points plotted correctly ± 1 full (2mm) square depending on sensible table (condone 1 addition error) B1 (dep) for points joined by curve or line segments provided no gradient is negative - ignore any part of graph outside range of their points. (SC:B1 if 4 or 5 points plotted not at end but consistent within each interval and joined)
	(c)	100 – 42	58	2	M1 (ft dep on graph being cf) for reading from graph at 18 or 19, can be implied by answer in range 40 to 46 A1 for answer in range 56 to 60 or ft for 100 – '42' ±1 full (2mm) square

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
1	(a)	900 × 1.70 =	1530	2	M1 900 × 1.7(0) or digits 153(0) seen A1 cao
	(b)	160 ÷ 1.70 =	£94.12 or £94.11	2	M1 160 ÷ 1.7(0) or digits 941(....) seen A1 cao
2	(a)	4.7 ÷ 5.9 = 0.796610169	0.7966..	2	B2 for 0.7966 or better (B1 for 0.8, 0.80, 0.79, 0.796, 0.797 or digits 59 seen)
	(b)	0.82, 0.8, 0.85, 0.66, 0.875 0.66, 0.8, 0.82, 0.85, 0.875 2/3, 4/5, 0.82, 85%, 7/8	2/3, 4/5, 0.82, 85%, 7/8	2	B2 correct order (oe decimals in order) (B1 correct order reversed, or one error in ordered listing) with or without decimal equivalents. NB Accept 0.67 or 0.66
3		**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	3	B3 all 6 correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
4	(a)	–3 (–1) (1) 3 5 7	**–3**, 3, **5**, **7**	2	B2 all correct (B1 2,3 correct)
	(b)	(-2,-3),(-1,-1),(0,1),(1,3),(2,5),(3,7)	line	2	B2 cao for line from $x = -2$ to $x = 3$ (B1 plotting at least 5 points correctly or single line passing through (0, 1) or single line of gradient 2) The six possible points are: (-2,-3), (-1,-1), (0,1), (1,3), (2,5), (3,7)
	(c)(i)		–2	2	B1 $y = -2$ or ft from line segment
	(ii)		2.5		B1 $x = 2.5$ or ft from line segment

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
5	(a) (i)	180 – 54 (=126) "126" ÷ 2	63	3	M1 for (180–54) ÷ 2 A1 cao
	(ii)		Reason		B1 (indep) angles in triangle add to 180 OR equal angles in isosceles triangle OR equal angles and 2 sides the same (B0 if any incorrect reasoning given eg parallel, equilateral triangle)
	(b)	180 – "x"	117	1	B1 117 or ft 180 – "x" if $x < 90$
6		15/120×360 = 45 Plain 40/120×360 = 120 Cheese & Onion 55/120×360 = 165 Salt & Vinegar 10/120 × 360 = 30 Beef	45°, 120°, 165°, 30°	4	M1 evidence of method for at least one angle (could be implied by one correct angle of four on pie chart or in the table) A2 All four angles drawn ±2° tolerance, any order (A1 at least 2 angles correctly drawn ±2°, or all 4 angles in the table) B1 (dep on at least 1 angle drawn correctly, and exactly 4 sectors) for labels (flavour or frequency; initials will do) NB: Ignore the table if the pie chart provides the marks.
7	(a)	260 – 50 = 210 210 ÷ 35 =	6	3	M1 for 260–50 or 210 seen. M1 for "260-50" ÷ 35 or 210÷35 A1 cao
	(b)		P=35h + 50	3	B3 for P=35h+50 or P=35×h+50 oe (B2 for correct RHS or P=h + 50 ×35 or P=35h+k where k is numerical oe) (B1 for P = some other linear expression in h, OR h + 50 ×35 OR 35h seen) NB: P=h scores no marks; ignore £ signs. SC B2 for $h = \frac{P-50}{35}$
8	(a)		Elevation	2	B2 for 4 vertical squares. Accept 4 by 1 rectangle. (B1 for 4 vertical squares with one square added or one parallelogram added at the top, or 3 vertical squares, or 4 horizontal squares)

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
(b)		Plan	2	B2 for 2 adjacent squares, vertical or horizontal. Accept 2 by 1 rectangle. (B1 for 3 adjacent horizontal or vertical squares or a rectangle with sides in the ratio 2:1)
9 (a)		Points	1	B1 all three points ±1 full square.
(b)		Negative	1	B1 Negative. Ignore other descriptors unless contradictory.
(c)		lobf	1	B1 A single straight line drawn to cross between (5,30), (5,40) and (40,0), (40,15); accept freehand if considered to be straight.
(d)(i) (ii)		18–25 30–40	2	B1 18g–25g inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient. B1 30–40 min inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient.
10		300, 3, 75, 150	3	B3 for 4 correct (B2 for 2 or 3 correct) (B1 for 1 correct)
11 (a)	$6x - 7 + 7 = 38 + 7$ $6x = 45$	7.5	2	M1 $6x = 45$ or +7 both sides A1 7.5 oe; accept 45/6
(b)	$5y - 2 = 10$ or $20y - 8 = 40$ $5y = 12$ $20y = 48$	$2\frac{2}{5}$	3	M1 $20y–8(=40)$ or $\frac{4(5y-2)}{4} = \frac{40}{4}$ or $5y–2=10$ M1 (indep) for correct rearrangement into the form $ay=b+c$ or better (eg $20y=40+8$ or $5y=10+2$, using own terms) A1 for $2\frac{2}{5}$, 2.4 oe
12	5 miles = 8 km 70mph÷5×8 = 122 km/h OR 120km/h ÷ 8 × 8 = 75 mph Faster than 70 mph	70mph (Great Britain) (112 km)	3	M1 5 miles = 8 km; OR 70 mph is about 100 km/h OR 1km=0.6(25) miles OR 1mile=1.6km oe M1 70 ÷ 5 × 8 (=112) or 120 ÷ 8 × 5 (=75) oe A1 (dep on at least M1) GB or 70 mph Refer to both answer line and working. NB GB or 70 mph without working scores 0 marks.

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
13 (a)	$\pi \times 0.65$	2.04–2.05	2	M1 for $\pi \times 0.65$ or 3.14×0.65 or 3.142×0.65 oe A1 2.04–2.05 SC Award B1 for 2.0 seen (not 2)
(b)	1000 ÷ "(a)"	487–491	2	M1 for 1000 (or 100) ÷ "(a)" A1 for 487–491
14	$45.00 + 45.00 \times \frac{15}{100} =$ $45.00 + 6.75 =$	51.75	3	M2 for $45.00 + 45.00 \times \frac{15}{100}$ oe or 45.00×1.15 oe OR $45.00 + 6.75$ OR complete method or 5175 seen. (M1 for $45.00 \times \frac{15}{100}$ oe OR 6.75 seen OR 675 seen OR correct method for calculating 15% of 45) A1 cao SC Award B2 for an answer of 38.25
15		273	2	M1 for 728 ÷ 8 or 728 ÷ "3+5" or 91 A1 cao SC B1 for 455, or for 273:455
16	3 24 3.7 46.9(53) 4 60 3.8 51.0(72) 3.1 26.6(91) 3.9 55.4(19) 3.2 29.5(68) 3.21 29.8(66…) 3.3 32.6(37) 3.22 30.1(66…) 3.4 35.9(04) 3.23 30.4(68..) 3.5 39.3(75) 3.24 30.7(72...) 3.6 43.0(56) 3.25 31.0(78...) or 31	3.2	4	B2 for trial between 3.2 and 3.3 inclusive (B1 for trial between 3 and 4 inclusive) B1 for different trial between 3.21 and 3.25 inclusive B1 (dep on at least one previous B1) cao for 3.2 as final answer. NB: embedded answers: –B1; award Bs for evaluations rounded or truncated to at least 1 dp or for 31

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
17	$45^2 + 34^2 =$ $2025 + 1156 = 3181$ $\sqrt{3181} = 56.4$	56	4	M1 for $45^2 + 34^2$ M1 (dep) for $\sqrt{(2025 + 1156)}$ A1 for 56.4 … B1 for rounding their diagonal to the nearest integer (dep on evidence from decimal) NB 56 as the final answer gets full marks. NB Scale drawings result in 0 marks.
18	$2000 \times (1.055)^3$ Interest = 2348.48 – 2000 =	348.48	3	M1 for $5.5/100 \times 2000$ (oe) or 330 or 2330 or 110 or 2110 M1 (dep) for $5.5/100 \times (2000 +$ "110" + "116.05") or 122.4… seen A1 cao (accept only 348.48 or 348.49) **OR** M2 for $2000 \times (1.055)^3$ or 2348.48(…) or 2348.49 seen (M1 for $2000 \times (1.055)^n$, $n \neq 3$) A1 for 348.48 or 348.49 [SC: B2 for 2348.48 - 2348.49]
19		Line	2	B2 line fully within tramlines, crossing AB and CD (B1 a straight line which crosses AB within the tramline, and also crosses CD) NB: Accept dotted or dashed lines, but not curves; accept freehand if considered to be straight. SC B1 for a perpendicular bisector of AB that is at least half way from AB to CD within the tramlines
20 (a)		$90<t\leq100$	1	B1 for $90<t\leq100$; accept 90-100.
(b)	$65\times12=780$ $75\times22=1650$ $85\times23=1955$ $95\times24=2280$ $105\times19=1995$ $8660/100=$	86.6	4	M1 for use of fx with x consistent within intervals (including end points). Allow one slip. M1 (dep) for use of midpoints M1 (dep on 1st M1) for use of $\sum fx/100$ or $\sum fx/\sum f$ A1 86.6

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
21	(a)	$8 \times \frac{10}{4}$	20	2	M1 $\frac{10}{4}$ or $\frac{4}{10}$ or 0.4 or 2.5 oe seen A1 cao NB ratios get M0 unless of the form 1:*n* OR M1 $\frac{8}{4}, \frac{4}{8}$ oe seen A1 cao
	(b)	$15 \times \frac{4}{10}$	6	2	M1 $15 \times \frac{4}{10}$ oe A1 cao
22	(a)	$x^2 - 4x + 3x - 12 = x^2 - x - 12$	$x^2 - x - 12$	2	M1 for exactly 4 terms correct ignoring signs (eg x^2, 4*x*, 3*x*, 12) or 3 correct terms out of 4 terms with correct signs (eg 3 out of 4 of x^2, –4*x*, +3*x*,–12) A1 cao
	(b)	$(x+2)(x+5)$	$(x+2)(x+5)$	2	B2 cao (B1 for exactly one of $(x+2)$, $(x+5)$)
	(c)	$6=15+4q-20$ $6-15=4(q-5)$ $p-3t=4q-4t$ $6-3\times5=4(q-5)$	2 ¾	3	M1 for correct substitution of *p* and *t*. M1 for correct expansion of $4(q-t)$ oe (eg $4q-20$, $4q-4t$) A1 11/4 or 2 ¾ or 2.75 OR M1 for correct substitution of *p* and *t*. M1 for $\frac{p-3t}{4} = q - t$ oe A1 11/4 or 2 ¾ or 2.75

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
23	(a)	$9.9\times10^8 - 6.0\times10^7$	9.3×10^8	2	M1 for $99\times10^7 - 6\times10^7$ or $9.9\times10^8 - 0.6\times10^8$ or conversion of either to an ordinary number, or 930000000 or 93×10^7 or 9.3×10^n where n is any positive integer A1 cao
	(b)	$\frac{6.0-4.5}{4.5}\times100 = \frac{1.5}{4.5}\times100 =$ or $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}\times100$	33.3%	3	M2 $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}\times100$ oe (M1 for $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}$ or $\frac{6.0\times10^7 - 4.5\times10^7}{6.0\times10^7}\times100$ oe A1 cao OR M2 $\frac{6.0\times10^7}{4.5\times10^7}\times100$ -100 = (M1 $\frac{6.0\times10^7}{4.5\times10^7}\times100$ or 133.33(%)) A1 cao NB Accept any of the above expressions without any reference to 10^7.
24		LCM (40,24) = 120 Bread buns 120÷40 Burgers 120÷24 OR Bread buns: 40 is $2\times2\times2$ (×5) Burgers: 24 is $2\times2\times2$ (×3)	Bread buns 3 Burgers 5	3	M1 attempt to find LCM by eg lists of multiples, or summing of 40 and 24. A1 identify 120 (as LCM) A1 cao (both) OR M1 expansion of either number into its prime factors in a factor tree or 8×5 or 8×3 A1 both expansions correct A1 cao (both) SC B2 if answers given the wrong way around

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
25	8 × 41 = 328 2 × 29 = 58 328 – 58 = 270 270 ÷ 6 = 45	45	3	M1 for either 8 × 41 (=328) or 2 × 29 (=58) M1 (dep) "328" – "58" (=270) A1 cao NB 328 and/or 58 on the answer line gets M1 (implied); 270 on the answer line gets M2 (implied)

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
1 (a)	$1 - (0.2 + 0.3 + 0.1)$	0.4	2	M1 for $1 - (0.2 + 0.3 + 0.1)$ A1 for 0.4 oe, accept $\frac{0.4}{1}$
(b)	0.2×200	40	2	M1 for 0.2×200 A1 cao NB $\frac{40}{200}$ is M1 A0, 40 out of 200 is M1 A1
2	$650 - 430 = 220$ 1 choc ice costs 110p $650 - 5 \times 110 = 100$p	50	3	M1 for $650 - 430$ or 220 or 110 oe seen M1 for $650 - 5 \times \frac{'220'}{2}$ or $430 - 3 \times \frac{'220'}{2}$ oe A1 for 50p or £0.50 or £0.5 *Alternative scheme* $2x + 5y = 650$ $2x + 3y = 430$ oe M1 for subtracting two simultaneous equations to eliminate x (lollies)(2 or 3 terms correct) M1 for $650 - 5 \times 'y'$ or $430 - 3 \times 'y'$ oe A1 for 50p or £0.50 or £0.5 *Alternative scheme* M1 for $3\times (2x + 5y = 650)$ evaluated and $5 \times (2x + 3y = 430)$ evaluated oe (5 or 6 terms correct) M1 for subtraction of equations to eliminate y (choc ices)(2 or 3 terms correct ft) A1 for 50p or £0.50 or £0.5

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
3		question + response boxes oe	2	1st aspect: One question (eg 'how long does it take you to travel to school?' or 'What time did you leave home to get to school?'); ignore other questions. 2nd aspect: Response list (at least two), not overlapping. 3rd aspect: Some mention of units (eg minutes) in either question or responses B2 for all three aspects, or B1 for just one aspects.
4	2 [(3×1) + (4×1)] + (3×6) + (1×6) + (2×6) + (4×6) + (1×6) + (5×6)	110 cm^2	4	M1 for attempt to find the area of one face M1 for at least 6 faces with intention to add A1 cao B1 (indep) for cm^2 (with or without numerical answer)
5	$\left(\frac{-6+4}{2}\right), \left(\frac{5+3}{2}\right)$	(–1, 4)	2	B2 cao [B1 for (–1,*a*) or (b,4) or (4,-1)]
6		Box plot	2	3 aspects: 1st aspect – vertical line for median 2nd aspect – box using correct quartiles 3rd aspect – whiskers (could be single line) drawn with correct end points B2 for fully correct box plot (B1 for 1 aspect)
7			2	M1 for a relevant pair of intersecting arcs A1 for line drawn within guidelines, at least 3cm in length, accept broken line [SC: B1 for line drawn within guidelines if M0]

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
8	(a)	e.g. 2 | 126 3 | 63 3 | 21 7	$2\times3\times3\times7$	2	M1 for a systematic method of at least 2 correct divisions by a prime number oe factor trees; can be implied by digits 2, 3, 3, 7 on answer line. A1 for $2\times3^2\times7$ or $2\times3\times3\times7$
	(b)	$2\times3\times7$	42	2	B2 cao (B1 for 6, 14, 21 or $2\times3\times7$)
9	(a)		$-1,0,1,2,3$	2	B2 cao (-1 each error or omission)
	(b)(i)		$x\geq\frac{7}{2}$	3	M1 for $2x\geq7$, condone use of = sign or wrong equality A1 for $x\geq\frac{7}{2}$ oe as final answer
	(ii)		4		SC:B1 for 3.5 or $\frac{7}{2}$ seen if M0 B1 ft from $x\geq$ "$\frac{7}{2}$" or $x>$"$\frac{7}{2}$"
10	(a)(i)		7^5	3	B1 cao
	(ii)		7^4		B2 cao (B1 for sight of 7^5 or 7^{2+3} or 7×7^3 or $7^1\times7^3$ or $7^2\times7^2$ or 7^{2+3-1})
	(b)		$\frac{1}{2}$	1	B1 for $\frac{1}{2}$ or 0.5 or 2^{-1}
11	(a)	$5n=m+21$	$n=\frac{m+21}{5}$	2	M1 for $5n=m+21$ or for attempt to divide three terms by 5 A1 $n=\frac{m+21}{5}$ oe

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
(b)	$4p-8q=3p+2$ $p-8q=2$ $p=8q+2$	$p=8q+2$	3	M1 for $4p-aq$ or $\frac{3}{4}p+b$ where a is an integer and b is a number M1 (dep) for taking one term correctly to LHS or RHS of expression A1 $p=8q+2$ oe
12		$y=\frac{1}{2}x+3$	2	B2 for $y = ½x + 3$ oe (B1 for $y = ½x + c$, $c \neq 7$ or $y = mx + 3$ oe or $½x + 3$ or $M = ½x + 3$)
13	$\frac{8}{3}\times\frac{5}{4}=\frac{8\times5}{3\times4}=\frac{40}{12}$	$3\frac{1}{3}$	3	B1 for $\frac{8}{3}$ oe or $\frac{5}{4}$ oe M1 (dep on B1) for multiplying numerator and denominator of "$\frac{8}{3}$" and "$\frac{5}{4}$" A1 for $3\frac{1}{3}$ oe mixed number or $\frac{10}{3}$ *OR* B1 for 1.25 and 2.67 or 2.66(…) M1(dep on B1) for correct method of multiplication A1 for $3.\dot{3}$
14	$4x+2y=8$ $4x-10y=20$ $12y=-12$ $y=-1$ $4x+2(-1)=8$ $x=2.5$	$x=2.5$ $y=-1$	3	M1 for correct process to eliminate either x or y (condone one arithmetical error) M1 (dep) for substituting found value into either equation A1 for $x = 2.5$, $y = -1$ [SC: B1 for $x = 2.5$ or $y = -1$ if M0]

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
15		Interior angle of hexagon = 180 – (360 ÷ 6) = 120 360 – (90 + 120)	150	4	Alternative 1 M1 for 360 ÷ 6 A1 for 60 M1 (dep on M1) for "60" + 90 A1 cao Alternative 2 M1 for 360 ÷ 6 A1 for 60 M1(dep on M1) for 360 – (2 × "60" + 90) A1 cao Alternative 3 M1 for (6 – 2) × 180 ÷ 6 A1 for 120 M1(dep on M1) for 360 – (90 + "120") A1 cao
16	(a)		(16), 50, 82, 96, 100	1	B1 cao
	(b)	Cumulative freq. diag. curve/ segments	Cum. freq graph	2	B1 ft for 4 or 5 points plotted correctly ± 1 full (2mm) square depending on sensible table (condone 1 addition error) B1 (dep) for points joined by curve or line segments provided no gradient is negative - ignore any part of graph outside range of their points. (SC:B1 if 4 or 5 points plotted not at end but consistent within each interval and joined)

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
(c)	$100 - 42$	58	2	M1 (ft dep on graph being cf) for reading from graph at 18 or 19, can be implied by answer in range 40 to 46 A1 for answer in range 56 to 60 or ft for 100 –'42' ±1 full (2mm) square
17 (a)		0.6 and 0.7,0.3,0.7	2	B1 for 0.6 on LH branch B1 for 0.7, 0.3 and 0.7 on RH branches
(b)	0.4×0.3	0.12	2	M1 for 0.4×0.3 A1 0.12 oe
(c)	$0.4\times0.7+0.6\times0.3$	0.46	3	M1 for '0.4×0.7' or '0.6×0.3' M1 for addition of two products from correct branches A1 0.46 oe Alternative M2 for an attempt to evaluate 1-(0.3×0.4 +'0.6×0.7') A1 cao
18	$x = 0.4545...$ so $100x = 45.4545...$ $99x = 45$ $x = \frac{45}{99} = \frac{15}{33}$	proof	3	M1 for $100x = 45.45\ldots$ or $10000x = 4545.45\ldots$ M1 (dep) for subtraction of both sides A1 for $\frac{15}{33}$ from correct proof
19	$\left(\sqrt{3}\right)^2 + \sqrt{3}\sqrt{2} - \sqrt{3}\sqrt{2} - \left(\sqrt{2}\right)^2$ $= 3 - 2$	1	2	B2 cao (B1 for $\sqrt{3}\sqrt{3} + \sqrt{3}\sqrt{2} - \sqrt{3}\sqrt{2} - \sqrt{2}\sqrt{2}$ oe, $\sqrt{3}\sqrt{3} - \sqrt{2}\sqrt{2}$ oe, or for 2,3, $\sqrt{4}$ $\sqrt{6}$ $\sqrt{9}$ seen)
20 (a)	$\frac{96}{24}$ or 4 $\sqrt{4}$ or 2	8	3	M1 for $\frac{96}{24}$ or $\frac{24}{96}$ or 4 or $\frac{1}{4}$ oe M1 for $\sqrt{\frac{96}{24}}$ or $\sqrt{\frac{24}{96}}$ or $\sqrt{'4'}$ or $\frac{1}{\sqrt{'4'}}$ or 2 or $\frac{1}{2}$ oe A1 cao

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
	(b)	12×2^3	96	2	M1 for '2'^{3} or 8 A1 cao
21	(a)	$x\times3-x\times2x^2$	$3x-2x^3$	2	B2 cao (B1 for a two term expression with either $3x$ or $2x^3$)
	(b)		$4x(3y+x)$	2	M1 for taking out a factor of x, $2x$, 2, 4 or $4x$ A1 cao
	(c)	$\frac{5a}{b^2}$	$\frac{5a}{b^2}$	2	B2 for $\frac{5a}{b^2}$ or $5ab^{-2}$ (accept $\frac{5a}{1b^2}$) (B1 for either dealing with the numbers or dealing with the powers of a)
	(d)	$\frac{\cancel{x-3}}{\cancel{(x-3)}(x+3)}$	$\frac{1}{x+3}$	2	M1 for $(x-3)(x+3)$ A1 cao
22	(a)		$\mathbf{a}+\mathbf{b}$	1	B1 $\mathbf{a}+\mathbf{b}$ oe
	(b)	$ED=\mathbf{a}$ $DX=-2\mathbf{b}+2\ AC=2\mathbf{a}$ (So, $DX=2ED$)	$2\mathbf{a}$	3	M1 for $(DX=)\ DA+AX$ A1 for $(DX=)\ -2\boldsymbol{b}+2$ ("$\boldsymbol{a}+\boldsymbol{b}$") A1 $2\boldsymbol{a}$ from fully correct proof
23		$\pi(2x)^2h=\frac{4}{3}\pi(3x)^3$ $h=\frac{\frac{4}{3}\pi(3x)^3}{\pi(2x)^2}$	$9x$	3	M1 for $\pi(2x)^2h=\frac{4}{3}\pi(3x)^3$ (condone absence of brackets) M1 (dep) for valid algebra that gets to $h=ax$ (condone one error in powers of numerical constants) A1 cao

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
24	(i)	$n^2+(n+1)^2=2(n^2+n)+1$	$2n^2+2n+1$	4	M1 for at least 3 terms correct from $n^2+n+n+1$ A1 for $2n^2+2n+1$ oe
	(ii)		$2(n^2+n)$ is always even so $2(n^2+n)+1$ is always odd		M1 for recognizing $2n^2$ is always even A1ft complete proof for their quadratic *Alternative method* M1 for recognizing that if n^2 is odd then $(n+1)^2$ is even or vice versa A1 for complete proof
25			42 318	2	B1 for answer in range 36 – 48 B1 for answer in range 312 – 324
26	(a)	$(x-3)^2-3^2+15$	$p=3, q=6$	2	B2 for $p=3$ and $q=6$ (B1 for $p=3$ OR $q=6$) SC: award B2 for $(x-3)^2+6$ if p and q are not identified
	(b)		Sketch	2	B1 for U shaped curve B1 ft for TP in first quadrant (ft if TP not in first quadrant)
27	(a)	Graph translated 2 units upwards through points (–4, 2), (–2, 4), (0,2) and (3,5)	Sketch	2	M1 for a vertical translation A1 curve through points (–4, 2), (–2, 4), (0,2) and (3,5) ± ½ square
	(b)	Graph reflected in x-axis through points (–4,0), (–2, –2), (0, 0) and (3, -3)	Sketch	2	M1 for reflection in x-axis or y-axis A1 curve through points (–4,0), (–2, –2), (0, 0) and (3, -3) ± ½ square

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
1		273	2	M1 for 728 ÷ 8 or 728 ÷ "3+5" or 91 A1 cao SC B1 for 455 or 273:455
2		$3n - 1$	2	B2 for $3n - 1$ oe (B1 for $3n + k$ where $k \neq -1$ but k could be 0)
3	3 24 3.7 46.9(53) 4 60 3.8 51.0(72) 3.1 26.6(91) 3.9 55.4(19) 3.2 29.5(68) 3.21 29.8(66…) 3.3 32.6(37) 3.22 30.1(66…) 3.4 35.9(04) 3.23 30.4(68..) 3.5 39.3(75) 3.24 30.7(72...) 3.6 43.0(56) 3.25 31.0(78...) or 31	3.2	4	B2 for trial between 3.2 and 3.3 inclusive (B1 for trial between 3 and 4 inclusive) B1 for different trial between 3.21 and 3.25 inclusive B1 (dep on at least one previous B1) cao for 3.2 NB: embedded answers: –B1; award Bs for evaluations rounded or truncated to at least 1 dp or for 31
4	2 is the only even prime number and the product of 2 odd numbers is odd	Yes	2	B2 for 'yes' and '2 is the only even prime number and the product of two odd numbers is odd' oe (B1 for 'yes' and either '2 is the only even prime number' oe **or** 'the product of two odd numbers is odd' oe)
5	$45^2 + 34^2 =$ $2025 + 1156 = 3181$ $\sqrt{3181} = 56.4$	56	4	M1 for $45^2 + 34^2$ M1 for $\sqrt{(2025 + 1156)}$ A1 for 56.4 … B1 for rounding their diagonal to the nearest integer (dep on evidence from decimal) NB 56 with no incorrect working as the final answer gets full marks. NB Scale drawings result in 0 marks.

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
6	2000 × $(1.055)^3$ Interest = 2348.48 – 2000 =	£348.48	3	M1 for 5.5/100 × 2000 (oe) or 330 or 16.5/100×2000 or 2330 or 110 or 2110 M1 (dep) for 5.5/100 × (2000 + "110" + "116.05") or 122.4… seen A1 cao (accept only 348.48 or 348.49) **OR** M2 for 2000 × $(1.055)^3$ or 2348.48(…) or 2348.49 seen (M1 for 2000 × $(1.055)^n$, $n \neq 3$) A1 for 348.48 or 348.49 [SC: B2 for 2348.48 - 2348.49]
7 (a)		Line	2	B2 line fully within tramlines, crossing AB and CD (B1 a straight line which crosses AB within the tramline, and also crosses CD) NB: Accept dotted or dashed lines, but not curves; accept freehand if considered to be straight. SC B1 for the perpendicular bisector of *AB* reaching halfway or more from *AB*
(b)		Region		B2 correct arc ±2mm and shaded within. Allow dotty or continous arc. (B1 inaccurate arc and shaded or accurate arc unshaded)
8	65×12=780 75×22=1650 85×23=1955 95×24=2280 105×19=1995 8660/100=	86.6	4	M1 for use of fx with x consistent within intervals (including end points). Allow one slip even if outside interval M1 (dep) for use of midpoints M1 (dep on 1st M1) for use of $\sum fx/100$ or $\sum fx/\sum f$ A1 86.6

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
9		Distance ÷ time: 1400 ÷ 2 h 20 min 20 mins is $\frac{1}{3}$ hour 1400 × 3 ÷ 7 =	600 kph	3	B1 20 mins as $\frac{1}{3}$ hour or as 0.33……hour M1 for distance ÷ time eg 1400 ÷ "2h 20 min" A1 cao OR B1 2 hour 20min = 140 (min) M1 Speed = $\frac{1400}{140}$ = (10 km per minute) A1 cao
10	(a)	$(x+18)+2x+(2x+7)=180$	Equation	2	B2 for $(x+18)+2x+(2x+7)=180$ oe (B1 for $(x+18)+2x+(2x+7)$)
	(b)	$5x+25 = 180$ $5x = 155$	31	2	M1 for simplifying to at least $5x+25=180$ or 360 (may be earned in (a)) A1 for $x=31$
11	(a)	(–14) –4 (0) 4 14	–4, 4, 14	2	B2 for all 3 values correct (B1 for 1 or 2 values correct)
	(b)		curve	2	B1 for all 5 points plotted correctly ± ½ square (ft from table if at least B1 awarded in (a)) B1 (indep) ft for any smooth curve through their points

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
12	(a)	$8 \times \frac{10}{4}$	20	2	M1 $\frac{10}{4}$ or $\frac{4}{10}$ or 0.4 or 2.5 oe seen A1 cao NB ratios get M0 unless of the form 1:*n* OR M1 $\frac{8}{4}, \frac{4}{8}$ oe seen A1 cao
	(b)	$15 \times \frac{4}{10}$	6	2	M1 $15 \times \frac{4}{10}$ oe A1 cao
13	(a)	$9.9\times10^8 - 6.0\times10^7$	$\pm 9.3\times10^8$	2	M1 for $99\times10^7 - 6.0\times10^7$ or $9.9\times10^8 - 0.60\times10^8$ or correct conversion of either to an ordinary number or 930000000 or 9.3×10^n where *n* is any positive integer A1 cao
	(b)	$\frac{6.0-4.5}{4.5} \times 100 = \frac{1.5}{4.5} \times 100 =$ or $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7} \times 100$	33.3%	3	M2 $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7} \times 100$ oe (M1 for $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}$ or $\frac{6.0\times10^7 - 4.5\times10^7}{6.0\times10^7} \times 100$ oe A1 33.3 – 33.4 OR M2 $\frac{6.0\times10^7}{4.5\times10^7} \times 100$ -100 or 33.3% (M1 $\frac{6.0\times10^7}{4.5\times10^7} \times 100$ or 133.3%) A1 33.3-33.4 NB Accept any of the above expressions without any reference to 10^7.

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
14	LCM (40,24) = 120 Bread buns 120÷40 Burgers 120÷24 OR Bread buns: 40 is 2 × 2 × 2 (×5) Burgers: 24 is 2 × 2 × 2 (×3)	Bread Buns 3 Burgers 5	3	M1 attempt to find LCM by eg lists of multiples, or summing of 40s and summing of 24s, with at least 3 numbers in each list A1 identify 120 as LCM A1 cao (both) **OR** M1 expansion of either number into its prime factors in a factor tree or 8×5 or 8×3 A1 both expansions correct A1 cao (both) SC B2 if answers given the wrong way around
15	$\sin 32 = \frac{AB}{12}$ $AB = 12 \times \sin 32$ $AB = 6.35903\ldots$	6.36	3	M1 $\sin 32 = \frac{AB}{12}$ (accept Sin $\frac{AB}{12}$) M1 $12 \times \sin 32$ or $12 \times 0.5299..$ A1 accept 6.359 – 6.360 SC Gradians 5.78(1…) Radians 6.62 Get M1M1A0 **OR** Use of Sine Rule $\frac{\sin 32}{AB} = \frac{\sin 90}{12}$ or $\frac{AB}{\sin 32} = \frac{12}{\sin 90}$ M1 $AB = \frac{12 \times \sin 32}{\sin 90}$ M1 AB = 6.359 – 6.36 A1 SC Gradians 5.85(…) Radians 7. 40(…) M1M1A0

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
16		$8 \times 41 = 328$ $2 \times 29 = 58$ $328 - 58 = 270$ $270 \div 6 = 45$	45	3	M1 for either 8×41 (=328) or 2×29 (=58) M1 (dep) "328" – "58" (=270) A1 cao NB 328 and /or 58 on the answer line gets M1 (implied); 270 on the answer line gets M2 (implied)
17	(a)	$x^2 - 4x + 3x - 12 = x^2 - x - 12$	$x^2 - x - 12$	2	M1 for exactly 4 terms correct ignoring signs (x^2, $4x$, $3x$, 12) or 3 out of 4 terms with correct signs (x^2, $-4x$, $+3x$, -12) A1 cao
	(b)	$6x^2 - 8x + 15x - 20 = 6x^2 + 7x - 20$	$6x^2 + 7x - 20$	2	M1 for exactly 4 terms correct ignoring signs ($6x^2$, $8x$, $15x$, 20) or 3 out of 4 terms with correct signs ($6x^2$, $-8x$, $+15x$, -20) A1 cao
	(c)	$(x+2)(x+5)$	$(x+2)(x+5)$	2	B2 cao (B1 for exactly one of $(x + 2)$, $(x + 5)$)
	(d)		$12p^8q^3$	2	B2 cao (B1 for any 2 out of 3 terms correct in a product or 3 terms correct in a sum or part product)
	(e)	$6 = 15 + 4(q - 5)$ $6 = 15 + 4q - 20$ $11 = 4q$	$2\frac{3}{4}$	3	M1 for correct substitution of p and t. M1 for correct expansion of $4(q\text{-}t)$ oe (eg $4q$-20, $4q$-$4t$) A1 11/4 or 2 ¾ or 2.75 OR M1 for correct substitution of p and t. M1 for $\frac{p-3t}{4} = q - t$ oe A1 11/4 or 2 ¾ or 2.75
18	(a)	$T=kx$; $150=6k$; $k=25$	$T=25x$	3	M1 for $T=kx$, k algebraic M1 subs $T=150$ and $x=6$ into $T = kx^n$ $(n \neq 0)$ A1 for $T=25x$ oe SC B1 $T \propto 25x$ oe

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
	(b)	$T = 25 \times 15 =$	375	1	B1 ft on k , $k \neq 1$
	(c)	$600 = 25x$; $x = 600 \div 25 =$	24	1	B1 ft on k , $k \neq 1$
19		$\frac{120}{570} \times 50 = 10.5263$ $\frac{250}{570} \times 50 = 21.9298$ $\frac{200}{570} \times 50 = 17.5438$	10 22 18	3	M1 method shown eg $\frac{120}{120+250+200} \times 50$ or one of 10.5(263) 21.9(298), 17.5(438) **OR** 570 ÷ 50 = 11.4 and one of 120 ÷ 11.4 =10.526…) , 270 ÷ 11.4 (=21.9298…) , 200 ÷ 11.4 (=17.5438…) A1 for 10.5(263), 21.9(298), 17.5(438) or 11, 22, 18 B1 correction to add to 50: 10, 22, 18 or 11, 22, 17
20		$x(x+1) + 4(2x–3) = (2x–3)(x+1)$ $x^2 + x + 8x – 12 = 2x^2 + 2x – 3x – 3$ $x^2 – 10x + 9 = 0$ $(x – 9)(x – 1) = 0$	$x = 1, 9$	5	M1 for multiplying through by common denominator $(2x–3)(x+1)$ M1 (dep)for either $x^2 + x + 8x – 12$ or $2x^2 + 2x – 3x – 3$ oe A1 for correct quadratic (= 0) M1 for a correct method to solve 3 term quadratic A1 cao for both solutions
21		$AB^2 = 8^2 + 9^2 – 2\times8\times9\times\cos40$ $AB^2 = 64 + 81 – 144\times\cos40$ $AB^2 = 145 – 144 \times 0.766$ $AB^2 = 145 – 110.31… = 34.6896$ $AB = \sqrt{34.6796} = 5.8897877$	5.89	3	M1 Subs in Cos Rule: $8^2 + 9^2 – 2\times8\times9\times\cos40$ M1 correct order of evaluation of $8^2 + 9^2 – 2\times8\times9\times\cos40$ A1 cao 5.88 – 5.89 SC: Award B2 for one of $AB^2 = 241.03…$ or $AB = 15.525…$ (radians) $AB^2 = 28.50…$ or $AB = 5.33…$ (gradians)
22	(a)		24 30	2	B1 24 cao B1 30 cao
	(b)	Column $60 \leq w < 75$ at 10 Column $75 \leq w < 95$ at 4	10 4	2	B1 10 and correct width (tol $\pm\frac{1}{2}$ small square) B1 4 and correct width (tol $\pm\frac{1}{2}$ small square)

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
23 (a)		100.5	1	B1 cao
(b)		10.515	1	B1 cao
(c)	$\frac{100.5}{10.515}$	9.5577746	2	M1 for greatest distance divided by least time Where 100 < greatest distance ≤ 100.5, 10.51 ≤ least time < 10.52 A1 for 9.555 – 9.56
(d)	$\frac{99.5}{10.525}$	9.45368..	2	M1 for least distance divided by greatest time Where 99.5 ≤ least distance < 100, 10.52 < greatest time ≤10.53 A1 for 9.45 – 9.455
24	$\frac{4}{7}\times\frac{4}{7}\times\frac{3}{7} + \frac{3}{7}\times\frac{3}{7}\times\frac{4}{7} =$ $\frac{48+36}{343} = \frac{84}{343}$ But there are three ways this can be achieved: BBG, BGB, GBB So the probability is $\frac{84}{343}\times 3$ OR $1-\frac{64}{343}-\frac{27}{343}$ NB: 84/343 = 0.244897; 252/343 = 0.73469 $\frac{4}{7} = 0.57(142...)$, $\frac{3}{7} = 0.42(857...)$	$\frac{252}{343}$	3	M1 for $\frac{4}{7}\times\frac{4}{7}\times\frac{3}{7}$ or $\frac{3}{7}\times\frac{3}{7}\times\frac{4}{7}$ oe or $\left(\frac{4}{7}\right)^3$ oe or $\left(\frac{3}{7}\right)^3$ oe or $\frac{91}{343}$ or 0.10(49…) or 0.13(99….) M1 (indep) for identification of all 6 outcomes (M2 for $1-\left[\left(\frac{4}{7}\right)^3+\left(\frac{3}{7}\right)^3\right]$) oe A1 $\frac{252}{343}$, $\frac{36}{49}$, 0.73(469…) oe

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
25	(a)	$\frac{1}{3} \times \pi \times 5^2 \times 8 = \pi \times 25 \times 8 \div 3 =$ 209.4395	209–210	2	M1 for $\frac{1}{3} \times \pi \times 5^2 \times 8$ A1 for 209–210
	(b)	Base radius = $\frac{216}{360} \times 15 = 9$ Height = $\sqrt{(15^2 - 9^2)} =$	12	4	M1 for 216 ÷ 360 A1 for 9 M1 for $\sqrt{(15^2 - \text{“9”}^2)}$, where “9” < 15 A1 cao

PMT

edexcel

Mark Scheme (Results)

Summer 2012

GCSE Mathematics (Linear) 1MA0
Foundation (Non-Calculator) Paper 1F

PMT

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at www.edexcel.com.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2012
Publications Code UG032619
All the material in this publication is copyright
© Pearson Education Ltd 2012

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labeling conventions.

ii) *select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect canceling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

12 Parts of questions
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers
Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

Guidance on the use of codes within this mark scheme
M1 – method mark A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
1	(a)		380	1	B1 cao
	(b)		6.2	1	B1 cao
	(c)		Arrow at 34	1	B1 cao
2	(a)		8	1	B1 for 8 ± 0.2
	(b)		35	1	B1 for 35 ± 2°
	(c)		Circle drawn	1	B1 for all parts within ± 2mm, (use overlay)
3	(a)		4, 7, 4, 3, 2	2	M1 for at least 3 correct tallies **or** at least 3 correct frequencies A1 for all frequencies correct
	(b)		7	1	B1 for 7 **or** ft from frequencies in (a) or tallies if no frequencies
	(c)		Diagram drawn	3	M1 for bar chart or other suitable chart with at least 3 correct heights for their scale (can f.t.) A1 for all 5 bars correctly labelled and vertical axis correctly scaled A1 for fully correct **or** ft frequencies in (a) **OR** M1 for pictogram with at least 3 correct rows (can f.t.) A1 for correct labels on all 5 rows and correctly key A1 for fully correct **or** ft frequencies in (a) **OR** M1 for pie chart with at least 3 correct sectors ± 2° (can f.t.) A1 for all 5 sectors correctly labelled A1 for fully correct **or** ft frequencies in (a)

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
4		£1.18 + 94p = £2.12 £5 – £2.12 – 30p = £2.58 £2.58 ÷ 2 =	1.29	3	M2 for (5 – 1.18 – 0.94 – 0.30) ÷ 2 oe **or** digits 129 (M1 for 1.18 + 0.94 **or** 2.12 seen **or** 1.18 + 0.94 + 0.30 oe **or** 2.42 seen **or** 5 – 1.18 – 0.94 oe **or** 2.88 seen **or** (5 – 1.18 – 0.94) ÷ 2 **or** 1.44 seen **or** 5 – 1.18 – 0.94 – 0.30 oe **or** 2.58 seen) A1 cao **NOTE**: Accept working in £ or pence
5	(a)(i)		(2, 3)	2	B1 cao
	(ii)		(–3, 1)		B1 cao
	(b)		Point plotted at (3, –4)	1	B1 cao
6	(a)		– 5	1	B1 cao
	(b)		6	1	B1 for 6 or – 6
	(c)		3	1	B1 cao
7			(P, B), (P, S), (P, L) (M, B), (M, S), (M, L) (H, B), (H, S), (H, L)	2	M1 for any 3 combinations with no incorrect combinations A1 for all 9 combinations with no duplicates or extras
8	(a)		Walk	1	B1 cao
	(b)	24 ÷ 4 =	6	2	M1 for 24 ÷ 4 oe or ¼ oe seen A1 cao

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
9	(a)		Isosceles triangle	1	B1 for isosceles triangle
	(b)		Rectangle with area 12 cm^2	2	M1 for rectangle drawn A1 cao
10	(a)		A marked at 0	1	B1 for A marked at 0 (within overlay)
	(b)		B marked at 1/4	1	B1 for B marked at 1/4 (within overlay)
11	(a)		9	1	B1 cao
	(b)		33	2	M1 for 5 × 5 **or** 25 seen in the working **or** 2 × 2 × 2 **or** 8 seen in the working A1 cao
12	(a)		20	2	M1 3 × 3 × 3 oe seen or drawn or 27 seen or use of 3 layers A1 cao
	(b)			2	B2 for correct view (B1 for  or )
13	(a)(i)		07 29	2	B1 for 07 29
	(ii)		36		B1 for 36 or ft difference between (i) and 06 53
	(b)		07 51	1	B1 cao
	(c)		09 55	1	B1 for 09 55 or 9 55 or five to ten

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
14		2 + 8 + 2 + 8 = 20 20 ÷ 4 =	5	4	M2 for 2 + 8 + 2 + 8 oe **or** 20 seen **or** (2 + 8) ÷ 2 oe (M1 for the sum of 3 sides of the rectangle) M1 (dep) for the sum of 3 or 4 sides of the rectangle ÷ 4 **or** an attempt to evaluate (2 + 8) ÷ 2 oe to get the length of one side A1 cao **SC**: B1 for an answer of 4 coming from $\sqrt{2\times 8}$ oe
15	(a)		4	1	B1 cao
	(b)	9.5 – 4.75 = **OR** 9.5 ÷ 2 =	4.75	2	M1 for 9.5 – 4.75 **or** 9.5 ÷ 2 **or** 4.75 – 9.5 A1 cao
	(c)		6	1	B1 cao
	(d)	12 × 4 =	48	2	M1 for × 4 seen **or** identifying + 0.5 for every 2 inches **or** 12 + 12 + 12 + 12 oe **or** build up method eg 12, 24, 36, 48 allow one error A1 cao
16	(a)		trapezium	1	B1 for trapezium or isosceles trapezium
	(b)			2	B2 for correct tessellation (at least 5 more shapes) (B1 for at least 4 shapes (including initial shape) correctly tessellating)

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
17*		S: 35 ÷ 100 × 40 = 14 W: 40 ÷ 8 × 3 = 15 **OR** D: 16 ÷ 40 (× 100) = 0.4 (40%) W: 3 ÷ 8 (× 100) = 0.375 (37.5%) **OR** D: $\frac{16}{40} = \frac{80}{200}$ S: $\frac{35}{100} = \frac{70}{200}$ W: $\frac{3}{8} = \frac{75}{200}$	Debbie and correct calculations	4	**Compares Marks out of 40 or fractions with denominator of 40** M1 for 35 ÷ 100 × 40 oe **or** 14 seen (or 14/40 seen) M1 for 40 ÷ 8 × 3 **or** 15 seen (or 15/40 seen) A1 for 14 and 15 **or** $\frac{14}{40}$ and $\frac{15}{40}$ C1 (dep on M1) for correct conclusion for their working QWC with 3 comparable marks: Decision and justification should be clear with working clearly presented and attributable. **OR Decimals (or Percentages)** M1 for 16 ÷ 40 (× 100) oe **or** 0.4 (or 40) seen M1 for 3 ÷ 8 (× 100) oe **or** 0.375 (or 37.5) seen A1 for 0.4 and 0.375 (**or** 40 and 37.5) C1 (dep on M1) for correct conclusion for their working QWC: with 3 comparable decimals (or percentages: Decision and justification should be clear with working clearly presented and attributable. **OR Compares Fractions with denominator other than 40** M1 for attempt to convert all to fractions with a common denominator other than 40 M1 for at least 1 correct A1 for $\frac{80}{200}$ and $\frac{70}{200}$ and $\frac{75}{200}$ oe C1 (dep on M1) for correct conclusion for their working QWC with 3 comparable fractions: Decision and justification should be clear with working clearly presented and attributable.

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
18*	(a)		10	1	B1 cao
	(b)	 **Miles** 0 10 20 30 40 50 **Ed** 0 15 30 45 60 75 **Bill** 10 20 30 40 50 60	Ed is cheaper up to 20 miles, Bill is cheaper for more than 20 miles	3	M1 for correct line for Ed intersecting at (20,30) ±1 sq tolerance **or** $10 + x = 1.5x$ oe C2 (dep on M1) for a correct full statement ft from graph eg. Ed cheaper up to 20 miles **and** Bill cheaper for more than 20 miles (C1 (dep on M1) for a correct conclusion ft from graph eg. cheaper at 10 miles with Ed ; eg. cheaper at 50 miles with Bill eg. same cost at 20 miles; eg for £5 go further with Bill **or** A general statement covering short **and** long distances eg. Ed is cheaper for shorter distances **and** Bill is cheaper for long distances) **OR** (continued on next page)

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
18 (contd)					**OR** M1 for correct method to work out Ed's delivery cost for at least 2 values of n miles where $0 < n \leq 50$ **or** for correct method to work out Ed and Bill's delivery cost for n miles where $0 < n \leq 50$ C2 (dep on M1) for 20 miles linked with £30 for Ed and Bill with correct full statement eg. Ed cheaper up to 20 miles **and** Bill cheaper for more than 20 miles (C1 (dep on M1) for a correct conclusion eg. cheaper at 10 miles with Ed; eg. cheaper at 50 miles with Bill eg. same cost at 20 miles; eg for £5 go further with Bill **or** A general statement covering short **and** long distances eg. Ed is cheaper for shorter distances **and** Bill is cheaper for long distances) SC: B1 for correct full statement seen with no working eg. Ed cheaper up to 20 miles **and** Bill cheaper for more than 20 miles QWC Decision and justification should be clear with working clearly presented and attributable

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
19		1,96 × 2.25 = 4.41 **OR** 4.23 ÷ 9 = 0.47 1.96 ÷ 4 = 0.49 **OR** 4.23 × 4 = 16.92 1.96 × 9 = 17.64 **OR** 4.23 ÷ 9 = 0.47 0.47 × 4 = 1.88 **OR** 1.96 ÷ 4 = 0.49 0.49 × 9 = 4.41 **OR** 9 ÷ 4.23 = 2.12 4 ÷ 1.96 = 2.04	Pack of 9	3	M2 for a fully correct method to enable a conclusion eg 1.96 × 2¼ **OR** M1 for 4.23 ÷ 9 or 423 ÷ 9 **or** 0.47 seen **or** 47 seen M1 for 1.96 ÷ 4 or 196 ÷ 4 or 0.49 seen or 49 seen **OR** M1 for 4.23 × 4 **or** 423 × 4 **or** 16.92 seen **or** 1692 seen M1 for 1.96 × 9 **or** 196 × 9 **or** 17.64 seen **or** 1764 seen **OR** M1 for 4.23 ÷ 9 **or** 423 ÷ 9 **or** 0.47 seen **or** 47 seen M1 for 0.47 × 4 **or** 47 × 4 **or** 1.88 seen **or** 188 seen **OR** M1 for 1.96 ÷ 4 **or** 196 ÷ 4 **or** 0.49 seen **or** 49 seen M1 for 0.49 × 9 **or** 49 × 9 **or** 4.41 seen **or** 441 seen **OR** M1 for 9 ÷ 4.23 **or** 2.12(...) seen **or** 2.13 seen M1 for 4 ÷ 1.96 **or** 2.04(...) seen A1 for Pack of 9 and fully correct calculations **NOTE:** B0 for an answer of 9 not supported by working.
20	(a)		6	1	B1 cao
	(b)		44	1	B1 cao
	(c)		31	2	M1 for 60 – 29 **or** 29 – 60 **or** any correct method that is attempting to find the difference between 29 and 60 (allow 1 arithmetic error) A1 cao

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
21*		Angle *DBC* = (180 – 50) ÷ 2 Base angles of isosceles triangle are equal Angle *ABD* = 180 – 65 Angles on a straight line add up to 180 *x* = 180 – 20 – 115 Angles in a triangle add up to 180 **OR** Angle *DBC* = (180 – 50) ÷ 2 Base angles of isosceles triangle are equal *x* = 65 – 20 Exterior angle of triangle is equal to sum of interior opposite angles **OR** Angle *DCB* = (180 – 50) ÷ 2 Base angles of isosceles triangle are equal *x* = 180 – 50 – 20 – 65 Angles in a triangle add up to 180	45 with reasons	4	M1 for (180 – 50) ÷ 2 oe **or** 65 seen M1 for 180 – 20 – (180 – "65") **or** "65" – 20 or 180 – 50 – 20 –'65' oe C2 for *x* identified as 45 with full reasons QWC: Reasons clearly laid out with correct geometrical language used (C1 (dep on M1) for one reason QWC: Reasons clearly laid out with correct geometrical language used) NOTE: *x* = 45 with no working or without any correct angles marked on the diagram cannot score.

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
22	(a)	360 ÷ 60 = 6 300 ÷ 60 = 5 6 × 5 =	Yes and 30	3	M1 for dividing side of patio by side of paving slab eg 360 ÷ 60 **or** 300 ÷ 60 **or** 3.6 ÷ 0.6 **or** 3 ÷ 0.6 **or** 6 and 5 seen **or** 6 divisions seen on length of diagram **or** 5 divisions seen on width of diagram M1 for correct method to find number of paving slabs eg (360 ÷ 60) × (300 ÷ 60) oe **or** 6 × 5 **or** 30 squares seen on diagram (units may not be consistent) A1 for Yes **and** 30 (or 2 extra) with correct calculations **OR** M1 for correct method to find area of patio **or** paving slab eg 360 × 300 **or** 108000 seen **or** 60 × 60 **or** 3600 seen **or** 3.6 × 3 **or** 10.8 seen **or** 0.6 × 0.6 **or** 0.36 seen M1 for dividing area of patio by area of a paving slab eg. (3.6 × 3) ÷ (0.6 × 0.6) oe (units may not be consistent) A1 for Yes **and** 30 (or 2 extra) with correct calculations **OR** M1 for method to find area of patio and area of 32 slabs eg. 60 × 60 × 32 **or** 360 × 300 M1 for method to find both areas eg. 60 × 60 × 32 **and** 360 × 300 (units may not be consistent) A1 for Yes **and** 115200 **and** 108000 **OR** Yes **and** 11.52 **and** 10.8 **NB :** Throughout the question, candidates could be working in metres or centimetres

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
22	(b)	1726 25890 27616 **8 6 3** **2** 2 4 1 8 9 **3** **7** 1 6 1 2 6 **2** 6 1 6 800 60 3 30 24000 1800 90 2 1600 120 6 24000 + 1800 + 90 + 1600 + 120 + 6 = 27616	276.16	3	M1 for complete correct method with relative place value correct. Condone 1 multiplication error, addition not necessary. **OR** M1 for a complete grid. Condone 1 multiplication error, addition not necessary. **OR** M1 for sight of a complete partitioning method, condone 1 multiplication error. Final addition not necessary. A1 for digits 27616 A1 ft (dep on M1) for correct placement of decimal point after addition (of appropriate values) (SC: B1 for attempting to add 32 lots of 8.63)
23	(a)		30	2	M1 for 25 ÷ 10 **or** 2.5 seen **or** 10 ÷ 25 **or** 0.4 seen **or** 12+12+6 oe **or** a complete method eg. 25×12÷10 oe A1 cao
	(b)	1000 ÷ 200 × 12	60	2	M1 for 500 ÷ 50 **or** 1000 ÷ 200 **or** 500 ÷ 10 **or** correct scale factor clearly linked with one ingredient eg 10 with sugar or 5 with butter or flour or 50 with milk **or** an answer of 120 or 600 A1 cao

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
24		Acton after 24, 48, 72, 96, .. Barton after 20, 40, 60, 80, .. LCM of 20 and 24 is 120 9:00 am + 120 minutes **OR** Acton after 24, 48, 1h 12 min... Barton after 20, 40, 1 h LCM is 2 hours 9:00 am + 2 hours **OR** Times from 9:00 am when each service leaves the bus station Acton at 9:24, 9:48, 10:12.. Barton at 9:20, 9:40, 10:00.. **OR** 20 = 2 × 2 × 5 24 = 2 × 2 × 2 × 3 2 × 2 × 2 × 3 × 5 = 120	11:00 am	3	M1 for listing multiples of 20 and 24 with at least 3 numbers in each list ; multiples could be given in minutes or in hours and minutes (condone one addition error in total in first 3 numbers in lists) A1 identify 120 (mins) **or** 2 (hours) as LCM A1 for 11:00 (am) **or** 11(am) **or** 11 o'clock **OR** M1 for listing times after 9am when each bus leaves the bus station, with at least 3 times in each list (condone one addition error in total in first 3 times after 9am in lists) A1 for correct times in each list up to and including 11:00 A1 for 11:00 (am) **or** 11(am) **or** 11 o'clock **OR** M1 for correct method to write 20 and 24 in terms of their prime factors 2, 2, 5 and 2, 2, 2, 3 (condone one error) A1 identify 120 as LCM A1 for 11:00 (am) **or** 11(am) **or** 11 o'clock

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
25	(a)		$6y - 15$	1	B1 cao
	(b)		$4x(2x + y)$	2	B2 cao (B1 for $x(8x + 4y)$ **or** $2x(4x + 2y)$ **or** $4(2x^2 + xy)$ **or** $4x(ax + by)$ where a, b are positive integers **or** $ax(2x + y)$ where a is a positive integer **or** $4x(2x \quad y)$)
	(c)	$10t = gh$ $h = \frac{10t}{g}$	$\frac{10t}{g}$	2	M1 for clear intention to multiply both sides of the equation by 10 (eg. ×10 seen on both sides of equation) **or** clear intention to divide both sides of the equation by g (eg. ÷ g seen on both sides of equation) **or** $10t = gh$ **or** $\frac{t}{g} = \frac{h}{10}$ **or** fully correct reverse flow diagram eg. ← ×10 ← ÷g ← A1 for $\frac{10t}{g}$ oe

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
26	(a)	$2 \times 5 \times 2 = 20$ $300 \div 20 =$	15	3	M2 for $300 \div (2 \times 5 \times 2)$ oe (M1 for $2 \times 5 \times 2$ **or** 20 seen **or** $300 \div (2 \times 5)$ **or** 30 seen A1 cao
	(b)	$c = \frac{30 \times 40}{150}$ =	8	2	M1 for $\frac{30 \times 40}{150}$ **or** 1200 seen A1 cao
27		$3x - 15 = 2x + 24$ $x = 39$ **OR** $2x + 3x - 15 + 2x + 2x + 24 = 360$ $9x + 9 = 360$ $9x = 351$ $x = 39$ **OR** $2x + 2x + 24 = 180$ $4x + 24 = 180$ $4x = 156$ $x = 39$ **OR** $2x + 3x - 15 = 180$ $5x - 15 = 180$ $5x = 195$ $x = 39$	39	3	M1 for forming an appropriate equation eg $3x - 15 = 2x + 24$ **or** $2x + 3x - 15 + 2x + 2x + 24 = 360$ oe **or** $2x + 2x + 24 = 180$ oe **or** $2x + 3x - 15 = 180$ oe **or** $2x + 3x - 15 = 2x + 2x + 24$ M1 (dep) for correct operation(s) to isolate x and non-x terms in an equation to get $ax = b$ A1 cao **OR** M2 for $\frac{351}{9}$ **or** $\frac{195}{5}$ **or** $\frac{156}{4}$ oe A1 cao

PMT

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481
Email publication.orders@edexcel.com

Order Code UG032619 Summer 2012

For more information on Edexcel qualifications, please visit our website
www.edexcel.com

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual
Llywodraeth Cynulliad Cymru
Welsh Assembly Government
Rewarding Learning

PMT



Mark Scheme (Results)

Summer 2015

Pearson Edexcel GCSE
In Mathematics B (2MB01)
Higher (Calculator) Unit 3



PMT

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2015
Publications Code UG042121
All the material in this publication is copyright
© Pearson Education Ltd 2015

PMT

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Note that in some cases a correct answer alone will not score marks unless supported by working; these situations are made clear in the mark scheme. Examiners should be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will award marks for the quality of written communication (QWC).
The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labelling conventions.

ii) *select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

PMT

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Partial answers shown (usually indicated in the ms by brackets) can be awarded the method mark associated with it (implied).
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks; transcription errors may also gain some credit. Send any such responses to review for the Team Leader to consider.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded (embedded answers).

12 Parts of questions
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers
Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

14 The detailed notes in the mark scheme, and in practice/training material for examiners, should be taken as precedents over the above notes.

Guidance on the use of codes within this mark scheme
M1 – method mark for appropriate method in the context of the question A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

PMT

5MB3H/01 June 2015				
Question	**Working**	**Answer**	**Mark**	**Notes**
1		71.40	2	M1 for 41.65÷7 A1 for 71.4(0)
2		4	3	M1 $\frac{4.5}{100} \times 300$ (=13.5) or $\frac{104.5}{100} \times 300$ (=313.5) oe M1 50 ÷ "13.5" (=3.7) or at least 3 repeated addition of "13.5" A1 cao SC B1 for $1.045^n \times 300$

5MB3H/01 June 2015				
Question	**Working**	**Answer**	**Mark**	**Notes**
3 (a)		1.75	2	M1 for intention to multiply brackets or for intention to divide all terms by 4 as the first step A1 for 1.75 oe
(b)		p > 7.5	2	M1 for correct method to isolate *p* or intention to divide all terms by 2 as the first step (condone the use of '=' in method) A1 for *p* > 7.5 oe
(c)		3, -5	3	M2 for $(x - 3)(x + 5)$ (M1 for $(x \pm 3)(x \pm 5)$) A1 cao 3 and –5 OR M1 for $\frac{-2 \pm \sqrt{2^2 - 4 \times 1 \times -15}}{2 \times 1}$ M1 for $\frac{-2 \pm \sqrt{64}}{2}$ A1 for 3 and – 5 cao

PMT

5MB3H/01 June 2015				
Question	**Working**	**Answer**	**Mark**	**Notes**
*4		275g (£7.95) pack with reason	4	M1 for one of 439 ÷ 150 (= 2.92), 639 ÷ 200 (= 3.19), 795 ÷ 275 (= 2.89) OR one of 150 ÷ 439 (= 0.341), 200 ÷ 639 (= 0.31), 275 ÷ 795 (= 0.345) OR any other calculation that could lead to a comparative figure M1 for calculations that could lead to comparative figures for 2 jars M1 for calculations that could lead to comparative figures for 3 jars eg all 3 from the above lists C1 for correct comparative figures for all 3 jars leading to a correctly stated comparison: 275g best value
5 (a)		7, -2, -1	2	B2 for all three correct values 7, -2, -1 (B1 for two correct values 7, -2 or -1)
(b)		Correct curve	2	B2 for fully correct curve (B1 ft for at least 5 points plotted correctly)
6		6	4	M1 for $7x + 22$ or $(5x+2)$ or $7b + 22p$ or $5b + 2p$ M1 for forming equation $7x + 22 = 2(5x + 2)$ M1 for correct intent to isolate x on one side A1 cao

PMT

5MB3H/01 June 2015				
Question	**Working**	**Answer**	**Mark**	**Notes**
7		28	3	M1 for 240 × 1.2 (=288) M1 for "288" ÷ 10 (=28.8) A1 cao OR M1 for 10 ÷ 1.2 (=8.33) M1 240 ÷ "8.33" (=28.8) A1 cao
8		Correct position marked	3	M1 for perpendicular bisector of the line AC M1 for arc of circle centre D radius 7cm A1 for correct point
*9		No with comparison of correct figures.	3	M1 for correct method to find frame dimensions 15+5+5 (=25) and 10+5+5 (=20) M1 for finding relationship between length and width for both eg 15 ÷ 10 (=1.5) and 25 ÷ 20 (= 1.25) or for correct use of a scale factor to find a comparable length C1 for correct conclusion following comparison of 2 correct comparable figures
10		Rotation, 90° clockwise centre (1,4)	3	B1 for rotation B1 for 90° clockwise or 270° anticlockwise B1 for (1,4) NB Award B0 if more than one transformation given

PMT

5MB3H/01 June 2015				
Question	**Working**	**Answer**	**Mark**	**Notes**
11		3.4	3	M1 for $\pi \times 3.5^2 \times 10$ (=384.85 or =122.5π) or $\pi \times 6^2$ (=113.1 or =36π) M1 (dep) for "384.85" ÷ "113.1" or 122.5(π) ÷ 36(π) A1 for 3.4 – 3.5 OR M1 for $\left(\frac{12}{7}\right)^2$ (=2.9) or $\left(\frac{7}{12}\right)^2$ (=0.34) M1 for $10 \div "\left(\frac{12}{7}\right)^2"$ or $10 \times "\left(\frac{7}{12}\right)^2"$ A1 for 3.4 – 3.5
12		7.5×10^9	2	M1 for 2.1×10^7 or 21000000 or sight of figures 75 A1 for 7.5×10^9
13		29.6	4	M1 for $8^2 + 5^2$ or 64 + 25 or 89 M1 (dep) $\sqrt{"8^2"+"5^2"}$ (=9.4) M1 for "9.4..." × π A1 for 29.5 – 29.65
*14		49	3	M1 for 180 - 56 - 75 A1 for 49 C1 for alternate segment theorem and angles on a straight line add up to 180° OR alternate segment theorem and angles in a triangle add up to 180° Appropriate to methods shown

PMT

5MB3H/01 June 2015				
Question	**Working**	**Answer**	**Mark**	**Notes**
15		0.84	3	M2 for 2×0.75^3 (M1 for 2×0.75 (=1.5) or 2×0.25 (=0.5) and 2 – "0.5") A1 for 0.84 − 0.844
16	$y(x+5) = 3x$ $yx+5y = 3x$ $5y = 3x - yx$ $5y = x(3 - y)$	$x = \frac{5y}{3-y}$	3	M1 for intention to multiply by $x + 5$ M1 for intention to isolate yx and $3x$ on one side to get $3x - xy$ oe A1 for $x = \frac{5y}{3-y}$ or $\frac{-5y}{y-3}$
17		85.6	4	M1 for 360 ÷ 5 (=72) M1 (dep) for $\frac{1}{2} \times 6^2 \times \sin 72$ (=17.12) M1 for completing full method to find total area of pentagon A1 for 85.5 − 85.6 OR M1 for 360 ÷ 10 (=36) or $\frac{1}{2}$(180 − 360÷5) (=54) M1(dep) for eg $6 \times \sin 36 \times 6 \times \cos 36$ (=17.12) or $\frac{1}{2} \times 6 \times \sin 54 \times 6 \times \cos 54$ (=8.55) M1 for completing full method to find total area of pentagon A1 for 85.5 − 85.6

PMT

5MB3H/01 June 2015				
Question	**Working**	**Answer**	**Mark**	**Notes**
18		$y = \frac{375}{x^2}$	3	M1 for $y \propto \frac{1}{x^2}$ or $y = \frac{k}{x^2}$ or $k = yx^2$ M1 for $5^2 \times 15$ (=375) A1 for $y = \frac{375}{x^2}$ or $y = \frac{k}{x^2}$ and $k = 375$
19		300	3	B1 for 45 or 0.15 M1 for 45 ÷ 0.15 A1 for 300 cao
20 (a)		$p = 4$, $q = -10$	3	M1 for sight of $(x - 4)^2$ or $p = 4$ M1 for $(x - 4)^2 - 16 + 6$ A1 for $p = 4$, $q = -10$ OR M1 for $x^2 - 2px + p^2 + q$ or $-2p = -8$ or $p^2 + q = 6$ M1 for $-2p = -8$ and $p^2 + q = 6$ A1 for $p = 4$, $q = -10$
(b)		(4, -10)	1	B1 ft
21 (a)		Correct graph	2	M1 for any translation along x axis A1 correct position
(b)		$y = -g(x)$	1	B1 for $y = -g(x)$ oe from correct interpretation of diagram

PMT

5MB3H/01 June 2015				
Question	**Working**	**Answer**	**Mark**	**Notes**
22		6.2	5	M1 for a method to find an angle *RAB* = 70, *ABR* = 50, *BRA* = 60 or *TAR* = 20 M1 for substitution into sine formula $\frac{AR}{\sin"50"} = \frac{12}{\sin"60"}$ M1 for use of sine rule to find *AR*, *AR* = $\frac{12}{\sin"60"} \times \sin"50"$ (=10.61) M1 for substitution into cosine formula $TR^2 = 5^2 + "10.61"^2 - 2 \times 5 \times "10.61" \times \cos 20$ (=37.92) A1 for 6.15 - 6.2
23		$\frac{3}{4}(\mathbf{b} + \mathbf{a})$	4	B1 for $\mathbf{b} - \mathbf{a}$ or $\frac{1}{2}(\mathbf{b} - \mathbf{a})$ or $\mathbf{a} - \mathbf{b}$ or $\frac{1}{2}(\mathbf{a} - \mathbf{b})$ M1 for $\overrightarrow{OM} = \mathbf{a} + \frac{1}{2}(\mathbf{b} - \mathbf{a})$ or $\mathbf{b} + \frac{1}{2}(\mathbf{a} - \mathbf{b})$ oe M1 for $\overrightarrow{ON} = \frac{3}{2}(\mathbf{a} + \frac{1}{2}(\mathbf{b} - \mathbf{a}))$ or $\frac{3}{2}(\mathbf{b} + \frac{1}{2}(\mathbf{a} - \mathbf{b}))$ A1 for $\frac{3}{4}(\mathbf{b} + \mathbf{a})$ oe (simplified)

PMT

Modifications to the mark scheme for Modified Large Print (MLP) papers.

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q4		Diagrams have been removed.	M1 for one of 439 ÷ 150 (= 2.92), 639 ÷ 200 (= 3.19), 795 ÷ 275 (= 2.89) OR one of 150 ÷ 439 (= 0.341), 200 ÷ 639 (= 0.31), 275 ÷ 795 (= 0.345) OR any other calculation that could lead to a comparative figure M1 for calculations that could lead to comparative figures for 2 jars M1 for calculations that could lead to comparative figures for 3 jars eg all 3 from the above lists C1 for correct comparative figures for all 3 jars leading to a correctly stated comparison: 275g best value
Q5	(a)	Wording added: 'There are three spaces to fill.' Table produced in vertical format.	B2 for all three correct values 7, -2, -1 (B1 for two correct values 7, -2 or -1)
Q5	(b)	1½ cm grid provided.	B2 for fully correct curve (B1 ft for at least 5 points plotted correctly)

PMT

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q8		Size of diagram kept as original. Scale put on diagram as well as in text.	M1 for perpendicular bisector of the line AC M1 for arc of circle centre D radius 7cm A1 for correct point
Q9		Diagram has been enlarged. Dotty shading. Measurement labels moved to be more accessible for candidates.	M1 for correct method to find frame dimensions 15+5+5 (=25) and 10+5+5 (=20) M1 for finding relationship between length and width for both eg 15 ÷ 10 (=1.5) and 25 ÷ 20 (= 1.25) or for correct use of a scale factor to find a comparable length C1 for correct conclusion following comparison of 2 correct comparable figures
Q10		The grid has been enlarged. Shapes are labelled triangle A and triangle B. Wording added: 'It shows two triangles on the grid, triangle A and triangle B.' Dotty shading	B1 for rotation B1 for 90° clockwise or 270° anticlockwise B1 for (1,4) NB Award B0 if more than one transformation given

PMT

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q11		The diagram has been enlarged. 2 models provided for all candidates. A diagram is also provided for MLP.	M1 for $\pi \times 3.5^2 \times 10$ (=384.85 or =122.5π) or $\pi \times 6^2$ (=113.1 or =36π) M1 (dep) for "384.85" ÷ "113.1" or 122.5(π) ÷ 36(π) A1 for 3.4 – 3.5 OR M1 for $\left(\frac{12}{7}\right)^2$ (=2.9) or $\left(\frac{7}{12}\right)^2$ (=0.34) M1 for 10 ÷ "$\left(\frac{12}{7}\right)^2$" or 10 × "$\left(\frac{7}{12}\right)^2$" A1 for 3.4 – 3.5
Q13		The diagram has been enlarged.	M1 for $8^2 + 5^2$ or 64 + 25 or 89 M1 (dep) $\sqrt{"8^2"+"5^2"}$ (=9.4) M1 for "9.4..." × π A1 for 29.5 – 29.65
Q14		The diagram has been enlarged. Wording added: 'Angle CAB is marked x'.	M1 for 180 - 56 - 75 A1 for 49 C1 for alternate segment theorem and angles on a straight line add up to 180° OR alternate segment theorem and angles in a triangle add up to 180° Appropriate to methods shown

PMT

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q16		MLP: *x* changed to *e*. *y* changed to *f*.	M1 for intention to multiply by $e + 5$ M1 for intention to isolate yx and $3x$ on one side to get $3e - ef$ oe A1 for $x = \frac{5f}{3-f}$ or $\frac{-5f}{f-3}$
Q17		The diagram has been enlarged.	M1 for $360 \div 5$ (=72) M1 (dep) for $\frac{1}{2} \times 6^2 \times$ sin"72" (=17.12) M1 for completing full method to find total area of pentagon A1 for 85.5 − 85.6 OR M1 for $360 \div 10$ (=36) or $\frac{1}{2}(180 - 360 \div 5)$ (=54) M1(dep) for eg 6 × sin"36" × 6 × cos"36" (=17.12) or $\frac{1}{2}$ 6 × sin"54" × 6 × cos"54" (=8.55) M1 for completing full method to find total area of pentagon A1 for 85.5 − 85.6
Q19		MLP and braille: a changed to x. b changed to f.	B1 for 45 or 0.15 M1 for $45 \div 0.15$ A1 for 300 cao

PMT

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q21	(a)	The grid has been enlarged.	M1 for any translation along x axis A1 correct position
Q21	(b)	The grid has been enlarged.	B1 for $y = -g(x)$ oe from correct interpretation of diagram
Q22		The diagram has been enlarged. TR is joined by a dashed line.	M1 for a method to find an angle, $RAB = 70$, $ABR = 50$, $BRA = 60$ or $TAR = 20$ M1 for substitution into sine formula $\frac{AR}{\sin"50"} = \frac{12}{\sin"60"}$ M1 for use of sine rule to find AR, $AR = \frac{12}{\sin"60"} \times \sin"50"$ (=10.61) M1 for substitution into cosine formula $TR^2 = 5^2 + "10.61"^2 - 2 \times 5 \times "10.61" \times \cos 20$ (=37.92) A1 for 6.15 - 6.2
Q23		The diagram has been enlarged. OM is joined by a dashed line.	B1 for $\mathbf{b} - \mathbf{a}$ or $\frac{1}{2}(\mathbf{b} - \mathbf{a})$ or $\mathbf{a} - \mathbf{b}$ or $\frac{1}{2}(\mathbf{a} - \mathbf{b})$ M1 for $\overrightarrow{OM} = \mathbf{a} + \frac{1}{2}(\mathbf{b} - \mathbf{a})$ or $\mathbf{b} + \frac{1}{2}(\mathbf{a} - \mathbf{b})$ oe M1 for $\overrightarrow{ON} = \frac{3}{2}(\mathbf{a} + \frac{1}{2}(\mathbf{b} - \mathbf{a}))$ or $\frac{3}{2}(\mathbf{b} + \frac{1}{2}(\mathbf{a} - \mathbf{b}))$ A1 for $\frac{3}{4}(\mathbf{b} + \mathbf{a})$ oe (simplified)

PMT

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom

PMT

edexcel

Mark Scheme (Results)

November 2015

Pearson Edexcel GCSE
In Mathematics A (1MA0)
Higher (Non-Calculator) Paper 1H

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

November 2015
Publications Code UG042920
All the material in this publication is copyright
© Pearson Education Ltd 2015

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Note that in some cases a correct answer alone will not score marks unless supported by working; these situations are made clear in the mark scheme. Examiners should be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will award marks for the quality of written communication (QWC).
The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labelling conventions.

ii*) select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Partial answers shown (usually indicated in the ms by brackets) can be awarded the method mark associated with it (implied).
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks; transcription errors may also gain some credit. Send any such responses to review for the Team Leader to consider.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 **Linear equations**
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded (embedded answers).

12 **Parts of questions**
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 **Range of answers**
Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

14 The detailed notes in the mark scheme, and in practice/training material for examiners, should be taken as precedents over the above notes.

Guidance on the use of codes within this mark scheme
M1 – method mark for appropriate method in the context of the question A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

1MA0 1H November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
1			69	4	M1 for finding 15% of £720 (=108) M1 (dep) for finding total of £720 plus interest (or 115% etc) (=828) M1 (dep on previous M1) dividing by 12 A1 cao OR M1 finding 720 ÷ 12 (=60) M1 (dep) finding 15% of "60" (=9) M1 (dep on previous M1) for adding, e.g. 60 + 9 A1 cao
2	(i)		19.44	2	B1 cao
	(ii)		19 440		B1 cao
3	(a)		$6n + 5$	2	B2 for $6n + 5$ (B1 for $6n + k$ where k is an integer or absent)
	(b)		No with explanation	2	M1 for "$6n + 5$" = 121 or any other valid method, e.g. counting on 6s (to get to 119 or more) A1 for No with complete explanation, e.g. $6n$=116 will not give a whole number
4	(a)		60	2	M1 for 200 × 0.3 oe A1 cao
	(b)		0.1	2	M1 subtracting sum of probabilities from 1, e.g. 1−(0.3+0.2+0.4) A1 cao

1MA0 1H November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
5			20	3	M1 for 330÷120 (=2.75) or 200÷60 (=3 1/3) or 450 ÷180 (=2.5) M1 for 450÷180 (=2.5) AND 8×"2.5" A1 cao OR M1 for 120÷8 (=15) or 60÷8 (=7.5) or 180÷8 (=22.5) M1 for 330÷(120÷8) [=22] or 200÷(60÷8) [=26.6..] or 450÷(180÷8) A1 cao OR M1 for multiples of 120:60:180 M1 for multiplication linked to 450 and 8+8+4 A1 cao
6			40°	4	M1 for angle FBC=70 or CFG = x or ABF = 110 may be seen in diagram M1 for angle CBF = BFC =70 or 90 – ½ x A1 for 40 supported by working C1 (dep on M2) for all reasons and linked to appropriate working, e.g. Alternate angles are equal; Allied angles / Co-interior angles add up to 180°; Base angles of an isosceles triangle are equal; angles in a triangle add to 180°, angles on a straight line equals 180°
7	(a)		explanations	2	B2 for two aspects from: no time frame; responses vague; no "never" box (B1 for one correct aspect)
	(b)		question response boxes+	2	B1 for a question with a time frame (may appear with response boxes) B1 for at least 3 correctly labelled response boxes (non-overlapping and exhaustive) Do not accept inequality symbols.

1MA0 1H November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
8		2p 1p ½ p Tot Sat 7 16 (31) 54 Sun (15) 14 17 (46) Tot (22)(30) 48 (100)	14	4	M1 for total Sat bottles 100 – 46 (=54) or for total ½ pint bottles 100 – 22 – 30 (=48) or (total 2 pint bottles on Sat) 22 – 15 (=7) M1 for total Sun bottles of ½ pint "48" – 31 (=17) or for total Sat bottles of 1 pint: "54" – 31 – (22 −15) (=16) M1 for 46 – 15 – "17" or for 30 – "16" A1 cao NB: any of the above figures could be shown in a 2-way table
*9			NO with evidence	4	M1 for 50 × 40 × 30 (=60000) M1 for "60000" ÷ 3000 (=20) M1 for "20" × £3.50 C1 eg for 70 and comparison resulting in NO OR M1 for £60 ÷ 3.50 (=17 bottles) M1 for "17" × 3000 (=51000) M1 for 50 × 40 × 30 (=60000) C1 eg for 51000 and 60000 and comparison resulting in NO

1MA0 1H November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
10	(a)		$x^2 + 2x$	1	B1 cao
	(b)		$3y + 4x + 2$	2	M1 for a method to expand a bracket, e.g. $3y + 6$ or $4x - 4$ A1 cao
	(c)	$2t^2 + 10t - 3t - 15$	$2t^2 + 7t - 15$	2	M1 for 4 terms correct ignoring signs or 3 out of no more than 4 terms with signs correct unless ambiguous A1 cao
	(d)		$4a(2a + 3)$	2	M1 for $4a(na+c)$ or $2a(4a+6)$ or $a(8a+12)$ [n,c integers, $c \neq 0$] A1 cao
	(e)		$(y + 1)(y - 2)$	2	M1 for $(y \pm 1)(y \pm 2)$ unless ambiguous A1 cao
11	(a)		150	2	M1 for $180 - (360 - 330)$ or $180 - 30$ or $330 - 180$ or a complete diagram showing the bearing of 330° A1 cao
	(b)		11 40	4	M1 for $200 \div 120$ (=1 2/3 h) M1 for conversion between hours and minutes A1 for 1 h 40 min or 100 minutes B1 (ft dep on M1) for 11 40

1MA0 1H November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
12	(a)		2, 0, 0, 6	2	B2 for 2, 0, 0, 6 (B1 for at least two of 2, 0, 0, 6); could be taken from graph
	(b)		Correct curve	2	M1 (ft) for at least 5 points plotted correctly A1 for a fully correct curve
	(c)		−0.6, 3.6	2	M1 (ft if M1 awarded in (b) and at least B1 in (a)) for indicating a point or line drawn at $y=4$, or one solution given A1 (ft) for both solutions
13			20	3	M1 for 30×14 (=420) or 18×10 (=180) M1 for $30 \times 14 - 18 \times 10$ or "420" – "180" (=240) A1 cao
14			126	4	M1 for method to find exterior or interior angle of octagon M1 for method to find exterior or interior angle of pentagon M1 for complete method A1 cao

1MA0 1H November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
15	(a)		19, 36, 51, 63, 73, 80	1	B1 cao
	(b)		cf graph	2	M1 for at least 5 of the 6 points plotted at each upper end of the interval (not joined) or 5 of the 6 points plotted consistently within interval (not upper end) and joined (dep on a cf table with no more than one arithmetic error) A1 correct graph
	*(c)		comparable value and conclusion	3	M1 for indication of a reading taken from a cf graph using weight = 3.4 kg or find UQ from 60 A1 for value given between 55 & 57 or 3.6 & 3.8 C1 (dep on at least M1) for conclusion (justified)
16			13.75	5	M1 for finding perimeter of rectangle e.g. $5x + 5 + 5x + 5 + 4x + 4x$ $(= 18x + 10)$ M1 for finding perimeter of trapezium e.g. $9x - 2 + 7x - 2 + 10x$ $(=26x - 4)$ M1 for equation e.g. $26x - 4 = 18x + 10$ (or $8x = 14$) A1 for finding the value of x as 1.75 B1 ft for subs of x into $5x + 5$ and evaluated (=13.75)
17			$x = 3\frac{1}{3}$ $y = -2$	4	M1 for a correct process to eliminate either variable (condone one arithmetic error) or to rearrange and substitute for elimination A1 cao for either x or y M1 (dep on M1) for correct substitution of found value into one of the equation or appropriate method after starting again (condone one arithmetic error) A1 cao

1MA0 1H November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
18			$2\sqrt{5}$	2	M1 for multiplication of denominator and numerator by $\sqrt{5}$ A1 cao
19			756π	5	M1 for $\frac{1}{3}\pi r^2 \times 10$ $(=270\pi)$ A1 for $r = 9$ M1 (dep on M1) for $\frac{1}{2} \times \frac{4}{3}\pi \times "9"^3$ $(= 486\pi)$ M1 for 270π + "486π" oe A1 cao
*20			Proof	5	M1for finding one other vector expressed as **a** and/or **b** M1 for method to find one of $\overrightarrow{DM}$, $\overrightarrow{MA}$ or $\overrightarrow{DA}$ eg $\overrightarrow{DM} = -\mathbf{b} + \frac{1}{2}(3\mathbf{b} + \mathbf{a})$ oe, $\overrightarrow{MA} = \frac{1}{2}(3\mathbf{b} + \mathbf{a}) + \mathbf{a}$ oe or $\overrightarrow{DA} = 2\mathbf{b} + 2\mathbf{a}$ oe M1 for method to find two of $\overrightarrow{DM}$, $\overrightarrow{MA}$ or $\overrightarrow{DA}$ A1 for two of $\overrightarrow{DM} = \frac{1}{2}(\mathbf{a} + \mathbf{b})$, $\overrightarrow{MA} = 1.5(\mathbf{a} + \mathbf{b})$, $\overrightarrow{DA} = 2(\mathbf{a} + \mathbf{b})$ ie simplified but oe C1 (dep on working shown) for conclusion relating to correct working

1MA0 1H November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
21	(a)		$\frac{9x-8}{x(2-x)}$	3	M1 for method to use a common denominator, e.g. $\frac{5(x)-4(2-x)}{x(2-x)}$ M1 (dep on M1) for correct expansion of brackets and combination of numerators e.g. $5x-8+4x$ $(=9x-8)$ A1 for $\frac{9x-8}{x(2-x)}$ or $\frac{9x-8}{2x-x^2}$
	(b)		$y = \frac{2-2t}{t+3}$	4	M1 for intention to multiply both sides by $y + 2$ as a first step e.g. $t \times y + 2 = 2 - 3y$ M1 for intention to correctly isolate their y terms on one side and the other terms on the other side, e.g. $ty+3y=2-2t$ M1 for intention to factorise, e.g. $y(t+3)$ $(=2-2t)$ A1 for $y = \frac{2-2t}{t+3}$ oe
*22			Similarity and proof	5	B1 for method matching a pair of opposite angles, e.g. if $EAB = x$, $BDE = 180-x$, $EAB + BDE = 180$ B1 for linking angles between quad and triangle, e.g. if $BDE = 180-x$ then $BDC = x$ B1 for stating or implying $ACE = BCD$ (same angle) C1 for Opposite angles of a cyclic quadrilateral add up to 180° or statement linking three angles for similarity C1 for complete proof

Modifications to the mark scheme for Modified Large Print (MLP) papers.

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 1MA0_1H			
Question		**Modification**	**Notes**
Q06		Diagram is enlarged.	Standard mark scheme
Q09		Model is provided for all candidates. Diagram also provided for MLP.	Standard mark scheme
Q10	(a)	MLP only: x changed to y Diagram is enlarged.	Standard mark scheme
Q12	(a) (b)	Wording added “There are four spaces to fill.” 1.5 cm grid..Small squares removed.	Standard mark scheme
Q14		Diagram enlarged.	Standard mark scheme
Q15	(c)	Frequencies changed to 25, 20, 15, 10, 5, 5. Grid: 1.5 cm 3.4 kg changed to 3.5 kg.	Standard mark scheme
Q16		“trapezium” and “rectangle” put inside the shapes. MLP only: x changed to y.	Standard mark scheme
Q19		Model provided for all candidates. Diagram also provided for MLP.	Standard mark scheme

PAPER: 1MA0_1H			
Question		**Modification**	**Notes**
Q20		Diagram is enlarged. Lower case vectors enlarged.	Standard mark scheme
Q22		Diagram is enlarged.	Standard mark scheme

PMT

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom

Question 17

This question was very poorly answered, many simply using a scale factor of 2.25 (9 ÷ 4) in an attempt to make progress. Some looked as though they wanted to use the formulae for surface area and volume of a cylinder, but never really made any meaningful start. A few students correctly found the ratio of lengths but could go no further.

Question 18

Many students found this question challenging and the modal score is zero. There were many blank responses.

Those who could use the cosine rule generally obtained the first two marks however, there was often confusion over the identity of angle BCA. Of those that correctly found angle BCA very few were able to apply the sine rule accurately and gain full marks for this question. A small number of students lost the final accuracy mark due to rounding earlier in their solution.

The most common errors seen involved using the trigonometry ratios for right-angled triangles.

Question 19

There were some excellent responses to this question from the most able students taking this paper but these were few and far between. Able students who were unable to get as far as the given result were sometimes able to score some marks for successfully finding expressions for the area of part of a hexagon or for a whole hexagon, whether it be *ABCDEF* or *FGHIJK*. Where this applied, students usually used trigonometry to find the height of a constituent triangle and then its area by using $\frac{1}{2} \times$ base $\times$ height or they found an area using the formula $\frac{1}{2}ab\sin C$.

Some weaker students used the given result and tried to justify it by using substitution or by algebraic manipulation whilst other students seemed to think that the question revolved around finding angles of a polygon. No marks could be awarded for these approaches.

Question 10

This question involving similar right-angled triangles was well answered by a minority of students. There were two main routes used by students who scored full marks. Some students used a scale factor together with Pythagoras' rule while other students used trigonometry to find the size of angle *B* then went on to use trigonometry again to find the length of side *CB*. Of those students who could only be awarded partial credit for their attempts, most scored the mark for using Pythagoras' rule correctly to find the length of *EB*. Unfortunately, they often then used a scale factor of 1.5 instead of 2.5 or failed to make any further progress.

Question 17

A similar solids question, but with an added layer of difficulty. It was pleasing to see so many candidates get some credit. This was normally for either finding the volume scale factor (by dividing the two volumes) or for a process to find the height of **B**, or in a good number of cases, both scoring 2. However, very few candidates were able to take this further. Typically candidates tried to use a scale factor of 27 when working with areas and lengths and gained no further credit e.g. 10.8 ÷ 27. There were also occasions where more marks could have been scored had candidates been able to deal with the half in the area of a triangle. A shame that such a low level skill was denying candidates high grade marks.

Question 5

This question involving similar triangles was well answered by a majority of candidates.
Often there was little or no working seen. Where working was seen, the most common approach was to find the scale factor ($\frac{20}{5}$) of the enlargement, mapping the smaller triangle onto the bigger triangle, then use it to find the lengths required. This approach was usually successful.
It was rare to see the alternative method of using the ratio of two known sides, for example $\frac{AC}{BC} = \frac{5}{4}$. Some candidates tried to solve the problem by using "Pythagoras' rule".
The most common error seen in part (a) was for candidates to use a scale factor of $\frac{22}{4}$.
In part (b) the most frequently incorrect answer seen was 4.4.

Question 23

Many candidates were caught out but the relatively familiar figures of 1, 8 and 27, and thought they had to work with cubing numbers. However, candidates had to find the length scale factor of the 3 triangles from the given ratio as 1:3:6 to gain the first mark, whereas many worked with the 1:2:3. Once they had the correct ratio of lengths they could then work out the correct ratio of areas for the 3 triangles to get 1:9:36. From here they had to work out the ratio of the areas of the 3 shapes, 2 of which were trapezia not triangles. There were a number of excellent responses, some working as above and others working nicely with algebra. There were also a number of candidates that gave lengths to the parallel sides and perpendicular height and worked correctly with areas.

Question 11

There were very few fully correct answers to this question. Lower attaining students did not appreciate the need to use 50^2 or 100^2 in one way or another to take into account area and unit conversions. These students often restricted their response to working out

$\frac{48\times50}{100}$, resulting in the answer 24. However, examiners were able to give partial credit to the relatively small proportion of students who realised the need to use $1\ m^2 = 10\ 000\ cm^2$, often expressed in the form of dividing by 100 twice. Very few students identified the need to use the scale factor of 2500 (or 50×50). Some students successfully chose dimensions for the room on the plan, for example 8 cm and 6 cm then converted these to metres before recalculating the area.

Question 15

The hint in the question given by the ordering of the vertices was lost on most students and this question was very poorly done. Most students were unable to identify the need to compare the pairing of AD and AC with the pairing of AE and AB or equivalent. Many students stopped after finding a scale factor between AD and AB (eg, $54 \div 32$) and the scale factor between AE and AC (eg $80 \div 21.6$). As they got different values, they stopped. There were many erroneous attempts involving "Pythagoras' rule" or areas of triangles. A small number of students who did get two figures to compare did not get the final mark because they did not write a conclusion. Students are advised that

drawing two separate triangles and marking in the lengths of the sides may help them to identify and use the similarity of the triangles.

Question 19

Overall, this question was answered poorly with the majority of students unable to find an appropriate strategy to work out the ratio $AB : AD$. Some students made a good start by using similar triangles to form an equation involving AB, e.g. $\frac{AB}{4} = \frac{25}{AB}$, or an equation involving the scale factor, e.g. 4 × scale factor = 25 ÷ scale factor. Many of those who formed a correct equation were able to complete the process to find the required ratio.
Most students, however, were unable to identify the need to compare the sides of the two similar triangles, *ABC* and *DAB*. Many used the ratio 4 : 21 incorrectly and answers of 4 : 25 were common. Students are advised that drawing two separate triangles and marking in lengths on the sides may help them to use the similarity of the triangles. A small number of students used Pythagoras to find the perpendicular height of the triangles and then *AB*. Some students assumed that Pythagoras could be applied to triangle *ACD* to find *AC* but it is not a right-angled triangle.

Question 17

This question was attempted by most, but only the stronger students were able to score. The first mark was for working with the volume scale factors of **L** and **M** to find the length scale factors, and the second similarly to work with the area scale factors of **M** and **P** to find the length scale factors. Both of these processes required students to take some form of root, and it is perhaps not surprising that students were better when the square root was involved. Many of the students who were able to find the 2 separate length scale factors or ratios, were then unsure as to how to combine them and scored no further credit. Those who did know, typically scored both the 3rd and 4th marks. The A mark was 'or equivalent' and we saw numerous acceptable possibilities, including 9.6 : 12 : 4 which had non-integer values.

Question 12

It was pleasing to see some fully correct solutions but, overall, this question was answered very poorly with relatively few learners showing that they understood the relationships between lengths, areas and volumes in similar solids. Starting the process to find the ratio of the radius of **A** to the radius of **B** by finding $\sqrt[3]{64}$ or $\sqrt[3]{125}$ gained the first mark and some learners reached the ratio 4 : 5. The second mark was awarded for using the information about the radius of sphere **C** and working out $5 \div 2$ or for starting to work with area and working out 4^2. A process to find the ratio of the area of **A** to the area of **C**, such as $4^2 : 2.5^2$, was needed for the third mark but few were able to get this far. Some of those that did gave 16 : 6.25 as the final answer and were not awarded the accuracy mark because the question asked for an answer in the form $a : b$ where a and b are integers. A successful route used by a

few learners was to combine the ratios 4 : 5 and 2 : 1, giving the ratio of the radii as 8 : 10 : 5 from which they could easily see that the required ratio is $8^2 : 5^2$. Many learners could not make a correct start. It was common to see 125, the volume of sphere **B**, divided by 2 and then 64 : 62.5 given as the final answer. Since the question involved three spheres some learners attempted to use the formulae for the surface area or volume of a sphere.

Question 13

The majority of those who identified that the triangles were similar proceeded efficiently with answering this question and gained full marks. However, many incorrectly assumed that the lengths of AB and BC or the lengths of BC and BE were

equal. Of those who didn't initially recognise that the triangles were similar, some did successfully navigate to a correct answer using trigonometry. The majority of learners appeared confident in using the formula for area of a trapezium, with only a small number choosing to split it into a rectangle and triangle.

Question 8

In part (a), a proof that triangle *ABC* is similar to triangle *ADE* was required and this appeared to be particularly challenging with many responses unsure of what was required. Those responses that showed some understanding of what they needed to do often gained the first mark for identifying angle *A* as common or for stating that angle *AED* = angle *ACB* or angle *ADE* = angle *ABC* and giving an appropriate reason. This mark could not be awarded when reasons were incomplete or poorly expressed. Reasons given to explain why angle *ADE* = angle *ABC*, for example, often referred only to parallel lines and did not mention corresponding angles. Equal pairs of angles were often identified with no reasons given and so no marks could be awarded. To complete the proof, it was necessary to identify a second pair of equal angles with an appropriate reason and give a statement that the angles in each triangle are the same. Many responses tried to prove that the two triangles are similar by using the conditions for congruency and it was common to see working that ended with SAS or ASA or RHS. These responses could gain at most one mark. Many responses contained only numerical work, often involving the scale factor, and these gained no marks.

In part (b), many responses were able to score all three marks for working out the length of *EC* as 6 cm. Finding the scale factor earned the first mark and a process to find the length of *AE*, such as 18 ÷ 1.5, earned the second mark. Some responses did not gain the final mark because 12 was given as the final answer. Giving the scale factor in ratio form earned the first mark but a common mistake was to then divide 18 in the ratio 2 : 3 which resulted in answers of 7.2 or 10.8. A common incorrect approach that gained no marks was 30 – 20 = 10 followed by 18 – 10 = 8. There were many incorrect attempts that involved Pythagoras' rule or areas of triangles and trapeziums.

Question 16

This similar solid question was one learners found challenging. The first mark was for beginning to work with scale factors and was normally awarded for finding the volume scale factor of 15. Unfortunately, many then didn't know how to find the linear scale factor and tried to use the volume scale factor with the lengths.
The strongest responses were able to take the cube root to find the length scale factor and then use this with the given height to find the missing height and an answer of 4.9.
Credit could have been given in the second process mark for equating scale factors, but this was rarely seen. Some premature rounding affected learners' final answers particularly if they worked with 1/15 as the scale factor rather than 15.
Part (b) effectively asked for the exact value of the volume scale factor. Unfortunately, many learners did not understand the real meaning of 'exact value' and gave a rounded decimal. This alone, without a correct expression as a surd or with indices, could not gain the mark.

Pearson Edexcel

Examiners' Report
Principal Examiner Feedback

Summer 2022

Pearson Edexcel GCSE (9 – 1)
In Mathematics (1MA1)
Higher (Calculator) Paper 3H

Edexcel and BTEC Qualifications
Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2022
Publications Code 1MA1_2H_2206_ER
All the material in this publication is copyright
© Pearson Education Ltd 2022

GCSE Mathematics (1MA1/3H)
Principal Examiner Feedback – Higher Paper 3

Introduction

The time allowed for the examination appears to have been sufficient for most students to complete this paper. The great majority of students seemed well suited to entry at the higher tier.

Most students entered for this examination presented their working in a clear and logical way. This helped examiners to award any credit due where final answers were not correct. Only a very small proportion of low attaining students presented weak scripts. Questions where working was often not so well organized included question 7 (algebraic expression for an area) and question 13 (manipulation of column vectors).

The paper gave the opportunity for students of all abilities to demonstrate positive achievement. All questions were accessible to some students but, as you would expect, there were relatively few students able to work confidently on all questions. In particular, questions 1 (Pythagoras' rule), 2 (formulae), 3 (ratio), 4 (best buy) and 11 (product rule for counting) were answered well by a large majority of students whereas questions, 10 (reverse percentage), 13 (column vectors), 14(a) (factorisation) and 15 (circle theorems) proved more of a challenge to the students in the attainment range they were aimed at.

REPORT ON INDIVIDUAL QUESTIONS

Question 1

This question provided a good start to the paper. Most students gained the full 2 marks. However, a significant number of students did not apply Pythagoras' rule correctly and instead added 4^2 and 8.5^2. These students could not be awarded any marks. It was surprising to see many students forget to take a square root and give the answer 56.25. A brief recap of the diagram might have alerted some students to there being an error in their working. In a small number of cases, trigonometry was attempted but the method was often incomplete and so no marks could be awarded.

Question 2

There was a disappointing proportion of fully correct answers to part (a) of this question. It has to be said that there was a great deal of sloppiness in students' notation when making the substitution and far too many students failed to write the −3 in brackets. They very often wrote $4 \times -3^2 - 11$, with most of these students then going on to use their calculators with little understanding of how to deal with the square of –3. This resulted in many answers of −47 instead of the correct 25. Students are advised to put brackets round any negative numbers when working with algebraic expressions

involving substitution both when writing down their working and when inputting any expression into their calculator.
Part (b) of the question was done quite well. Students usually changed the subject by subtracting 4 from both sides first and then dividing by 3. Only a small minority of students divided by 3 as their first step. Students who did do this often forgot to divide the 4 by 3. Students are reminded that they should give a complete answer on the answer line, including the "$p =$". A number of students just wrote "$\frac{d-4}{3}$" on the answer line.

Question 3

A majority of students scored full marks on this question. Many students in this higher tier paper used an algebraic approach, formulating and solving an equation, then using the solution to find a ratio and the value of p. Students usually worked accurately, but there were a number of students who wrote down a correct equation but collected terms in x to get $4x$ instead of the correct $5x$. Other students, having got the correct equation, $5x - 6 = 54$, then made an error in the first step to solve it, writing $5x = 48$.

A significant proportion of students adopted a purely numerical approach, finding, by trial and improvement, three numbers which satisfied the criteria given in the question, then using them to form a ratio. Some students found the number of counters for Rick and for Tony but did not form a ratio. There were also some other students who did form a ratio but did not write it in the ratio $1 : p$.

Question 4

A good discriminator between lower attaining students, this question attracted many fully correct solutions. Despite there being a number of possible routes that students could take, there were also many incorrect or incomplete solutions. The most common approach was to find the cost of 15 rolls of wallpaper at each of the shops, though credit was given for comparing any number of rolls, for example 1 roll or 5 rolls. A high proportion of students compared £180 with £184.80 and reached a correct conclusion to score all 4 marks. However, some students worked out the discount and took that to be the cost of the wallpaper at Style Papers.

A common error seen was, for students who did not appear to have access to a calculator, to use a build-up method to find 12%, for example by writing down 10%, 1% and 1%. Students who did this often made an error with the arithmetic and lost marks because they did not show the calculation, for example the addition of the three percentages or how they worked out the 10% or 1%. Students are reminded that when showing their calculation of a percentage, they should show a full method, so for example instead of writing 12% of £70 = £8.40, they should write $\frac{12}{100} \times 70 = 8.40$ or $0.12 \times 70 = 8.40$. This is so that in the event of an incorrect answer, marks for a correct method can be awarded.

Question 5

This question was quite well answered and descriptions of the mistakes made when the frequency polygon was drawn were generally clearly described. There were two mistakes to identify and students needed to describe them both to score the full 2 marks. Examiners were surprised that more students gained the mark for stating that the point plotted at (50, 5) should have been plotted at (45, 5) than gained the mark for stating that 40 had been omitted from the scale on the frequency axis. Any statement equivalent to one of these, and which was unambiguously stated, earned a mark. The most commonly seen unacceptable statements included not joining the polygon to (0, 0) and not joining the ends of the polygon.

Question 6

The majority of students made a good attempt at this problem on distance, time and average speed. Many students produced a concise, accurate and complete solution. Lower attaining students sometimes lost accuracy in their working because they used 0.6, 0.66 or another decimal approximation when coverting 40 minutes to hours. These students were still able to be given 3 of the 4 marks available if they had used and shown correct processes in their solutions. A significant number of students appeared to ignore the units in the question and were very fortunate to get a correct answer because the nature of the problem depended on the ratio of the times taken, rather than the times themselves. Of those students who answered the question incorrectly, many did so because they did not know the formula and divided the speed by the time, instead of multiplying them.

Question 7

This question was a good discrminator. There were a number of routes which could be followed in order to show the given result. The most popular method was to use the formula for the area of the trapezium *QUVR*. Once a student had found the length of *QU* (= $x + 5$), they could make substitutions into the formula then simplify it to show $A = 2x^2 + 20x$. Many students found this to be quite straightforward.

A second commonly seen and generally successful method was to use
Area of trapezium *QUVR* = Area of rectangle *TUVS* − Area of trapezium *TQRS*.
Perhaps the most straightforward way was to find the area of the rectangle with length and width $4x$ and 5 then add the area of a triangle of base x and height $4x$. It was surprising how few students took this approach.

The presentation of students answers was often not very clear with examiners having to search the working space for relevant working. However, many students did use algebra in a confident way to show the given result. The absence of brackets to make working clear sometimes led to a student losing marks but examiners were sympathetic where intentions were clear.

Question 8

Many students found this question to be straightforward. They often used a triangle where readings were easy to take and gave a value for the gradient of the graph within the accepted range, more often than not 0.14. For some other students who drew a small triangle on the line getting an accurate answer was less likely. However, if the value given for the gradient was outside the range 0.135 to 0.145, examiners awarded both marks in part (a) provided a correct method was clearly shown. A small number of students used "increase in x ÷ increase in y" or counted squares rather than using the scales on the axes. These students could not be credited with any marks in part (a).

Part (b) was answered less well and many incorrect or ambiguous statements were seen. An answer equivalent to "cost per unit of electricity" was expected but many students merely described what the graph, not the gradient, showed and statements such as "cost of how many units used" were often seen.

Question 9

Just over a quarter of all students completed part (a) of this question successfully to gain all 3 marks. A commonly seen final answer was 10^{120}. This could only score 1 mark. However, about two thirds of all students got as far as simplifying the expression to $\sqrt{10^{120}}$. Some of these students left this as their answer while other students went on to evaluate it incorrectly.

Answers to part (b) were often marred by a lack of clarity. Answers such as "Liam should have multiplied the numbers inside and outside the brackets" were commonplace but could not usually be rewarded with the mark available. Another common incorrect response was for a student to state that the 12 should be squared. More successful responses usually referred to the answer Liam should get, that is 12^{100} or that Liam should have multiplied 50 and 2, not worked out 50^2.

Question 10

This question was not well done. Too many students showed the misunderstanding that, to reverse a 10% decrease, they had to increase the value of the car by 10%, so they multiplied by 1.1 rather than using the correct method of dividing by 0.9. It was relatively unusual to award 1 or 2 marks for responses. Students usually either showed a good understanding or no understanding of the processes required to solve the problem.

Question 11

This question was generally well answered with most students scoring the full 2 marks. Where examiners saw errors, most either involved adding 16, 5 and 3 instead of multiplying, or multiplying the 16, 5 and 3 but then adjusting their answer by, for

example dividing by 3 or dividing by 2, as they mistakenly believed that the 240 included repeats. This latter approach was given some credit for the 16 × 5 × 3.

Question 12

In answering this question, most students showed a good understanding of how to apply trigonometrical ratios to this 2 stage problem. However, a significant proportion of them made errors in manipulating the ratio used in the second stage of the problem.

It was envisaged that students would use the trigonometry of right angled triangles to solve this problem and most did. However, a large number of students used the sine rule to get their answer. Examiners were surprised how few students realised that triangle *ACD* was isosceles and so they were not able to write down the length of AC without further calculation. However, they usually used tan 45 = $\frac{AC}{8}$ correctly to get 8 cm for the length of *AC*. Most students were then able to write down a relationship which would enable them to find the length of *AB*, and this was awarded some credit. However, too many of these students did not rearrange their equation correctly and instead of getting *AB* = $\frac{8}{\sin 20}$, wrote down *AB* = 8 × sin20, leading to an incorrect answer.

Question 13

This question was successfully answered by just under a half of all students who took this paper. Nearly all students scored a mark for finding 3**a** as a column vector. However, students often failed to write their responses in a clear, logical manner. Few students manipulated the equation to get $-2\mathbf{b} = \begin{pmatrix} 8 \\ 17 \end{pmatrix} - 3\mathbf{a}$ or $2\mathbf{b} = 3\mathbf{a} - \begin{pmatrix} 8 \\ 17 \end{pmatrix}$ before substituting for **a**. This might have led to a clearer approach and the avoidance of errors with signs. As it was, examiners were often left not knowing whether the student was trying to find the value of −2**b** or 2**b** with a consequent loss of marks for those students who did not get a fully correct final answer. Students who got one correct component of **b** were given the benefit of the doubt and rewarded with two of the three marks available.

Question 14

A minority of students scored full marks for their responses to this question. However, most students were able to gain some marks and the question was a good discriminator between students who could manipulate algebraic expressions with accuracy and those who made, sometimes unnecessary, errors.

A fully factorised expression was asked for in answering part (a) of this question. It was disappointing to see so few students factorise the expression fully and it was much more usual for students to gain just one of the two marks available for getting as far as, for example, $4(p^2 - 9)$ or $(2p - 6)(2p + 6)$.

Students usually scored well in part (b) which required the expansion of a product of three linear expressions to give a fully simplified cubic expression. Students generally showed their working in an organized way or used a grid to show terms in their products. Errors were usually restricted to incorrect terms or difficulties in dealing with the signs when collecting terms together, rather than a flawed strategy although some students omitted terms from their expansion. For students who did not give a fully correct answer, it was commonplace for them to earn 2 or 3 of the available marks in part (b).

Question15

There were some excellent complete, concise and clear proofs to this question involving the application of circle theorems. However, these were only seen in a minority of cases and students need to be aware of the need to state results clearly using the statements indicated on the mark scheme as guidance. There were many students who could not give reasons or who gave them with a lack of clarity. These students often gained some credit for matching up at least two pairs of equal angles. A clear indication of this on the diagram was accepted by examiners.

Of students who completed the proof successfully, most used statements of the same segment theorem together with a pair of vertically opposite angles. Students should note it was not enough to state "same segment theorem" – they needed to say what the theorem states, for example "angles in the same segment are equal". Some students used the result that angles in a triangle sum to 180° as a reason to support the equality of their third pair of angles. This was, of course, acceptable. A small number of students used the result that the angle at the centre of a circle is twice the angle at the circumference to show that angle PQX = angle PRS and that angle QPR = angle QSR, another acceptable approach.

About one in ten students obtained full marks for their proof.

Question 16

This question discriminated well between more able students sitting this paper with many students being awarded each of the 0, 1, 2 or 3 marks available. Some students did not attempt the question and many lower attaining students restricted their answer to working out the value of p by substituting $e = 6.8$ and $f = 0.05$. This gained no credit. Of those students who did score marks, many of them scored just one mark for writing down at least one of the bounds for e or f. Students who substituted bounds for e and f into the expression for p were split between those who calculated
$2[\text{UB of } e] \div [\text{LB of } f]$, the correct expression and those who calculated
$2[\text{UB of } e] \div [\text{UB of } f]$. Some students who had the correct fraction, failed to take the square root or made an error in taking the square root and so lost the final accuracy mark.

There was a significant number of students who calculated [UB of $2e$] ÷ [LB of f], that is $\frac{13.65}{0.045}$ instead of $\frac{13.7}{0.045}$. These students could not be awarded full marks even though their final answer worked out to be in the range 17.4 to 17.5

Question 17

In part (a) of this question, lower attaining students often drew a bar chart and did not recognise the need to find and use frequency densities. Some students used the midpoints of the class intervals erroneously together with the frequencies to get values which they then used to draw a diagram. Having said that, there was a good proportion of students who did successfully calculate frequency densities and use these to draw a histogram. It was interesting to note that a significant number of students drew the first four bars correctly but then made an error drawing the bar representing the interval $3.0 < d \leq 5.0$

Part (b) of the question was less well done with only a small proportion of students able to use the ratio of the areas of the two bars concerned to find a correct expression. Examiners did award one mark to a significant number of students who showed that they needed to compare the heights, widths and areas of the two bars concerned.

Question 18

It is encouraging to report that the majority of students gained some credit for their attempt at this question. The length of *TC* was often found successfully and a good proportion of students were also able to find the length of at least one of *SD* and *TD*. Only the higher attaining students were able to identify the angle which was required and put everything together in order to find the size of this angle, that is the size of angle *DTS*. Errors commonly seen included, substituting 12 and 14 incorrectly into the formula for the area of the cross section and making an error in the use of Pythagoras' rule to find the length of *DT*.

Question 19

Fully correct answers to this question were seen infrequently. However, a significant proportion of students taking this paper were able to score at least one mark for using a correct common denominator and writing at least one of the three terms with that common denominator, usually the first fraction as $\frac{3x(x-2)}{x^2-4}$. Only the best students were able to work accurately in order to gain more credit for their responses. The question discriminated well between high ability students aiming to get one of the two top grades. For many students, this question proved to be the most challenging question on the paper. In particular mistakes were often made with signs when the fraction $\frac{(x+2)(2x+1)}{x^2-4}$ was subtracted from the fraction $\frac{3x(x-2)}{x^2-4}$ or in writing $-(x^2-4)$ as $-x^2-4$.

Question 20

Though a significant number of students did not attempt this question, many of those students who did attempt it scored full marks for a complete and correct solution. The most common error seen was for students to write down a correct equation, $29\,600 = 24\,000a + 800$ but then make an error in solving the equation. It was not unusual for students to divide 29600 by 24800 as a first step, then subtract 800. Some students who found a correct value for "a" went on to use the iterative relationship correctly but only went as far as finding the profit made by the shop in the year 2020.

Question 21

Higher attaining students usually made a good start on this question on probability and identified the non-replacement nature of this problem. They usually presented at least one correct product of three fractions with denominators 9, 8 and 7 respectively. Only a small proportion of students gave all four possible ways of getting an even sum with the resultant correct answer, $\frac{11}{21}$. Students who drew a tree diagram were generally more successful at finding all 4 correct combinations.

Very few students approached the problem by finding the probability that the result was an odd number then subtracted from 1. Those students who did do this were rewarded appropriately.

Question 22

This question was accessible to and discriminated well between the most high attaining students sitting this paper. These students often gave a complete and fully correct solution and scored full marks. They usually used the approach of eliminating y to get an equation in x. This was more straightforward to deal with than the corresponding equation in y. The most common errors made by students included expanding $(2x - 5)^2$ incorrectly, often as $4x^2 + 25$, and making errors when re-arranging their quadratic equation to get all terms on one side of the equation. Some students who did get the correct quadratic equation to solve, made errors when trying to factorise and so restricted themselves to the award of 2 marks.

Summary

Based on their performance on this paper, students are offered the following advice:

- make sure you put negative numbers in brackets when substituting them into algebraic expressions or when using a calculator.
- check all your working for careless errors and to make sure you have not miscopied numbers or algebraic expressions either given in the question or in a previous line of your working.
- practise the addition and subtraction of algebraic fractions.
- check your working when expanding brackets and collecting terms, especially when finding the product of three linear expressions or when removing brackets after a subtraction of one expression from another.
- ensure you have a good understanding of how to use multipliers in problems concerning reverse percentages.
- practise drawing histograms and using them to solve problems.

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom