Chhetri Academy

chhetriacademy.com

GCSE MATHS

Geometry and measures

Edexcel 1MA1 | Subtopic collection Topic 49

Vectors

Higher tier

Answers and examiner comments

39	2004-2024	Higher
question groups	available papers	tier

Using this booklet

1. Answer numbers match the companion question booklet.
2. Follow the original question number on marking tables.
3. Examiner comments are included where available.

Name _____ **Date** _____

Some entries use full source marking material in an appendix. Any unavailable mark schemes are identified beside the relevant answer.

Original question 17

The full source mark scheme begins on page 75. Use the original paper code, tier and question number shown above.

Original question 22

No matching mark scheme was available in the supplied archive for this question.

Original question 22

The full source mark scheme begins on page 126. Use the original paper code, tier and question number shown above.

Original question 20

No matching mark scheme was available in the supplied archive for this question.

23		$360-(120+140+58)$	42	2	M1 $360-"(120+140+58)"$ or equivalent) or for (a + 58 + 120 + 140 = 360) oe seen A1 cao [Note: The subtraction MUST be from 360]

1380/1F

22	(a)		$\mathbf{b} - \mathbf{a}$	1	B1 cao
	(b)	$OP = OA + AP$ $= OA + \frac{2}{3}\ AB = \underline{\mathbf{a}} + \frac{2}{3}(\underline{\mathbf{b}} - \underline{\mathbf{a}})$	$\frac{1}{3}\mathbf{a} + \frac{2}{3}\mathbf{b}$	3	M1 for $\vec{OP} = \vec{OA} + \vec{AP}$ or $\vec{OP} = \vec{OB} + \vec{BP}$ M1 for $\vec{AP} = k\,(\mathbf{b} - \mathbf{a})$ ft from (a) with $0 < k < 1$ or $\vec{AP} = \frac{2}{3}\vec{AB}$ or $\vec{BP} = k\,(\mathbf{a} - \mathbf{b})$ ft from (a) with $0 < k < 1$ or $\vec{BP} = \frac{1}{3}\vec{BA}$ A1 for $\frac{1}{3}\mathbf{a} + \frac{2}{3}\mathbf{b}$ oe (must be in its simplest form)

1380 3H

1380/3H

Question	Working	Answer	Mark	Notes
27 (a)	$\overrightarrow{OP} = \mathbf{a} + \mathbf{b}$ $\overrightarrow{OM} = \frac{1}{2}\overrightarrow{OP}$	$\frac{1}{2}(\mathbf{a} + \mathbf{b})$	2	M1 for $\overrightarrow{OP} = \overrightarrow{OT} + \overrightarrow{TP}$ **or** $\overrightarrow{OM} = \frac{1}{2}\overrightarrow{OP}$ **or** $\overrightarrow{OM} = \frac{1}{2}\overrightarrow{OT} + \frac{1}{2}\overrightarrow{TP}$ **or** $\overrightarrow{OP} = \mathbf{a} + \mathbf{b}$ A1 for $\frac{1}{2}(\mathbf{a} + \mathbf{b})$ oe SC : B1 for $\mathbf{a} + \mathbf{b} \div 2$
(b)	$\overrightarrow{TO} + \overrightarrow{OM}$ $-\mathbf{a} + \frac{1}{2}(\mathbf{a} + \mathbf{b})$	$-\frac{1}{2}\mathbf{a} + \frac{1}{2}\mathbf{b}$	2	M1 for $-\mathbf{a}$ + "$\frac{1}{2}(\mathbf{a} + \mathbf{b})$" oe or $\overrightarrow{TM} = \overrightarrow{TO} + \overrightarrow{OM}$ or $\overrightarrow{TM} = \overrightarrow{TP} + \overrightarrow{PM}$ A1 ft

26	(a)	$\overrightarrow{AB} = \overrightarrow{AO} + \overrightarrow{OB}$	$-2\mathbf{a} + 3\mathbf{b}$	1	B1 for $-2\mathbf{a} + 3\mathbf{b}$ or $3\mathbf{b} - 2\mathbf{a}$
	(b)	$\overrightarrow{OP} = 2\mathbf{a} + \frac{2}{5}(3\mathbf{b} - 2\mathbf{a})$ $= \frac{6}{5}\mathbf{a} + \frac{6}{5}\mathbf{b}$ $= \frac{6}{5}(\mathbf{a} + \mathbf{b})$ parallel	$\frac{6}{5}(\mathbf{a} + \mathbf{b})$ is parallel to $\mathbf{a} + \mathbf{b}$	3	M1 for $2\mathbf{a} \pm \frac{2}{5}$('$3\mathbf{b} - 2\mathbf{a}$') OR $3\mathbf{b} \pm \frac{3}{5}$ ('$2\mathbf{a}$ - $3\mathbf{b}$') A1 for $\frac{6}{5}\mathbf{a} + \frac{6}{5}\mathbf{b}$ oe A1 for $\frac{6}{5}(\mathbf{a} + \mathbf{b})$ is parallel to $\mathbf{a} + \mathbf{b}$ oe

1380_3H

Question		Working	Answer	Mark	Notes
26	(a)		$\mathbf{b} - \mathbf{a}$	1	B1 for $\mathbf{b} - \mathbf{a}$ or $-\mathbf{a} + \mathbf{b}$
	(b)	$\overrightarrow{OP} = \overrightarrow{OA} + \overrightarrow{AP}$ $\overrightarrow{AP} = \frac{3}{4} \times (\mathbf{b} - \mathbf{a})$ $\overrightarrow{OP} = \mathbf{a} + \frac{3}{4} \times (\mathbf{b} - \mathbf{a})$ OR $\overrightarrow{OP} = \overrightarrow{OB} + \overrightarrow{BP}$ $\overrightarrow{BP} = \frac{1}{4} \times (\mathbf{a} - \mathbf{b})$ $\overrightarrow{OP} = \mathbf{b} + \frac{1}{4} \times (\mathbf{a} - \mathbf{b})$	$\frac{1}{4}(\mathbf{a} + 3\mathbf{b})$	3	B1 for $\frac{3}{4} \times$ '$(\mathbf{b} - \mathbf{a})$' M1 for $(\overrightarrow{OP} =) \overrightarrow{OA} + \overrightarrow{AP}$ or $(\overrightarrow{OP} =) \overrightarrow{OA} + \frac{3}{4}\overrightarrow{AB}$ or $\mathbf{a} \pm \frac{3}{4} \times$ '$(\mathbf{b} - \mathbf{a})$' A1 for $\frac{1}{4}(\mathbf{a} + 3\mathbf{b})$ or $\frac{1}{4}\mathbf{a} + \frac{3}{4}\mathbf{b}$ **OR** B1 for $\frac{1}{4} \times$ '$(\mathbf{a} - \mathbf{b})$' M1 for $(\overrightarrow{OP} =) \overrightarrow{OB} + \overrightarrow{BP}$ or $(\overrightarrow{OP} =) \overrightarrow{OB} + \frac{1}{4}\overrightarrow{BA}$ or $\mathbf{b} \pm \frac{1}{4} \times$ '$(\mathbf{a} - \mathbf{b})$' A1 for $\frac{1}{4}(\mathbf{a} + 3\mathbf{b})$ or $\frac{1}{4}\mathbf{a} + \frac{3}{4}\mathbf{b}$

15	(a)	i $CA = 2OA$ ii $BA = BO + OA = -\mathbf{b} + \mathbf{a}$ iii $BC = BO + OC = -\mathbf{b} - \mathbf{a}$	$2\mathbf{a}$ $\mathbf{a} - \mathbf{b}$ $-\mathbf{a} - \mathbf{b}$	3	B1 for $2\mathbf{a}$ oe B1 for $\mathbf{a} - \mathbf{b}$ oe B1 for $-\mathbf{a} - \mathbf{b}$ oe
	(b)	i $AX = AO + OX$ $= -\mathbf{a} + 2\mathbf{a} - \mathbf{b} = \mathbf{a} - \mathbf{b}$ ii $AX = BA$ so AX is parallel to BA; A is on both AX and BA, so B, A, X are all on a straight line	$\mathbf{a} - \mathbf{b}$ explanation	3	M1 for $AX = AO + OX$ A1 for $\mathbf{a} - \mathbf{b}$ oe B1 (dep on M1) for explanation eg $BX = BO + OX = -\mathbf{b} + 2\mathbf{a} - \mathbf{b} = 2(\mathbf{a} - \mathbf{b})$

5MB3H_01

Question		Working	Answer	Mark	Notes
28	(a)		$\mathbf{a} - 3\mathbf{b}$	1	B1 for $\mathbf{a} - 3\mathbf{b}$ oe
	(b)			4	M1 for (**NC** =) $2\mathbf{a} - 2\mathbf{b}$ oe M1 for (**NM** =) $\mathbf{b} + \frac{1}{2}"(\mathbf{a} - 3\mathbf{b})"$ A1 for $\frac{1}{2}(\mathbf{a} - \mathbf{b})$ oe and $2\mathbf{a} - 2\mathbf{b}$ oe C1 for **NC** is a multiple of **NM** (+ common point) **OR** M1 for (**NC** =) $2\mathbf{a} - 2\mathbf{b}$ oe M1 for (**MC** =) $\frac{1}{2}"(\mathbf{a} - 3\mathbf{b})" + \mathbf{a}$ A1 for $\frac{3}{2}(\mathbf{a} - \mathbf{b})$ oe and $2\mathbf{a} - 2\mathbf{b}$ oe C1 for **NC** is a multiple of **MC** (+ common point) **OR** M1 for (**NM** =) $\mathbf{b} + \frac{1}{2}"(\mathbf{a} - 3\mathbf{b})"$ M1 for (**MC** =) $\frac{1}{2}"(\mathbf{a} - 3\mathbf{b})" + \mathbf{a}$ A1 for $\frac{1}{2}(\mathbf{a} - \mathbf{b})$ oe and $\frac{3}{2}(\mathbf{a} - \mathbf{b})$ oe C1 for **NM** is a multiple to **MC** (+ common point)

18	(a)	$\begin{pmatrix}4\\8\end{pmatrix}-\begin{pmatrix}2\\4\end{pmatrix}$	$\begin{pmatrix}2\\4\end{pmatrix}$	2	M1 $\overrightarrow{OQ}-\overrightarrow{OP}$ in co-ordinates or vectors or $\begin{pmatrix}2\\y\end{pmatrix}$ or $\begin{pmatrix}x\\4\end{pmatrix}$ A1 cao [SC If no marks then B1 $\begin{pmatrix}4\\2\end{pmatrix}$ or $\begin{pmatrix}-2\\-4\end{pmatrix}$

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
18	(b)	$M = (3, 6)$ $N = (4, 8) + ½ (6, –4) = (7, 6)$ $\overrightarrow{MN} = \begin{pmatrix}7\\6\end{pmatrix} - \begin{pmatrix}3\\6\end{pmatrix}$ OR $\overrightarrow{MN} = ½\overrightarrow{PR}$ $\overrightarrow{PR} = \begin{pmatrix}6\\–4\end{pmatrix} + \begin{pmatrix}2\\4\end{pmatrix} = \begin{pmatrix}8\\0\end{pmatrix}$ OR $\overrightarrow{MN} = ½\overrightarrow{PQ} + ½\overrightarrow{QR}$ $\overrightarrow{MN} = ½ \begin{pmatrix}2\\4\end{pmatrix} + ½ \begin{pmatrix}6\\–4\end{pmatrix}$	$\begin{pmatrix}4\\0\end{pmatrix}$	3	B1 $M = (3, 6)$ M1 $N = (4, 8) + ½ (6, –4)$ or $(7, 6)$ or $\overrightarrow{MN} = \begin{pmatrix}7\\6\end{pmatrix} - \begin{pmatrix}3\\6\end{pmatrix}$ A1 cao OR B1 $\overrightarrow{MN} = ½\overrightarrow{PR}$ M1 ft $\overrightarrow{PR} = \begin{pmatrix}6\\–4\end{pmatrix} + \begin{pmatrix}'2'\\'4'\end{pmatrix}$ or $\begin{pmatrix}8\\0\end{pmatrix}$ A1 cao OR B1 $\overrightarrow{MN} = ½\overrightarrow{PQ} + ½\overrightarrow{QR}$ M1 ft $\overrightarrow{MN} = ½ \begin{pmatrix}'2'\\'4'\end{pmatrix} + ½ \begin{pmatrix}6\\–4\end{pmatrix}$ A1 cao

Question		Working	Answer	Mark	Notes
26	(a)		$6\mathbf{b} - 3\mathbf{a}$	1	B1 for $6\mathbf{b} - 3\mathbf{a}$ oe
	(b)			4	M1 for $\overrightarrow{AX} = \frac{1}{3}\overrightarrow{AB}$ or $\frac{1}{3}$ '$(6\mathbf{b} - 3\mathbf{a})$' or ft to $2\mathbf{b} - \mathbf{a}$ M1 for $\overrightarrow{OY} = \overrightarrow{OB} + \overrightarrow{BY} = 6\mathbf{b} + 5\mathbf{a} - \mathbf{b}$ $(= 5\mathbf{b} + 5\mathbf{a})$ oe M1 for $\overrightarrow{OX} = 3\mathbf{a} +$ '$2\mathbf{b} - \mathbf{a}$' $= 2\mathbf{a} + 2\mathbf{b}$ oe **Or** $\overrightarrow{OX} = 6\mathbf{b} - \frac{2}{3}$ '$(6\mathbf{b} - 3\mathbf{a})$' $(= 2\mathbf{a} + 2\mathbf{b})$ oe C1 for $\frac{2}{5}\overrightarrow{OY} = \frac{2}{5} \times 5(\mathbf{a} + \mathbf{b}) = 2(\mathbf{a} + \mathbf{b}) = \overrightarrow{OX}$

4.



18.



22



25(a)



Original question 23

The full source mark scheme begins on page 178. Use the original paper code, tier and question number shown above.

Question		Working	Answer	Mark	Notes
27	(a)		**a - b**	1	B1 for **a - b** oe
	(b)		$\frac{2}{5}\mathbf{a} + \frac{3}{5}\mathbf{b}$	3	M1 for a correct vector statement for $\overrightarrow{NR}$ eg. ($\overrightarrow{NR}$ =) $\overrightarrow{NQ} + \overrightarrow{QR}$ or ($\overrightarrow{NR}$ =) $\overrightarrow{NS} + \overrightarrow{SR}$ M1 for $\frac{2}{5}SQ$ (+ QR) or $\frac{3}{5}QS$ (+ SR) (SQ, QR, QS, SR may be written in terms of **a** and **b)** A1 for $\frac{2}{5}(\mathbf{a} - \mathbf{b}) + \mathbf{b}$ oe or $\frac{3}{5}(\mathbf{b} - \mathbf{a}) + \mathbf{a}$ oe

Question		Working	Answer	Mark	Notes
20	(a)		$4\mathbf{b} - 2\mathbf{a}$	1	B1 for $4\mathbf{b} - 2\mathbf{a}$ oe
	(b)	$\overrightarrow{ON} = \overrightarrow{OA} + \overrightarrow{AN}$, $2\mathbf{a} + \frac{3}{4}(4\mathbf{b} - 2\mathbf{a})$ $2\mathbf{a} + 3\mathbf{b} - 1.5\mathbf{a}$	$3\mathbf{b} + 0.5\mathbf{a}$	3	M1 for a correct vector for $\overrightarrow{ON}$, e.g. $(\overrightarrow{ON} =) \overrightarrow{OA} + \overrightarrow{AN}$, may be written in terms of **a** and **b** e.g. $(\overrightarrow{ON} =) 2\mathbf{a} + \frac{3}{4}(4\mathbf{b} - 2\mathbf{a})$ M1 for $(\overrightarrow{AN} =) \frac{3}{4}(4\mathbf{b} - 2\mathbf{a})$ oe or $(\overrightarrow{NB} =) \frac{1}{4}(4\mathbf{b} - 2\mathbf{a})$ oe or $(\overrightarrow{NA} =) \frac{3}{4}(2\mathbf{a} - 4\mathbf{b})$ oe or $(\overrightarrow{BN} =) \frac{1}{4}(2\mathbf{a} - 4\mathbf{b})$ oe A1 for $3\mathbf{b} + 0.5\mathbf{a}$

PAPER: 1MA0_1H

Question		Working	Answer	Mark	Notes
24	(a)	$\overrightarrow{AB} = -\mathbf{a} + \mathbf{b}$ $\overrightarrow{ON} = \overrightarrow{OA} + \frac{2}{3}\overrightarrow{AB}$ $\overrightarrow{ON} = \mathbf{a} + \frac{2}{3}(-\mathbf{a} + \mathbf{b})$ $= \frac{1}{3}\mathbf{a} + \frac{2}{3}\mathbf{b}$ OR $\overrightarrow{ON} = \overrightarrow{OB} + \frac{1}{3}\overrightarrow{BA}$ $\overrightarrow{ON} = \mathbf{b} + \frac{1}{3}(-\mathbf{b} + \mathbf{a})$ $= \frac{1}{3}\mathbf{a} + \frac{2}{3}\mathbf{b}$	$\frac{1}{3}\mathbf{a} + \frac{2}{3}\mathbf{b}$	3	M1 for correct vector equation involving $\overrightarrow{ON}$, eg. $\overrightarrow{ON} = \overrightarrow{OA} + \overrightarrow{AN}$, may be written, partially or fully, in terms of **a** and **b**, e.g. $(\overrightarrow{ON} =)$ $\mathbf{a} + \frac{2}{3}\overrightarrow{AB}$ M1 for showing answer requires $\overrightarrow{AN} = \frac{2}{3}\overrightarrow{AB}$ or $\overrightarrow{BN} = \frac{1}{3}\overrightarrow{BA}$ A1 $\frac{1}{3}\mathbf{a} + \frac{2}{3}\mathbf{b}$ oe
	(b)	$\overrightarrow{OD} = \overrightarrow{OA} + \overrightarrow{AC} + \overrightarrow{CD}$ $= \mathbf{a} + \mathbf{b} + \mathbf{b}$ $= \mathbf{a} + 2\mathbf{b}$ $\overrightarrow{OD} = 3(\frac{1}{3}\mathbf{a} + \frac{2}{3}\mathbf{b})$ $\overrightarrow{OD} = 3\overrightarrow{ON}$	Proof	3	M1 for a correct vector statement for $\overrightarrow{OD}$ or $\overrightarrow{ND}$ in terms of **a** and **b**, e.g. $\overrightarrow{OD} = \mathbf{a} + \mathbf{b} + \mathbf{b}$ oe or $\overrightarrow{ND} = \frac{2}{3}(-\mathbf{b} + \mathbf{a}) + \mathbf{b} + \mathbf{b}$ oe A1 for correct and fully simplified vectors for $\overrightarrow{ON}$ (may be seen in (a)) **and** for $\overrightarrow{OD}$ $(= \mathbf{a} + 2\mathbf{b})$ or $\overrightarrow{ND}$ $(= \frac{2}{3}\mathbf{a} + \frac{4}{3}\mathbf{b})$ C1 (dep on A1) for statement that $\overrightarrow{OD}$ or $\overrightarrow{ND}$ is a multiple of $\overrightarrow{ON}$ (+ common point)

Modifications to the mark scheme for Modified Large Print (MLP) papers.

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5º
Measurements of length: ±5 mm

PAPER: 1MA0_1H			
Question		**Modification**	**Notes**
2		*x* axis 2cm for 10,*y* axis 2cm for 2 ½ . Crosses changed to filled in circles. Right axis labelled.	Standard mark scheme
3		Model as well as diagram	Standard mark scheme
4	(a)	*x* changed to *e*. *y* changed to *f*.	M1 5*e* or 5 seen A1 cao
	(c)	*x* changed to *y*	Standard mark scheme
	(d)	*x* changed to *y*	Standard mark scheme

PAPER: 1MA0_1H			
Question		**Modification**	**Notes**
6	(a)	Move shape A two squares to the right. Cut left axis at -5. 2cm grid.	B1 for correct shape in correct position
	(b)	2cm grid. *x* axis -6 to +7. *y* axis -3 to +7.	Standard mark scheme
Q11		Remove 501 – 750g row. .	Standard mark scheme
Q12		2 cm grid. Axes drawn but not labelled	Standard mark scheme
Q18		Change line so it goes through (20, 0) (50, 20) (60, 30) (70, 40) (80,60) (100, 80). *x* axis 2cm for 10, *y* axis 2cm for 5.	B1 for 70 (accept 68 to 72) M1 for reading a value from graph at time = 60 (=30, accept 28 to 32) M1 for '30' ÷ 80 × 100 (= 37.5) or 25 ÷ 100 × 80 (= 20) C1 (dep on M2) for correct decision based on their figures OR M1 for 25 ÷ 100 × 80 (= 20) M1 for reading a value from graph at cf = 20 (=50, accept 48 to 52) C1 (dep on M2) for correct decision based on their figures
	(c)	28 seconds changed to 30 seconds. 96 seconds changed to 100 seconds. Boxplot *x* axis 3cm for 10 in an intermediate line.	B1 for ends of whiskers at 30 and 100 with a box B1 ft for median at '70' inside a box B1 for ends of box at 50 (accept 52 to 54) and 80 (accept 78 to 82)

PAPER: 1MA0_1H			
Question		**Modification**	**Notes**
Q23		Triangle A (the image) moved down 1 square. Triangle B (the enlargement) given scale factor - $\frac{1}{2}$ centre (0, -1). Wording: 'Describe fully the single transformation that will map Triangle A onto Triangle B'.	B1 for enlargement scale factor $-\frac{1}{2}$ B1 for (centre) (0, –1)
Q24		Vectors increased in size.	Standard mark scheme

19	(a)		$\mathbf{b} - \mathbf{a}$	1	B1 for $\mathbf{b} - \mathbf{a}$ oe
	(b)		$\frac{4}{5}\mathbf{a} + \frac{1}{5}\mathbf{b}$ oe	3	M1 for a fully correct vector statement for $\overrightarrow{OX}$ eg.($\overrightarrow{OX}$ =) $\overrightarrow{OA}$ (or $\mathbf{a}$) + $\overrightarrow{AX}$ (or $\frac{1}{5}\overrightarrow{AB}$) or ($\overrightarrow{OX}$ =) $\overrightarrow{OB}$ (or $\mathbf{b}$) + $\overrightarrow{BX}$ (or $\frac{4}{5}\overrightarrow{BA}$ or $-\frac{4}{5}\overrightarrow{AB}$) M1 (indep) for $\pm\frac{1}{5}(\mathbf{b} - \mathbf{a})$ or $\pm\frac{4}{5}(\mathbf{a} - \mathbf{b})$, ft their (a) A1 for $\mathbf{a} + \frac{1}{5}(\mathbf{b} - \mathbf{a})$ or $\mathbf{b} + \frac{4}{5}(\mathbf{a} - \mathbf{b})$ oe

PAPER: 5MB3H_01

19	(a)		$\mathbf{b} - \mathbf{a}$	1	B1 for $\mathbf{b} - \mathbf{a}$ oe
	(b)		$\frac{4}{5}\mathbf{a} + \frac{1}{5}\mathbf{b}$ oe	3	M1 for a fully correct vector statement for $\overrightarrow{OX}$ eg.($\overrightarrow{OX}$ =) $\overrightarrow{OA}$ (or $\mathbf{a}$) + $\overrightarrow{AX}$ (or $\frac{1}{5}\overrightarrow{AB}$) or ($\overrightarrow{OX}$ =) $\overrightarrow{OB}$ (or $\mathbf{b}$) + $\overrightarrow{BX}$ (or $\frac{4}{5}\overrightarrow{BA}$ or $-\frac{4}{5}\overrightarrow{AB}$) M1 (indep) for $\pm\frac{1}{5}(\mathbf{b} - \mathbf{a})$ or $\pm\frac{4}{5}(\mathbf{a} - \mathbf{b})$, ft their (a) A1 for $\mathbf{a} + \frac{1}{5}(\mathbf{b} - \mathbf{a})$ or $\mathbf{b} + \frac{4}{5}(\mathbf{a} - \mathbf{b})$ oe

PAPER: 5MB3H_01

Original question 24

The full source mark scheme begins on page 193. Use the original paper code, tier and question number shown above.

Paper: 5MB3H_01

Question		Working	Answer	Mark	Notes
19			proof	4	B1 for $\overrightarrow{AM}$ = 0.5**b** or $\overrightarrow{MC}$ = 0.5**b** or $\overrightarrow{BC}$ = **a** or $\overrightarrow{CX}$ = **a** or $\overrightarrow{BX}$ = 2**a** Note: This could be shown on the diagram or in a correct vector expression M1 for a correct relevant vector expression for $\overrightarrow{OM}$ or $\overrightarrow{MX}$ or $\overrightarrow{OX}$ eg $\overrightarrow{OM}$=$\overrightarrow{OA}$ + $\overrightarrow{AM}$ or $\overrightarrow{OX}$= **b** + 2**a** A1 for any two from $\left(\overrightarrow{OM}\right)$ = **a** + 0.5**b**, $\left(\overrightarrow{MX}\right)$ = **a** + 0.5**b** and $\left(\overrightarrow{OX}\right)$ = **b** + 2**a** B1 for a fully correct proof, eg." $\overrightarrow{OX}$ = 2$\overrightarrow{OM}$ so the vectors are parallel and have a common point *O* " OR (geometric proof) M1 for <OAM = <MCX or OA = CX and AM = CM A1 for <OAM = <MCX with reason (alternate angles) and OA = CX and AM = CM B1 for ΔOAM ≡ ΔXCM with reason, eg SAS B1 for correct proof, eg <AMO = <CMX with reason (vertically opposite angles)

27		$\overrightarrow{AB} = \overrightarrow{AO} + \overrightarrow{OB}$ $= \mathbf{a} + \mathbf{b}$ $\overrightarrow{AC} = \frac{7}{2}\overrightarrow{AB}$ $\overrightarrow{OC} = \overrightarrow{OA} + \overrightarrow{AC}$ $= 2\mathbf{a} + \mathbf{b} + \frac{7}{2}(\mathbf{a} + \mathbf{b})$	$\frac{11}{2}\mathbf{a} + \frac{9}{2}\mathbf{b}$	4	M1 for $\overrightarrow{AB} = \overrightarrow{AO} + \overrightarrow{OB}$ $(= -(2\mathbf{a} + \mathbf{b}) + (3\boldsymbol{a} + 2\boldsymbol{b}))$ or $\mathbf{a} + \mathbf{b}$ M1 for $\overrightarrow{AC} = \frac{7}{2}\overrightarrow{AB}$ or $\overrightarrow{BC} = \frac{5}{2}\overrightarrow{AB}$, may be in terms of **a** and **b** M1 (dep M2) for complete method to find $\overrightarrow{OC}$ in terms of **a** and **b** A1 for $\frac{11}{2}\mathbf{a} + \frac{9}{2}\mathbf{b}$ or equivalent simplest form (SCB2 for $\frac{11}{2}\mathbf{a} + \frac{23}{2}\mathbf{b}$ or $\frac{11}{2}\mathbf{a} + \frac{19}{2}\mathbf{b}$)

PAPER: 1MA0_2H

23		$\frac{3}{4}(\mathbf{b} + \mathbf{a})$	4	B1 for $\mathbf{b} - \mathbf{a}$ or $\frac{1}{2}(\mathbf{b} - \mathbf{a})$ or $\mathbf{a} - \mathbf{b}$ or $\frac{1}{2}(\mathbf{a} - \mathbf{b})$ M1 for $\overrightarrow{OM} = \mathbf{a} + \frac{1}{2}(\mathbf{b} - \mathbf{a})$ or $\mathbf{b} + \frac{1}{2}(\mathbf{a} - \mathbf{b})$ oe M1 for $\overrightarrow{ON} = \frac{3}{2}(\mathbf{a} + \frac{1}{2}(\mathbf{b} - \mathbf{a}))$ or $\frac{3}{2}(\mathbf{b} + \frac{1}{2}(\mathbf{a} - \mathbf{b}))$ A1 for $\frac{3}{4}(\mathbf{b} + \mathbf{a})$ oe (simplified)

Modifications to the mark scheme for Modified Large Print (MLP) papers.

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q4		Diagrams have been removed.	M1 for one of 439 ÷ 150 (= 2.92), 639 ÷ 200 (= 3.19), 795 ÷ 275 (= 2.89) OR one of 150 ÷ 439 (= 0.341), 200 ÷ 639 (= 0.31), 275 ÷ 795 (= 0.345) OR any other calculation that could lead to a comparative figure M1 for calculations that could lead to comparative figures for 2 jars M1 for calculations that could lead to comparative figures for 3 jars eg all 3 from the above lists C1 for correct comparative figures for all 3 jars leading to a correctly stated comparison: 275g best value
Q5	(a)	Wording added: 'There are three spaces to fill.' Table produced in vertical format.	B2 for all three correct values 7, -2, -1 (B1 for two correct values 7, -2 or -1)
Q5	(b)	1½ cm grid provided.	B2 for fully correct curve (B1 ft for at least 5 points plotted correctly)

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q8		Size of diagram kept as original. Scale put on diagram as well as in text.	M1 for perpendicular bisector of the line AC M1 for arc of circle centre D radius 7cm A1 for correct point
Q9		Diagram has been enlarged. Dotty shading. Measurement labels moved to be more accessible for candidates.	M1 for correct method to find frame dimensions 15+5+5 (=25) and 10+5+5 (=20) M1 for finding relationship between length and width for both eg 15 ÷ 10 (=1.5) and 25 ÷ 20 (= 1.25) or for correct use of a scale factor to find a comparable length C1 for correct conclusion following comparison of 2 correct comparable figures
Q10		The grid has been enlarged. Shapes are labelled triangle A and triangle B. Wording added: 'It shows two triangles on the grid, triangle A and triangle B.' Dotty shading	B1 for rotation B1 for 90° clockwise or 270° anticlockwise B1 for (1,4) NB Award B0 if more than one transformation given

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q11		The diagram has been enlarged. 2 models provided for all candidates. A diagram is also provided for MLP.	M1 for $\pi \times 3.5^2 \times 10$ (=384.85 or =122.5π) or $\pi \times 6^2$ (=113.1 or =36π) M1 (dep) for "384.85" ÷ "113.1" or 122.5(π) ÷ 36(π) A1 for 3.4 – 3.5 OR M1 for $\left(\frac{12}{7}\right)^2$ (=2.9) or $\left(\frac{7}{12}\right)^2$ (=0.34) M1 for 10 ÷ "$\left(\frac{12}{7}\right)^2$" or 10 × "$\left(\frac{7}{12}\right)^2$" A1 for 3.4 – 3.5
Q13		The diagram has been enlarged.	M1 for $8^2 + 5^2$ or 64 + 25 or 89 M1 (dep) $\sqrt{"8^2"+"5^2"}$ (=9.4) M1 for "9.4..." × π A1 for 29.5 – 29.65
Q14		The diagram has been enlarged. Wording added: 'Angle CAB is marked x'.	M1 for 180 - 56 - 75 A1 for 49 C1 for alternate segment theorem and angles on a straight line add up to 180° OR alternate segment theorem and angles in a triangle add up to 180° Appropriate to methods shown

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q16		MLP: x changed to e. y changed to f.	M1 for intention to multiply by $e + 5$ M1 for intention to isolate yx and $3x$ on one side to get $3e - ef$ oe A1 for $x = \frac{5f}{3-f}$ or $\frac{-5f}{f-3}$
Q17		The diagram has been enlarged.	M1 for $360 \div 5$ (=72) M1 (dep) for $\frac{1}{2} \times 6^2 \times \sin 72°$ (=17.12) M1 for completing full method to find total area of pentagon A1 for 85.5 – 85.6 OR M1 for $360 \div 10$ (=36) or $\frac{1}{2}(180 - 360 \div 5)$ (=54) M1(dep) for eg $6 \times \sin 36° \times 6 \times \cos 36°$ (=17.12) or $\frac{1}{2} \times 6 \times \sin 54° \times 6 \times \cos 54°$ (=8.55) M1 for completing full method to find total area of pentagon A1 for 85.5 – 85.6
Q19		MLP and braille: a changed to x. b changed to f.	B1 for 45 or 0.15 M1 for $45 \div 0.15$ A1 for 300 cao

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q21	(a)	The grid has been enlarged.	M1 for any translation along x axis A1 correct position
Q21	(b)	The grid has been enlarged.	B1 for $y = -g(x)$ oe from correct interpretation of diagram
Q22		The diagram has been enlarged. TR is joined by a dashed line.	M1 for a method to find an angle, $RAB = 70$, $ABR = 50$, $BRA = 60$ or $TAR = 20$ M1 for substitution into sine formula $\frac{AR}{\sin"50"} = \frac{12}{\sin"60"}$ M1 for use of sine rule to find AR, $AR = \frac{12}{\sin"60"} \times \sin"50"$ (=10.61) M1 for substitution into cosine formula $TR^2 = 5^2 + "10.61"^2 - 2 \times 5 \times "10.61" \times \cos 20$ (=37.92) A1 for 6.15 - 6.2
Q23		The diagram has been enlarged. OM is joined by a dashed line.	B1 for $\mathbf{b} - \mathbf{a}$ or $\frac{1}{2}(\mathbf{b} - \mathbf{a})$ or $\mathbf{a} - \mathbf{b}$ or $\frac{1}{2}(\mathbf{a} - \mathbf{b})$ M1 for $\overrightarrow{OM} = \mathbf{a} + \frac{1}{2}(\mathbf{b} - \mathbf{a})$ or $\mathbf{b} + \frac{1}{2}(\mathbf{a} - \mathbf{b})$ oe M1 for $\overrightarrow{ON} = \frac{3}{2}(\mathbf{a} + \frac{1}{2}(\mathbf{b} - \mathbf{a}))$ or $\frac{3}{2}(\mathbf{b} + \frac{1}{2}(\mathbf{a} - \mathbf{b}))$ A1 for $\frac{3}{4}(\mathbf{b} + \mathbf{a})$ oe (simplified)

Original question 20

The full source mark scheme begins on page 213. Use the original paper code, tier and question number shown above.

Question	Working	Answer	Mark	Notes
23 (a)(i)		$\mathbf{a} + \mathbf{b}$	2	B1 for $\mathbf{a} + \mathbf{b}$ oe
(ii)		$-\mathbf{a} + 3\mathbf{b}$		B1 for $-\mathbf{a} + 3\mathbf{b}$ oe
(b)		$\frac{3}{4}\mathbf{a} + \frac{3}{4}\mathbf{b}$	2	M1 for $\overrightarrow{OP} + \frac{1}{4}\overrightarrow{PR}$ or $\overrightarrow{OR} + \frac{3}{4}\overrightarrow{RP}$ (may be in terms of **a** and **b)** A1 for $\frac{3}{4}\mathbf{a} + \frac{3}{4}\mathbf{b}$ or $\frac{3}{4}(\mathbf{a} + \mathbf{b})$
*(c)		$OS = \frac{3}{4}OT$	2	C2 (dep A1) for S divides OT in the ratio 3:1 oe or $OS = \frac{3}{4}OT$ oe (C1 (dep A1) for S lies on OT or that OT and PR intersect at S oe)

Original question 21

The full source mark scheme begins on page 229. Use the original paper code, tier and question number shown above.

Question	Working	Answer	Mark	Notes
23		Correct conclusion from correct working	4	B1 for $\overrightarrow{AB} = -5\mathbf{a} + 2\mathbf{b}$ or $\overrightarrow{BA} = 5\mathbf{a} - 2\mathbf{b}$ M1 for a correct vector statement for $\overrightarrow{OT}$ eg $\overrightarrow{OA} + \overrightarrow{AT}$ or $\overrightarrow{OB} + \overrightarrow{BT}$ or $\overrightarrow{OA} + \frac{5}{6}\overrightarrow{AB}$ or $\overrightarrow{OB} + \frac{1}{6}\overrightarrow{BA}$, may be written partially or fully in terms of **a** and **b** M1 for $5\mathbf{a} + \frac{5}{6}(-5\mathbf{a} + 2\mathbf{b})$ oe or $2\mathbf{b} + \frac{1}{6}(5\mathbf{a} - 2\mathbf{b})$ oe A1 for $\frac{5}{6}(\mathbf{a} + 2\mathbf{b})$ is parallel to $\mathbf{a} + 2\mathbf{b}$

PAPER: 1MA0/2H				
Question	**Working**	**Answer**	**Mark**	**Notes**
*24	$(n^2 + 4n + 4) - (n^2 + 2n + 1)$ $\frac{2n+3}{2n^2+3n}$ $\frac{2n+3}{n(2n+3)}$	Proof	4	M1 for correct method to expand $(n + 2)^2$ or $(n + 1)^2$ M1 for correct simplification of numerator M1 for factorisation of $2n^2 + 3n$ **or** for clearing the fractions on both sides correctly C1 for complete and correct proof OR M1 for $\{(n+2)-(n+1)\}\{(n+2)+(n+1)\}$ M1 for $1 \times (2n+3)$ M1 for factorisation of $2n^2 + 3n$ **or** for clearing the fractions on both sides correctly C1 for complete and correct proof OR M1 for $n\{(n+2)^2-(n+1)^2\} = (2n^2 + 3n) \times 1$ M1 for $n(n+2)^2 - n(n+1)^2$ or for correct expansion of $(n+2)^2-(n+1)^2$ M1 for correct expansion of $n\{(n+2)^2-(n+1)^2\}$ C1 for complete and correct proof (must include statement recognising the equality of LHS and RHS)

Modifications to the mark scheme for Modified Large Print (MLP) papers.

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 1MA0_2H			
Question		**Modification**	**Notes**
1		Diagram enlarged. 4th horizontal line added to the bottom of the stem and leaf diagram. Key moved above and to the left of the diagram. Wording added 'It shows the basis of a stem and leaf diagram.	Standard mark scheme
3		Diagram enlarged. Wording added 'Angle $B\,O\,C = 4x + 31$'. Wording added 'Angle $A\,O\,C = 6x - 21$'.	Standard mark scheme
4		Diagram enlarged. Spike removed from the spinner and the spinner has been straightened.	Standard mark scheme
7		Grid enlarged.	Standard mark scheme
9		Diagram enlarged. Wording added '$CB = 200$ cm'. Wording added '$DC = 60$ cm'.	Standard mark scheme

PAPER: 1MA0_2H			
Question		**Modification**	**Notes**
10		Pictures have been removed.	Standard mark scheme
11		Diagram enlarged.	Standard mark scheme
12	(b)	MLP only: x changed to y.	B2 for $3y^2 + 4y$ or $y(3y + 4)$ (B1 for $y^2 - 2y$ or $2y^2 + 6y$ or $3y^2 + ny$ or $py^2 + 4y$)
	(d)	'a' changed to 's' and 'b' changed to 't' for all candidates.	B2 for $9st(2 + 3t)$ (B1 for $9s\,(2t + 3t^2)$ or $9t(2s + 3st)$ or $st(18 + 27t)$ or $3st(6 + 9t)$ or $3s(6t + 9t^2)$ or $3t(6s + 9st)$ or $9st$(a two term algebraic expression))
13	(a)	Wording added 'There are five spaces to fill.'	Standard mark scheme
	(b) to (d)	Diagram enlarged. Axes labels have been moved to the top of the vertical axis and to the left of the horizontal axis. Right axis has been labelled.	Standard mark scheme
14		Diagram enlarged.	Standard mark scheme
15		Diagram enlarged. Wording 'AB = 15 m' changed to 'BA = 15 metres'. Wording 'BC = 24 m' changed to 'CB = 24 metres'. Wording added 'Angle CBD = angle BDA = 90°.'	Standard mark scheme

PAPER: 1MA0_2H			
Question		**Modification**	**Notes**
18		Diagram enlarged.	Standard mark scheme
19	(b)	Braille only: e changed to m.	M1 for $(m \pm 3)(m \pm 4)$ A1 for $(m - 3)(m + 4)$
21		Simplified diagram provided for MLP candidates.	Standard mark scheme
22	(a) (b)	Grid enlarged. Grid enlarged.	Standard mark scheme Standard mark scheme
23		Diagram enlarged. Cross at T changed to a solid dot.	Standard mark scheme

Question	Working	Answer	Mark	Notes
24 (a)		$3\mathbf{a} + 3\mathbf{b}$	2	M1 for $\vec{AB} = 6\mathbf{b} - 6\mathbf{a}$ oe or $\vec{BA} = 6\mathbf{a} - 6\mathbf{b}$ oe or $(\vec{OM}) = \frac{1}{2}(6\mathbf{a} + 6\mathbf{b})$ oe A1 cao
*(b)	Method 1 Show that $\vec{AG}$ is parallel to $\vec{AN}$	Shown	4	M1 for a method to find $\vec{OG}$ or $\vec{GM}$ in terms of **a** and **b** eg $\vec{OG} = \frac{2}{3}$ "$(3\mathbf{a} + 3\mathbf{b})$" $(= 2\mathbf{a} + 2\mathbf{b})$ eg $\vec{GM} = \frac{1}{3}$ "$(3\mathbf{a} + 3\mathbf{b})$" $(= \mathbf{a} + \mathbf{b})$ M1 for a method to find $\vec{AN}$ in terms of **a** and **b** eg $-6\mathbf{a} + 6\mathbf{b} \div 2$ $(= 3\mathbf{b} - 6\mathbf{a})$ M1 for a method to find $\vec{AG}$ in terms of **a** and **b** eg $-6\mathbf{a}$ + "$2\mathbf{a} + 2\mathbf{b}$" $(= 2\mathbf{b} - 4\mathbf{a})$ C1 for correct simplified expressions in terms of **a** and **b** for $\vec{AG}$ and $\vec{AN}$ followed by conclusion OR
	Method 2 Show that $\vec{AG}$ is parallel to $\vec{GN}$			M1 for a method to find $\vec{OG}$ in terms of **a** and **b** eg $\frac{2}{3}$ "$(3\mathbf{a} + 3\mathbf{b})$" $(= 2\mathbf{a} + 2\mathbf{b})$ M1 for a method to find $\vec{GN}$ in terms of **a** and **b** eg $-$"$(2\mathbf{a} + 2\mathbf{b})$" $+ 6\mathbf{b} \div 2$ $(= \mathbf{b} - 2\mathbf{a})$ M1 for a method to find $\vec{AG}$ in terms of **a** and **b** eg $\frac{1}{2}(6\mathbf{b} - 6\mathbf{a}) - \frac{1}{3}$ "$(3\mathbf{a} + 3\mathbf{b})$" $(= 2\mathbf{b} - 4\mathbf{a})$ C1 for correct simplified expressions in terms of **a** and **b** for $\vec{AG}$ and $\vec{GN}$ followed by conclusion

8a



8b



13a



3a



16


200
180
160
140
120
100
80
60
40
20
0
20
40
60
80
100
120
x

11


P
S (104, 105.5)
R
(106, 105)
A
Q
(106, 103)

Modifications to the mark scheme for Modified Large Print (MLP) papers.

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5º
Measurements of length: ±5 mm

PAPER: 1MA0_1H			
Question		**Modification**	**Mark scheme notes**
1	(a)	MLP only: *x* changed to *e*.	Standard mark scheme but *x* changed to *e* for MLP
3		Number 6 on the frequency column of the table changed to number 5. Diagram enlarged. Right axis has been labelled. Axes labels have been moved to the left of the horizontal axis and above the vertical axis.	Standard mark scheme
4		MLP only: *x* changed to *y*.	Standard mark scheme but *x* changed to *y* for MLP
5		Diagram enlarged. Angle sizes moved outside of the angle arcs; arcs have been made smaller.	Standard mark scheme
7		See below	
8	(a)	Question reversed. Triangle S has been rotated by anticlockwise 90° at the origin and has been labelled triangle T. Question wording changed to 'It shows triangle S and triangle T given on a grid. Describe fully the single transformation that maps triangle S onto triangle T.' 3 answer lines have been provided. Dotty shading has been used.	New mark scheme: B1 for Rotation B1 for 90° anticlockwise about origin [(0,0)]
8	(b)	Diagram enlarged. Shading has changed to dotty shading. Shapes labelled 'triangle Q' and 'triangle P'. Wording added 'It shows triangle Q and triangle P given on a grid.' Question wording 'shape' changed to 'triangle'. *x* axis cut so it finishes at 10. The grid to the left and below the axis is removed.	Standard mark scheme
12		Model provided for all candidates. Diagram enlarged and also provided for MLP. Arrow heads removed and a dot has been added to the centre.	Standard mark scheme
13		In (a) Table has been turned to vertical format and left aligned. Wording added 'There are three spaces to fill.'. In (b) Diagram enlarged.	Standard mark scheme
15		Diagram enlarged. Arrow removed from 6 cm. Arrow and line for 10 cm now a dashed line.	Standard mark scheme

<table>
<tr><th colspan="3">PAPER: 1MA0_1H</th></tr>
<tr><th colspan="2">Question</th><th>Modification</th><th>Mark scheme notes</th></tr>
<tr><td>16</td><td>(a)</td><td>Numbers on the frequency table changed to: 40, 70, 50, 20, 10, 10.</td><td>(a) award B1 for 40, 110, 160, 180, 190, 200
In (b) standard mark scheme (with the above figures) but apply extended tolerance to the plotting of their points.</td></tr>
<tr><td>16</td><td>(c)</td><td>Diagram enlarged. Right axis has been labelled. Axes labels have been moved to the left of the horizontal axis and above the vertical axis.
On larger print versions the horizontal axis is labelled 0, 20, 40, etc. due to lack of space
The number 54 has been changed to 50.</td><td>For (c): M1 indicates using diagram with a mark of 50 eg draws line from 50 to their cumulative frequency graph, or sght of 160 eg 200 − 160
A1 for 40 or ft on a cumulative frequency graph</td></tr>
<tr><td>17</td><td></td><td>MLP only: x changed to y.</td><td>Standard mark scheme with x replaced with y</td></tr>
<tr><td>19</td><td></td><td>Diagram enlarged. Points on the Male box plot moved to: 10, 30, 40, 60, 100. Points on the Female box plot moved to: 0, 20, 40, 70, 110. Horizontal axis label moved to the left of the axis.

</td><td>Standard mark scheme for the award of marks, but not the different figures they might use as shown below.
<table><tr><td></td><td>Male</td><td>Female</td></tr><tr><td>Median</td><td>40</td><td>40</td></tr><tr><td>Range</td><td>90</td><td>110</td></tr><tr><td>IQR</td><td>30</td><td>50</td></tr></table></td></tr>
<tr><td>22</td><td></td><td>See below</td><td></td></tr>
<tr><td>24</td><td></td><td>Diagram enlarged. Crosses at G and N changed to a solid dots.</td><td>Standard mark scheme.</td></tr>
</table>

PAPER: 1MA0_1H			
Question		**Modification**	**Mark scheme notes**
7		 Diagram changed as shown. Model provided for all candidates. Diagram also enlarged and provided for MLP. Cross section of the prism also provided. Key provided for the cross section of the prism. Braille only: will label the cross section R P Q. Cross section of prism h = height 40 cm 	Standard mark scheme

		arc has been made smaller. An enlarged part of the diagram has also been provided and labelled 'Diagram (ii)'. Wording added 'Diagram (ii) shows an enlarged section of part of the diagram.'	

Question 22

Diagram (i)



Diagram (ii)



Question	Answer	Mark	Mark scheme	Additional guidance
10 (a)	Diagram	B1	for correct vector drawn including arrow	May be drawn anywhere on the grid. Condone missing label
(b)	$\begin{pmatrix} 3 \\ -4 \end{pmatrix}$	M1	for **a** + 2**b** drawn with resultant vector **or** for writing **a** and **b** as column vectors **and** attempt to add **a** + 2**b** , eg $\begin{pmatrix} 1 \\ 2 \end{pmatrix} + 2 \times \begin{pmatrix} 1 \\ -3 \end{pmatrix}$ **or** $\begin{pmatrix} 1+2 \\ c \end{pmatrix}$ **or** $\begin{pmatrix} d \\ 2+-6 \end{pmatrix}$ **or** $\begin{pmatrix} -4 \\ 3 \end{pmatrix}$	Accept consistent incorrect notation for M1
		A1	cao	

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
21	Proof	M1	for $\overrightarrow{DQ}$ = ½ (**b** − **a**) oe or $\overrightarrow{EQ}$ = ½ (**a** − **b**) oe	Vectors could be written on the diagram
		M1	for $\overrightarrow{PQ}$ = ½ **a** + $\overrightarrow{DQ}$ **or** ½ **a** + ½ (**b** − **a**) oe **or** $\overrightarrow{PQ}$ = − ½ **a** + **b** + $\overrightarrow{EQ}$ **or** − ½ **a** + **b** + ½ (**a** − **b**) oe	
		B1	for $\overrightarrow{PQ}$ = ½ **b**	
		C1	for complete proof with statement, eg $FE = 2PQ$ **or** FE is a multiple of PQ **or** **b** = 2(½ **b**)	

6	$\begin{pmatrix}-9\\14\end{pmatrix}$	M1 A1	for $2\begin{pmatrix}3\\4\end{pmatrix}-3\begin{pmatrix}5\\-2\end{pmatrix}$ **or** $\begin{pmatrix}6\\8\end{pmatrix}$ **and** $\begin{pmatrix}15\\-6\end{pmatrix}$ **or** $\begin{pmatrix}-9\\y\end{pmatrix}$ **or** $\begin{pmatrix}x\\14\end{pmatrix}$ cao	May be seen in two separate calculations eg 2×3 + –3×5 and 2×4 + –3×–2 Condone incorrect notation if method is clear for this mark only

Question	Answer	Mark	Mark scheme	Additional guidance
18	$\frac{2}{5}\mathbf{a}+\mathbf{b}$	P1	for relationship involving D eg $\overrightarrow{OD}=\frac{2}{5}\overrightarrow{OB}$ or $\overrightarrow{DB}=\frac{3}{5}\overrightarrow{OB}$ **or** for relationship involving E eg $\overrightarrow{BE}=\frac{1}{5}\overrightarrow{BC}$ or $\overrightarrow{EC}=\frac{4}{5}\overrightarrow{BC}$	
		P1	for relationship involving D in terms of **a** and **b** eg $\overrightarrow{OD}=\frac{2}{5}(\mathbf{a}+\mathbf{b})$ or $\overrightarrow{DB}=\frac{3}{5}(\mathbf{a}+\mathbf{b})$ **or** for relationship involving E in terms of **a** and **b** eg $\overrightarrow{BE}=\frac{1}{5}(-\mathbf{b}-\mathbf{a}+3\mathbf{b})$ oe or $\overrightarrow{EC}=\frac{4}{5}(-\mathbf{b}-\mathbf{a}+3\mathbf{b})$ oe **or** $\overrightarrow{BC}=2\mathbf{b}-\mathbf{a}$ oe or $\overrightarrow{CB}=\mathbf{a}-2\mathbf{b}$ oe	
		P1	(dep P2) for expression for $\overrightarrow{DE}$ in terms of **a** and **b** eg $\overrightarrow{DE}=\frac{3}{5}(\mathbf{a}+\mathbf{b})+\frac{1}{5}(-\mathbf{b}-\mathbf{a}+3\mathbf{b})$	
		A1	for $\frac{2}{5}\mathbf{a}+(1)\mathbf{b}$ or $\frac{1}{5}(2\mathbf{a}+5\mathbf{b})$	

Paper: 1MA1/3H

Question	Answer	Mark	Mark scheme	Additional guidance
13	$\begin{pmatrix}-1\\4\end{pmatrix}$	M1	for $3\begin{pmatrix}2\\-3\end{pmatrix} - 2\mathbf{b} = \begin{pmatrix}8\\-17\end{pmatrix}$ oe or for $\begin{pmatrix}6\\-9\end{pmatrix}$	
		M1	for $2\mathbf{b} = \begin{pmatrix}3\times 2-8\\3\times -3--17\end{pmatrix} \left(=\begin{pmatrix}-2\\8\end{pmatrix}\right)$ oe or $-2\mathbf{b} = \begin{pmatrix}8-3\times 2\\-17-3\times -3\end{pmatrix} \left(=\begin{pmatrix}2\\-8\end{pmatrix}\right)$ oe or for one element correct, –1 or 4	Must see "$2\mathbf{b}$ = ..." or "$-2\mathbf{b}$ = **...**" to award 2 marks One correct element scores 2 marks
		A1	cao (if M0 scored, SC B1 for $\begin{pmatrix}1\\-4\end{pmatrix}$)	

Question	Answer	Mark	Mark scheme	Additional guidance
24	Proof	C1	for $\overrightarrow{CE} = 2\mathbf{a} - \mathbf{b}$ oe	Vectors may be seen on diagram. Award marks provided not ambiguous. For the award of the first 3 marks, expressions for vectors $\overrightarrow{CE}$, $\overrightarrow{EP}$, $\overrightarrow{CP}$, $\overrightarrow{CF}$ and $\overrightarrow{DP}$ may not be simplified
		C1	for $\overrightarrow{EP} = 2\mathbf{a} - \mathbf{b}$ oe or for $\overrightarrow{CP} = 4\mathbf{a} - 2\mathbf{b}$ oe	
		C1	for $\overrightarrow{CF} = \mathbf{a} - \mathbf{b}$ oe or for $\overrightarrow{DP} = 2\mathbf{a} - 2\mathbf{b}$ oe	
		C1	for $\overrightarrow{CF} = \mathbf{a} - \mathbf{b}$ and $\overrightarrow{DP} = 2\mathbf{a} - 2\mathbf{b}$ **(**or 2**(a – b))** leading to conclusion	

Question	Answer	Mark	Mark scheme	Additional guidance
20	$\frac{3}{5}\mathbf{a}+\frac{2}{5}\mathbf{b}$	P1	for process to find $\overrightarrow{OR}=\mathbf{a}-\mathbf{b}$ or $\overrightarrow{RO}=\mathbf{b}-\mathbf{a}$	
		P1	for process to find $\overrightarrow{MR}=\frac{3}{5}(\mathbf{a}-\mathbf{b})$ or $\overrightarrow{MO}=\frac{2}{5}(\mathbf{b}-\mathbf{a})$ or $\overrightarrow{RM}=\frac{3}{5}(\mathbf{b}-\mathbf{a})$ or $\overrightarrow{OM}=\frac{2}{5}(\mathbf{a}-\mathbf{b})$	
		P1	for complete process to find $\overrightarrow{MT}$ in terms of **a** and **b,** eg $\frac{3}{5}(\mathbf{a}-\mathbf{b})+\mathbf{b}$ oe or $\frac{2}{5}(\mathbf{b}-\mathbf{a})+\mathbf{a}$ oe	
		A1	for $\frac{3}{5}\mathbf{a}+\frac{2}{5}\mathbf{b}$ or $\frac{1}{5}(3\mathbf{a}+2\mathbf{b})$ or $\frac{3\mathbf{a}+2\mathbf{b}}{5}$	Accept decimals instead of fractions

Question	Answer	Mark	Mark scheme	Additional guidance
22	$b = 3a$	P1	for $a\binom{2}{6} + b\binom{8}{2} = k\binom{13}{6}$ oe or for setting up a linear equation in a or b eg $2a + 8b = 13k$ or $6a + 2b = 6k$, $k \neq 0, 1$ **or** for $2a + 8b = 13$ and $6a + 2b = 6$ **or** for $\frac{2a+8b}{6a+2b} = \frac{13}{6}$ oe	Accept any non zero value substituted for k
		P1	for process to solve the simultaneous equations to get $b = \frac{3k}{2}$ and $a = \frac{k}{2}$ or $b = \frac{3}{2}$ and $a = \frac{1}{2}$ **or** both $2a + 8b = 13k$ and $6a + 2b = 6k$ with process to eliminate k	
		A1	for $b = 3a$ oe	

Paper: 1MA1/3H

Question	Answer	Mark	Mark scheme	Additional guidance
18	Proof	B1	for $\overrightarrow{EH} = -\mathbf{b} + \mathbf{a}$ **or** $\overrightarrow{HE} = -\mathbf{a} + \mathbf{b}$ **or** $\overrightarrow{HG} = \mathbf{a} + \mathbf{c}$ **or** $\overrightarrow{GH} = -\mathbf{c} - \mathbf{a}$	
		B1	for $\overrightarrow{BC} = -2\mathbf{b} + 2\mathbf{a} + 2\mathbf{c}$ **or** $\overrightarrow{CB} = 2\mathbf{b} - 2\mathbf{a} - 2\mathbf{c}$	
		B1	for $\overrightarrow{EF} = \mathbf{b} + \frac{1}{2}(-2\mathbf{b} + 2\mathbf{a} + 2\mathbf{c})$ $(= \mathbf{a} + \mathbf{c})$ **or** $\overrightarrow{FE} = \frac{1}{2}(2\mathbf{b} - 2\mathbf{a} - 2\mathbf{c}) - \mathbf{b}$ $(= -\mathbf{a} - \mathbf{c})$ **or** $\overrightarrow{FG} = \frac{1}{2}(-2\mathbf{b} + 2\mathbf{a} + 2\mathbf{c}) - \mathbf{c}$ $(= -\mathbf{b} + \mathbf{a})$ **or** $\overrightarrow{GF} = \mathbf{c} + \frac{1}{2}(2\mathbf{b} - 2\mathbf{a} - 2\mathbf{c})$ $(= -\mathbf{a} + \mathbf{b})$	
		A1	for completing proof, $\overrightarrow{EH} = \overrightarrow{FG}$ **or** $\overrightarrow{HG} = \overrightarrow{EF}$ (so *EFGH* is a parallelogram)	

Question	Answer	Mark	Mark scheme	Additional guidance
19 (a)	$\frac{1}{18}(8k\mathbf{b}+10\mathbf{b}-\mathbf{a})$	P1	for a correct expression for $\overrightarrow{CB}$ or $\overrightarrow{BC}$ eg, $\overrightarrow{CB}=-k\mathbf{b}-\mathbf{a}+\mathbf{b}$ or $\overrightarrow{BC}=-\mathbf{b}+\mathbf{a}+k\mathbf{b}$	All vectors must be clearly identified
		P1	for a correct expression for $\overrightarrow{CN}$ or $\overrightarrow{BN}$ or $\overrightarrow{NC}$ or $\overrightarrow{NB}$ eg, $\overrightarrow{CN}=\frac{5}{9}(-k\mathbf{b}-\mathbf{a}+\mathbf{b})$ or $\overrightarrow{BN}=\frac{4}{9}(-\mathbf{b}+\mathbf{a}+k\mathbf{b})$ or $\overrightarrow{NC}=\frac{5}{9}(-\mathbf{b}+\mathbf{a}+k\mathbf{b})$ or $\overrightarrow{NB}=\frac{4}{9}(-k\mathbf{b}-\mathbf{a}+\mathbf{b})$	This mark implies the previous one
		P1	for a correct unsimplified expression for $\overrightarrow{MN}$ eg $\frac{1}{2}\mathbf{a}+k\mathbf{b}+\frac{5}{9}(-k\mathbf{b}-\mathbf{a}+\mathbf{b})$ oe or $-\frac{1}{2}\mathbf{a}+\mathbf{b}+\frac{4}{9}(-\mathbf{b}+\mathbf{a}+k\mathbf{b})$ oe	
		A1	for $\frac{1}{18}(8k\mathbf{b}+10\mathbf{b}-\mathbf{a})$ oe eg $\frac{5}{9}\mathbf{b}+\frac{4}{9}k\mathbf{b}-\frac{1}{18}\mathbf{a}$	Must have a maximum of 3 vector terms, **a**, **b**, and k**b**
(b)	No, with explanation	C1	No with supporting reason ft (a) **Acceptable reasons:** No, since $\frac{1}{18}(8k\mathbf{b}+10\mathbf{b}-\mathbf{a})$ is not a multiple of **b** No, as N is not the midpoint of BC No, they are not multiples of each other No, does not factorise to **b** Not a multiple of OB OB doesn't have an **a** **Not acceptable reasons:** Yes… No, they don't share the same multiples OB doesn't go into MN	

Paper: 1MA1/2H

Paper: 1MA1/3H

Question	Answer	Mark	Mark scheme	Additional guidance
8	Vector $\binom{5}{8}$ drawn	M1	for $2 \times 3 + -1$ (= 5) or $2 \times 2 + 4$ (= 8) seen as a calculation **or** for $\binom{2\times3}{2\times2} + \binom{-1}{4}$ **or** for $\binom{5}{b}$ or $\binom{a}{8}$ **or** for $\binom{3}{2}$ drawn or $\binom{6}{4}$ drawn or $\binom{-1}{4}$ drawn	May be in the correct position in a column vector or coordinate. Condone missing arrows
		M1	for $\binom{5}{8}$ **or** for $\binom{5}{8}$ drawn with no arrow or incorrect arrow **or** for $\binom{5}{b}$ drawn with arrow or $\binom{a}{8}$ drawn with arrow, where $b \neq 8$ and $a \neq 5$	
		A1	cao	For this mark the drawn vector must include an arrow showing direction. Need not be labelled but do not award if there is any ambiguity

Paper: 1MA1/3H

edexcel

GCSE
Edexcel GCSE
Mathematics A 1387

July 2004

advancing learning, changing lives

Mark Scheme

Edexcel GCSE
Mathematics A 1387

London Qualifications is one of the leading examining and awarding bodies in the UK and throughout the world. It incorporates all the qualifications previously awarded under the Edexcel and BTEC brand. We provide a wide range of qualifications including general (academic), vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, our centres receive the support they need to help them deliver their education and training programmes to learners.

For further information please call Customer Services on 0870 240 9800, or visit our website at www.edexcel.org.uk

July 2004

Publications Code UG015616

All the material in this publication is copyright
© London Qualifications Ltd 2004

CONTENTS

Notes on Marking Principles....2
Paper 5514....4
Paper 5515....8
Paper 5516.... 11
Paper 5517.... 15
Paper 5518.... 19
Paper 5519.... 23

NOTES ON MARKING PRINCIPLES

NOTES ON MARKING PRINCIPLES

1 Three types of mark are available

M marks - awarded for correct working seen or implied.
A marks - conditional accuracy marks which are awarded for accurate working following the award of M marks.
B marks - unconditional accuracy marks (independent of M).

2 Abbreviations

cao - correct answer only.
ft - follow through.
$\surd$ - Denotes a "follow through" answer.

SC - special case.
isw - ignore subsequent working.
oe - or equivalent (and appropriate).
NB: a candidate cannot benefit from both isw and ft.

3 If no working shown, then

correct answers normally score full marks
incorrect (even though nearly correct) answers score no marks.

4 Marking instructions

Misread loses A marks (and sometimes B marks) on that part but ft can be allowed on subsequent parts. M marks can still be earned. If in doubt contact your team leader.
If there is a wrong answer in the answer space DO CHECK the working in the body of the script.

- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.
- If there is a wrong answer in the answer space DO CHECK the working in the body of the script.
- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.

5 Style of marking

Answer correct: tick and write part mark in margin NEXT TO BRACKETED MARK.
Answer incorrect: cross, but show M, A or B marks if any earned in body of script and transfer the total of these to the margin next to the bracket mark.
Total for each double page at bottom right page (except for back if used), FINAL TOTAL IN RELEVANT BOX ON FRONT COVER.
Nought in margin for fully incorrect question or page AND FOR NO ATTEMPT.
Where no attempt has been made a line should be put in the answer space and zero in the margin next to the bracketed mark
CHOICE OF METHOD – No marks unless one answer is in answer space – then mark that.
CROSSED OUT WORK - if not replaced this should be marked (if legible).
There must always be a mark next to bracketed mark in the margin/

6 Follow Through Marks

Follow throughs are guided by two principles:

(a) Follow throughs which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous, do not award.
(b) Follow throughs which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

7 Probability

Probability answers must be given as fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least two decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.

6 Linear Equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

8 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

PAPER 5501					
No		**Working**	**Answer**	**Mark**	**Notes**
1			K: 1020 L: 8.06	2	B1 for 1020 or 1,020 B1 for 8.06 Accept £8.06p and £8,06
2	(a)		28, 33	2	B1, B1 (B1 ft for "28" + 5 if both numbers >25)
	(b)			1	B1 for add 5, +5, for going up in 5's the difference is 5 oe
	(c)			1	B1 for they end in 3 and 8, it ends in 7; or refers to 383 and 388 are in sequence
3	(a)(i)		$\frac{1}{2}$	2	B2 for ½ accept half (B1 for an equivalent unsimplified fraction eg 4/8 or 50% or 0.5)
	(ii)		2 rectangles shaded	1	B1 for correct shading (any 2 rectangles)
	(b)(i)		40	4	B1 for 40 cao
	(ii)		150 000		B1 for 150 000 cao (accept 150,000 not 150.000)
	(iii)		6.55		B1 for 6.55 cao not $6.5^{½}$
	(iv)		$\frac{3}{8}$		B1 for $\frac{3}{8}$ oe accept 0.375
	(c)		Cross 3cm from *A*	1	B1 mark a cross 3 cm (±2 mm) from *A*
4	(a)(i)		metres	2	B1 for metres (m)
	(ii)		grams		B1 for grams (g)
	(b)		miles	1	B1 for miles
5	(a)		(3, 2)	1	B1 for (3, 2)
	(b)(i)		*Q* at (0, 3)	1	B1 for *Q* plotted correctly on *y*-axis at (0, 3) ±2mm
	(ii)		*R* at (−2, −3)	1	B1 for *R* plotted correctly at (−2, −3) ±2mm
6	(a)(i)		30	2	B1 for 30
	(ii)		25		B1 for 25
	(b)(i)		4 shapes	2	B1 for drawing 4 shapes
	(ii)		1½ shapes		B1 for drawing 1½ shapes

PAPER 5501					
	No	Working	Answer	Mark	Notes
7	(a)		30	1	B1 for 30
	(b)	$20 \times 3 + 5$	65	2	M1 for $20 \times 3 + 5$ oe A1 for 65
8	(a)		Draws perp.	1	B1 for correctly drawing perp must touch line or cut line *AB* ± 2mm
	(b)		Sketches a cylinder	1	B1 for sketching cylinder
9	(a)	1974 6580 8554	855.4	3	M1 for complete method with relative place value correct, condone 1 error in multiplication A2 cao (A1 for digits 8554 seen or A1 for "855.4" dependent on 1 arithmetic error)
	(b)	990 ÷ 69 = 14.3…. or 14 rem 24	14	4	Method 1 - Everything excluding long division M2 for a valid method with no errors …need to see 966 or 1035 (M1 for a valid method, including estimation, that reaches an answer between 901 and 1099 inclusive. Ignore errors.) A2 for 14 with a fully correct method (A1 (dep on 1st M1) for an answer between 14 and 15 inclusive. Ignore errors. Cannot have used estimation.) Note: Estimation can only score a maximum of M1 unless accompanied by another method or 966 or 1035 see (award M2) Method 2 – long division method M2 for 990 ÷ 69 = integer answer between 12 and 19 inclusive and rem 30 seen (M1 for 990 ÷ 69 with 1 in tens column.) A2 for 14 with fully correct method (A1 (dep on 1st M1) for an answer between 14 and 15 inclusive.)

PAPER 5501					
	No	Working	Answer	Mark	Notes
10	(a)(i)		7	2	B1 for 7
	(ii)		−10		B1 for −10
	(b)(i)		6	2	B1 for 6 (accept −6)
	(ii)		8		B1 for 8 (accept −8)
	(c)		−7	1	B1 for −7 cao
11	(a)		B and D	2	B1, B1 (–1 each extra)
	(b)(i)		A	2	B1 for A
	(ii)		3		B1 for 3
12	(a)	800 − 144	656	2	M1 for at least 1 digit correct and in correct position needs to be 3 digit number A1 for 656
	(b)	144 is less than 200 so Trudy is wrong		2	B1 for 200 B1ft for 'correct' explanation based on cand's "200"
	(c)	$\frac{45\times800}{100}$	360	2	M1 for 45 × 800 ÷ 100 oe A1 for 360
	(d)	$\frac{176}{800}\times100$	22	2	M1 for $\frac{176}{800}\times100$ oe A1 for 22
13	(a)(i)		25	4	B1 for 25
	(ii)		28		B1 for 28
	(iii)		5 and 20		B1 for 5 and 20
	(iv)		26 and 33		B1 for 26 and 33
	(b)	$2^3 = 8$ or $2^3 = 2 \times 2 \times 2$ (which is not 6)		1	B1 for valid 'explanation'
14			10 m^2	2	B1 for 10 B1 indep for m^2.

PAPER 5501				
No	**Working**	**Answer**	**Mark**	**Notes**
15 (a)	$\frac{11}{12}-\frac{10}{12}$	$\frac{1}{12}$	2	M1 for correctly writing both fractions to a common denominator A1 for $\frac{1}{12}$ oe
(b)	$\frac{70\times400}{200}$	140	2	B2 for 140 (accept 136) (B1 for sight of any two of 70, 400 or 200)
16 (a)		$2y$	1	B1 for $2y$ or $2\times y$, $y2$, $y\times2$
(b)		$3p^2$.	1	B1 for $3p^2$ or $3\times p^2$ or $p^2\times3$ or $p^2\,3$
(c)		$x(x-3)$	2	B2 for $x(x-3)$ (B1 for x (x…….)) $\frac{25}{60}$ wrongly cancelled gets B1 ISW
17 (a)		2 23 9 34 15 2 9 26 17 25 18 60	3	B3 for all correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
(b)		$\frac{25}{60}$	1	B1 for $\frac{25}{60}$ or $\frac{5}{12}$ oe $\frac{25}{60}$ wrongly cancelled gets B1 ISW

PAPER 5501					
	No	**Working**	**Answer**	**Mark**	**Notes**
18	(a)(i)		60	2	B1 for 60
	(ii)				B1 for all **angles equal** so equilateral triangle oe
	(b)(i)		130	3	B1 for 130
	(ii)				B1 for isosceles triangle oe or 2 angles equal accept $\angle Q = \angle R$ B1 for angles on a straight **line** add up to **180°**oe (180° could be in working)
	(c)		64	1	B1 for 64
19	(a)		1	1	B1 for 1
	(b)		3	1	B1 for 3
	(c)	$9\times1+3\times2+5\times3+3\times4$ (= 42) "42" ÷ 20	2.1	3	M1 for 9×1, 3×2, 5×3, 3×4 or for 42 seen M1 (dep) for "42" ÷ 20 A1 for 2.1 or $2\frac{1}{10}$ or $2\frac{2}{20}$
20	(a)		09 05	1	B1 for 09 05 oe
	(b)		7	1	B1 for 7
	(c)		10	1	B1 for 10
	(d)	7km in 20 mins	21	3	B1 for 20 (minutes) oe or $\times3$ seen M1 for $\frac{"7"}{"20"}$ A1 cao [SC: "7" × 3 seen gets B1 M1]

PAPER 5501				
No	**Working**	**Answer**	**Mark**	**Notes**
21	Vertices of new triangle at (2, −2) (2, 6) and (6, −2)	Correct triangle drawn	3	B3 cao (B2 for either correct triangle in wrong translated position or for 2 vertices correct) (B1 a right angled triangle with horizontal length 4 or vertical length 8 in the same orientation as the shaded triangle) SC B2 for s + 3 totally correct
22 (a) (b) (c)	 8560 ÷ (10 × 10)	8 Pentagon 85.6	1 1 2	B1 for 8 B1 for pentagon M1 for 8560 ÷ (10 × 10) oe A1 for 85.6
23	$2x + x + 100 + 47 = 360$ $2x + x = 360 - 100 - 47$ $x = 71$ Largest angle = $2x$ =	142	4	M1 for $2x + x + 100 + 47 = 360$ or 360 – 147 or 213 seen M1 dep for correctly separating x-terms and non x terms or "360 – 147" ÷ 3 A1 for $x = 71$provided M2 awarded A1 ft for 142

PAPER 5502					
	No	**Working**	**Answer**	**Mark**	**Notes**
1	(a)		Diagram	1	B1 cao
	(b)		22, 26	1	B1 cao
	(c)		46	1	B1 cao
2	(a)	£5.95 + £1.62 + 25p + 2 × 48p = £8.78	1.22	3	M1 for 3 of £5.95, £1.62, 25p, 48p seen with intention to add or 170 seen A1 for £8.78 A1 ft from " 8.78"
	(b)	160 ÷ 12 = 13.(333…)	14	2	M1 for 160 ÷ 12, or 13 or better or 12 × 13 or 12 × 14 A1 14 cao
	(c)	"14" × £12.20 = £170.8**0**	170.80	2	M1 for "14" × 12.20 A1 for 170.8**0** or 158.6**0** if 13 in (b) NB: 170.8 or 158.6 gets M1 A0
3	(a)(i)		1459	1	B1 cao
	(ii)		9541	1	B1 cao
	(b)		9 + 5 = 14	1	B1 cao
	(c)		0	1	B1 cao
4	(a)		1123	1	B1 cao
	(b)		1125	1	B1 cao
	(c)		43	1	B1 cao
	(d)	67 – "43"	24	2	M1 for 67 – "43" A1 ft from "43"
	(e)(i)		80	2	B1 cao
	(ii)		56 – 58		B1 answer in range 56 to 58 inc
	(f)		Completes graph	2	B2 for (Apr , 70) and (May , 60 < p < 65) plotted and joined (B1 for one point plotted)

PAPER 5502					
	No	Working	Answer	Mark	Notes
5	(a)(i)		4	1	B1 cao
	(ii)		10	1	B1 cao
	(b)		28	2	B2 for 28 (B1 for 29 or 14×2)
6	(a)		$\frac{3}{4}$	1	B1 accept 0.75, 75%
	(b)	$\frac{3}{4} \times 28 = 21$	15.54	3	M1 for $\frac{3}{4} \times 28$ or 21 seen M1 for "21" $\times$ 74 or 28×74 or 7×74 A1 cao
	(c)(i)		44 – 46	2	B1 44 – 46 inc
	(ii)		5.2 – 5.4		B1 5.2 – 5.4 inc
7	(a)		Correct shape	1	B1
	(b)		Correct shape	1	B1
8	(a)		Hexagon	1	B1
	(b)		Right	1	B1 cao
	(c)		Drawing	2	B2 for 7 additional hexagons with at least 2 points at which 3 hexagons meet (B1 for one point at which 3 hexagons meet)
9	(a)		$\frac{3}{4}$	2	M1 $\frac{80-20}{80}$ or $\frac{60}{80}$ or 0.75 (oe) A1 cao Do not accept equivalents
	(b)		15	1	B1 cao
	(c)		Iron and Cook set	1	B1 for both Accept mixer and fryer
	(d)		Microwave	1	B1 cao
	(e)		Bar chart	2	B1 one column (mixer): 70, 90 B1 second column (fryer): 70, 85 [SC: B1 columns correct but reversed]

PAPER 5502					
No		**Working**	**Answer**	**Mark**	**Notes**
10	(a)		Acute	1	B1
	(b)		Reflex	1	B1
	(c)		reason	1	B1 reason eg 120 + 230 = 350 ≠ 360
11	(a)		$\frac{1}{2}$	1	B1 for $\frac{1}{2}$ oe
	(b)		reason	1	B1 for reason e.g could get 30 heads
12	(a)	7.5×4	30	2	M1 for 7.5×4 or $7\frac{1}{2} \times 4$ A1 cao
	(b)	$\sqrt{(42.25)}$	6.5	2	B2 for 6.5 (B1 for 42.25)
13	(a)		$6x$	1	B1 accept $6 \times x$, $x6$ oe
	(b)		$x - 4$	1	B1 cao
	(c)		$12(x - 4)$	1	B1 for $12(x - 4)$ or $12x - 48$ or $12 \times x - 48$ oe
	(d)		$18x - 48$	2	M1 for "$6x$" + "$12(x-4)$" or "$6x$" + "$12x - 48$" oe A1 ft $18x - 48$, $2(9x - 24)$, $3(6x - 16)$, $6(3x - 8)$
14	(a)		90	2	B2 for 90 ± 2 (B1 for 9 ± 0.2)
	(b)		317	1	B1 for 317 ± 4
15	(a)	Accurate drawing of triangle		2	B1 cao for 38° (±2°) B1 cao (indep) for BC drawn 7.3 cm (±2mm) and completing the triangle.
	(b)	Measure angle A	63°	1	B1 for 63° (±2°) or ft their diagram (±2°)

PAPER 5502					
	No	Working	Answer	Mark	Notes
16	(a)	0 × 1 = 0 1 × 1 = 1 2 × 4 = 8 3 × 8 = 24 4 × 8 = 32 5 × 7 = 35	100	3	M1 col 1 × col 2 (at least 3 shown) M1 (dep) sum of totals A1 cao SC: M1 M1 A0 for 101
	(b)	Either 35.50 ÷ 1.42 = 25; 26.99 – 25 = 1.99 Cheaper in the USA Or 26.99 × 1.42 = 38.33; 38.33 – 35.50 = 2.83 Cheaper in the USA		3	M1 35.50 ÷ 1.42 A1 25 seen B1 ft (dep on M1) £1.99 with conclusion (must have units) **OR** M1 26.99 × 1.42 A1 38.32 or 38.33… seen B1 ft (dep on M1) $2.83 or $2.82 with conclusion (must have units) must be to 2dp
17	(a)		-1, (1), (3), 5, 7, 9	2	B2 cao (B1 for 2 values)
	(b)		Graph	2	B1 ft for plotting points ±1/2 square B1 cao for line between $x = -2$ and $x = 3$
	(c)(i)		0.4	2	B1 for 0.4 or ft from single straight line with positive gradient
	(ii)		1.2		B1 for 1.2 or ft from single straight line with positive gradient

PAPER 5502					
	No	Working	Answer	Mark	Notes
18	(a)		Plots	1	B1 cao
	(b)		description	1	B1 dynamic relationship or "positive" (correlation)
	(c)		line of best fit	1	Line within overlay region, and to the extent of.
	(d)(i)	280 g →	reading	2	B1 ft from single straight line of positive gradient (±1/2 square)
	(ii)	120 pages →	reading		B1 ft from single straight line of positive gradient (±1/2 square)
19			200 150 225 150 10	3	B3 cao (B2 for three correct, or B1 for one correct)
20	(a)			2	B1 B1 for correct sketch – ignore additional internal lines B1 B0 for rectangle or right-angled triangle or correct shape without line NB 3-D sketch gets B0 B0
	(b)			2	B1 B1 for correct plan – ignore internal lines – accept a rotated plan, not reflected B1 B0 for a single rectangle drawn NB 3-D sketch gets B0 B0
21	(a)	163.25 – 35.50 = 127.75 127.75 ÷ 18.25 = 7	8	3	M1 163.25 – 35.50 (or sight of 127.75) M1 (dep) "127.75" ÷ 18.25 A1 cao SC: M2 for 7 days
	(b)	1.175 × £64 or $\frac{17.5}{100}$ × 64 or £11.20 seen	75.20	2	M1 1.175 × 64, 0.175 × 64 oe or 11.20 or 75.2 seen or 6.40, 3.20 and 1.60 seen A1 cao

PAPER 5502					
	No	Working	Answer	Mark	Notes
22	(a)		6	1	B1 cao
	(b)		$t^2 - 2t$	1	B1 oe
	(c)		$3(y - 4)$	1	B1 Accept $3 \times (y - 4)$ or $3(y - 4$
23	(a)		Reason	1	B1 Makes some mention of bias either directly or making reference to an insufficient or biased range of responses
	(b)		Reason	1	B1 (a) an insufficient range of responses (b) No mention of money (c) No time frame in the question (d) Misunderstanding of "A lot" and "Not much"

PAPER 5503					
No		**Working**	**Answer**	**Mark**	**Notes**
1			10.	1	B1 for 10
2	(a)(i)		$3e + 2f$	2	B1 for $3e + 2f$
	(ii)		$3p^2$		B1 for $3p^2$
	(b)	$5 \times -3 + 1$	-14	2	M1 for $5 \times -3 + 1$ A1 for -14
3		1974 $\frac{6580}{8554}$	855.4	3	M1 for complete correct method with relative place value correct, condone 1 error in multiplication. A2 cao (A1 for digits 8554 seen or A1 ft for "855.4" dependent on one arithmetic error only)
4	(a)		2 23 9 34 15 2 9 26 17 25 18 60	3	B3 all correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
	(b)		$\frac{25}{60}$	1	B1 for $\frac{25}{60}$ or $\frac{5}{12}$ oe
5	(a)	$\frac{45}{100} \times 800$	360	2	M1 for $\frac{45}{100} \times 800$ oe A1 for 360
	(b)	$\frac{176}{800} \times 100$	22	2	M1 for $\frac{176}{800} \times 100$ oe A1 for 22

PAPER 5503					
	No	Working	Answer	Mark	Notes
6	(a)	$\frac{11}{12}-\frac{10}{12}$	$\frac{1}{12}$	2	M1 for correctly writing both fractions to a common denominator A1 for $\frac{1}{12}$ oe
	(b)	$\frac{70\times400}{200}$	140	2	B2 for 140 (accept 136) (B1 for sight of any two of 70, 400 or 200)
7	(a)(i)		60	2	B1 for 60
	(ii)				B1 for all angles equal so equilateral triangle oe
	(b)(i)		130	3	B1 for 130
	(ii)				B1 for isosceles triangle oe B1 for angles on a straight line add up to 180°oe
	(c)		64	1	B1 for 64
8		$9\times1+3\times2+5\times3+3\times4$ (=42) "42" ÷ 20	2.1	3	M1 for 9×1, 3×2, 5×3, 3×4 or for 42 seen M1 (dep) "42" ÷20 A1 for 2.1 or $2\frac{1}{10}$ or $2\frac{2}{20}$
9	(a)		09 05	1	B1 for 09 05 oe
	(b)		7	1	B1 for 7
	(c)		10	1	B1 for 10
	(d)	7 km in 20 minutes	21	3	B1 for 20 (minutes) oe or $\times3$ seen M1 for $\frac{"7"}{"20"}$ A1 cao SC: "7" $\times3$ gets B1 M1

PAPER 5503					
No		**Working**	**Answer**	**Mark**	**Notes**
10		Vertices of new triangle at (2, −2) (2, 6) and (6, −2)	Correct triangle drawn	3	B3 cao (B2 for either correct triangle in wrong translated position or for 2 vertices correct) (B1 for a right-angled triangle with horizontal length 4 or vertical length 8 in the same orientation as shaded triangle) SC B2 for scale factor 3, totally correct
11		$2^2 + 3 =$	7 which is not even	2	B2 (B1 for correctly evaluating $n^2 + 3$ with a prime number value for n.)
12	(a)	360 ÷ 5	72	2	M1 for 360 ÷ 5 oe A1 for 72
	(b)	8560 ÷ (10 × 10)	85.6	2	M1 for 8560 ÷ (10 × 10) oe A1 for 85.6
	(c)(i)	Least length = 100.5	100.5	2	B1 for 100.5
	(ii)	Greatest length = 101.5	101.5		B1 for 101.5 or 101.499 or better
13		$2x + x + 100 + 47 = 360$ $2x + x = 360 - 100 - 47$ $x = 71$ Largest angle = $2x$ =	142	4	M1 for $2x + x + 100 + 47 = 360$ or 360 – 147 or 213 seen M1 (dep) for correctly separating x-terms and non x terms or $\frac{"360-147"}{3}$ A1 for $x = 71$ A1 ft (provided M2 awarded) for "142"
14	(a)(i)		2.21	2	B1 for 2.21
	(ii)		0.013		B1 for 0.013
	(b)	LCM = 3 × 13 × 17 = 3 × 221	663	2	M1 for 3 × 13 × 17 oe A1 for 663

PAPER 5503				
No	**Working**	**Answer**	**Mark**	**Notes**
15		Tick boxes 1,3 & 6	3	B1 for $\frac{\pi abc}{2d}$; B1 for $2a^2$; B1 for $2\left(c^2+d^2\right)$ (−B1 for each additional expression ticked (>3) to a minimum of 0)
16	200×0.2	40	2	M1 for 200×0.2 or $\frac{40}{200}$ seen A1 for 40
17 (a)	eg 2 | 108 2 | 54 3 | 27 3 | 9 3	$2^2 \times 3^3$	3	M2 for full systematic method of at least 4 divisions by prime numbers oe factor trees; condone 1 calculation error (M1 for 108 written as a correct product(s) with only one non-prime or equivalent division or a full process with 2 calculation errors.) A1 for $2^2 \times 3^3$ (accept $2 \times 2 \times 3 \times 3 \times 3$)
(b)		12	1	B1 for 12
18	Perpendicular from P to intersecting arcs (within tramlines); perpendicular at least 2cm long		2	M1 relevant pair of arcs crossing within tramlines A1 SC M1 AO for a full construction of a perpendicular to AB not passing through P.
19	15×10	150 cm^3.	3	M1 for 15×10 A1 for 150 B1 for cm^3.

PAPER 5503					
No		**Working**	**Answer**	**Mark**	**Notes**
20			$y = 2x + 6$	2	B2 for $y = 2x + 6$ (B1 for $y = 2x + k$, $k \neq 1$ or for $y = mx + 6$, m $\neq 0$ or for $2x + 6$)
21		$6\frac{2}{5} = \frac{32}{5}$ Area of triangle = $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ (=2) Length of a side of sq. = $\sqrt{18 \times "2"}$ (=6) Perimeter of square = 4 × 6	24	5	B1 for $6\frac{2}{5} = \frac{32}{5}$ oe or $3\frac{1}{5} = \frac{16}{5}$ or $\frac{30}{8} + \frac{2}{8}$ oe (or implied by area of triangle=2) M1 for $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ oe M1 for area of square = 18 × product of two lengths A1 for $\sqrt{18 \times "2"}$ A1 for 24
22	(a)		$x(x-3)$	2	B2 for $x(x-3)$ (B1 for x (…….))
	(b)		k^3.	1	B1 for k^3.
	(c)(i)	$4x + 20 + 3x - 21$	$7x - 1$	4	M1 for **three** of 4 terms $4x + 20 + 3x - 21$ (or better) A1 for $7x - 1$
	(ii)	$x^2 + 3yx + 2xy + 6y^2$	$x^2 + 5xy + 6y^2$		M1 for **three** of 4 terms $x^2 + 3yx + 2xy + 6y^2$ A1 for $x^2 + 5xy + 6y^2$
	(d)		$(p+q)(p+q+5)$	1	B1 for $(p+q)(p+q+5)$

PAPER 5503					
	No	Working	Answer	Mark	Notes
23	(a)		32	1	B1 for 32 (accept 31.5 to 33.5 inclusive)
	(b)			3	B1 for ends of whiskers at 9 and 57, (with a box) B1 for ends of box at 16 and 45 (± 0.5) B1 for median marked at 32 or a complete box and whisker diagram drawn with a median inside the box
	(c)	Median(B)>Median(G); on average boys take longer IQR(B)>IQR(G); times for boys have a greater spread		2	B1 eg for comparison of medians B1 eg for comparison of (interquartile) ranges
24	(a)	$6y = 15 - 5x$	$y = \frac{15-5x}{6}$	2	M1 for either $6y = 15 - 5x$ or -6 $y = 5x - 15$ or for $\frac{5x}{6} + y = \frac{15}{6}$ or a correct ft on sign error to $y=$ A1 for $y = \frac{15-5x}{6}$ oe
	(b)	$6k + 5(-21) = 15$	20	2	M1 for substitution of $x = -21$ (or $x = 21$) into given equation or candidate's answer to (a) A1 for $k = 20$
	(c)(i)		Region R indicated	3	B2 correct region shaded (accept unshaded if R clear) (B1 shaded or R region satisfies 3 of the 4 given inequalities with same boundaries)
	(ii)		(1,1)		B1 for (1,1)

PAPER 5503					
	No	Working	Answer	Mark	Notes
25	(a)		60	1	B1 for 60 B1 for *BDC* = 25
	(b)	eg Angle *BDC* = 25 *ADB* = 60 − 25	35	2	[Award the mark for equivalent approaches] B1 ft for (a) − 25
	(c)	Ben is correct; angle *DAB* = 65 + 25 = 90 and since angle in a semi-circle is 90°, *BD* must be a diameter		1	B1 for full valid justification

PAPER 5504					
	No	Working	Answer	Mark	Notes
1	(a)	Accurate drawing of triangle		2	B1 cao for 38º (±2º) B1 cao (indep) for BC drawn 7.3 cm (±2mm) and completing the triangle.
	(b)	Measure angle A	63º	1	B1 for 63º (±2º) or ft their diagram (±2º)
2	(a)	0 × 1 = 0 1 × 1 = 1 2 × 4 = 8 3 × 8 = 24 4 × 8 = 32 5 × 7 = 35	100	3	M1 col 1 × col 2 (at least 3 shown) could be implied by answers. M1 (dep) sum of totals A1 cao SC: M1 M1 A0 for 101
	(b)	Either $35.50 ÷ 1.42 = £25; £26.99 - £25 = £1.99 Cheaper in the USA Or £26.99 × 1.42= $38.33; $38.33 - $35.50 = $2.83 Cheaper in the USA		3	M1 35.50 ÷ 1.42 A1 25 seen B1 ft (dep on M1) £1.99 with conclusion; units needed OR: M1 26.99 × 1.42 A1 38.33 or 38.32…seen B1 ft (dep on M1) $2.83 or $2.82 with conclusion; units needed
3	(a)		-1, (1), (3), 5, 7, 9	2	B2 cao (B1 for 2 values)
	(b)		Graph	2	B1 ft for plotting points ±1/2 square B1 cao for line between $x = -2$ and $x = 3$
	(c)(i)		0.4	2	B1 0.4 or ft single straight line with positive gradient
	(ii)		1.2		B1 1.2 or ft single straight line with positive gradient

PAPER 5504					
	No	Working	Answer	Mark	Notes
4	(a)			2	B1 Rectangle drawn B1 Triangle attached Ignore additional internal lines; 3D sketch gets B0
	(b)			2	B1 single Rectangle drawn B1 Attached smaller rectangle drawn Accept rotated plan, Ignore internal lines down; Nets get B0
5	(a)	£163.25 - £35.50 = £127.75 £127.75 ÷ £18.25 = 7	8	3	M1 163.25 – £35.50 (or sight of £127.75) M1 (dep) "127.75" ÷ 18.25 A1 cao SC: M2 for 7 days
	(b)	1.175 × £64 or $\frac{17.5}{100}$ × 64 or £11.20 seen	75.20	2	M1 1.175 × 64.00 0.175 × 64 oe or 11.20 or 75.2 seen OR 6.40, 3.20, and 1.60 seen A1 cao
	(c)	54 × £2.38 =	128.52	2	M1 54 × £2.38 A1 cao
6		Tessellation	Drawing	2	B2 seven additional hexagons, with at least 2 points at which 3 hexagons meet (B1 one point at which 3 hexagons meet).
7			200 150 225 150 10	3	B3 cao (B2 for three correct, or B1 for one correct)
8	(a)(i)	Bearing measured	70	2	B1 70 ± 4
	(ii)	Bearing measured	317		B1 317 ± 4
	(b)	Locus (circle) drawn & shaded		2	B1 circle ±2 mm centre Manchester B1 shading with accurate or approximate circle within tolerance

PAPER 5504					
	No	Working	Answer	Mark	Notes
9	(a)		Plots	1	B1 cao
	(b)		Description	1	B1 dynamic relationship or "positive" (correlation)
	(c)		Line of best fit	1	Line within overlay region, and to the extent of.
	(d)(i)	280 g →	Reading	2	B1 ft from single straight line of positive gradient (±1/2 square) B1 ft from single straight line of positive gradient (±1/2 square)
	(ii)	120 pages →	Reading		
10	(a)	$\frac{1}{6}, \frac{3}{8}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}$		2	B2 for all correct (B1 for 1 error or all correct but wrong order, or use of a common denominator decimals).
	(b)	3/5, 65%, 2/3, 0.72, 3/4		2	B2 for all correct (B1 for 1 error or all correct but wrong order or conversions to decimals oe)
11	(a)		182, 178, 180, 184	2	M1 mean of any three consecutive months, eg (147 + 161 + 238) ÷ 3 oe A1 cao
	(b)	Sale price = 80% Fun Friday price = 70% of 80% = 56% (oe)		2	B1 B1 for a fully correct explanation involving a worked example (oe) B1 a partially complete explanation

PAPER 5504					
	No	Working	Answer	Mark	Notes
12	(a)		Reason	1	B1 makes a mention of bias, either directly, or making reference to an insufficient or biased range of responses.
	(b)(i)		Reason	1	B1 make some mention of any one of (a) an insufficient range of responses (b) no mention of money (c) no time frame in the question (d) misunderstanding of "a lot" or "not much"
	(ii)		Question	2	B2 B1 for each of the following upto B2 (a) an improved question eg time frame made clear (b) response boxes (imperfect) (c) response boxes no errors OR For suggesting a generally improved question (a) a question clearly in the context of changes to the canteen (b) at least 3 boxes showing a full range of responses
13	(a)	$20y - 18y = 16 - 9$ oe $2y = 7$	$3\frac{1}{2}$ oe	3	M1 $20y - 18y = 16 - 9$ oe M1 $2y = 7$ A1 cao
	(b)	$40 - x = 3(4 + x)$ $40 - x = 12 + 3x$ $40 - 12 = x + 3x$ $4x = 28$	7	3	M1 multiplying through by 3: $3 \times \frac{40 - x}{3} = 3 \times 4 + 3 \times x$ A1 $40 - 12 = x + 3x$ A1 cao
14	(a)		$12(x\text{-}4)$	1	B1 for $12(x - 4)$ or $12x - 48$ or $12 \times x - 48$ oe $x = 12\ (x - 4)$ gets B0
	(b)		$18x\text{-}48$	2	M1 $6x +$ "$12(x - 4)$" or $6x +$ "$12x - 48$" A1 ft a linear expression $18x - 48$, $2(9x - 24)$, $3(6x - 16)$, $6(3x - 8)$

PAPER 5504					
No		**Working**	**Answer**	**Mark**	**Notes**
15		x $x^3 - 2x$ 4.1 60.7(21) 4.2 65.6(88) 4.3 70.9(07) Accept answer to 1 dp 4.22 66.7(114) or more 4.23 67.2(269) 4.24 67.7(45) 4.25 68.2(656)	4.2	4	B2 for trial $4.1 \leq x \leq 4.3$ evaluated (B1 for trial $4 < x < 5$ evaluated) B1 for different trial $4.225 \leq x \leq 4.25$ evaluated B1 (dep on at least one previous B1) for 4.2 cao
16	(a)		1.0×10^{-9}	1	B1 accept 1×10^{-9} or just 10^{-9}
	(b)	$1 \div 5 \times 10^{-9}$	2×10^{8}	2	M1 for $1 \div$ ("5×10^{-9}") or digit 2 with zeros only seen Condone omission of bracket for M1. A1 cao
17	(a)	$\frac{6.27 \times 4.52}{4.81 + 9.63} = \frac{28.3404}{14.44} = 1.962631579$	1.9626…	2	B2 for 1.9626… (B1 for 28.34 or 14.44)
	(b)		1.96	1	B1 ft from (a) as rounded to 1dp or 2dp. Do not accept 2, 2.00, but accept 2.0
18			$d = 4n + 6$	2	B2 $d = 4n + 6$ oe (B1 $d = 4n + k$, k an integer or 0, $4n + 6$, $n = 4n + 6$)
19		Vol = $\pi \times 3.8^2 \times 2.5 = \pi \times 14.44 \times 2.5$ = 45.36… × 2.5 = 113.411 Mass = "113" × 1.5 = 170.1165	170	4	M1 for $\pi \times r^2 \times 2.5$ where r = is 3.8 or 7.6 A1 if $r = 3.8$ M1 for "113" × 1.5 A1 for 169.5 – 170.3 cao

PAPER 5504					
	No	Working	Answer	Mark	Notes
20	(a)	$10^2 + 6^2$ or 136 $\sqrt{(100+36)}$ or $\sqrt{136} = 11.66\ldots$	11.7	3	M1 $10^2 + 6^2$ or 136 seen M1 (dep) $\sqrt{100+36}$ or $\sqrt{136}$ A1 11.66 – 11.7
	(b)	$\cos x = \frac{8}{10}$ or 0.8 $x = \cos^{-1} 0.8 = 36.869^\circ$	36.9	3	M1 for $\cos = \frac{8}{10}$, cos = 0.8 (oe) M1 (dep) for $\cos^{-1}$ (oe) A1 for 36.86 – 36.9
21		$18x - 6y = 99$ $\quad 12x - 4y = 66$ $8x + 6y = 18$ $\quad 12x + 9y = 27$ $26x = 117$ $\quad -13y = 39$	$x = 4\frac{1}{2}$ $y = -3$	4	M1 correct process to eliminate either x or y (condone one error) A1 cao for non – eliminated one. M1 (dep on 1st M1) for correct substitution of their found value. A1 cao (need both)
22		$\frac{133}{72} = 1.8472, \frac{160}{82} = 1.9512$ OR $\frac{72}{133} = 0.54135, \frac{82}{160} = 0.5125$ OR $\frac{160}{133} = 1.203...., \frac{82}{72} = 1.1388....$ OR $\frac{133}{160} = 0.83125...., \frac{72}{82} = 0.878$	1.84..≠ 1.95.. 1.20..≠ 1.13..	3	M1 for $\frac{133}{72}$ (= 1.8472…) oe Accept 1.8, 1.85 M1 for $\frac{160}{82}$ (=1.9512…) oe consistent pairing Accept 2.0, 1.9 **OR** M1 for $\frac{160}{133}$ (= 1.203…) oe M1 for $\frac{82}{72}$ (=1.1388) oe A1 for enough decimal places to show that the ratios are not equal; since the scale factors are different the shapes cannot be similar. NB Do Not need conclusion

PAPER 5504					
	No	Working	Answer	Mark	Notes
23	(a)		0.4, 0.6 0.4, 0.6, 0.4	2	B1 for LHS: (0.6), 0.4 B1 for RHS: 0.6, 0.4, 0.6, 0.4
	(b)	(30 × 42) – (25 × 42.8) = 1260 – 1070 = 190 190 ÷ 5 =	38	3	M1 for (30 × 42) – (25 × 42.8) or 1260 – 1070 or 190 seen M1 (dep) for "190" ÷ 5 A1 cao
24	(a)	£12000 × 0.25 = £3000; £12000 – £3000 = £9000 £9000 × 0.25 = £2250; £9000 – £2250 = £6750 £6750 × 0.25 = £1687.50; £6750 – £1687.50 =	£5062.50	3	M1 for 12000 × 0.75 (=9000) oe or £3000 or £23437.50 seen M1 (dep) for at least two further depreciation calculations (complete steps) A1 cao **OR** M2 for 12000 × $(0.75)^3$ or 5062.50 seen (M1 for 12000 × $(0.75)^n$, n = 2 or 4)
	(b)	0.8 × 0.8 × 0.8 × 0.8 (oe)	0.4096	2	M1 0.8^4 (oe) A1 cao

PAPER 5505					
	No	Working	Answer	Mark	Notes
1	(a)(i) (ii) (b)	 LCM = 3 × 13 × 17 = 3 × 221	2.21 0.013 663	2 2	B1 for 2.21 B1 for 0.013 M1 for 3 × 13 × 17 oe A1 for 663
2			Tick boxes 1,3 & 6	3	B1 for $\frac{\pi abc}{2d}$; B1 for $2a^2$; B1 for $2(c^2 + d^2)$ (−B1 for each additional expression ticked (>3) to a minimum of 0)
3	(a) (b)	200 × 0.2 0.2 + 0.4	40 0.6	2 2	M1 for 200 × 0.2 or $\frac{40}{200}$ seen A1 for 40 M1 for 0.2 + 0.4 A1 for 0.6
4	(a) (b)	eg 2 | 108 2 | 54 3 | 27 3 | 9 3 12	$2^2 \times 3^3$ 12	3 1	M2 for full systematic method of at least 4 divisions by prime numbers oe factor trees; condone 1 calculation error (M1 for 108 written as a correct product with only one non-prime or equivalent division or a full process with 2 calculation errors.) A1 for $2^2 \times 3^3$ (accept 2 × 2 × 3 × 3 × 3) B1 for 12
5		Perpendicular from P to intersecting arcs (within tramlines); perpendicular at least 2 cm long		2	M1 relevant pair of arcs crossing within tramlines A1 SC M1A0 for full construction of a line perpendicular to AB not through P
6		15 × 10	150 cm^3.	3	M1 for 15 × 10 A1 for 150 B1 for cm^3

PAPER 5505					
	No	Working	Answer	Mark	Notes
7	(a)		k^3.	1	B1 for k^3.
	(b)(i)	$4x + 20 + 3x - 21$	$7x - 1$	4	M1 for **three** of 4 terms $4x + 20 + 3x - 21$ (or better) A1 for $7x - 1$
	(ii)	$x^2 + 3yx + 2xy + 6y^2$	$x^2 + 5xy + 6y^2$		M1 for **three** of 4 terms $x^2 + 3yx + 2xy + 6y^2$ A1 for $x^2 + 5xy + 6y^2$
	(c)		$(p+q)(p+q+5)$	1	B1 for $(p+q)(p+q+5)$
	(d)		m^8	1	B1 for m^8.
	(e)		$6r^3t^6$	2	B2 for $6r^3t^6$ (B1 for $.....r^3t^6$ or for $6....t^6$)
8	(i)	Least length = 100.5	100.5	2	B1 for 100.5
	(ii)	Greatest length = 101.5	101.5		B1 for 101.5; ACCEPT 101.499 or better
9		$6\frac{2}{5} = \frac{32}{5}$ Area of triangle = $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ (= 2) Length of a side of sq. = $\sqrt{18 \times "2"}$ (=6) Perimeter of square = 4 × 6	24	5	B1 for $6\frac{2}{5} = \frac{32}{5}$ oe or $3\frac{1}{5} = \frac{16}{5}$ oe or $\frac{30}{8} + \frac{2}{8}$ oe (or implied by area of triangle=2) M1 for $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ oe M1 for (area of square) = 18 × product of two lengths A1 = $\sqrt{18 \times "2"}$ A1 for 24

PAPER 5505					
	No	Working	Answer	Mark	Notes
10	(a)	$6y = 15 - 5x$	$y = \frac{15-5x}{6}$	2	M1 for either $6y = 15 - 5x$ or for $\frac{5x}{6} + y = \frac{15}{6}$ or $-6y = 5x - 15$ or a correct ft on sign error to $y =$ A1 for $y = \frac{15-5x}{6}$ oe
	(b)	$6k + 5(-21) = 15$	20	2	M1 for subst. of $x = -21$ (or $x = 21$) into given eq^n or answer to (a) A1 for $k = 20$
	(c)(i)		Region R indicated	3	B2 correct region shaded (accept unshaded if R clear) (B1 shaded (R) region satisfies 3 of the 4 given inequalities with same boundaries)
	(ii)		(1,1)		B1 for (1,1)
11	(a)		$y = 2x + 6$	2	B2 for $y = 2x + 6$ (B1 for $y = 2x + k$, $k \neq 1$ or for $y = mx + 6$, $m \neq 0$ or for $2x+6$)
	(b)	Grad of $AB = 2$; Grad of $BC = \frac{-1}{"2"}$	$y = -\frac{1}{2}x + 6$	2	M1 for Grad of $BC = \frac{-1}{"2"}, -\frac{1}{2}$ or grad of $BC = -\frac{1}{\text{grad of AB}}$ A1 ft for $y = -\frac{1}{2}x + 6$ oe ft on wrong coeff. of x in (a)
	(c)	Eg A rectangle is always a cyclic quadrilateral because the opposite angles of a rectangle always add up to 180° (90+90)		1	B1 for valid explanation. (eg lines from the pt of int. of diagonals of rect to all 4 vertices are equal(radii))
12		Vertices of new triangle at $(-1, -4)$ $(-4, 2)$ and $(2, 2)$	Correct triangle drawn	3	B3 cao (B2 for 3 correct vertices no triangle or triangle with 2 correct vertices (B1 for triangle with two of six co-ordinates correct from using P as centre or any isosceles triangle with base 6 and height 6)

PAPER 5505					
	No	Working	Answer	Mark	Notes
13	(a)		32	1	B1 for 32 (accept 31.5 to 33.5 inclusive)
	(b)			3	B1 for ends of whiskers at 9 and 57 (with a box) B1 for ends of box at 16 and 45/46 ($\pm$ 0.5) B1for median marked at "32"or complete box and whisker diagram drawn with a median inside the box
	(c)	Median(B)>Median(G); on average boys take longer IQR(B)>IQR(G); times for boys have a greater spread		2	B1 eg for comparison of medians (ft on diagrams) B1 eg for comparison of (interquartile) ranges (ft on diagram)
14		Reading top to bottom frequencies are 20; 18; 45; 52		2	B2 all correct (B1 for one frequency correct)
15	(a)	$1600 = p\,q^0$.; $400 = pq^2$ $q^2 = 0.25$	$p = 1600$ $q = 0.5$	3	M1 for either $400 = pq^2$ or $1600 = pq^0$ B1 for $p = 1600$ A1 for $q = 0.5$
	(b)	$V = pq^{-2}$	6400	2	M1 for recognition that t = –2 A1 for 6 400

PAPER 5505					
	No	Working	Answer	Mark	Notes
16	(a)	(I) $AB = AC$ (triangle ABC is isosceles) (II) PB = PC tangents (from a point to a circle are) equal (III) AP = AP (common side) so the 2 triangles are congruent ,SSS.		3	B3 for I, II, III with congruency reason (B2 for any two of I, II or III) (B1 for any one of the I, II or III)
	(b)	$BPC = 20°$ PBC (or PCB) = 90-1/2 "20" (= 80°) $BAC = PBC$ = "80"	50°	4	B4 for 50° (B3 for $BAC = 80°$) (B2 for $PBC = 80°$ or $PCB = 80°$) (B1 for $APC = 10°$ or $BPC = 20°$ or a middle angle = 90°) SC if clear numerical slip seen eg "PBC = 180 – 90 – 10 = 70" then goes on to get correct ft angle $ABC = 55$ deduct 1 from total so this cand would get B4–1 = B3
17	(a)	$\overrightarrow{OP} = \overrightarrow{OA} + \overrightarrow{AP}$ $= \overrightarrow{OA} + \frac{2}{3}(6\mathbf{c} - 6\mathbf{a})$ $= 6\mathbf{a} + 4\mathbf{c} - 4\mathbf{a}$	$2\mathbf{a} + 4\mathbf{c}$	3	M1 for $\overrightarrow{OP} = \overrightarrow{OA} + \overrightarrow{AP}$ or any correct vector journey involving $\overrightarrow{OP}$ M1 for $\overrightarrow{AP} = \frac{2}{3}(6\mathbf{c} - 6\mathbf{a})$ oe or $\overrightarrow{CP} = \frac{1}{3}(-6\mathbf{c} + 6\mathbf{a})$ oe or reverse vectors A1 for $2\mathbf{a} + 4\mathbf{c}$ oe (accept unsimplified)
	(b)	Eg $\overrightarrow{OM} = \overrightarrow{OC} + \overrightarrow{CM} = 6\mathbf{c} + 3\mathbf{a}$ $\overrightarrow{OM} = 1.5\overrightarrow{OP}$	$\overrightarrow{OM} = 1.5\overrightarrow{OP}$ so OPM is a straight line	2	B1 for $\overrightarrow{OM} = 6\mathbf{c} + \frac{1}{2}(6\mathbf{a})$ or $\overrightarrow{PM} = 2\mathbf{c} + \mathbf{a}$ unsimplified or reverse vectors B1 for a fully correct proof.

PAPER 5505					
	No	Working	Answer	Mark	Notes
18	(a)		4	1	B1 for 4 condone $\pm$ 4
	(b)		2	1	B1 for 2 condone ± 2
	(c)	$\sqrt{160} = 4\sqrt{10}$; $\left[\frac{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)-\sqrt{2}\times\sqrt{5}}{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)}\right]\times 100$ $\left[\frac{6\sqrt{10}-\sqrt{10}}{6\sqrt{10}}\right]\times 100$	$\frac{500}{6}$	4	B1 for **either** $\sqrt{160} = 4\sqrt{10}$ or $\sqrt{8} = 2\sqrt{2}$ **or** $\sqrt{20} = 2\sqrt{5}$ M1 for $\left[\frac{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)-\sqrt{2}\times\sqrt{5}}{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)}\right]$ oe $(\times 100)$ B1 for either $6\sqrt{10} - \sqrt{10}$ or $6\sqrt{10}$ A1 for $\frac{500}{6}$ (accept 83.3 if no obvious earlier error)

PAPER 5505					
	No	Working	Answer	Mark	Notes
19	(a)(i)		$(2x-7)(x-14)$	3	M1 x^2 term and constant term (± 98) obtained or $2x(x-14) - 7(x-14)$ or $x(2x-7) - 14(2x-7)$ A1 for $(2x-7)(x-14)$
	(ii)		$x = \frac{7}{2}$; $x = 14$		B1ft ft (i) provided of form $(2x\pm a)(x\pm b)$
	(b)(i)		$\frac{7}{n+7}$	3	B1 for $\frac{7}{n+7}$ oe
	(ii)	$\frac{7}{n+7} = \frac{2}{5} \Rightarrow 2(n+7) = 5 \times 7$ $2n = 21$	$n=10.5$ is not possible since n has to be an integer		M1 for $2(n+7) = 5\times 7$ or $n+7=5\times 3.5$ (can be implied) ft (b)(i) fractional in terms of n and < 1 A1 ft for $n=10.5$ not possible (since n not integer) oe
	(c)	$2\times\left(\frac{n}{n+7}\right)\times\left(\frac{7}{n+7}\right) = \frac{4}{9}$ $14n\times 9 = 4(n+7)^2$ $14n\times 9 = 4(n^2+14n+49)$ $4n^2+56n+196-126n = 0$	$2n^2 - 35n + 98 = 0$	5	M1 for $\left(\frac{n}{n+7}\right)\times\left(\frac{7}{n+7}\right)$ seen M1 for $2\times\left(\frac{n}{n+7}\right)\times\left(\frac{7}{n+7}\right)$ oe $(=\frac{4}{9})$ M1(dep on 1st M) elimination of fractions within an equation B1 3 terms correct in expansion of $(n+7)^2 = n^2 + 7n + 7n + 49$ A1 full valid completion to printed answer

PAPER 5505					
No		**Working**	**Answer**	**Mark**	**Notes**
	(d)	$\frac{7}{n+7}\times\frac{7}{n+7}=\frac{7}{21}\times\frac{7}{21}=$	$\frac{1}{9}$	2	M1 for $\frac{7}{n+7}\times\frac{7}{n+7}$ or better or ft [answer (b)(i)]^2 or $1-\frac{4}{9}-(\frac{n}{n+7})^2$ A1 for $\frac{1}{9}$ oe cao
20	(a)(i)		$y=1+\sin x$	2	B1 for $y=1+\sin x$
	(ii)		$y=2\sin x$		B1 for $y=2\sin x$ SC both (i) f(x) + 1, (ii) 2f(x) B1
	(b)	Stretch parallel to y-axis scale factor 3 Stretch parallel to x-axis scale factor $\frac{1}{2}$		3	M1 for 'stretch' A1 for Stretch parallel to y-axis scale factor 3 oe A1 for Stretch parallel to x-axis scale factor $\frac{1}{2}$ oe SC if M0 award BI for "sf 3 vertically" **and** "sf $\frac{1}{2}$ horizon."

PAPER 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
1		Question	2	B1 for each of the following upto B2 (a) an improved question eg time frame made clear (b) response boxes (imperfect) (c) response boxes no errors OR For suggesting a generally improved question (a) a question clearly in the context of changes to the canteen (b) at least 3 boxes showing a full range of responses
2	Locus (circle) drawn & shaded		2	B1 circle centre Manchester ±2mm B1 shading with accurate or approximate circle within tolerance
3 (a)		182, 178, 180, 184	2	M1 mean of any three consecutive months, eg (147 + 161 + 238) ÷ 3 oe A1 cao
(b)	Sale price = 80% Fun Friday price = 70% of 80% = 56% (oe)		2	B1 B1 for a fully correct explanation involving a worked example (oe) B1 a partially complete explanation
4	x $x^3 - 2x$ 4.1 60.7(21) 4.2 65.6(88) 4.3 70.9(07) 4.22 66.7(114) 4.23 67.2(269) 4.24 67.7(45) 4.25 68.2(656)	4.2	4	B2 for trial $4.1 \leq x \leq 4.3$ evaluated (B1 for trial $4 < x < 5$ evaluated) B1 for different trial $4.225 \leq x \leq 4.25$ evaluated B1 (dep on at least one previous B1) for 4.2 cao

PAPER 5506					
	No	Working	Answer	Mark	Notes
5	(a)		1.0×10^{-9}	1	B1 accept 1×10^{-9} or just 10^{-9}
	(b)	$1 \div (5 \times 10^{-9})$	2×10^{8}	2	M1 for $1 \div$ ("5×10^{-9}") or digit 2 with zeros only seen Condone omission of bracket for M1. A1 cao
6	(a)	$\frac{6.27 \times 4.52}{4.81 + 9.63} = \frac{28.3404}{14.44} = 1.962631579$	1.9626…	2	B2 for 1.9626… (B1 for 28.34… or 14.44)
	(b)		1.96	1	B1 ft from (a) as rounded to 1dp or 2dp. Do not accept 2, 2.00, but accept 2.0
7			$d = 4n + 6$	2	B2 $d = 4n + 6$ oe (B1 $d = 4n + k$, k an integer $\neq 6$, $4n + 6$, $n = 4n + 6$)
8		Vol = $\pi \times 3.8^2 \times 2.5 = \pi \times 14.44 \times 2.5$ $= 45.36\ldots \times 2.5 = 113.411$ Mass = "113" × 1.5 = 170.1165	170	4	M1 for $\pi \times r^2 \times 2.5$ where r = is 3.8 or 7.6 A1 if $r = 3.8$ M1 for "113" × 1.5 A1 for 169.5 – 170.3 cao

PAPER 5506					
	No	Working	Answer	Mark	Notes
9	(a)	$10^2 + 6^2$ or 136 $\sqrt{(100 + 36)}$ or $\sqrt{136} = 11.66\ldots$	11.7	3	M1 for $10^2 + 6^2$ or 136 seen M1 (dep) $\sqrt{100+36}$ or $\sqrt{136}$ A1 11.66 – 11.7
	(b)	Cos x = 8/10 or 0.8 $x = \cos^{-1} 0.8 = 36.869^o$	36.9	3	M1 for cos = $\frac{8}{10}$, cos = 0.8(oe) M1 (dep) for $\cos^{-1}$ (oe) A1 for 36.86 – 36.9 M1 Use of sine rule and x found M1 for $x = 90 - \sin^{-1}$("0.8") A1 for 36.86 – 36.9
10		$18x - 6y = 99$ $\quad 12x - 4y = 66$ $8x + 6y = 18$ $\quad 12x + 9y = 27$ $26x = 117$ $\quad -13y = 39$	$x = 4\ ½$ $y = –3$	4	M1 correct process to eliminate either x or y (condone one error) A1 cao for non – eliminated one. M1 (dep on 1st M1) for correct substitution of their found value. A1 cao (need both)

PAPER 5506					
	No	Working	Answer	Mark	Notes
11		$\frac{133}{72} = 1.8472\,,\ \frac{160}{82} = 1.9512$ OR $\frac{72}{133} = 0.54135, \frac{82}{160} = 0.5125$ OR $\frac{160}{133} = 1.203...., \frac{82}{72} = 1.1388....$ OR $\frac{133}{160} = 0.83125...., \frac{72}{82} = 0.878$	1.84..≠ 1.95.. 1.20..≠ 1.13..	3	M1 for $\frac{133}{72}$ (= 1.8472…) oe Accept 1.8, 1.85 M1 for $\frac{160}{82}$ (=1.9512…) oe consistent pairing Accept 2.0, 1.9 **OR** M1 for $\frac{160}{133}$ (= 1.203…) oe M1 for $\frac{82}{72}$ (=1.1388) oe A1 for enough decimal places to show that the ratios are not equal; since the scale factors are different the shapes cannot be similar. NB Do Not need conclusion
12	(a)	£12000 × 0.25 = £3000; £12000 – £3000 = £9000 £9000 × 0.25 = £2250; £9000 – £2250 = £6750 £6750 × 0.25 = £1687.50; £6750 – £1687.50 =	£5062.50	3	M1 for 12000 × 0.75 (=9000) oe or £3000 or £23437.50 seen M1 (dep) for at least two further depreciation calculations (complete steps) A1 cao **OR** M2 for 12000 × $(0.75)^3$ or 5062.50 seen (M1 for 12000 × $(0.75)^n$, $n = 2$ or 4)
	(b)	0.8 × 0.8 × 0.8 × 0.8 (oe)	0.4096	2	M1 0.8^4 (oe) A1 cao

PAPER 5506					
No		**Working**	**Answer**	**Mark**	**Notes**
13		$r^2 = \frac{3\times10}{\pi h} = \frac{3\times10}{\pi 1.5} = \frac{3\times10}{4.712..} = 6.36, 6.37$ $r = \sqrt{6.366\ldots}$	2.52-2.54	3	M1 for correct rearrangement to give $r^2 = \frac{3V}{\pi h}$ or $30 \div 4.712..$ or $6.36 - 6.37$ *Allow 0.3, 0.33 for* $\frac{1}{3}$ M1 (dep) for $\sqrt{}$ A1 cao 2.52 – 2.54
14	(a)		0.4 0.6,0.4, 0.6,0.4	2	B1 for LHS: (0.6), 0.4 B1 for RHS: 0.6, 0.4, 0.6, 0.4
	(b)	0.6×0.6	0.36	2	M1 $0.6 \times$ "0.6" [0 < "0.6" < 1] A1 cao
	(c)	$(30 \times 42) - (25 \times 42.8) = (1260 - 1070 = 190$ $190 \div 5 =$	38	3	M1 for $(30 \times 42) - (25 \times 42.8)$ or $1260 - 1070$ or 190 seen M1(dep) for "190" ÷ 5 A1 cao 38

PAPER 5506					
	No	**Working**	**Answer**	**Mark**	**Notes**
15	(a)	$S = \frac{k}{f^2};\ 125 = \frac{k}{8^2};\ k = 8000$	$S = \frac{8000}{f^2}$	3	M1 for $S = \frac{k}{f^2}$ M1 for $125 = \frac{k}{8^2}$ A1 cao These marks can be awarded if the full formula appears in part (b), rather than in part (a). [SC: $S = \frac{125}{64} f^2$ M1 M0 A0, $S = 1.95(3125) f^2$ M1 M0 A0 $S = \frac{1000}{f}$ M1 M0 A0]
	(b)		500	1	B1 cao

PAPER 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
16	$0.5 \times 3.2 \times 8.4 \times \text{Sin } B = 10$ $\sin B = 0.74404\ldots$ 48.077 $AC^2 = 3.2^2 + 8.4^2 - 2 \times 3.2 \times 8.4 \times \cos B$ $AC^2 = 44.8815\ldots..$ $AC = 6.69\ (936\ldots)$ Perimeter = 18.3 Use the altitude AD, $\frac{h \times 8.4}{2} = 10 \Rightarrow h = (2.381)$ $BD = \sqrt{3.2^2 - h^2} = 2.139$ $DC = 6.261$ $AC = \sqrt{2.38^2 + '6.261'^2} = 6.69(936)$ Perimeter = 18.3	18.3	6	M1 for $0.5 \times 3.2 \times 8.4 \times \sin B\ (= 10)$ A1 for $\sin B = 0.74(404\ldots)$ or $B = 47.7 - 48.1$ M1 for $3.2^2 + 8.4^2 - 2 \times 3.2 \times 8.4 \times \cos$ "48.077" M1(dep) for AC^2 = "44.8(815)"….. with correct order of evaluation A1 $AC = 6.69 - 6.7$ A1 18.29 – 18.3 **for** $\frac{h \times 8.4}{2} = 10 \Rightarrow h = (2.381)$ M1 for $BD^2 = 3.2^2 - "2.381"^2$ A1 $BD = 2.1 - 2.2$ M1 (dep) $AC^2 = "2.381"^2 + "6.261"^2$ A1 $AC = 6.69 - 6.7$ A1 18.29 – 18.3

PAPER 5506					
	No	Working	Answer	Mark	Notes
17	(a)	$(x + 8)^2 = x^2 + (x + 5)^2$ $x^2 + 16x + 64 = 2x^2 + 10x + 25$		4	B1 for angle $OTA = 90°$ (implied by use of Pythagoras with OA as hypotenuse) M1 $(x + 8)^2 = x^2 + (x + 5)^2$ oe M1 for correct squaring of x+8 or x+5 A1 for completion following correct working
	(b)	$(+6 + \sqrt{(6^2 - - 4\times39)})/2$ $= (6 \pm \sqrt{(36 + 156)})/2$ $= (6 \pm \sqrt{192})/2 = (6 + 13.856)/2$	9.93	3	M1 for substitution into quadratic formula, allow sign errors in b and c M1 for $x = (6 \pm \sqrt{192})/2$ [+ alone will do] A1 for 9.92 – 9.93
					M1 for $(x-3)^2-9-39$ M1 for $x = 3 \pm \sqrt{48}$ [+ alone will do it] A1 for 9.92 – 9.93 T&I= 9.93 gets M1,M0,A0
18		$EB = 60 \times \tan 30°$ $BD = \sqrt{(60^2 + 60^2)}$ $\tan BDE = 34.64 \div 84.85$	22.2°	4	M1 for $EB = 60\times\tan30$ M1 for $BD = \sqrt{(60^2 + 60^2)}$ M1 for $\tan BDE =$ "34.64" ÷ "84.85" A1 22.17 – 22.21
		$EB = 60 \times \tan 30°$ (= 34.64) $ED^2 = 60^2 + \left(\frac{60}{\cos 30}\right)^2$ $ED = \sqrt{8400} = (91.65)$ $\text{Angle} = \sin^{-1}\left(\frac{EB}{\sqrt{8400}}\right) = 22.2$			M1 for $EB = 60 \times \tan 30°$ oe M1 for fully correct method for ED M1 for $\sin BDE = \left(\frac{'34.84'}{'\sqrt{8400}'}\right)$ (oe) A1 22.17 – 22.21

PAPER 5506					
	No	Working	Answer	Mark	Notes
19	(a)	$x^2 + 6^2 = 25$ $x^2 = -11$		2	M1 substitutes y=6 to get x^2+36 = 25 A1 deduces x^2=25-36<0; impossible, (so line does not intersect curve)
	(b)	$y^2 = (2x - 2)^2$ $x^2 + (2x - 2)^2 = 25$ $x^2 + 4x^2 - 8x + 4 = 25$ $5x^2 - 8x + 4 - 25 = 0$ $5x^2 - 8x - 21 = 0$ $(5x + 7)(x - 3) = 0$ or $x = \frac{-(-8) \pm \sqrt{(-8)^2 - 4 \times 5 \times (-21)}}{10} = \frac{8 \pm \sqrt{484}}{10}$ So $x = 3$, or $x = -1.4$ Subs into either equation gives: $y = 4$, or $y = -4.8$	$x = 3$ $y = 4$ $x = -1.4$ $y = -4.8$	6	M1 $y^2=(2x-2)^2$ seen or implied A1 $x^2 + 4x^2 - 8x + 4 = 25$ (oe expanded form) M1 correct attempt to solve 3 term quadratic equation A1 $x = 3$, $x = -1.4$ M1 (dep on previous Ms) Sub one value of x into either equation gives: A1 $y = 4$, $y = -4.8$ NB: incorrectly matched pairs loses the final A1.

PAPER 5506					
	No	Working	Answer	Mark	Notes
20	(a)	400.005 ÷ 59.95 = 6.672310 6.672310 ÷ 1000 × 3600 = 24.02032 399.995 ÷ 60.05 = 6.661032 6.661032 ÷ 1000 × 3600 = 23.97972	24.020 23.980	5	B1 for 400.005 or 59.95 seen oe M1 for "400.005" ÷ "59.95" where "400.005" ∈ [400.005,400.5] and "59.95" ∈ [59.5,59.95] oe B1 for 399.995 or 60.05 seen oe M1 for "399.995" ÷ "60.05" where "399.995" ∈ [399.5,399.95] and "60.05" ∈ [60.05,60.5] oe A1 23.979-23.980 and 24.020-24.0204
	(b)	24.0 because to 1dp the answers are the same		1	B1cao for 24.0 with reason
	(c)	177 × 50/477 = 18.553 → 19 → 18 111 × 50/477 = 11.635 → 12 86 × 50/477 = 9.0147 → 9 82 × 50/477 = 8.595 → 9 21 × 50/477 = 2.201 → 2	18 12 9 9 2	3	M1 for $\frac{50}{"477"} \times 177$ or 111 or 86 or 82 or 21 A1 for all integers or better answers, at least 3 correct A1 cao
21	(a)	$40 - x = 3(4 + x)$ $40 - x = 12 + 3x$ $40 - 12 = x + 3x$ $4x = 28$	7	3	M1 multiplying through by 3: $3 \times \frac{40 - x}{3} = 3 \times 4 + 3 \times x$ A1 $40 - 12 = x + 3x$ A1 cao
	(b)	$\frac{2x(2x - 3)}{(2x - 3)(2x + 3)} = \frac{2x}{2x + 3}$	$\frac{2x}{2x + 3}$	3	B1 for $(2x\text{-}3)(2x\text{+}3)$ B1 for $2x(2x\text{-}3)$ or $(2x + 0)(2x + 3)$ B1 cao

PAPER 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
22	Distance from x axis is y. Distance from (0, 2) is $\sqrt{(x^2+(y-2)^2)}$ $y^2 = x^2 + (y-2)^2$ $y^2 = x^2 + y^2 - 4y + 4$ $0 = x^2 - 4y + 4$ $4y = x^2 + 4$ and finish		4	B1 for $(x-0)^2 + (y-2)^2$ or $\sqrt{((x-0)^2+(y-2)^2)}$ oe seen B1 for $y = \sqrt{(x-0)^2+(y-2)^2}$ or $y^2 = (x-0)^2 + (y-2)^2$ oe B1 $(y-2)^2 = y^2 - 4y + 4$ seen B1 for $4y = x^2 + 4$ and finish

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone: 01623 467467
Fax: 01623 450481
Email: publications@linneydirect.com

Order Code UG015616 July 2004

For more information on Edexcel and BTEC qualifications please contact
Customer Services on 0870 240 9800
or http://enquiries.edexcel.org.uk
or visit our website: www.edexcel.org.uk

London Qualifications Limited, trading as Edexcel. Registered in England and Wales No. 4496750
Registered Office: Stewart House, 32 Russell Square, London WC1B 5DN



edexcel

Edexcel GCSE

Mathematics A 1387

June 2007

advancing learning, changing lives

Mark Scheme

Edexcel GCSE

Mathematics A 1387

Table Of Contents

Notes on Marking Principles	3 - 4
5521 / 01 Mark Scheme	5 - 11
5521 / 02 Mark Scheme	12 – 17
5523 / 03 Mark Scheme	18 – 27
5523 / 04 Mark Scheme	28 – 35
5525 / 05 Mark Scheme	36 – 43
5525 / 06 Mark Scheme	44 - 52

NOTES ON MARKING PRINCIPLES

1 **Types of mark**

M marks: method marks
A marks: accuracy marks
B marks: unconditional accuracy marks (independent of M marks)

2 Abbreviations

cao – correct answer only
ft – follow through
isw – ignore subsequent working
SC: special case
oe – or equivalent (and appropriate)
dep – dependent
indep - independent

3 No working

If no working is shown then correct answers normally score full marks
If no working is shown then incorrect (even though nearly correct) answers score no marks.

4 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.
If there is no answer on the answer line then check the working for an obvious answer.

5 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

6 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

7 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

8 Linear equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

9 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

Remember: if you are having difficulty making a decision on how you should mark a candidate response contact your Team Leader for advice, or send the item to review.

Paper 5521_01					
No		Working	Answer	Mark	Notes
1		Cat |||| /||| 8 Dog |||| | 6 Fish || 2 Hamster |||| 4	8, 6, 2, 4	3	M1 for attempt to tally or one frequency correct in either column A1 for 1 frequency correct or all tallies correct in correct column A1 for all frequencies correct (accept if /20)
2			27, 35, 42, 67, 118	1	B1 cao
3	(a)		Diameter drawn	1	B1 for a diameter
	(b)		Right angle marked	1	B1 *R* marked correctly
	(c)		Rectangle drawn	1	B1 for a rectangle
4	(a)		40	1	B1 cao
	(b)		50	1	B1 cao
	(c)		3 full loaves 4 full loaves + 1 half loaf	2	B1 for 3 full loaves B1 for 4 full loaves + 1 half loaf
5	(a)		7252	1	B1 cao
	(b)		Three thousand and eighty six	1	B1 accept 3 thousand and eighty six (condone 0 hundred)
	(c)		4600	1	B1 accept 4600
	(d)		200	1	B1 for 200 or 2 hundred or 100 or hundred
6	(i)		Cube	2	B1 for 'cube' (accept 'cuboid') ignore spelling
	(ii)		Cylinder		B1 for 'cylinder' ignore spelling
7		5×100	500	2	B2 for 490 or 500 or 510 (B1 for either 5 or 5.0 or 100 seen)

Paper 5521_01					
No		Working	Answer	Mark	Notes
8	(a)		8.4 cm or 84 mm	2	B2 allow ±2mm (B1 for 8.4 or 84, B1 for appropriate unit)
	(b)		37°	1	B1 allow ±2°
9	(a)		Carbon black	1	B1 accept 'black carbon' accept 26%
	(b)		0.1(0)	1	B1 cao
	(c)		0.04	1	B1 cao
	(d)	$\frac{26}{100}$	$\frac{13}{50}$	2	M1 for $\frac{26}{100}$ A1 cao
10	(a)		97	1	B1 cao
	(b)		London Reading	1	B1 cao
	(c)	41 + 57 + 58	156	3	M1 for two of 41, 57, 58 M1(dep) for '41' + '57' + '58' A1 cao
11	(a)		3	1	B1 cao allow ± 0.2
	(b)		–5	1	B1 cao allow ±0.2

Paper 5521_01					
No		**Working**	**Answer**	**Mark**	**Notes**
12	(a)		Centimetres (cm) miles litres (*l*)	3	B3 (B1 for each correct answer) accept abbreviations
	(b)		300	1	B1 cao
	(c)	'1500>1400' or '1.5>1.4'	Reason	1	B1 for No and '1500>1400' or '1.5>1.4'
13	(a)	Height of bars 12, 8, 6 lines drawn between points	Bars drawn	2	B2 for 3 bars correctly drawn (B1 for 2 bars correct)
	(b)		July and August	1	B1 oe
	(c)	24 – 4	20	1	B1 (Accept '4 to 24' oe)
	(d)	The temperatures are rising	Temp's rising oe	1	B1 for reason
14	(a) (i)		$(-4, 3)$	2	B1 cao
	(ii)		$(2, -1)$		B1 cao
	(b) (i)	*D* marked at $(-4, -1)$	Point marked on grid	2	B1 for point marked at $(-4, -1)$ cao
	(ii)		$(-4, -1)$		B1 ft
15	(a)			1	B1 cao
	(b)			1	B1 cao

Paper 5521_01				
No	**Working**	**Answer**	**Mark**	**Notes**
16 (a)		(Pat +) reason	1	B1 correct comment (Pat may be implied)
(b)	21 ÷ 3	7	1	B1 cao
17 (i)		S marked at 1	3	B1 for S within ½ cm of 1
(ii)		P marked at 0		B1 for P marked at 0 cao
(iii)		Q marked at "1/3"		B1 for Q marked at 1/3 ± 1 cm use overlay
18	6 + 6 + 3 or $2\frac{1}{2}\times 6$	15	3	M1 for realizing 6 glasses in one bottle M1 for realizing 3 glasses in ½ a bottle A1 cao (M2 for attempt to find $2\frac{1}{2}\times 6$) oe
19	15 and 16 parts shaded Alternative 1 $\frac{3}{4}=0.75,\ \frac{4}{5}=0.8$ Alternative 2 $\frac{3}{4}=\frac{15}{20},\ \frac{4}{5}=\frac{16}{20}$	$\frac{4}{5}$ + reason	3	M1 for correctly shading 15 parts for 3/4 M1 for correctly shading 16 parts for 4/5 A1 (dependent on M2) for selection of 4/5 Alternative 1 M1 for $\frac{3}{4}=0.75$ M1 for $\frac{4}{5}=0.8$ A1 (dep on M2) for selection of 0.8 or $\frac{4}{5}$ or $\frac{16}{20}$ Alternative 2 M1 for $\frac{3}{4}=\frac{15}{20}$ M1 for $\frac{4}{5}=\frac{16}{20}$ A1(dep on M2) for selection of $\frac{4}{5}$ or $\frac{16}{20}$

Paper 5521_01				
No	**Working**	**Answer**	**Mark**	**Notes**
20 (a)		−5 (−3) (−1) 1 3 5	2	B2 cao (B1 for any 2 or 3 correct)
(b)	Points plotted	Line	2	B2 for line from (−1, −5) to (4, 5) (B1 ft for plotting at least 5 "points")
21	Shapes shaded on grid	6 tessellating shapes	2	B2 for fully correct with 5 or more additional shapes, no gaps (B1 for 4 shapes tessellating with at least one shape inverted, with or without the given shape ignore extras)
22 (a)		$\frac{7}{20}$	1	B1 for $\frac{7}{20}$ oe
(b)	7 + 2 (or 20 – 11) are not lime flavour	$\frac{9}{20}$	1	B1 for $\frac{9}{20}$
(c)		0	1	B1 for 0, zero or nought ($\frac{0}{20}$ gets B0)
23 (a)		$4a$	1	B1 accept $4\times a$, $a \times 4$, $a4$
(b)		$12b$	1	B1 accept $12\times b$, $b \times 12$, $b12$
(c)		$2a + 6b$	2	B2 cao (B1 for $2a$ or $6b$ seen)
(d)		$x(x-6)$	2	B2 cao (B1 for $x(ax+b)$ where a, b are numbers not equal to 0 or $x - 6$ seen on its own, or as part of an expression)
24 (a)		40	1	B1 cao
(b)		45	1	B1 for 42 – 48 accept ¾ hour
(c)	40×2 or $\frac{40}{30}\times 60$ or $40\div\frac{1}{2}$	80	2	M1 for 40×2 or $\frac{40}{30}$ or $40\div\frac{1}{2}$ A1 cao NB $\frac{40}{45}\times 60$ gets M0 A0

Paper 5521_01					
No		**Working**	**Answer**	**Mark**	**Notes**
25	(a)		4560	1	B1 cao
	(b)		45.6	1	B1 cao
	(c)		2.4	1	B1 cao
26		$-7 - 3 = -10$ $2 \times -10 = -20$ $-20 \div 4$	-5	3	M1 for substitution of 2 and –7 into $p(q - 3)$ or sight of -20 or $-14 - 6$ M1dep for '–20' ÷ 4 A1 cao B1 SC for sight of –10 if M0 awarded
27	(a)		Reflection in *y*-axis	1	B1 cao
	(b)		Rotation by half turn about (0,0)	2	B2 cao (B1 for half turn not about (0,0).)
	(c)		Enlargement Scale factor 3 Centre (0,0)	3	B1 for 'enlargement' B1 for "scale factor 3" or 3 seen B1 for 'centre (0,0)
28	(a)		Reason	1	B1 for 'The frequencies are nearly equal' oe
	(b)	$1\times26+2\times26+3\times23+4\times25 = 247$ $247/100$	2.47	3	M1 for fx (attempting at least 2 relevant products) M1 for $\sum fx \div 100$ A1 2.47 cao

Paper 5521_01				
No	**Working**	**Answer**	**Mark**	**Notes**
29	5 × 5 × 6	150	4	M1 for attempt at 1 division (e.g. 40 ÷ 8), may be implied by marks or number on one edge of diagram or by 5 or 6 seen M1 for attempt at 3 divisions (40÷8, 40÷8, 60÷10), may be implied by marks or numbers on diagram or by 5,5 and 6 seen. M1 (dep on 1st M1) for "5" × "5" × "6" A1 cao Alternatively M1 for 40 × 40 × 60 **or** 8 × 8 × 10 **or** 96000 or 640 seen M1 for 40 × 40 × 60 **and** 8 × 8 × 10 or 96000 **and** 640 seen M1 (dep on 1st M1) for "(40 × 40 × 60)" ÷ "(8 × 8 × 10)" A1 cao SC:B1 for dividing area of one carton face by area of corresponding box face if M0

Paper 5521_02					
No		Working	Answer	Mark	Notes
1	(a)	Draw diagram.	Diagram	1	B1 cao
	(b)		13,16	1	B1 cao
	(c)		31	1	B1 cao
2	(a)		8	1	B1 cao
	(b)		14	2	B2 for 14 (B1 for 13 or 15)
	(c)		16	2	B2 for 16 (B1 for 15, 17 or 8)
3	(a)		4,7 drawn	2	B2 for car height 4 and bus height 7, (B1 for one correct)
	(b)		6	1	B1 cao
	(c)		Walk	1	B1
	(d)		27	1	B1 cao
4	(a)		40	1	B1 for 40–41 inclusive
	(b)		12	1	B1 for 11.5 – 12.5 inclusive
5	(a)		Row complete	2	B2 for 1+..+11; 36 (B1 for one of the 2 cells complete)
	(b)		Square	1	B1 "square"
6		One line of symmetry		1	B1 within 2mm of centre of base / 2mm of vertex

Paper 5521_02				
No	**Working**	**Answer**	**Mark**	**Notes**
7 (a) (i)		09 06	2	B1 (accept 9 06 oe)
(ii)		39		B1 cao
(b)		06 55	1	B1 (accept 6 55 oe)
(c)		2h 6min	1	B1 cao
(d)		15 min	2	M1 for 0906 – 0645 – "(c)" or 0906 – 0645 − 2hr 6min or 2hr 21min − "(c)" or 2hr 21min − 2hr 6 min or 141 − 126 or 20 − 5 A1 cao SC: B1 for 55 or 75 or 93 seen
8 (i)		8,10,12,20 or 30	3	B1 at least one of 8, 10, 12, 20, 30 (no extras)
(ii)		8,12 or 20		B1 at least one of 8,12, 20 (no extras)
(iii)		3 or 5		B1 3 or 5 or both (no extras)
9 (a)		C or G	1	B1 at least one of C or G (no extras)
(b)		A and F	1	B1 cao
(c)		2	1	B1 (accept –2)
10 (a)		$\frac{7}{10}$	1	B1 7/10 oe
(b)(i)		4 squares	2	B1 4 squares shaded
(ii)		80%		B1 80% or ft from unshaded part (no ft from 0% or 100%)
(c)(i)		2.5	2	B1 2.4–2.6 inclusive
(ii)		1.7		B1 1.6–1.8 inclusive
11 (a)		2	1	B1 for 2 or –2
(b)		14	1	B1 for 14 or –14

Paper 5521_02					
No		**Working**	**Answer**	**Mark**	**Notes**
12		2 × 8.50 = 17.00 3 × 4.50 = 13.50 Total = 30.50 50.00 – 30.50	19.5(0)(*p*)	3	M1 for adding 5 correct values or 2 × 8.50 + 3 × 4.50 (ignore units) or 30.5(0) or 3050 seen M1 dep for 50 − "30.50" (ignore units) (OR M1for adding at least 1 adult ticket and at least 1 child ticket **and** subtracting from 50) A1 cao SC: B1 for 24 or 37 or 2400 or 3700 seen
13	(a)		Hexagon	1	B1
	(b)(i)		120	2	B1 cao
	(ii)		Str line		B1 reference to a (straight) line and 180°
	(c)		Obtuse	1	B1 Accept "interior"
14	(a)	2, 2, 3, 3, 3, 4, 4, 4, 5, 6	3.5	2	M1 ordering the numbers (condone 1 error or omission) A1 cao
	(b)	36 ÷ 10	3.6	2	M1 sum of numbers ÷ 10 A1 cao SC B1 for 3r 6
	(c)	6 – 2	4	1	B1 cao
15	(a)		Paul	1	B1 cao
	(b)	36 ÷ 2 oe	18	1	B1 cao
	(c)	60/360 =	1/6	2	M1 60/360 oe A1 cao

Paper 5521_02					
No		**Working**	**Answer**	**Mark**	**Notes**
16		4.7 ÷ 5.9 = 0.796610169	0.7966..	2	B2 for 0.7966 or better (B1 for 0.8, 0.80, 0.79, 0.796, 0.797 or digits 59 seen
17		$6x - 7 + 7 = 38 + 7$ $6x = 45$	7.5	2	M1 $6x = 45$ or +7 both sides A1 7.5 oe; accept 45/6 oe
18		**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	3	B3 all six entries correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
19	(a)	900 × 1.70 =	1530	2	M1 900 × 1.7(0) or digits 153(0) seen A1 cao
	(b)	160 ÷ 1.70 =	£94.12 or £94.11	2	M1 160 ÷ 1.7(0) or digits 941(....) seen A1 cao
20	(a)(i)	180 – 54 (=126) "126" ÷ 2	63	3	M1 for (180–54) ÷ 2 A1 cao
	(ii)		Reason		B1 (indep) angles in triangle add to 180 OR equal angles in isosceles triangle OR equal angles and 2 sides the same (B0 if any incorrect reasoning given eg parallel, equilateral triangle)
	(b)	180 – "x"	117	1	B1 117 or ft 180 – "x" if $x < 90$
21	(a)	3 × 35 + 50	155	2	M1 for 3 × 35 + 50 or digits 155 seen A1 cao
	(b)	260 – 50 = 210 210 ÷ 35 =	6	3	M1 for 260–50 or 210 seen. M1 for "260-50" ÷ 35 or 210÷35 A1 cao SC B1 for starting at a number between 100 and 170 and adding at least two 35's and showing a total between 230 and 290 OR For adding at least three 35's, perhaps with other numbers, and showing a total between 180 and 240 (or between 230 and 290 if 50 is included in the sum)

Paper 5521_02				
No	**Working**	**Answer**	**Mark**	**Notes**
(c)		P=35h + 50	3	B3 for P=35h+50 or P=35×h+50 oe (B2 for correct RHS or P=h + 50 ×35 or P=35h+k where k is numerical oe) (B1 for P = some other linear expression in h, OR h + 50 ×35 OR 35h seen) NB: P=h scores no marks; ignore £ signs. SC B2 for $h = \frac{P-50}{35}$
22 (a)		Elevation	2	B2 for 4 vertical squares. Accept 4 by 1 rectangle. (B1 for 4 vertical squares with one square added or one parallelogram added at the top, or 3 vertical squares, or 4 horizontal squares)
(b)		Plan	2	B2 for 2 adjacent squares, vertical or horizontal. Accept 2 by 1 rectangle. (B1 for 3 adjacent horizontal or vertical squares or a rectangle with sides in the ratio 2:1)
23 (i) (ii) (iii)		5 9 6	3	B1 cao B1 cao B1 cao
24	$45.00 + 45.00 \times \frac{15}{100} =$ $45.00 + 6.75 =$	51.75	3	M2 for $45.00 + 45.00 \times \frac{15}{100}$ oe or 45.00×1.15 oe OR $45.00 + 6.75$ OR complete method or 5175 seen. (M1 for $45.00 \times \frac{15}{100}$ oe OR 6.75 seen OR 675 seen OR correct method for calculating 15% of 45) A1 cao SC Award B2 for an answer of 38.25

Paper 5521_02				
No	**Working**	**Answer**	**Mark**	**Notes**
25 (a)		Points	1	B1 all three points ±1 full square
(b)		Negative	1	B1 Negative (ignore additional descriptors unless contradictory)
(c)		lobf	1	B1 A single straight line drawn to cross between (5,30), (5,40) and (40,0), (40,15); accept freehand if considered to be straight.
(d)(i)		18–25	2	B1 18g–25g inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient.
(ii)		30–40		B1 30–40 min inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient
26		300 3 75 150	3	B3 for 4 correct answers (B2 for 2 or 3 correct answers) (B1 for 1 correct answer)
27	π × 0.65	2.04–2.05	2	M1 for π × 0.65 or 3.14 x 0.65 or 3.142 x 0.65 A1 for 2.04–2.05 SC Award B1 for 2.0 seen (not 2)
28	5 miles = 8 km 70mph÷5×8 = 112 km/h OR 120km/h ÷ 8 × 5 = 75 mph Faster than 70 mph	70mph (Great Britain) (112 km)	3	M1 5 miles = 8 km; OR 70 mph is about 100 km/h OR 1km=0.6(25) miles OR 1mile=1.6km oe M1 70 ÷ 5 × 8 (=112) or 120 ÷ 8 × 5 (=75) A1 (dep on at least M1) GB or 70 mph Refer to both answer line and working. NB GB or 70 mph without working scores 0 marks

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
1		Cuboid drawn	2	B2 for correct isometric drawing in any orientation (ignore points 'behind', mark 7 vertices only); accept lines drawn near to dots as long as there is no ambiguity. (B1 for one of the three faces drawn correctly **or** for an isometric drawing of any cuboid)
2	Different makes of car Tally Frequency	Make of car Tally Frequency	3	B1 for make of car or list of at least 3 different makes B1 for tally or tally marks B1 for frequency or totals
3		6 tessellating shapes	2	B2 for fully correct with 5 or more additional shapes, no gaps (B1 for 4 or more shapes tessellating, with at least one shape inverted, with or without the given shape, ignore extras)
4	24.90 ÷ 3 or 8.30 24.90 – '8.30' or 2 × 8.30	16.6(0)	3	M1 for 24.90 ÷ 3 or 8.30 M1 (dep) for 24.90 – “8.30” or 2 × “8.30” A1 for 16.60 or 16.6

Paper 5523_03

No	Working	Answer	Mark	Notes
5	315 × 24 24 × 315 1260 6300 7560 7200 7560 **3 1 5** 0 6 | 0 2 | 1 0 — **2** 1 2 | 0 4 | 2 0 — **4** 7 5 6 0 **300 10 5** 6000 | 200 | 100 — **20** 1200 | 40 | 20 — **4** 6000+200+100+1200+40+20 = 7560 **3 0.1 0.05** 60 | 2 | 1 — **20** 12 | 0.4 | 0.2 — **4** 60 + 2 + 1 + 12 + 0.4 + 0.2 = 75.6	75.6(0)	3	M1 for a complete method with relative place value correct. Condone 1 multiplication error, addition not necessary. **OR** M1 for a complete grid with not more than 1 multiplication error, addition not necessary. **OR** M1 for sight of a complete partitioning method, condone 1 multiplication error, final addition not necessary. A1 for 7560 or digits 756(0) A1 (dep on M1, but not previous A1) for correct placement of decimal point.

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
6	15 and 16 parts shaded Alternative 1 $\frac{3}{4} = 0.75$ or 75%, $\frac{4}{5} = 0.8$ or 80% Alternative 2 $\frac{3}{4} = \frac{15}{20}$, $\frac{4}{5} = \frac{16}{20}$	$\frac{4}{5}$ + reason	3	M1 for shading 15 parts for $\frac{3}{4}$ M1 for shading 16 parts for $\frac{4}{5}$ A1 (dep on M2) for selection of $\frac{4}{5}$ with correct shading Alternative 1 M1 for $\frac{3}{4} = 0.75$ or 75% M1 for $\frac{4}{5} = 0.8$ or 80% A1 (dep on M2) for selection of 0.8 or 80% or $\frac{4}{5}$ with correct decimals or percentages Alternative 2 M1 for $\frac{3}{4} = \frac{15}{20}$ oe M1 for $\frac{4}{5} = \frac{16}{20}$ oe A1 (dep on M2) for selection of $\frac{4}{5}$ or $\frac{16}{20}$ with equivalent fractions

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
7		$5 \times 5 \times 6$	150	4	M1 for attempt at 1 division (e.g. $40 \div 8$), may be implied by marks or number on one edge of diagram or by 5 or 6 seen M1 for attempt at 3 divisions ($40 \div 8$, $40 \div 8$, $60 \div 10$), may be implied by marks or numbers on diagram or by 5,5 and 6 seen. M1 (dep on 1st M1) for "5" × "5" × "6" A1 cao Alternatively M1 for $40 \times 40 \times 60$ **or** $8 \times 8 \times 10$ **or** 96000 or 640 seen M1 for $40 \times 40 \times 60$ **and** $8 \times 8 \times 10$ or 96000 **and** 640 seen M1 (dep on 1st M1) for "$(40 \times 40 \times 60)$" ÷ "$(8 \times 8 \times 10)$" A1 cao SC:B1 for dividing area of one carton face by area of corresponding box face if M0
8	(a)		$\frac{7}{20}$	1	B1 for $\frac{7}{20}$ oe
	(b)	7 + 2 (or 20 – 11) are not lime flavour	$\frac{9}{20}$	1	B1 for $\frac{9}{20}$ oe
	(c)		0	1	B1 for 0, zero or nought ($\frac{0}{20}$ gets B0)
9	(a)		$80x$	1	B1 for $80x$ (accept $80 \times x$, $x80$, $x \times 80$) seen
	(b)		$95y$	1	B1 for $95y$ (accept $95 \times y$, $y95$, $y \times 95$) seen
	(c)		$80x + 95y$	2	M1ft for adding "$80x$" and "$95y$" (algebraic expressions only) A1 for $80x + 95y$

Paper 5523_03

No	Working	Answer	Mark	Notes
10 (a)		40	1	B1 cao
(b)		45	1	B1 for 42 to 48 (accept 3/4 hour)
(c)	40×2 or $\frac{40}{30}\times60$ or $40\div\frac{1}{2}$	80	2	M1 for 40×2 or $\frac{40}{30}$ or $40\div\frac{1}{2}$ A1 cao NB $\frac{40}{45}\times 60$ gets M0 A0
11 (a)	$3\times3-4\times2$ or $9-8$	1	2	M1 for substitution of 3 and 2 into expression or 9 and 8 seen A1 cao
(b)	$-7-3=-10$ $2\times-10=-20$ $-20\div4$	-5	3	M1 for substitution of 2 and -7 into $p(q-3)$ or sight of -20 or $-14-6$ M1 (dep) for "-20" $\div 4$ A1 cao SC: B1 for -10 seen if M0
12 (a)	6 | 8 9 7 7 | 8 5 9 6 3 8 | 1 3 1 7 1 9 | 0 1	6 | 7 8 9 7 | 3 5 6 8 9 8 | 1 1 1 3 7 9 | 0 1	3	M1 for unordered diagram (condone one error) A1 cao B1 for key (eg 6 | 7 = 67)
(b)(i)		Explanation	2	B1 for '(order numbers and) select middle value' oe
(ii)		79		B1 cao

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
13	(a)		Reflection in y-axis	1	B1 for triangle with vertices at (-1, 1) (-3, 1) and (-1,4)
	(b)		Rotation by half turn about (0, 0)	2	B2 for triangle with vertices (-1, -1) (-3, -1) and (-1, -4) (B1 for half turn not about (0,0))
	(c)		Enlargement Scale factor 3 Centre (0, 0)	3	B1 for 'enlargement' B1 for 'scale factor 3' or 3 seen B1 for 'centre (0,0)' B0 for any combination of transformations
14	(a)		4560	1	B1 cao
	(b)		45.6	1	B1 cao
	(c)		2.4	1	B1 cao
15	(a)	$4a-2a+5b+b$	$2a+6b$	2	B2 cao (B1 for $2a$ or $6b$ seen)
	(b)		$x(x-6)$	2	B2 cao (B1 for $x(ax+b)$ where a, b are numbers not equal to zero or $x-6$ seen on its own, or part of an expression)
	(c)		$3x-2x^3$	2	B2 cao (B1 for $3x$ or $2x^3$)
	(d)		$4x(3y+x)$	2	B2 cao (B1 for $2(6xy+2x^2)$ or $4(3xy+x^2)$ or $x(12y+4x)$ or $2x(6y+2x)$ or $4x(\quad)$)

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
16	(a)	$1 - (0.2 + 0.3 + 0.1)$	0.4	2	M1 for $1 - (0.2 + 0.3 + 0.1)$ A1 for 0.4 oe, accept $\frac{0.4}{1}$
	(b)	0.2×200	40	2	M1 for 0.2×200 A1 cao NB $\frac{40}{200}$ is M1 A0, 40 out of 200 is M1 A1
17	(a) (i)	$180 - 2 \times 25$	130	3	M1 for $180 - 2 \times 25$ A1 cao
	(ii)		Reason		B1 for mentioning isosceles and equal (or base) angles or equal sides and equal (or base) angles
	(b)	$180 - 95$	85	1	B1 cao
18	(a) (i)		7^5	3	B1 cao
	(ii)		7^4		B2 cao (B1 for sight of 7^5 or 7^{2+3} or 7×7^3 or $7^1 \times 7^3$ or $7^2 \times 7^2$ or 7^{2+3-1})
	(b)		$\frac{1}{2}$	1	B1 for $\frac{1}{2}$ or 0.5 or 2^{-1}
19	(a)		3×10^7	1	B1 cao
	(b)		0.002	1	B1 cao

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
20		Box plot	2	3 aspects: 1st aspect – vertical line for median 2nd aspect – box using correct quartiles 3rd aspect – whiskers (could be single line) drawn with correct end points B2 for fully correct box plot (B1 for 1 aspect)
21 (a)	e.g. 2 | 126 3 | 63 3 | 21 7	$2\times3\times3\times7$	2	M1 for a systematic method of at least 2 correct divisions by a prime number oe factor trees; can be implied by digits 2, 3, 3, 7 on answer line. A1 for $2\times3^2\times7$ or $2\times3\times3\times7$
(b)	$2\times3\times7$	42	2	B2 cao (B1 for 6, 14, 21 or $2\times3\times7$)
22	$\frac{8}{3}\times\frac{5}{4}=\frac{8\times5}{3\times4}=\frac{40}{12}$	$3\frac{1}{3}$	3	B1 for $\frac{8}{3}$ oe improper fraction or $\frac{5}{4}$ oe improper fraction M1 (dep on B1) for multiplying numerator and denominator of "$\frac{8}{3}$" and "$\frac{5}{4}$" A1 for $3\frac{1}{3}$ oe mixed number or $\frac{10}{3}$ OR B1 for 1.25 and 2.67 or 2.66(…) M1 (dep on B1) for correct method of multiplication A1 for $3.\dot{3}$

Paper 5523_03				
No	**Working**	**Answer**	**Mark**	**Notes**
23			2	M1 for a relevant pair of intersecting arcs A1 for line drawn within guidelines, at least 3cm in length, accept broken line [SC: B1 for line drawn within guidelines if M0]
24 (a)		$-1, 0, 1, 2, 3$	2	B2 cao (-1 each error or omission)
(b)(i)		$x \geq \frac{7}{2}$	3	M1 for $2x \geq 7$, condone use of = sign or wrong equality A1 for $x \geq \frac{7}{2}$ oe as final answer SC:B1 for 3.5 or $\frac{7}{2}$ seen if M0 B1 ft from $x \geq$ "$\frac{7}{2}$"
(ii)		4		
25	$4x + 2y = 8$ $4x - 10y = 20$ $12y = -12$ $y = -1$ $4x + 2(-1) = 8$ $x = 2.5$	$x = 2.5$ $y = -1$	3	M1 for correct process to eliminate either x or y (condone one arithmetical error) M1 (dep) for substituting found value into either equation A1 for $x = 2.5$, $y = -1$ [SC: B1 for $x = 2.5$ or $y = -1$ if M0]

Paper 5523_03					
No		**Working**	**Answer**	**Mark**	**Notes**
26		Interior angle of hexagon = 180 – (360 ÷ 6) = 120 360 – (90 + 120)	150	4	Alternative 1 M1 for 360 ÷ 6 A1 for 60 M1 (dep on M1) for "60" + 90 A1 cao Alternative 2 M1 for 360 ÷ 6 A1 for 60 M1 (dep on M1) for 360 – (2 × "60" + 90) A1 cao Alternative 3 M1 for (6 – 2) × 180 ÷ 6 A1 for 120 M1 (dep on M1) for 360 – (90 + "120") A1 cao
27	(a)		(16), 50, 82, 96, 100	1	B1 cao
	(b)	Cumulative freq. diag. curve/ segments	Cum. freq graph	2	B1 for 4 or 5 points plotted correctly ± 1 full (2mm) square depending on sensible table (condone 1 addition error) B1 (dep) for points joined by curve or line segments provided no gradient is negative - ignore any part of graph outside range of their points. (SC:B1 if 4 or 5 points plotted not at end but consistent within each interval and joined)
	(c)	100 – 42	58	2	M1 (ft dep on graph being cf) for reading from graph at 18 or 19, can be implied by answer in range 40 to 46 A1 for answer in range 56 to 60 or ft for 100 – '42' ±1 full (2mm) square

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
1	(a)	900 × 1.70 =	1530	2	M1 900 × 1.7(0) or digits 153(0) seen A1 cao
	(b)	160 ÷ 1.70 =	£94.12 or £94.11	2	M1 160 ÷ 1.7(0) or digits 941(....) seen A1 cao
2	(a)	4.7 ÷ 5.9 = 0.796610169	0.7966..	2	B2 for 0.7966 or better (B1 for 0.8, 0.80, 0.79, 0.796, 0.797 or digits 59 seen)
	(b)	0.82, 0.8, 0.85, 0.66, 0.875 0.66, 0.8, 0.82, 0.85, 0.875 2/3, 4/5, 0.82, 85%, 7/8	2/3, 4/5, 0.82, 85%, 7/8	2	B2 correct order (oe decimals in order) (B1 correct order reversed, or one error in ordered listing) with or without decimal equivalents. NB Accept 0.67 or 0.66
3		**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	**55** 61 74 **190** 33 **17** **10** 60 **88** **78** 84 250	3	B3 all 6 correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
4	(a)	–3 (–1) (1) 3 5 7	**–3**, 3, **5**, **7**	2	B2 all correct (B1 2,3 correct)
	(b)	(-2,-3),(-1,-1),(0,1),(1,3),(2,5),(3,7)	line	2	B2 cao for line from $x = -2$ to $x = 3$ (B1 plotting at least 5 points correctly or single line passing through (0, 1) or single line of gradient 2) The six possible points are: (-2,-3), (-1,-1), (0,1), (1,3), (2,5), (3,7)
	(c)(i)		–2	2	B1 $y = -2$ or ft from line segment
	(ii)		2.5		B1 $x = 2.5$ or ft from line segment

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
5	(a) (i)	180 – 54 (=126) "126" ÷ 2	63	3	M1 for (180–54) ÷ 2 A1 cao
	(ii)		Reason		B1 (indep) angles in triangle add to 180 OR equal angles in isosceles triangle OR equal angles and 2 sides the same (B0 if any incorrect reasoning given eg parallel, equilateral triangle)
	(b)	180 – "x"	117	1	B1 117 or ft 180 – "x" if $x < 90$
6		15/120×360 = 45 Plain 40/120×360 = 120 Cheese & Onion 55/120×360 = 165 Salt & Vinegar 10/120 × 360 = 30 Beef	45°, 120°, 165°, 30°	4	M1 evidence of method for at least one angle (could be implied by one correct angle of four on pie chart or in the table) A2 All four angles drawn ±2° tolerance, any order (A1 at least 2 angles correctly drawn ±2°, or all 4 angles in the table) B1 (dep on at least 1 angle drawn correctly, and exactly 4 sectors) for labels (flavour or frequency; initials will do) NB: Ignore the table if the pie chart provides the marks.
7	(a)	260 – 50 = 210 210 ÷ 35 =	6	3	M1 for 260–50 or 210 seen. M1 for "260-50" ÷ 35 or 210÷35 A1 cao
	(b)		P=35h + 50	3	B3 for P=35h+50 or P=35×h+50 oe (B2 for correct RHS or P=h + 50 ×35 or P=35h+k where k is numerical oe) (B1 for P = some other linear expression in h, OR h + 50 ×35 OR 35h seen) NB: P=h scores no marks; ignore £ signs. SC B2 for $h = \frac{P-50}{35}$
8	(a)		Elevation	2	B2 for 4 vertical squares. Accept 4 by 1 rectangle. (B1 for 4 vertical squares with one square added or one parallelogram added at the top, or 3 vertical squares, or 4 horizontal squares)

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
(b)		Plan	2	B2 for 2 adjacent squares, vertical or horizontal. Accept 2 by 1 rectangle. (B1 for 3 adjacent horizontal or vertical squares or a rectangle with sides in the ratio 2:1)
9 (a)		Points	1	B1 all three points ±1 full square.
(b)		Negative	1	B1 Negative. Ignore other descriptors unless contradictory.
(c)		lobf	1	B1 A single straight line drawn to cross between (5,30), (5,40) and (40,0), (40,15); accept freehand if considered to be straight.
(d)(i) (ii)		18–25 30–40	2	B1 18g–25g inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient. B1 30–40 min inclusive OR if not in this range ft ±1 square dep on single straight line with negative gradient.
10		300, 3, 75, 150	3	B3 for 4 correct (B2 for 2 or 3 correct) (B1 for 1 correct)
11 (a)	$6x - 7 + 7 = 38 + 7$ $6x = 45$	7.5	2	M1 $6x = 45$ or +7 both sides A1 7.5 oe; accept 45/6
(b)	$5y - 2 = 10$ or $20y - 8 = 40$ $5y = 12$ $20y = 48$	$2\frac{2}{5}$	3	M1 $20y–8(=40)$ or $\frac{4(5y-2)}{4} = \frac{40}{4}$ or $5y–2=10$ M1 (indep) for correct rearrangement into the form $ay=b+c$ or better (eg $20y=40+8$ or $5y=10+2$, using own terms) A1 for $2\frac{2}{5}$, 2.4 oe
12	5 miles = 8 km 70mph÷5×8 = 122 km/h OR 120km/h ÷ 8 × 8 = 75 mph Faster than 70 mph	70mph (Great Britain) (112 km)	3	M1 5 miles = 8 km; OR 70 mph is about 100 km/h OR 1km=0.6(25) miles OR 1mile=1.6km oe M1 70 ÷ 5 × 8 (=112) or 120 ÷ 8 × 5 (=75) oe A1 (dep on at least M1) GB or 70 mph Refer to both answer line and working. NB GB or 70 mph without working scores 0 marks.

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
13 (a)	$\pi \times 0.65$	2.04–2.05	2	M1 for $\pi \times 0.65$ or 3.14×0.65 or 3.142×0.65 oe A1 2.04–2.05 SC Award B1 for 2.0 seen (not 2)
(b)	1000 ÷ "(a)"	487–491	2	M1 for 1000 (or 100) ÷ "(a)" A1 for 487–491
14	$45.00 + 45.00 \times \frac{15}{100} =$ $45.00 + 6.75 =$	51.75	3	M2 for $45.00 + 45.00 \times \frac{15}{100}$ oe or 45.00×1.15 oe OR $45.00 + 6.75$ OR complete method or 5175 seen. (M1 for $45.00 \times \frac{15}{100}$ oe OR 6.75 seen OR 675 seen OR correct method for calculating 15% of 45) A1 cao SC Award B2 for an answer of 38.25
15		273	2	M1 for 728 ÷ 8 or 728 ÷ "3+5" or 91 A1 cao SC B1 for 455, or for 273:455
16	3 24 3.7 46.9(53) 4 60 3.8 51.0(72) 3.1 26.6(91) 3.9 55.4(19) 3.2 29.5(68) 3.21 29.8(66…) 3.3 32.6(37) 3.22 30.1(66…) 3.4 35.9(04) 3.23 30.4(68..) 3.5 39.3(75) 3.24 30.7(72...) 3.6 43.0(56) 3.25 31.0(78...) or 31	3.2	4	B2 for trial between 3.2 and 3.3 inclusive (B1 for trial between 3 and 4 inclusive) B1 for different trial between 3.21 and 3.25 inclusive B1 (dep on at least one previous B1) cao for 3.2 as final answer. NB: embedded answers: –B1; award Bs for evaluations rounded or truncated to at least 1 dp or for 31

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
17	$45^2 + 34^2 =$ $2025 + 1156 = 3181$ $\sqrt{3181} = 56.4$	56	4	M1 for $45^2 + 34^2$ M1 (dep) for $\sqrt{(2025 + 1156)}$ A1 for 56.4 … B1 for rounding their diagonal to the nearest integer (dep on evidence from decimal) NB 56 as the final answer gets full marks. NB Scale drawings result in 0 marks.
18	$2000 \times (1.055)^3$ Interest = 2348.48 – 2000 =	348.48	3	M1 for 5.5/100 × 2000 (oe) or 330 or 2330 or 110 or 2110 M1 (dep) for 5.5/100 × (2000 + "110" + "116.05") or 122.4… seen A1 cao (accept only 348.48 or 348.49) **OR** M2 for $2000 \times (1.055)^3$ or 2348.48(…) or 2348.49 seen (M1 for $2000 \times (1.055)^n$, $n \neq 3$) A1 for 348.48 or 348.49 [SC: B2 for 2348.48 - 2348.49]
19		Line	2	B2 line fully within tramlines, crossing AB and CD (B1 a straight line which crosses AB within the tramline, and also crosses CD) NB: Accept dotted or dashed lines, but not curves; accept freehand if considered to be straight. SC B1 for a perpendicular bisector of AB that is at least half way from AB to CD within the tramlines
20 (a)		$90<t\leq100$	1	B1 for $90<t\leq100$; accept 90-100.
(b)	65×12=780 75×22=1650 85×23=1955 95×24=2280 105×19=1995 8660/100=	86.6	4	M1 for use of fx with x consistent within intervals (including end points). Allow one slip. M1 (dep) for use of midpoints M1 (dep on 1st M1) for use of $\sum fx/100$ or $\sum fx/\sum f$ A1 86.6

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
21	(a)	$8 \times \frac{10}{4}$	20	2	M1 $\frac{10}{4}$ or $\frac{4}{10}$ or 0.4 or 2.5 oe seen A1 cao NB ratios get M0 unless of the form 1:*n* OR M1 $\frac{8}{4}, \frac{4}{8}$ oe seen A1 cao
	(b)	$15 \times \frac{4}{10}$	6	2	M1 $15 \times \frac{4}{10}$ oe A1 cao
22	(a)	$x^2 - 4x + 3x - 12 = x^2 - x - 12$	$x^2 - x - 12$	2	M1 for exactly 4 terms correct ignoring signs (eg x^2, 4*x*, 3*x*, 12) or 3 correct terms out of 4 terms with correct signs (eg 3 out of 4 of x^2, –4*x*, +3*x*,–12) A1 cao
	(b)	$(x+2)(x+5)$	$(x+2)(x+5)$	2	B2 cao (B1 for exactly one of $(x+2)$, $(x+5)$)
	(c)	$6=15+4q-20$ $6-15=4(q-5)$ $p-3t=4q-4t$ $6-3\times5=4(q-5)$	2 ¾	3	M1 for correct substitution of *p* and *t*. M1 for correct expansion of $4(q-t)$ oe (eg $4q-20$, $4q-4t$) A1 11/4 or 2 ¾ or 2.75 OR M1 for correct substitution of *p* and *t*. M1 for $\frac{p-3t}{4} = q - t$ oe A1 11/4 or 2 ¾ or 2.75

Paper 5523_04					
No		**Working**	**Answer**	**Mark**	**Notes**
23	(a)	$9.9\times10^8 - 6.0\times10^7$	9.3×10^8	2	M1 for $99\times10^7 - 6\times10^7$ or $9.9\times10^8 - 0.6\times10^8$ or conversion of either to an ordinary number, or 930000000 or 93×10^7 or 9.3×10^n where n is any positive integer A1 cao
	(b)	$\frac{6.0-4.5}{4.5}\times100 = \frac{1.5}{4.5}\times100 =$ or $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}\times100$	33.3%	3	M2 $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}\times100$ oe (M1 for $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}$ or $\frac{6.0\times10^7 - 4.5\times10^7}{6.0\times10^7}\times100$ oe A1 cao OR M2 $\frac{6.0\times10^7}{4.5\times10^7}\times100$ -100 = (M1 $\frac{6.0\times10^7}{4.5\times10^7}\times100$ or 133.33(%)) A1 cao NB Accept any of the above expressions without any reference to 10^7.
24		LCM (40,24) = 120 Bread buns 120÷40 Burgers 120÷24 OR Bread buns: 40 is $2\times2\times2$ (×5) Burgers: 24 is $2\times2\times2$ (×3)	Bread buns 3 Burgers 5	3	M1 attempt to find LCM by eg lists of multiples, or summing of 40 and 24. A1 identify 120 (as LCM) A1 cao (both) OR M1 expansion of either number into its prime factors in a factor tree or 8×5 or 8×3 A1 both expansions correct A1 cao (both) SC B2 if answers given the wrong way around

Paper 5523_04				
No	**Working**	**Answer**	**Mark**	**Notes**
25	8 × 41 = 328 2 × 29 = 58 328 – 58 = 270 270 ÷ 6 = 45	45	3	M1 for either 8 × 41 (=328) or 2 × 29 (=58) M1 (dep) "328" – "58" (=270) A1 cao NB 328 and/or 58 on the answer line gets M1 (implied); 270 on the answer line gets M2 (implied)

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
1 (a)	$1 - (0.2 + 0.3 + 0.1)$	0.4	2	M1 for $1 - (0.2 + 0.3 + 0.1)$ A1 for 0.4 oe, accept $\frac{0.4}{1}$
(b)	0.2×200	40	2	M1 for 0.2×200 A1 cao NB $\frac{40}{200}$ is M1 A0, 40 out of 200 is M1 A1
2	$650 - 430 = 220$ 1 choc ice costs 110p $650 - 5 \times 110 = 100$p	50	3	M1 for $650 - 430$ or 220 or 110 oe seen M1 for $650 - 5 \times \frac{'220'}{2}$ or $430 - 3 \times \frac{'220'}{2}$ oe A1 for 50p or £0.50 or £0.5 *Alternative scheme* $2x + 5y = 650$ $2x + 3y = 430$ oe M1 for subtracting two simultaneous equations to eliminate x (lollies)(2 or 3 terms correct) M1 for $650 - 5 \times 'y'$ or $430 - 3 \times 'y'$ oe A1 for 50p or £0.50 or £0.5 *Alternative scheme* M1 for $3\times (2x + 5y = 650)$ evaluated and $5 \times (2x + 3y = 430)$ evaluated oe (5 or 6 terms correct) M1 for subtraction of equations to eliminate y (choc ices)(2 or 3 terms correct ft) A1 for 50p or £0.50 or £0.5

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
3		question + response boxes oe	2	1st aspect: One question (eg 'how long does it take you to travel to school?' or 'What time did you leave home to get to school?'); ignore other questions. 2nd aspect: Response list (at least two), not overlapping. 3rd aspect: Some mention of units (eg minutes) in either question or responses B2 for all three aspects, or B1 for just one aspects.
4	2 [(3×1) + (4×1)] + (3×6) + (1×6) + (2×6) + (4×6) + (1×6) + (5×6)	110 cm^2	4	M1 for attempt to find the area of one face M1 for at least 6 faces with intention to add A1 cao B1 (indep) for cm^2 (with or without numerical answer)
5	$\left(\frac{-6+4}{2}\right), \left(\frac{5+3}{2}\right)$	(–1, 4)	2	B2 cao [B1 for (–1,*a*) or (b,4) or (4,-1)]
6		Box plot	2	3 aspects: 1st aspect – vertical line for median 2nd aspect – box using correct quartiles 3rd aspect – whiskers (could be single line) drawn with correct end points B2 for fully correct box plot (B1 for 1 aspect)
7			2	M1 for a relevant pair of intersecting arcs A1 for line drawn within guidelines, at least 3cm in length, accept broken line [SC: B1 for line drawn within guidelines if M0]

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
8	(a)	e.g. 2 | 126 3 | 63 3 | 21 7	$2\times3\times3\times7$	2	M1 for a systematic method of at least 2 correct divisions by a prime number oe factor trees; can be implied by digits 2, 3, 3, 7 on answer line. A1 for $2\times3^2\times7$ or $2\times3\times3\times7$
	(b)	$2\times3\times7$	42	2	B2 cao (B1 for 6, 14, 21 or $2\times3\times7$)
9	(a)		$-1,0,1,2,3$	2	B2 cao (-1 each error or omission)
	(b)(i)		$x\geq\frac{7}{2}$	3	M1 for $2x\geq7$, condone use of = sign or wrong equality A1 for $x\geq\frac{7}{2}$ oe as final answer
	(ii)		4		SC:B1 for 3.5 or $\frac{7}{2}$ seen if M0 B1 ft from $x\geq$ "$\frac{7}{2}$" or $x>$"$\frac{7}{2}$"
10	(a)(i)		7^5	3	B1 cao
	(ii)		7^4		B2 cao (B1 for sight of 7^5 or 7^{2+3} or 7×7^3 or $7^1\times7^3$ or $7^2\times7^2$ or 7^{2+3-1})
	(b)		$\frac{1}{2}$	1	B1 for $\frac{1}{2}$ or 0.5 or 2^{-1}
11	(a)	$5n=m+21$	$n=\frac{m+21}{5}$	2	M1 for $5n=m+21$ or for attempt to divide three terms by 5 A1 $n=\frac{m+21}{5}$ oe

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
(b)	$4p-8q=3p+2$ $p-8q=2$ $p=8q+2$	$p=8q+2$	3	M1 for $4p-aq$ or $\frac{3}{4}p+b$ where a is an integer and b is a number M1 (dep) for taking one term correctly to LHS or RHS of expression A1 $p=8q+2$ oe
12		$y=\frac{1}{2}x+3$	2	B2 for $y = ½x + 3$ oe (B1 for $y = ½x + c$, $c \neq 7$ or $y = mx + 3$ oe or $½x + 3$ or $M = ½x + 3$)
13	$\frac{8}{3}\times\frac{5}{4}=\frac{8\times5}{3\times4}=\frac{40}{12}$	$3\frac{1}{3}$	3	B1 for $\frac{8}{3}$ oe or $\frac{5}{4}$ oe M1 (dep on B1) for multiplying numerator and denominator of "$\frac{8}{3}$" and "$\frac{5}{4}$" A1 for $3\frac{1}{3}$ oe mixed number or $\frac{10}{3}$ *OR* B1 for 1.25 and 2.67 or 2.66(…) M1(dep on B1) for correct method of multiplication A1 for $3.\dot{3}$
14	$4x+2y=8$ $4x-10y=20$ $12y=-12$ $y=-1$ $4x+2(-1)=8$ $x=2.5$	$x=2.5$ $y=-1$	3	M1 for correct process to eliminate either x or y (condone one arithmetical error) M1 (dep) for substituting found value into either equation A1 for $x = 2.5$, $y = -1$ [SC: B1 for $x = 2.5$ or $y = -1$ if M0]

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
15		Interior angle of hexagon = 180 – (360 ÷ 6) = 120 360 – (90 + 120)	150	4	Alternative 1 M1 for 360 ÷ 6 A1 for 60 M1 (dep on M1) for "60" + 90 A1 cao Alternative 2 M1 for 360 ÷ 6 A1 for 60 M1(dep on M1) for 360 – (2 × "60" + 90) A1 cao Alternative 3 M1 for (6 – 2) × 180 ÷ 6 A1 for 120 M1(dep on M1) for 360 – (90 + "120") A1 cao
16	(a)		(16), 50, 82, 96, 100	1	B1 cao
	(b)	Cumulative freq. diag. curve/ segments	Cum. freq graph	2	B1 ft for 4 or 5 points plotted correctly ± 1 full (2mm) square depending on sensible table (condone 1 addition error) B1 (dep) for points joined by curve or line segments provided no gradient is negative - ignore any part of graph outside range of their points. (SC:B1 if 4 or 5 points plotted not at end but consistent within each interval and joined)

Paper 5525_05				
No	**Working**	**Answer**	**Mark**	**Notes**
(c)	$100 - 42$	58	2	M1 (ft dep on graph being cf) for reading from graph at 18 or 19, can be implied by answer in range 40 to 46 A1 for answer in range 56 to 60 or ft for 100 –'42' ±1 full (2mm) square
17 (a)		0.6 and 0.7,0.3,0.7	2	B1 for 0.6 on LH branch B1 for 0.7, 0.3 and 0.7 on RH branches
(b)	0.4×0.3	0.12	2	M1 for 0.4×0.3 A1 0.12 oe
(c)	$0.4\times0.7+0.6\times0.3$	0.46	3	M1 for '0.4×0.7' or '0.6×0.3' M1 for addition of two products from correct branches A1 0.46 oe Alternative M2 for an attempt to evaluate 1-(0.3×0.4 +'0.6×0.7') A1 cao
18	$x = 0.4545...$ so $100x = 45.4545...$ $99x = 45$ $x = \frac{45}{99} = \frac{15}{33}$	proof	3	M1 for $100x = 45.45\ldots$ or $10000x = 4545.45\ldots$ M1 (dep) for subtraction of both sides A1 for $\frac{15}{33}$ from correct proof
19	$\left(\sqrt{3}\right)^2 + \sqrt{3}\sqrt{2} - \sqrt{3}\sqrt{2} - \left(\sqrt{2}\right)^2$ $= 3 - 2$	1	2	B2 cao (B1 for $\sqrt{3}\sqrt{3} + \sqrt{3}\sqrt{2} - \sqrt{3}\sqrt{2} - \sqrt{2}\sqrt{2}$ oe, $\sqrt{3}\sqrt{3} - \sqrt{2}\sqrt{2}$ oe, or for 2,3, $\sqrt{4}$ $\sqrt{6}$ $\sqrt{9}$ seen)
20 (a)	$\frac{96}{24}$ or 4 $\sqrt{4}$ or 2	8	3	M1 for $\frac{96}{24}$ or $\frac{24}{96}$ or 4 or $\frac{1}{4}$ oe M1 for $\sqrt{\frac{96}{24}}$ or $\sqrt{\frac{24}{96}}$ or $\sqrt{'4'}$ or $\frac{1}{\sqrt{'4'}}$ or 2 or $\frac{1}{2}$ oe A1 cao

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
	(b)	12×2^3	96	2	M1 for '2'^{3} or 8 A1 cao
21	(a)	$x\times3-x\times2x^2$	$3x-2x^3$	2	B2 cao (B1 for a two term expression with either $3x$ or $2x^3$)
	(b)		$4x(3y+x)$	2	M1 for taking out a factor of x, $2x$, 2, 4 or $4x$ A1 cao
	(c)	$\frac{5a}{b^2}$	$\frac{5a}{b^2}$	2	B2 for $\frac{5a}{b^2}$ or $5ab^{-2}$ (accept $\frac{5a}{1b^2}$) (B1 for either dealing with the numbers or dealing with the powers of a)
	(d)	$\frac{\cancel{x-3}}{\cancel{(x-3)}(x+3)}$	$\frac{1}{x+3}$	2	M1 for $(x-3)(x+3)$ A1 cao
22	(a)		**a** + **b**	1	B1 **a** + **b** oe
	(b)	ED = **a** DX = –2**b** + 2 AC = 2**a** (So, DX = 2ED)	2**a**	3	M1 for (DX=) DA + AX A1 for (DX=) –2***b*** + 2 ("***a*** + ***b***") A1 2***a*** from fully correct proof
23		$\pi(2x)^2h=\frac{4}{3}\pi(3x)^3$ $h=\frac{\frac{4}{3}\pi(3x)^3}{\pi(2x)^2}$	$9x$	3	M1 for $\pi(2x)^2h=\frac{4}{3}\pi(3x)^3$ (condone absence of brackets) M1 (dep) for valid algebra that gets to $h=ax$ (condone one error in powers of numerical constants) A1 cao

Paper 5525_05					
No		**Working**	**Answer**	**Mark**	**Notes**
24	(i)	$n^2+(n+1)^2=2(n^2+n)+1$	$2n^2+2n+1$	4	M1 for at least 3 terms correct from $n^2+n+n+1$ A1 for $2n^2+2n+1$ oe
	(ii)		$2(n^2+n)$ is always even so $2(n^2+n)+1$ is always odd		M1 for recognizing $2n^2$ is always even A1ft complete proof for their quadratic *Alternative method* M1 for recognizing that if n^2 is odd then $(n+1)^2$ is even or vice versa A1 for complete proof
25			42 318	2	B1 for answer in range 36 – 48 B1 for answer in range 312 – 324
26	(a)	$(x-3)^2-3^2+15$	$p=3, q=6$	2	B2 for $p=3$ and $q=6$ (B1 for $p=3$ OR $q=6$) SC: award B2 for $(x-3)^2+6$ if p and q are not identified
	(b)		Sketch	2	B1 for U shaped curve B1 ft for TP in first quadrant (ft if TP not in first quadrant)
27	(a)	Graph translated 2 units upwards through points (–4, 2), (–2, 4), (0,2) and (3,5)	Sketch	2	M1 for a vertical translation A1 curve through points (–4, 2), (–2, 4), (0,2) and (3,5) ± ½ square
	(b)	Graph reflected in x-axis through points (–4,0), (–2, –2), (0, 0) and (3, -3)	Sketch	2	M1 for reflection in x-axis or y-axis A1 curve through points (–4,0), (–2, –2), (0, 0) and (3, -3) ± ½ square

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
1		273	2	M1 for 728 ÷ 8 or 728 ÷ "3+5" or 91 A1 cao SC B1 for 455 or 273:455
2		$3n - 1$	2	B2 for $3n - 1$ oe (B1 for $3n + k$ where $k \neq -1$ but k could be 0)
3	3 24 3.7 46.9(53) 4 60 3.8 51.0(72) 3.1 26.6(91) 3.9 55.4(19) 3.2 29.5(68) 3.21 29.8(66…) 3.3 32.6(37) 3.22 30.1(66…) 3.4 35.9(04) 3.23 30.4(68..) 3.5 39.3(75) 3.24 30.7(72...) 3.6 43.0(56) 3.25 31.0(78...) or 31	3.2	4	B2 for trial between 3.2 and 3.3 inclusive (B1 for trial between 3 and 4 inclusive) B1 for different trial between 3.21 and 3.25 inclusive B1 (dep on at least one previous B1) cao for 3.2 NB: embedded answers: –B1; award Bs for evaluations rounded or truncated to at least 1 dp or for 31
4	2 is the only even prime number and the product of 2 odd numbers is odd	Yes	2	B2 for 'yes' and '2 is the only even prime number and the product of two odd numbers is odd' oe (B1 for 'yes' and either '2 is the only even prime number' oe **or** 'the product of two odd numbers is odd' oe)
5	$45^2 + 34^2 =$ $2025 + 1156 = 3181$ $\sqrt{3181} = 56.4$	56	4	M1 for $45^2 + 34^2$ M1 for $\sqrt{(2025 + 1156)}$ A1 for 56.4 … B1 for rounding their diagonal to the nearest integer (dep on evidence from decimal) NB 56 with no incorrect working as the final answer gets full marks. NB Scale drawings result in 0 marks.

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
6	$2000 \times (1.055)^3$ Interest = 2348.48 – 2000 =	£348.48	3	M1 for 5.5/100 × 2000 (oe) or 330 or 16.5/100×2000 or 2330 or 110 or 2110 M1 (dep) for 5.5/100 × (2000 + "110" + "116.05") or 122.4… seen A1 cao (accept only 348.48 or 348.49) **OR** M2 for $2000 \times (1.055)^3$ or 2348.48(…) or 2348.49 seen (M1 for $2000 \times (1.055)^n$, $n \neq 3$) A1 for 348.48 or 348.49 [SC: B2 for 2348.48 - 2348.49]
7 (a)		Line	2	B2 line fully within tramlines, crossing AB and CD (B1 a straight line which crosses AB within the tramline, and also crosses CD) NB: Accept dotted or dashed lines, but not curves; accept freehand if considered to be straight. SC B1 for the perpendicular bisector of *AB* reaching halfway or more from *AB*
(b)		Region		B2 correct arc ±2mm and shaded within. Allow dotty or continous arc. (B1 inaccurate arc and shaded or accurate arc unshaded)
8	65×12=780 75×22=1650 85×23=1955 95×24=2280 105×19=1995 8660/100=	86.6	4	M1 for use of fx with x consistent within intervals (including end points). Allow one slip even if outside interval M1 (dep) for use of midpoints M1 (dep on 1st M1) for use of $\sum fx/100$ or $\sum fx/\sum f$ A1 86.6

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
9		Distance ÷ time: 1400 ÷ 2 h 20 min 20 mins is $\frac{1}{3}$ hour 1400 × 3 ÷ 7 =	600 kph	3	B1 20 mins as $\frac{1}{3}$ hour or as 0.33……hour M1 for distance ÷ time eg 1400 ÷ "2h 20 min" A1 cao OR B1 2 hour 20min = 140 (min) M1 Speed = $\frac{1400}{140}$ = (10 km per minute) A1 cao
10	(a)	$(x+18)+2x+(2x+7)=180$	Equation	2	B2 for $(x+18)+2x+(2x+7)=180$ oe (B1 for $(x+18)+2x+(2x+7)$)
	(b)	$5x+25 = 180$ $5x = 155$	31	2	M1 for simplifying to at least $5x+25=180$ or 360 (may be earned in (a)) A1 for $x=31$
11	(a)	(–14) –4 (0) 4 14	–4, 4, 14	2	B2 for all 3 values correct (B1 for 1 or 2 values correct)
	(b)		curve	2	B1 for all 5 points plotted correctly ± ½ square (ft from table if at least B1 awarded in (a)) B1 (indep) ft for any smooth curve through their points

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
12	(a)	$8 \times \frac{10}{4}$	20	2	M1 $\frac{10}{4}$ or $\frac{4}{10}$ or 0.4 or 2.5 oe seen A1 cao NB ratios get M0 unless of the form 1:*n* OR M1 $\frac{8}{4}, \frac{4}{8}$ oe seen A1 cao
	(b)	$15 \times \frac{4}{10}$	6	2	M1 $15 \times \frac{4}{10}$ oe A1 cao
13	(a)	$9.9\times10^8 - 6.0\times10^7$	$\pm 9.3\times10^8$	2	M1 for $99\times10^7 - 6.0\times10^7$ or $9.9\times10^8 - 0.60\times10^8$ or correct conversion of either to an ordinary number or 930000000 or 9.3×10^n where *n* is any positive integer A1 cao
	(b)	$\frac{6.0-4.5}{4.5} \times 100 = \frac{1.5}{4.5} \times 100 =$ or $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7} \times 100$	33.3%	3	M2 $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7} \times 100$ oe (M1 for $\frac{6.0\times10^7 - 4.5\times10^7}{4.5\times10^7}$ or $\frac{6.0\times10^7 - 4.5\times10^7}{6.0\times10^7} \times 100$ oe A1 33.3 – 33.4 OR M2 $\frac{6.0\times10^7}{4.5\times10^7} \times 100$ -100 or 33.3% (M1 $\frac{6.0\times10^7}{4.5\times10^7} \times 100$ or 133.3%) A1 33.3-33.4 NB Accept any of the above expressions without any reference to 10^7.

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
14	LCM (40,24) = 120 Bread buns 120÷40 Burgers 120÷24 OR Bread buns: 40 is $2 \times 2 \times 2$ (×5) Burgers: 24 is $2 \times 2 \times 2$ (×3)	Bread Buns 3 Burgers 5	3	M1 attempt to find LCM by eg lists of multiples, or summing of 40s and summing of 24s, with at least 3 numbers in each list A1 identify 120 as LCM A1 cao (both) **OR** M1 expansion of either number into its prime factors in a factor tree or 8×5 or 8×3 A1 both expansions correct A1 cao (both) SC B2 if answers given the wrong way around
15	$\sin 32 = \frac{AB}{12}$ $AB = 12 \times \sin 32$ $AB = 6.35903\ldots$	6.36	3	M1 $\sin 32 = \frac{AB}{12}$ (accept Sin $\frac{AB}{12}$) M1 $12 \times \sin 32$ or $12 \times 0.5299..$ A1 accept 6.359 – 6.360 SC Gradians 5.78(1…) Radians 6.62 Get M1M1A0 **OR** Use of Sine Rule $\frac{\sin 32}{AB} = \frac{\sin 90}{12}$ or $\frac{AB}{\sin 32} = \frac{12}{\sin 90}$ M1 $AB = \frac{12 \times \sin 32}{\sin 90}$ M1 AB = 6.359 – 6.36 A1 SC Gradians 5.85(…) Radians 7. 40(…) M1M1A0

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
16		$8 \times 41 = 328$ $2 \times 29 = 58$ $328 - 58 = 270$ $270 \div 6 = 45$	45	3	M1 for either 8×41 (=328) or 2×29 (=58) M1 (dep) "328" – "58" (=270) A1 cao NB 328 and /or 58 on the answer line gets M1 (implied); 270 on the answer line gets M2 (implied)
17	(a)	$x^2 - 4x + 3x - 12 = x^2 - x - 12$	$x^2 - x - 12$	2	M1 for exactly 4 terms correct ignoring signs (x^2, $4x$, $3x$, 12) or 3 out of 4 terms with correct signs (x^2, $-4x$, $+3x$, -12) A1 cao
	(b)	$6x^2 - 8x + 15x - 20 = 6x^2 + 7x - 20$	$6x^2 + 7x - 20$	2	M1 for exactly 4 terms correct ignoring signs ($6x^2$, $8x$, $15x$, 20) or 3 out of 4 terms with correct signs ($6x^2$, $-8x$, $+15x$, -20) A1 cao
	(c)	$(x+2)(x+5)$	$(x+2)(x+5)$	2	B2 cao (B1 for exactly one of $(x + 2)$, $(x + 5)$)
	(d)		$12p^8q^3$	2	B2 cao (B1 for any 2 out of 3 terms correct in a product or 3 terms correct in a sum or part product)
	(e)	$6 = 15 + 4(q - 5)$ $6 = 15 + 4q - 20$ $11 = 4q$	$2\frac{3}{4}$	3	M1 for correct substitution of p and t. M1 for correct expansion of $4(q\text{-}t)$ oe (eg $4q$-20, $4q$-$4t$) A1 11/4 or 2 ¾ or 2.75 OR M1 for correct substitution of p and t. M1 for $\frac{p - 3t}{4} = q - t$ oe A1 11/4 or 2 ¾ or 2.75
18	(a)	$T=kx$; $150=6k$; $k=25$	$T=25x$	3	M1 for $T=kx$, k algebraic M1 subs $T=150$ and $x=6$ into $T = kx^n$ $(n \neq 0)$ A1 for $T=25x$ oe SC B1 $T \propto 25x$ oe

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
	(b)	$T = 25 \times 15 =$	375	1	B1 ft on k , $k \neq 1$
	(c)	$600 = 25x$; $x = 600 \div 25 =$	24	1	B1 ft on k , $k \neq 1$
19		$\frac{120}{570} \times 50 = 10.5263$ $\frac{250}{570} \times 50 = 21.9298$ $\frac{200}{570} \times 50 = 17.5438$	10 22 18	3	M1 method shown eg $\frac{120}{120+250+200} \times 50$ or one of 10.5(263) 21.9(298), 17.5(438) **OR** 570 ÷ 50 = 11.4 and one of 120 ÷ 11.4 =10.526…) , 270 ÷ 11.4 (=21.9298…) , 200 ÷ 11.4 (=17.5438…) A1 for 10.5(263), 21.9(298), 17.5(438) or 11, 22, 18 B1 correction to add to 50: 10, 22, 18 or 11, 22, 17
20		$x(x+1) + 4(2x–3) = (2x–3)(x+1)$ $x^2 + x + 8x – 12 = 2x^2 + 2x – 3x – 3$ $x^2 – 10x + 9 = 0$ $(x – 9)(x – 1) = 0$	$x = 1, 9$	5	M1 for multiplying through by common denominator $(2x–3)(x+1)$ M1 (dep)for either $x^2 + x + 8x – 12$ or $2x^2 + 2x – 3x – 3$ oe A1 for correct quadratic (= 0) M1 for a correct method to solve 3 term quadratic A1 cao for both solutions
21		$AB^2 = 8^2 + 9^2 – 2\times8\times9\times\cos40$ $AB^2 = 64 + 81 – 144\times\cos40$ $AB^2 = 145 – 144 \times 0.766$ $AB^2 = 145 – 110.31… = 34.6896$ $AB = \sqrt{34.6796} = 5.8897877$	5.89	3	M1 Subs in Cos Rule: $8^2 + 9^2 – 2\times8\times9\times\cos40$ M1 correct order of evaluation of $8^2 + 9^2 – 2\times8\times9\times\cos40$ A1 cao 5.88 – 5.89 SC: Award B2 for one of $AB^2 = 241.03…$ or $AB = 15.525…$ (radians) $AB^2 = 28.50…$ or $AB = 5.33…$ (gradians)
22	(a)		24 30	2	B1 24 cao B1 30 cao
	(b)	Column $60 \leq w < 75$ at 10 Column $75 \leq w < 95$ at 4	10 4	2	B1 10 and correct width (tol $\pm\frac{1}{2}$ small square) B1 4 and correct width (tol $\pm\frac{1}{2}$ small square)

Paper 5525_06				
No	**Working**	**Answer**	**Mark**	**Notes**
23 (a)		100.5	1	B1 cao
(b)		10.515	1	B1 cao
(c)	$\frac{100.5}{10.515}$	9.5577746	2	M1 for greatest distance divided by least time Where 100 < greatest distance ≤ 100.5, 10.51 ≤ least time < 10.52 A1 for 9.555 – 9.56
(d)	$\frac{99.5}{10.525}$	9.45368..	2	M1 for least distance divided by greatest time Where 99.5 ≤ least distance < 100, 10.52 < greatest time ≤10.53 A1 for 9.45 – 9.455
24	$\frac{4}{7}\times\frac{4}{7}\times\frac{3}{7} + \frac{3}{7}\times\frac{3}{7}\times\frac{4}{7} =$ $\frac{48+36}{343} = \frac{84}{343}$ But there are three ways this can be achieved: BBG, BGB, GBB So the probability is $\frac{84}{343}\times 3$ OR $1-\frac{64}{343}-\frac{27}{343}$ NB: 84/343 = 0.244897; 252/343 = 0.73469 $\frac{4}{7} = 0.57(142...)$, $\frac{3}{7} = 0.42(857...)$	$\frac{252}{343}$	3	M1 for $\frac{4}{7}\times\frac{4}{7}\times\frac{3}{7}$ or $\frac{3}{7}\times\frac{3}{7}\times\frac{4}{7}$ oe or $\left(\frac{4}{7}\right)^3$ oe or $\left(\frac{3}{7}\right)^3$ oe or $\frac{91}{343}$ or 0.10(49…) or 0.13(99….) M1 (indep) for identification of all 6 outcomes (M2 for $1-\left[\left(\frac{4}{7}\right)^3+\left(\frac{3}{7}\right)^3\right]$) oe A1 $\frac{252}{343}$, $\frac{36}{49}$, 0.73(469…) oe

Paper 5525_06					
No		**Working**	**Answer**	**Mark**	**Notes**
25	(a)	$\frac{1}{3} \times \pi \times 5^2 \times 8 = \pi \times 25 \times 8 \div 3 =$ 209.4395	209–210	2	M1 for $\frac{1}{3} \times \pi \times 5^2 \times 8$ A1 for 209–210
	(b)	Base radius = $\frac{216}{360} \times 15 = 9$ Height = $\sqrt{(15^2 - 9^2)} =$	12	4	M1 for 216 ÷ 360 A1 for 9 M1 for $\sqrt{(15^2 - \text{“9”}^2)}$, where “9” < 15 A1 cao

PMT

edexcel

Mark Scheme (Results)

March 2013

GCSE Mathematics (2MB01) Higher
5MB3H (Calculator) Paper 01

PMT

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk for our BTEC qualifications.
Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

If you have any subject specific questions about this specification that require the help of a subject specialist, you can speak directly to the subject team at Pearson.
Their contact details can be found on this link: www.edexcel.com/teachingservices.

You can also use our online Ask the Expert service at www.edexcel.com/ask. You will need an Edexcel username and password to access this service.

Pearson: helping people progress, everywhere
Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

March 2013
Publications Code UG035055
All the material in this publication is copyright
© Pearson Education Ltd 2013

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labeling conventions.

ii) *select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect canceling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

PMT

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

12 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers

Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

Guidance on the use of codes within this mark scheme
M1 – method mark A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
*1			small and correct answers	3	M1 for 4.2 ÷ 1.75 (=2.4(0)) **or** 8.68 ÷ 3.5 (=2.48) M1 for 4.2 ÷ 1.75 (=2.4(0)) **and** 8.68 ÷ 3.5 (=2.48) C1 (dep on at least M1) for small and (£)2.4(0) and (£)2.48 **OR** M1 for 1.75 ÷ 4.2 (=0.41(6)) **or** 3.5 ÷ 8.68 (=0.40(3)) M1 for 1.75 ÷ 4.2 (=0.41(6)) **and** 3.5 ÷ 8.68 (=0.40(3)) C1 (dep on at least M1) for small and 0.41(6)(kg) and 0.40(3)(kg) **OR** M1 for 4.2 ÷ 1.75 (=2.4(0)) **or** 8.68 ÷ 3.5 (=2.48) M1 for (4.2 ÷ 1.75) × 3.5 (=8.4(0)) or (8.68 ÷ 3.5) × 1.75 (=4.34) C1 (dep on at least M1) for small and (£)8.4(0) or for small and (£)4.34 **OR** M1 for 1.75 ÷ 4.2 (=0.41(6)) or 3.5 ÷ 8.68 (=0.40(3)) M1 for (1.75 ÷ 4.2) × 8.68 (=3.61(6)) or (3.5 ÷ 8.68) × 4.2 (=1.69(3)) C1 (dep on at least M1) for small and 3.61(6)(kg) or 1.69(3)(kg) **OR** M2 for 8.68 ÷ 2 (=4.34) or 4.20 × 2 (=8.40) C1 (dep on at least M1) for small and (£)4.34 or (£8.40) **OR** Any other method considered equivalent. Figures may be truncated or rounded as long as their method is clear.

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
2		$7 \times 8^2 + 320 = 448 + 320$ 768p = 7.68	£7.68	3	M1 for 7×8^2 or 448 seen A1 for 768 B1 for correct monetary answer written correctly eg £7.68 or 768p
3			correct shaded region	2	B1 for arc of circle, centre oak tree, radius 4cm(±2mm) B1 for shaded region where the tree can be planted, bounded by sides of garden and arc of circle centre oak tree
4	(a)		−3, −2, −1,0,1	2	B2 for all 5 values and no others (B1 for 4 correct values and no others or -4,-3,-2,-1,0,1 or -3,-2,-1,0,1,2
	(b)		$-2 \leq x < 4$	2	B2 for $-2 \leq x < 4$ (B1 for $-2 \leq x$ or $x < 4$ or $-2 < x \leq 4$) [Note: accept the use of any letter other than x throughout and ignore any attempt to list integer values]
5			25.1-25.2	2	M1 for $2 \times 3.142 \times 4$ or 8π oe A1 for 25.1 – 25. 2
6	(a)		5, −4, −3	2	B2 for 5, −4 and −3 (B1 for 5 or −4 or −3)
	(b)		correct curve	2	B2 for fully correct curve (B1 ft for at least 5 points plotted correctly)

PMT

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
7			enlargement, scale factor 2, centre (6, −5)	3	B1 for enlargement B1 for (scale) factor 2 or ×2 or sf 2 B1 for (6, −5) NB: award 0 marks for an explanation that includes reference to more than one transformation.
*8		Cost of wages, rent and other expenses last year 92 000+10 800+7000=109 800 Cost of wages, rent and other expenses this year $1.075\times92\,000+\frac{7}{9}\times 10$ 800 + 7000÷2 = 98 900+8400+3500 = 110 800 Costs are more this year, so she cannot increase the amount she spends on goods	no, with working and reason	4	M1 for an attempt to calculate using 7.5% and 92000 (eg 7.5% of 92000 or increase of 7.5%) eg 92000 + 6900 or 98900 M1 for an attempt to find a fractional amount of 10800 (eg $\frac{7}{9}\times 10800$ or 8400 or $\frac{2}{9}\times 10800$ or 2400) M1 for complete method of increasing 92000 by 7.5%, finding $\frac{7}{9}$ of 10800, and halving 7000, with at least all of these three added together. C1 for statement eg "no" with both 110800 & 109800 seen, OR "no" with "1000 more" OR "no" and compares 181800 with 182800, oe.
9	(a)	$\sqrt{30}=5.4772255\ldots$ $2.5^2=6.25$ 5.4772255...÷ 6.25 =	0.876	2	M1 for $\sqrt{30}\div 6.25$ or $5.4(7...)\div 2.5^2$ A1 for any answer in the range 0.876 to 0.877
	(b)	4.5 × 1000 × 1000	4 500 000	2	M1 for complete method equivalent to 4.5 × 1000 × 1000 A1 for 4 500 000 oe

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
10		3 → 21 4 → 56 3.1 → 23.5(9) 3.2 → 26.3(6) 3.3 → 29.3(3) 3.4 → 32.5(0) 3.5 → 35.8(7) 3.6 → 39.4(5) 3.7 → 43.2(5) 3.8 → 47.2(7) 3.9 → 51.(5) 3.35 → 30.8(9)	3.3	4	B2 for a trial between 3.3 and 3.4 inclusive (B1 for a trial between 3 and 4 inclusive) B1 for a different trial between 3.3 and 3.4 exclusive B1 (dep on at least one previous B1) for 3.3 NB trials should be evaluated to at least 1dp truncated or rounded
11			45	3	M1 for 60000 × 0.75 or ÷1000 or digits 45 or the numbers 60 or 0.00075) M1 for the complete method 60 000 × 0.75 ÷ 1000 oe A1 cao
12	(a)		−7	2	M1 for an attempt to get *x*s on one side or number terms on one side A1 cao
	(b)		8.5	2	M1 for an attempt to divide each side by 4 or an attempt to expand the bracket eg $4 \times x - 4 \times 5$ A1 for 8.5 oe

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
*13		2500×0.04×2 =£200 2500×$(1.039)^2$ = £2698.80 £2698.80 – £2500 = £198.80 Scheme A gives £1.20 more	statement	4	M1 for a correct method to find 4% of 2500 or 3.9% of 2500 A1 for a correctly calculated amount 2700 or 2698.8(0) or 200 or 198.8(0); or percentage rate calculated over the 2 years for comparison: 3.93% or 3.976% oe M1 for a correct compound interest method using 3.9% and 2500 over 2 years C1 for statement of scheme A, with two correct comparable figures.
14		$3x - 2y = 7$ $7x + 2y = 13$ $10x = 20$ $x = 2$ $3\times2 - 2y = 7$ $-2y = 1$ $y = -0.5$	2, −0.5	3	M1 for a correct process to eliminate either x or y (allow one arithmetic error) M1 (dep) for correct substitution of their found variable or an otherwise correct method to eliminate the other variable A1 for 2 and −0.5 oe
15	(a)		3.7×10^{-3}	1	B1 cao
	(b)		49 000	1	B1 cao
	(c)	500 ÷ 250 000 [= 0.002] OR $(5 \times 10^2) \div (2.5 \times 10^5)$ [= $(5\div2.5)\times10^{2-5}$] OR $\frac{500}{250} \times \frac{1}{10^3}$ [= 2×10^{-3}] OR $\frac{500\div100}{250000\div100} = \frac{5}{2500}$ [= 0.002]	2×10^{-3}	2	M1 for an attempt to divide using two numbers that are in consistent form eg following an attempt to convert to ordinary numbers, or standard form numbers. A1 for 2×10^{-3}

PMT

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
16			12.7	3	M1 for 3.142 × 5 × 5 oe or 3.142 × 5 × 5 × '*h*' (=78.5 – 78.55) M1 for 1000 ÷ (3.142 × 5 × 5) A1 for 12.7 – 12.8 NB: multiples of π acceptable for M marks
17		$4x - 3 = 2x + 2y$ $2x - 3 = 2y$ $2x = 2y + 3$ $x = y + 1.5$	$x = y + 1.5$	3	M1 for attempt to expand brackets eg $2\times x+2\times y$ or divide through by 2 (each term). M1 for attempt to get x on one side of equation. A1 for $(x =) y + 1.5$, $\frac{2y + 3}{2}$, oe
18			(11, 21)	3	M1 for 5 – 2 or 9 – 3 OR 3 or 6 as long as these are not related to the 3 from *A* or multiple 3 or 9,18 M1 for 2 + 3 × "3" and 3 + 3 × "6" OR for 5 + 2 × "3" and 9 + 2 × "6" OR for 2 + 3 × "3" (=11) and (y=) 2 × '11' – 1 A1 cao SC: B1 if no method shown and answer shows x coordinate as 11 or y coordinate as 21.

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
19			4.6	5	M1 for $AC = \sqrt{(3^2+4^2)}$ (=5) or $DA = \sqrt{(5^2+2^2)}$ M1 for tan BAC = 0.75 oe or (BAC=) 36.8(6) or 36.9 M1 for tan DAC = 0.4 oe or (DAC=) 21.8(0) M1 for $\sqrt{'5'^2+2^2}$ × sin("36.8(6)"+"21.8(0)") or 5.385 × sin 58.7 A1 for 4.59 – 4.61 OR M1 for $AC = \sqrt{3^2+4^2}(=5)$ or $DP = \sqrt{(29 - AP^2)}$ M1 for cos BAC = 0.8 oe or (BAC=) 36.8(6) oe M1 for angle CDP = angle BAC M1 for (DP=) 3 + 2×cos CDP or (AP=) 4 – 2×sin CDP A1 for 4.59 – 4.61
20			0.5, −3	3	M1 for $(2x \pm 1)(x \pm 3)$ M1 for $(2x - 1)(x + 3)$ A1 cao OR M1 for $\frac{-5 \pm \sqrt{5^2 - 4 \times 2 \times -3}}{2 \times 2}$ allow sub. of $c = \pm 3$ M1 for $\frac{-5 \pm \sqrt{49}}{4}$ A1 cao NB: any other method put in review.

PMT

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
21		$1180.5 \div \frac{4}{3}\pi(6.15)^3$	1.21	4	M1 for 1180 ± 0.5 or 6.2 ± 0.05 M1 for 'max. mass' ÷ 'min. volume' where "min volume" is defined as any expression involving 6.15 M1 for $1180.5 \div \frac{4}{3}\pi(6.15)^3$ A1 for 1.21 – 1.22
22			proof	3	B1 for *AD* = *AB* and *AE* = *AG* B1 for angle *EAB* = angle *DAG* (= 90 + angle *DAE*) B1 (dep on previous B marks) for 2 sides and included angle (SAS) oe
23	(a)		$3\boldsymbol{a} + \boldsymbol{b}$	1	B1 oe
	(b)		3**c**	4	M1 for writing *QB* as a combination of vectors eg $\frac{1}{2}$ ('3**a** + **b**') + **b** or 3**a** + 2**b** – $\frac{1}{2}$ ('3**a** + **b**') or 2**a** + 2**b** − **c** oe M1 for equating **a** + **b** = 2**c** or writing *PQ* in terms of vectors **a** and **b** eg −**a** + $\frac{1}{2}$ ('3**a** + **b**') or 2**a** + **b** – $\frac{1}{2}$ ('3**a** + **b**') A1 for (*PQ*=) $\frac{1}{2}$**a** + $\frac{1}{2}$**b or** (*QB*=) $\frac{3}{2}$**a** + $\frac{3}{2}$**b** or (*QB*=) 2**c**−**a** + **a** + **c** A1 cao NB: accept equivalent expressions throughout

PMT

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481
Email publication.orders@edexcel.com
Order Code UG035055 March 2013

For more information on Edexcel qualifications, please visit our website
www.edexcel.com

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual
Llywodraeth Cynulliad Cymru
Welsh Assembly Government
Rewarding Learning

PMT

edexcel

Mark Scheme (Results)

Summer 2014

Pearson Edexcel GCSE
In Mathematics A (1MA0)
Higher (Non-Calculator) Paper 1H

PMT

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2014
Publications Code UG039399
All the material in this publication is copyright
© Pearson Education Ltd 2014

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labelling conventions.

ii) *select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

12 Parts of questions
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers
Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

Guidance on the use of codes within this mark scheme
M1 – method mark A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

PAPER: 1MA0_1H					
Question		**Working**	**Answer**	**Mark**	**Notes**
1	(a)		$\frac{2}{21}$	1	B1 for $\frac{2}{21}$
	(b)		$\frac{4}{15}$	2	M1 for attempting to use a suitable common denominator with at least one of the two fractions correct A1 for $\frac{4}{15}$ oe
2			0 | 5 9 1 | 3 5 6 8 9 9 2 | 1 2 3 3 5 7 8 9 3 | 1 2 4 4 | 0	3	B2 for fully correct diagram. Accept a stem of 10, 20, etc. (B1 for ordered with at most 2 errors or omissions or for correct unordered diagram) B1 for a correct key (units may be omitted) consistent with diagram.
3		<table><tr><td></td><td>Sq</td><td>G</td><td>S</td><td>Tot</td></tr><tr><td>F</td><td>2</td><td>4</td><td>15</td><td>21</td></tr><tr><td>M</td><td>6</td><td>14</td><td>9</td><td>29</td></tr><tr><td>Tot</td><td>8</td><td>18</td><td>24</td><td>50</td></tr></table>	4	4	M1 for a correct first step which results in a value that could be in the table: eg. 50 – 18 – 8 (= 24) or 50 – 21 (= 29) or 8 – 6 (= 2) M1 for a correct method to find a second value that could be in the table using their first value eg "29" – 9 – 6 (=14) or "24" – 9 (=15) M1 for a fully correct and complete method. A1 cao
4			25.60	4	M1 for a correct method to find $\frac{1}{3}$ of 24 (=8) or $\frac{2}{3}$ of 24 (=16) M1 for a correct method to find 60% (= 7.2) or 40% (= 4.8) of 12 or 60% (=14.4) or 40% (= 9.6) of 24 M1 (dep on at least M1) for a method to find the sum of their discounted adult ticket + 2 × their discounted child ticket A1 25.6(0)

PAPER: 1MA0_1H					
Question		**Working**	**Answer**	**Mark**	**Notes**
5			Question given	2	B1 for a suitable question which includes a time frame (the time frame could appear with the response boxes) B1 for at least 3 non-overlapping exhaustive response boxes with no use of inequality symbols
6	(a)		$2m^2 + 6m$	1	B1 for $2m^2 + 6m$
	(b)		$3xy(y - 2)$	2	B2 for $3xy(y - 2)$ (B1 for $3x(y^2 - 2y)$ or $3y(xy - 2x)$ or $xy(3y - 6)$ or $3xy$(a two term algebraic expression))
*7			3	4	M1 for a method to calculate at least one area eg 10×7 (=70) or 16×10 (=160) M1 for a method to find the total area (=124) M1 (dep on M1) for "124" ÷ 36 C1 (dep on M3) for 3 (pigs) clearly identified and supported by correct calculations Or M1 for an area of $36m^2$ drawn with dimensions shown M1 for 3 areas of $36m^2$ drawn with dimensions shown M1 (dep on M1) for method to find the area left (=16) C1 (dep on M3) for 3 (pigs) clearly identified and supported by correct calculations
8			Correct region	3	B1 for full line drawn 1.5 cm from edge of patio and parallel to it B1 for full arc of circle radius 3 cm centre the centre of the pond B1 ft for shading region to the right of their vertical line **and** outside the arc of their circle with correct centre

PAPER: 1MA0_1H					
Question		**Working**	**Answer**	**Mark**	**Notes**
9	(a)		Shape drawn	2	B2 for shape with vertices at (0, –1), (–1, – 3), (–2, –3), (–2, –1) (B1 for rotation of 180° about the wrong centre)
	(b)		Triangle drawn	2	B2 for triangle with vertices at (6, 9), (9, 9), (9, 3) (B1 for 2 vertices correct or enlargement sf 3 in wrong position or enlargement, centre (0, 0), but sf >1, ≠ 3)
*10		1.18 ÷ 4 = 0.295 (118 ÷ 4 = 29.5) 1.74 ÷ 6 = 0.29 (174 ÷ 6 = 29) 1.18 ÷ 2 = 0.59 1.74 ÷ 3 = 0.58 1.74 × 4 = 6.96 1.18 × 6 = 7.08 1.74 × 2 = 3.48 1.18 × 3 = 3.54 1.18÷2×3=1.77 1.74÷3×2=1.16 4÷1.18=3.3(…) 6÷1.74=3.4(…)	6 pints	3	M1 for division of price by quantity for both bottles or division of quantity by price for both bottles or complete method to find price of same quantity of milk A1 for two correct values that could be used for a comparison C1 ft (dep on M1) for comparison of their values with a correct conclusion.
11			36	3	M1 for a correct method to find either an interior or an exterior angle; eg. (180 × 3) ÷ 5 or 540 ÷ 5 (=108) or 360 ÷ 5 (=72) M1 (dep) for a complete method to find angle *CFD*. A1 cao

PAPER: 1MA0_1H					
Question		**Working**	**Answer**	**Mark**	**Notes**
12	*(a)		No, temp is 25°C	3	M1 for substitution of 77 into the RHS of the formula A1 for 25 cao or for 225/9 and 180/9 cao C1 (dep on M1) for conclusion (ft) following from working shown OR M1 for substitution of 20 into LHS of formula and correct process to find F A1 for 68 cao C1 (dep on M1) for conclusion (ft) following from working shown
	(b)		$F = \frac{9C}{5} + 32$	3	M1 for expansion of the brackets (eg $5 \times F - 5 \times 32$) or an attempt to multiply both sides by 9, or divide both sides by 5 as the first step. M1 (dep) for a correct second step A1 for $F = \frac{9C}{5} + 32$ oe
*13			Answer in range 35 – 50	4	M1 for a method to either find the exact or approximate number of seconds in one day, e.g. $24 \times 60 \times 60$ (=86400) or the number of minutes in 2014 seconds, e.g. $2014 \div 60$ or $2000 \div 60$ (≈30) M1 for a correct method to find the number of prizes; eg. '$24 \times 60 \times 60$' $\div$ 2014 oe or $60 \div$ "30" $\times 24$ oe B1 for rounding at least one appropriate value in the working to 1 sf, e.g. 24 rounded to 20 or 2014 rounded to 2000 or 86400 rounded to 90000 C1 (dep on M2) for answer in 35 – 50 clearly identified
14			6 hours	4	B1 for 5 miles = 8 km or equivalent statement or for $\frac{8}{5}$ or $\frac{5}{8}$ used correctly M1 for $50 \times r$ with $1.5 \leq r \leq 1.7$ oe or $480 \times s$ with $0.6 \leq s \leq 0.7$ oe M1 for $480 \div$ speed or distance $\div 50$ A1 oe

<table>
<tr><th colspan="6">PAPER: 1MA0_1H</th></tr>
<tr><th colspan="2">Question</th><th>Working</th><th>Answer</th><th>Mark</th><th>Notes</th></tr>
<tr><td>15</td><td>(a)</td><td></td><td>2, –1, 2, 7</td><td>2</td><td>B2 for all correct
(B1 for 2 or 3 correct)</td></tr>
<tr><td></td><td>(b)</td><td></td><td>Correct graph</td><td>2</td><td>M1 (dep on at least B1) for at least 6 points from their table plotted correctly
A1 cao for fully correct graph</td></tr>
<tr><td></td><td>(c)</td><td>$x^2 - 3x - 4 = 0$
$(x - 4)(x + 1) = 0$</td><td>–1, 4</td><td>2</td><td>M1 for line $y = x + 3$ drawn correctly or for reduction to correct 3 term quadratic (=0) and :
$(x \pm 1)(x \pm 4)$
or formula using $a = 1$, $b = -3$ and $c = -4$, allow one sign error in the formula,
or $\left(x - \frac{3}{2}\right)^2 = 4 + \left(\frac{3}{2}\right)^2$
A1 cao</td></tr>
<tr><td>16</td><td>(a)</td><td></td><td>0.75</td><td>2</td><td>M1 for “5.6” – “4.85” with at least one value correct
A1 cao</td></tr>
<tr><td></td><td>(b)</td><td></td><td>20</td><td>2</td><td>M1 for a complete method e.g. 80 ÷ 4
A1 cao</td></tr>
<tr><td></td><td>(c)</td><td><table><tr><th></th><th>1st half</th><th>2nd half</th></tr><tr><td>Med</td><td>5.3</td><td>4.75</td></tr><tr><td>Range</td><td>2.2</td><td>2.45</td></tr><tr><td>IQR</td><td>0.75</td><td>0.75</td></tr></table></td><td>2 comparisons</td><td>2</td><td>B1 ft from (a) for a correct comparison of a measure of spread
B1 for a correct comparison of medians (accept averages)
For the award of both marks at least one of the comparisons made must be in the context of the question.</td></tr>
</table>

PAPER: 1MA0_1H					
Question		**Working**	**Answer**	**Mark**	**Notes**
17	(a)		1	1	B1 cao
	(b)		$\frac{1}{100}$	1	B1 for $\frac{1}{100}$ or 0.01
	(c)		0.00273 27.3×10^{-3} 2.73×10^{3} 273×10^{2}	2	M1 for converting all numbers to same form with at least one conversion correct A1 for fully correct order with correct numbers in any correct form (SC B1 if one number incorrectly placed or all 4 numbers listed in reverse order)
18			$x = 7$ $y = -3$	3	M1 for correct process to eliminate one variable (condone one arithmetic error) M1 (dep) for substituting found value in one of the equations or appropriate method after starting again (condone one arithmetic error) A1 for $x = 7$ and $y = -3$
19			$y = \frac{1}{2}x - 5$	3	M1 for method to find gradient of L_1 e.g $\frac{6-3}{6-0}$ $\left(=\frac{1}{2}\right)$ M1 for $y =$ "$\frac{1}{2}$"$x + c$ or $y = mx - 5$ (c, m do not have to be numerical, or correct numerical values) or for $(L =)\frac{1}{2}x - 5$ A1 $y = \frac{1}{2}x - 5$ oe

PAPER: 1MA0_1H					
Question		**Working**	**Answer**	**Mark**	**Notes**
20			6	3	M1 for $\frac{15}{10}$ (=1.5) or $\frac{10}{15}$ (=0.66..) or $\frac{16}{10}$(=1.6) or $\frac{10}{16}$(=0.625) M1 for $\frac{15}{10}$ × 16 (=24) oe A1 cao OR M1 for $\frac{15}{16}$(=0.9375) or $\frac{16}{15}$ (=1.066…) or $\frac{16}{10}$(=1.6) or $\frac{10}{16}$(=0.625) M1 for $\frac{15}{16}$× 10 (=9.375) oe A1 20.625 oe
21			55	3	M1 for angle ABO = 90 or angle ADO = 90, or angle OBC = 15 or angle FDO = 90 or angle EBO = 90 (could be marked on the diagram) M1 for reflex angle BOD = 360 – (360 – 90 – 90 – 40) (= 220) or angle BCD = (360 – 90 – 90 – 40) ÷ 2 (= 70) or angle BDO or angle DBO = 90 – (180 – 40)/2 (= 20) or angle BOC = 180 – (15 + 15) (=150) A1 cao
22	(a)		$27x^6y^{12}$	2	B2 cao (B1 for any two terms correct in a product, e.g. $9x^6y^{12}$)
	(b)		$\frac{x-3}{2x-1}$	3	M1 for $(x-3)(x+3)$ M1 for $(2x-1)(x+3)$ A1 for $\frac{x-3}{2x-1}$ cao

PMT

PAPER: 1MA0_1H					
Question		**Working**	**Answer**	**Mark**	**Notes**
23	(a)		$\frac{3}{10}, \frac{6}{9}, \frac{3}{9}, \frac{7}{9}, \frac{2}{9}$	2	B1 for $\frac{3}{10}$ on LH yellow branch B1 for $\frac{6}{9}, \frac{3}{9}, \frac{7}{9}, \frac{2}{9}$ correct on tree diagram
	(b)		$\frac{48}{90}$	3	M1 for $\frac{7}{10} \times "\frac{3}{9}"$ or $"\frac{3}{10}" \times "\frac{7}{9}"$ or $"\frac{3}{10}" \times "\frac{2}{9}"$ M1 for $\frac{7}{10} \times "\frac{3}{9}" + "\frac{3}{10}" \times "\frac{7}{9}" + "\frac{3}{10}" \times "\frac{2}{9}"$ A1 for $\frac{48}{90}$ oe OR M1 for $\frac{7}{10} \times "\frac{6}{9}"$ M1 for $1 - \frac{7}{10} \times "\frac{6}{9}"$ A1 for $\frac{48}{90}$ oe

PAPER: 1MA0_1H					
Question		**Working**	**Answer**	**Mark**	**Notes**
*24			Proof	3	M1 for $\overrightarrow{MN} = \overrightarrow{MO} + \overrightarrow{ON}$ (= **n** – **m**) or $\overrightarrow{NM} = \overrightarrow{OM} + \overrightarrow{NO}$ (= **m** – **n**) or $\overrightarrow{AB} = \overrightarrow{AO} + \overrightarrow{OB}$ (= 2**n** – 2**m**) or $\overrightarrow{BA} = \overrightarrow{OA} + \overrightarrow{BO}$ (= 2**m** – 2**n**) M1 for $\overrightarrow{MN}$= **n** – **m** and $\overrightarrow{AB}$= 2**n** – 2**m** oe C1 (dep on M1, M1) for fully correct proof, with $\overrightarrow{AB}$= 2$\overrightarrow{MN}$ or $\overrightarrow{AB}$ is a multiple of $\overrightarrow{MN}$ [SC M1 for $\overrightarrow{MN}$ = 0.5**n** – 0.5**m** and $\overrightarrow{AB}$ = **n** – **m** C1 (dep on M1) for fully correct proof, with $\overrightarrow{AB}$= 2$\overrightarrow{MN}$ or $\overrightarrow{AB}$ is a multiple of of $\overrightarrow{MN}$]
25	(a)		$4\sqrt{3}$	2	M1 for $\frac{12}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}}$ A1 for $\frac{12\sqrt{3}}{3}$ oe with a rational denominator
	(b)		18	2	M1 for $\sqrt{2}\times\sqrt{2} + \sqrt{2}\times\sqrt{8} + \sqrt{8}\times\sqrt{2} + \sqrt{8}\times\sqrt{8}$ oe A1 cao OR M1 for $(\sqrt{2} + 2\sqrt{2})^2$ A1cao

PAPER: 1MA0_1H					
Question		**Working**	**Answer**	**Mark**	**Notes**
26	(a)		180, 0	1	B1 for 180, 0 Accept π, 0
	(b)		270, –1	1	B1 for 270, –1 accept $\frac{3\pi}{2}$, –1
	(c)		$a = 2$ $b = 3$ $c = 1$	3	B1 cao B1 cao B1 cao

Modifications to the mark scheme for Modified Large Print (MLP) papers.

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 1MA0_1H			
Question		**Modification**	**Notes**
Q08		Diagram made x 2 size. 3 metres changed to 6 metres. 6 metres changed to 12 metres.	B1 for full line drawn 3 cm from edge of patio and parallel to it B1 for full arc of circle radius 6 cm centre the centre of the pond B1 ft for shading region to the right of their vertical line **and** outside the arc of their circle with correct centre
Q09	(a) (b)	2 cm grid. A cut-out shape is provided. Axis removed. 2 cm plain grid. L shape put on grid. 3 squares down and an extra square across. Total 4 squares. Candidates asked to enlarge the shape by SF3	B2 for shape with vertices at (0, –1), (–1, – 3), (–2, –3), (–2, –1) (B1 for rotation of 180° about the wrong centre) B2 for a correct enlargement (B1 for an enlargement with a SF other than 3 or for 3 sides correct)
Q10		No pictures	M1 for division of price by quantity for both bottles or division of quantity by price for both bottles or complete method to find price of same quantity of milk A1 for two correct values that could be used for a comparison C1 ft (dep on M1) for comparison of their values with a correct conclusion.

PMT

PAPER: 1MA0_1H			
Question		**Modification**	**Notes**
Q14		No picture, just a box giving the distances	B1 for 5 miles = 8 km or equivalent statement or for $\frac{8}{5}$ or $\frac{5}{8}$ used correctly M1 for $50 \times r$ with $1.5 \leq r \leq 1.7$ oe or $480 \times s$ with $0.6 \leq s \leq 0.7$ oe M1 for 480 ÷ speed or distance ÷ 50 A1 oe
Q15		Grid x 2 – 2 cm squares. Top row removed.	B2 for all correct (B1 for 2 or 3 correct) M1 (dep on at least B1) for at least 6 points from their table plotted correctly A1 cao for fully correct graph M1 for line $y = x + 3$ drawn correctly or for reduction to correct 3 term quadratic (=0) and : $(x \pm 1)(x \pm 4)$ or formula using $a = 1$, $b = -3$ and $c = -4$, allow one sign error in the formula, or $\left(x - \frac{3}{2}\right)^2 = 4 + \left(\frac{3}{2}\right)^2$ A1 cao

PMT

PAPER: 1MA0_1H			
Question		**Modification**	**Notes**
Q16	(a)	Box plots: "first half" – relevant figures are 4.25, 4.75, 5.25, 5.5, 6.4.	M1 for "5.5" – "4.75" with at least one value correct A1 cao
	(c)	"Second half" figures are 3.6, 4.25, 4.75, 5, and 6. 2 cm squares	B1 ft from (a) for a correct comparison of a measure of spread B1 for a correct comparison of medians (accept averages) For the award of both marks at least one of the comparisons made must be in the context of the question.
Q19		2 cm grid	M1 for method to find gradient of L_1 e.g $\frac{6-3}{6-0}$ $\left(=\frac{1}{2}\right)$ M1 for $y =$ "$\frac{1}{2}$"$x + c$ or $y = mx - 5$ (c, m do not have to be numerical, or correct numerical values) or for $(L =)\frac{1}{2}x - 5$ A1 $y = \frac{1}{2}x - 5$ oe

PAPER: 1MA0_1H			
Question		**Modification**	**Notes**
Q20		Picture removed – box with the word "Photo inserted" Diagram labelled Diagram (i) and (ii)	M1 for $\frac{15}{10}$ (=1.5) or $\frac{10}{15}$ (=0.66..) or $\frac{16}{10}$(=1.6) or $\frac{10}{16}$(=0.625) M1 for $\frac{15}{10} \times 16$ (=24) oe A1 cao OR M1 for $\frac{15}{16}$(=0.9375) or $\frac{16}{15}$ (=1.066…) or $\frac{16}{10}$(=1.6) or $\frac{10}{16}$(=0.625) M1 for $\frac{15}{16} \times 10$ (=9.375) oe A1 20.625 oe
Q22	(b)	MLP only : x changed to y	M1 for $(y-3)(y+3)$ M1 for $(2y-1)(y+3)$ A1 for $\frac{y-3}{2y-1}$ cao
Q25		x 2 size, solid answer lines Braille only – Roman numerals inserted on the five answer lines	Standard mark scheme

PMT

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

PMT

edexcel

Mark Scheme (Results)

November 2015

Pearson Edexcel GCSE
In Mathematics A (1MA0)
Higher (Non-Calculator) Paper 1H

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

November 2015
Publications Code UG042920
All the material in this publication is copyright
© Pearson Education Ltd 2015

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Note that in some cases a correct answer alone will not score marks unless supported by working; these situations are made clear in the mark scheme. Examiners should be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will award marks for the quality of written communication (QWC).
The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labelling conventions.

ii*) select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Partial answers shown (usually indicated in the ms by brackets) can be awarded the method mark associated with it (implied).
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks; transcription errors may also gain some credit. Send any such responses to review for the Team Leader to consider.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded (embedded answers).

12 Parts of questions
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers
Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

14 The detailed notes in the mark scheme, and in practice/training material for examiners, should be taken as precedents over the above notes.

Guidance on the use of codes within this mark scheme
M1 – method mark for appropriate method in the context of the question A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

1MA0 1H November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
1			69	4	M1 for finding 15% of £720 (=108) M1 (dep) for finding total of £720 plus interest (or 115% etc) (=828) M1 (dep on previous M1) dividing by 12 A1 cao OR M1 finding 720 ÷ 12 (=60) M1 (dep) finding 15% of "60" (=9) M1 (dep on previous M1) for adding, e.g. 60 + 9 A1 cao
2	(i)		19.44	2	B1 cao
	(ii)		19 440		B1 cao
3	(a)		$6n + 5$	2	B2 for $6n + 5$ (B1 for $6n + k$ where k is an integer or absent)
	(b)		No with explanation	2	M1 for "$6n + 5$" = 121 or any other valid method, e.g. counting on 6s (to get to 119 or more) A1 for No with complete explanation, e.g. $6n$=116 will not give a whole number
4	(a)		60	2	M1 for 200 × 0.3 oe A1 cao
	(b)		0.1	2	M1 subtracting sum of probabilities from 1, e.g. 1−(0.3+0.2+0.4) A1 cao

1MA0 1H November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
5			20	3	M1 for 330÷120 (=2.75) or 200÷60 (=3 1/3) or 450 ÷180 (=2.5) M1 for 450÷180 (=2.5) AND 8×"2.5" A1 cao OR M1 for 120÷8 (=15) or 60÷8 (=7.5) or 180÷8 (=22.5) M1 for 330÷(120÷8) [=22] or 200÷(60÷8) [=26.6..] or 450÷(180÷8) A1 cao OR M1 for multiples of 120:60:180 M1 for multiplication linked to 450 and 8+8+4 A1 cao
6			40°	4	M1 for angle FBC=70 or CFG = x or ABF = 110 may be seen in diagram M1 for angle CBF = BFC =70 or 90 – ½ x A1 for 40 supported by working C1 (dep on M2) for all reasons and linked to appropriate working, e.g. Alternate angles are equal; Allied angles / Co-interior angles add up to 180°; Base angles of an isosceles triangle are equal; angles in a triangle add to 180°, angles on a straight line equals 180°
7	(a)		explanations	2	B2 for two aspects from: no time frame; responses vague; no "never" box (B1 for one correct aspect)
	(b)		question response boxes+	2	B1 for a question with a time frame (may appear with response boxes) B1 for at least 3 correctly labelled response boxes (non-overlapping and exhaustive) Do not accept inequality symbols.

1MA0 1H November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
8		2p 1p ½ p Tot Sat 7 16 (31) 54 Sun (15) 14 17 (46) Tot (22)(30) 48 (100)	14	4	M1 for total Sat bottles 100 – 46 (=54) or for total ½ pint bottles 100 – 22 – 30 (=48) or (total 2 pint bottles on Sat) 22 – 15 (=7) M1 for total Sun bottles of ½ pint "48" – 31 (=17) or for total Sat bottles of 1 pint: "54" – 31 – (22 −15) (=16) M1 for 46 – 15 – "17" or for 30 – "16" A1 cao NB: any of the above figures could be shown in a 2-way table
*9			NO with evidence	4	M1 for 50 × 40 × 30 (=60000) M1 for "60000" ÷ 3000 (=20) M1 for "20" × £3.50 C1 eg for 70 and comparison resulting in NO OR M1 for £60 ÷ 3.50 (=17 bottles) M1 for "17" × 3000 (=51000) M1 for 50 × 40 × 30 (=60000) C1 eg for 51000 and 60000 and comparison resulting in NO

1MA0 1H November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
10	(a)		$x^2 + 2x$	1	B1 cao
	(b)		$3y + 4x + 2$	2	M1 for a method to expand a bracket, e.g. $3y + 6$ or $4x - 4$ A1 cao
	(c)	$2t^2 + 10t - 3t - 15$	$2t^2 + 7t - 15$	2	M1 for 4 terms correct ignoring signs or 3 out of no more than 4 terms with signs correct unless ambiguous A1 cao
	(d)		$4a(2a + 3)$	2	M1 for $4a(na+c)$ or $2a(4a+6)$ or $a(8a+12)$ [n,c integers, $c \neq 0$] A1 cao
	(e)		$(y + 1)(y - 2)$	2	M1 for $(y \pm 1)(y \pm 2)$ unless ambiguous A1 cao
11	(a)		150	2	M1 for $180 - (360 - 330)$ or $180 - 30$ or $330 - 180$ or a complete diagram showing the bearing of 330° A1 cao
	(b)		11 40	4	M1 for $200 \div 120$ (=1 2/3 h) M1 for conversion between hours and minutes A1 for 1 h 40 min or 100 minutes B1 (ft dep on M1) for 11 40

1MA0 1H November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
12	(a)		2, 0, 0, 6	2	B2 for 2, 0, 0, 6 (B1 for at least two of 2, 0, 0, 6); could be taken from graph
	(b)		Correct curve	2	M1 (ft) for at least 5 points plotted correctly A1 for a fully correct curve
	(c)		−0.6, 3.6	2	M1 (ft if M1 awarded in (b) and at least B1 in (a)) for indicating a point or line drawn at $y=4$, or one solution given A1 (ft) for both solutions
13			20	3	M1 for 30 × 14 (=420) or 18×10 (=180) M1 for 30 × 14 – 18 × 10 or "420" – "180" (=240) A1 cao
14			126	4	M1 for method to find exterior or interior angle of octagon M1 for method to find exterior or interior angle of pentagon M1 for complete method A1 cao

1MA0 1H November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
15	(a)		19, 36, 51, 63, 73, 80	1	B1 cao
	(b)		cf graph	2	M1 for at least 5 of the 6 points plotted at each upper end of the interval (not joined) or 5 of the 6 points plotted consistently within interval (not upper end) and joined (dep on a cf table with no more than one arithmetic error) A1 correct graph
	*(c)		comparable value and conclusion	3	M1 for indication of a reading taken from a cf graph using weight = 3.4 kg or find UQ from 60 A1 for value given between 55 & 57 or 3.6 & 3.8 C1 (dep on at least M1) for conclusion (justified)
16			13.75	5	M1 for finding perimeter of rectangle e.g. $5x + 5 + 5x + 5 + 4x + 4x$ $(= 18x + 10)$ M1 for finding perimeter of trapezium e.g. $9x - 2 + 7x - 2 + 10x$ $(=26x - 4)$ M1 for equation e.g. $26x - 4 = 18x + 10$ (or $8x = 14$) A1 for finding the value of x as 1.75 B1 ft for subs of x into $5x + 5$ and evaluated (=13.75)
17			$x = 3\frac{1}{3}$ $y = -2$	4	M1 for a correct process to eliminate either variable (condone one arithmetic error) or to rearrange and substitute for elimination A1 cao for either x or y M1 (dep on M1) for correct substitution of found value into one of the equation or appropriate method after starting again (condone one arithmetic error) A1 cao

1MA0 1H November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
18			$2\sqrt{5}$	2	M1 for multiplication of denominator and numerator by $\sqrt{5}$ A1 cao
19			756π	5	M1 for $\frac{1}{3}\pi r^2 \times 10$ $(=270\pi)$ A1 for $r = 9$ M1 (dep on M1) for $\frac{1}{2} \times \frac{4}{3}\pi \times "9"^3$ $(= 486\pi)$ M1 for 270π + "486π" oe A1 cao
*20			Proof	5	M1for finding one other vector expressed as **a** and/or **b** M1 for method to find one of $\overrightarrow{DM}$, $\overrightarrow{MA}$ or $\overrightarrow{DA}$ eg $\overrightarrow{DM} = -\mathbf{b} + \frac{1}{2}(3\mathbf{b} + \mathbf{a})$ oe, $\overrightarrow{MA} = \frac{1}{2}(3\mathbf{b} + \mathbf{a}) + \mathbf{a}$ oe or $\overrightarrow{DA} = 2\mathbf{b} + 2\mathbf{a}$ oe M1 for method to find two of $\overrightarrow{DM}$, $\overrightarrow{MA}$ or $\overrightarrow{DA}$ A1 for two of $\overrightarrow{DM} = \frac{1}{2}(\mathbf{a} + \mathbf{b})$, $\overrightarrow{MA} = 1.5(\mathbf{a} + \mathbf{b})$, $\overrightarrow{DA} = 2(\mathbf{a} + \mathbf{b})$ ie simplified but oe C1 (dep on working shown) for conclusion relating to correct working

1MA0 1H November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
21	(a)		$\frac{9x-8}{x(2-x)}$	3	M1 for method to use a common denominator, e.g. $\frac{5(x)-4(2-x)}{x(2-x)}$ M1 (dep on M1) for correct expansion of brackets and combination of numerators e.g. $5x-8+4x$ $(=9x-8)$ A1 for $\frac{9x-8}{x(2-x)}$ or $\frac{9x-8}{2x-x^2}$
	(b)		$y=\frac{2-2t}{t+3}$	4	M1 for intention to multiply both sides by $y+2$ as a first step e.g. $t \times y+2=2-3y$ M1 for intention to correctly isolate their y terms on one side and the other terms on the other side, e.g. $ty+3y=2-2t$ M1 for intention to factorise, e.g. $y(t+3)$ $(=2-2t)$ A1 for $y=\frac{2-2t}{t+3}$ oe
*22			Similarity and proof	5	B1 for method matching a pair of opposite angles, e.g. if $EAB = x$, $BDE = 180-x$, $EAB + BDE = 180$ B1 for linking angles between quad and triangle, e.g. if $BDE = 180-x$ then $BDC = x$ B1 for stating or implying $ACE = BCD$ (same angle) C1 for Opposite angles of a cyclic quadrilateral add up to 180° or statement linking three angles for similarity C1 for complete proof

Modifications to the mark scheme for Modified Large Print (MLP) papers.

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 1MA0_1H			
Question		**Modification**	**Notes**
Q06		Diagram is enlarged.	Standard mark scheme
Q09		Model is provided for all candidates. Diagram also provided for MLP.	Standard mark scheme
Q10	(a)	MLP only: x changed to y Diagram is enlarged.	Standard mark scheme
Q12	(a) (b)	Wording added “There are four spaces to fill.” 1.5 cm grid..Small squares removed.	Standard mark scheme
Q14		Diagram enlarged.	Standard mark scheme
Q15	(c)	Frequencies changed to 25, 20, 15, 10, 5, 5. Grid: 1.5 cm 3.4 kg changed to 3.5 kg.	Standard mark scheme
Q16		“trapezium” and “rectangle” put inside the shapes. MLP only: x changed to y.	Standard mark scheme
Q19		Model provided for all candidates. Diagram also provided for MLP.	Standard mark scheme

PMT

PAPER: 1MA0_1H			
Question		**Modification**	**Notes**
Q20		Diagram is enlarged. Lower case vectors enlarged.	Standard mark scheme
Q22		Diagram is enlarged.	Standard mark scheme

PMT

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom

PMT

edexcel

Mark Scheme (Results)

Summer 2016

Pearson Edexcel GCSE
In Mathematics B (2MB01)
Higher (Calculator) Unit 3

PMT

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2016
Publications Code 5MB3H_01_1606_MS
All the material in this publication is copyright
© Pearson Education Ltd 2016

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Note that in some cases a correct answer alone will not score marks unless supported by working; these situations are made clear in the mark scheme. Examiners should be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will award marks for the quality of written communication (QWC).
The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labelling conventions.

ii) *select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 **With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Partial answers shown (usually indicated in the ms by brackets) can be awarded the method mark associated with it (implied).
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks; transcription errors may also gain some credit. Send any such responses to review for the Team Leader to consider.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 **Follow through marks**

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 **Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.

10 **Probability**

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded (embedded answers).

12 Parts of questions
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers
Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

Guidance on the use of codes within this mark scheme
M1 – method mark A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

PMT

PAPER: 5MB3H/01					
Question		**Working**	**Answer**	**Mark**	**Notes**
1*			Decision (supported)	3	M1 for correct first step e.g. 15 ÷ 3 M1 for complete method C1 for correct decision with accurate figures
2	(a)		Correct diagram	2	B2 cao (B1 for correct orientation)
	(b)		Enlargement, sf 4, centre (0, 1)	3	B1 for enlargement B1 for scale factor 4 B1 for centre (0, 1)
3			$w = \frac{d+5}{2}$	2	M1 for attempt to add 5 to both sides or divide both sides by 2 as a first step, eg $d + 5 = 2w$ oe or $\frac{d}{2} = w - \frac{5}{2}$ oe A1 for $w = \frac{d+5}{2}$ oe

PMT

PAPER: 5MB3H/01					
Question		Working	Answer	Mark	Notes
4			22.5	4	M1 for $4(x - 8) = 2x + 13$ M1 for expansion of bracket or division of all terms by 4, eg $4x - 32 = 2x + 13$, or $x - 8 = \frac{2x}{4} + \frac{13}{4}$ M1 for isolating x and number terms eg $2x = 45$, $\frac{x}{2} = \frac{45}{4}$ A1 for $\frac{45}{2}$ or 22.5 OR M1 for $(180 - 64) \div 2$ (= 58) M1 for $4(x - 8)$ = "58" or $2x + 13$ = "58" or "58" – 13 (= 45) M1 for isolating x and number terms eg $4x = 90$, $2x = 45$ or "45" ÷ 2 A1 for $\frac{45}{2}$ or 22.5 OR M1 for $64 + 4(x - 8) + 2x + 13$ M1 for $64 + 4(x - 8) + 2x + 13 = 180$ M1 for isolating x and number terms eg $6x = 135$ A1 for $\frac{45}{2}$ or 22.5
5			Correct region shaded	3	M1 for a circle centre M or N (accept arc of sufficient length to define the region) M1 for circle centre M radius 5 cm **and** circle centre N radius 3 cm (accept arc of sufficient length to define the region) A1 for correct region shaded

PMT

PAPER: 5MB3H/01					
Question		Working	Answer	Mark	Notes
*6			Conclusion (supported)	4	M1 for $\pi \times 120^2$ (= 45 216 – 45 249) M1 for "$\pi \times 120^2$" ÷ 1800 A1 for 25 – 25.2 C1 ft (dep on M2) for appropriate conclusion from their figures OR M1 for $\pi \times 120^2$ (= 45 216 – 45 249) M1 for 20 × 1800 A1 for 36 000 and 45 216 – 45 249 C1 ft (dep on M2) for appropriate conclusion from their figures OR M1 for $\pi \times 120^2$ (= 45 216 – 45 249) M1 for "$\pi \times 120^2$" ÷ 20 A1 for 2260 – 2263 C1 ft (dep on M2) for appropriate conclusion from their figures OR M1 for 1800 × 20 M1 for 36 000 ÷ π (=11 457 – 11 465) A1 for 107(.0…) C1 ft (dep on M2) for appropriate conclusion from their figures
7			161.50	5	M2 for a correct method to decrease 6720 by 20%, eg 6720 × 0.8 (= 5376) or 6720 × 0.2 (= 1344 and 6720 – 1344 (= 5376)) (M1 for a correct method to find 20% of 6720 eg 6720 × 0.2 or $\frac{20}{100} \times 6720$ (= 1344)) M1 for subtracting 1500 (= 3876) after a percentage calculation M1 "3876" ÷ 24 after the subtraction of 1500 A1 for 161.5(0)

PMT

PAPER: 5MB3H/01					
Question		**Working**	**Answer**	**Mark**	**Notes**
8		3 42 3.1 45.(291) 3.2 48.(768) 3.3 52.(437) 3.4 56.(304) 3.5 60.(375) 3.6 64.(656) 3.7 69.(153) 3.8 73.(872) 3.9 78.(819) 4 84 3.71 69.6(14811) 3.72 70.(078848) 3.73 70.5(45117) 3.74 71.(013624) 3.75 71.4(84375)	3.7	4	B2 for a trial between 3 and 4 exclusive (B1 for a trial at 3 or 4) B1 for a different trial of $3.7 < x \leq 3.75$ B1 (dep on at least one previous B1) for 3.7 NB. Trials should be evaluated to at least 2 sf truncated or rounded for values of x correct to 1 decimal place. Trials should be evaluated to at least 1 dp truncated or rounded for values of x correct to 2dp. **NB** No working scores 0 marks even if answer is correct
9			618	4	M1 for a method to find an area which is part of the cross section M1 (dep) for a complete method to find the total area of the cross section M1 (dep M1) for their cross sectional area × 12 A1 cao **OR** M1 for a method to find the volume of cuboid, 8 × 8 × 12 (= 768) M1 for a method to find the volume of the triangular prism, ½ ×5 × 5 × 12(= 150) M1 (dep M2) for a complete method for the volume of the prism A1 cao

5MB3H_01					
Question		Working	Answer	Mark	Notes
10	(a)		15.2	3	M1 for $12.7^2 + 8.3^2$ or $161.29 + 68.89$ or 230.18 M1 for √"$(12.7^2 + 8.3^2)$" or √"$161.29 + 68.89$" or $\sqrt{230.18}$ A1 for 15.2 or 15.17(168…)
	(b)		16.1	3	M1 for $\sin 40° = \frac{DE}{25}$ oe or $\frac{DE}{\sin 40} = \frac{25}{\sin 90}$ oe M1 for 25 sin 40° A1 for 16.1 or 16.06(969…)
11	(a)		4.081	2	M1 for 8.875 or $\frac{71}{8}$ or 2.979…or 4.08 A1 for 4.081 or 4.0809(5…)…
	(b)		4×10^{-3}	2	M1 for 4×10^n, $n \neq -3$ or $a \times 10^{-3}$, $a \neq 4$ or 0.004 or 0.4×10^{-2} or 1/250 A1 cao
12			$x < 2$	2	M1 for an attempt to isolate x and number terms or multiply all terms by 2, or for $\frac{3}{2}x < 3$ or $-\frac{3}{2}x < -3$ or $x = 2$ A1 cao
13	(a)		0, 6, 4, –6	2	B2 for all values correct (B1 for any one value correct)
	(b)		Graph drawn	2	M1 ft (provided B1 in (a) for at least 6 points plotted correctly from their table A1 cao for correct curve drawn from (–4, –6) to (3, –6)
	(c)		1.5 to 1.6 and –2.5 to –2.6	2	M1 for correct use $y = 2$ (may be implied by one correct estimate) or correct use of formula. A1 for 1.5 to 1.6 and –2.5 to –2.6

5MB3H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
14*		Sher Yas **Old 19300 12560** New 20265 13502 **Inc 965 942**	Correct comparison (supported)	3	M1 for a correct method to calculate the population at the beginning of the 10 year period for at least one place. eg $\frac{20265}{1.05}$ (= 19 300) oe or $\frac{13502}{1.075}$ (= 12 560) oe A1 for 965 (Sherbury) **and** 942(Yaston) C1 (dep on M1) ft for statement comparing increases leading to conclusion based on two comparable amounts eg increase in Sherbury is greater than increase in Yaston
15		$20x + 10y = 35$ $6x - 10y = -48$ $x = -\frac{1}{2}$ $y = \frac{9}{2}$ OR $12x + 6y = 21$ $12x - 20y = -96$ $26y = 117$ $y = \frac{9}{2}$ $x = -\frac{1}{2}$	$-\frac{1}{2}, \frac{9}{2}$	4	M1 for a correct process to eliminate either variable (condone one arithmetic error) A1 cao for either x or y M1 (dep on M1) for correct substitution of found value into one of the equations or appropriate method after starting again (condone one arithmetic error) A1 cao OR M1 for full method to rearrange and substitute to eliminate either variable (condone one arithmetic error) A1 cao for either x or y M1 (dep on M1) for correct substitution of found value into one of the equations or appropriate method after starting again (condone one arithmetic error) A1 cao **NB** Trial and improvement methods score 0 marks unless both x and y are correct
16			226	2	M1 for angle $ABC = 180 - 67$ (= 113) A1 cao OR M1 for obtuse angle $AOC = 2 \times 67$ (= 134) A1 cao

5MB3H_01					
Question		Working	**Answer**	**Mark**	**Notes**
17			– 2.82, 19.82	3	M1 for $\frac{--17 \pm \sqrt{(-17)^2 - 4\times1\times-56}}{2\times1}$ (condone one sign error) or $\left(x - \frac{17}{2}\right)^2$ oe M1 for $\frac{17 \pm \sqrt{513}}{2}$ or for method leading to $\frac{17}{2} \pm \sqrt{\frac{513}{4}}$ oe A1 for answers in the range 19.82 to 19.825; – 2.82 to –2.825
18			21.2	4	M1 for correct use of sine rule to find *AB* or *BC*, eg $\frac{6}{\sin 25} = \frac{AB}{\sin 120}$ or $\frac{6}{\sin 25} = \frac{BC}{\sin(180-120-25)}$ M1 for (*AB* =) $\frac{6}{\sin 25} \times \sin 120$ (= 12.2(951…)) or (*BC* =) $\frac{6}{\sin 25} \times \sin 35$ (= 8.1(431…) M1(dep M2) for method to find area, eg 0.5 × 6 × "*AB*" sin35 or 0.5 × 6 × "*BC*" sin 120 A1 for answer in the range 21.1 – 21.2 OR (with perpendicular from *A* meeting *BC* extended at a point *X*) M1 for method to find *AX* M1 for method to find *BX* M1(dep M2) for 0.5 × "*AX*" × "*BC*" A1 for answer in the range 21.1 – 21.2
*19			0.38 because the LB and UB agree to that number of figures	5	B1 for 10.75 or 10.85 or 10.84999… B1 for 75.055 or 75.065 or 75.064999… M1 for $\sqrt{\frac{10.85}{75.055}}$ as UB **OR** $\sqrt{\frac{10.75}{75.065}}$ as LB A1 (dep on all previous marks) for both 0.3802… and 0.3784 must clearly come from working with correct values C1 for 0.38 from 0.3784… and 0.3802 … **and** "both LB and UB round to 0.38" **NB** An answer of 038(932….) without working scores no marks

5MB3H_01					
Question		Working	**Answer**	**Mark**	**Notes**
20			7.35	3	M1 for $y \propto \frac{1}{\sqrt{x}}$ or $y = \frac{k}{\sqrt{x}}$ or $9 = \frac{k}{\sqrt{4}}$ M1 for $k = 9\sqrt{4}$ (=18) A1 for 7.35 or 7.348…
*21			$\frac{2}{3}$ (7**a** + 3**b**) and correct conclusion	4	M1 for correct vector for $\overrightarrow{OY}$, or $\overrightarrow{AP}$ eg $(\overrightarrow{OY}) = \overrightarrow{OA} + \overrightarrow{AY}$, or $\overrightarrow{OY} = \overrightarrow{OA} + \frac{2}{3}\overrightarrow{AP}$ may include terms in **a** and **b,** eg. 6**a** + $\overrightarrow{AY}$, 4**b** + 4**a** – **b** + $\overrightarrow{PY}$ or $\overrightarrow{AP} = \overrightarrow{AO} + \overrightarrow{OB} + \overrightarrow{BP}$ or –6**a** + 4**b** + 4**a** – **b** or – 2**a** + 3**b** M1 for $(\overrightarrow{AY}) = \frac{2}{3}\vec{AP}$ or ⅔(–6**a** + 4**b** + 4**a** – **b**) or ⅔(–2**a** + 3**b**) or $(\overrightarrow{PY} = \frac{1}{3}\overrightarrow{PA}$ or ⅓(– 4**a** + **b** – 4**b** + 6**a**) or ⅓(2**a** – 3**b**) M1 for correct expression for $\overrightarrow{OY}$ in terms of **a** and **b,** eg $\frac{14}{3}$ **a** + 2**b** C1 for $\frac{2}{3}$ (7**a** + 3**b**) and "$\overrightarrow{OY}$ is parallel to the vector 7**a** +3**b**" **oe**

PMT

Modifications to the mark scheme for Modified Large Print (MLP) papers.

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q2	(a)	Grid enlarged. Shading changed to dotty shading. Labels added to shapes. Image labelled shape A. Rotation given and labelled shape B. Question changed to 'Describe fully the single transformation which maps shape A to shape B. One row top and bottom, one column left and right has been removed. Three answer lines given.	2 cao (B1 for correct orientation in third quadrant or correct quarter turn)
Q2	(b)	Grid enlarged. Shading changed to dotty shading. Labels added next to the triangles 'triangle P' and 'triangle Q'. One column removed from the left.	B1 for enlargement B1 for scale factor 4 B1 for centre (0, 1)

PMT

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q4		Diagram enlarged. MLP only – x changed to y. Braille only – measurements given in the text.	M1 for $4(x - 8) = 2x + 13$ M1 for expansion of bracket or division of all terms by 4, eg $4x - 32 = 2x + 13$, $x - 8 = 2x/4 + 13/4$ M1 for isolating x and number terms eg $2x = 45$, $x/2 = 45/4$ A1 for 45/2 or 22.5 OR M1 for $(180 - 64) \div 2$ (= 58) M1 for $4(x - 8) = 58$ or $2x + 13 = 58$ M1 for isolating x and number terms eg $4x = 90$, $2x = 45$ A1 for 45/2 or 22.5 OR M1 for $64 + 4(x - 8) + 2x + 13$ M1 for $64 + 4(x - 8) + 2x + 13 = 180$ M1 for isolating x and number terms eg $6x = 135$ A1 for 45/2 or 22.5
Q5		Diagram kept the same size as original. Scale moved above the diagram and also put in the question paper.	B1 for circle centre M radius 5 cm B1 for circle centre N radius 3 cm B1 for correct region shaded

PMT

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q6		Diagram enlarged. Cross changed to a filled in circle. Arrow heads removed.	M1 for $\pi \times 120^2$ (= 45238….) M1 for "$\pi \times 120^2$" ÷ 1800 A1 for 25.0 – 25.2 C1 for "surface is large enough as 25 > 20" oe OR M1 for $\pi \times 120^2$ (= 45238….) M1 for 20 × 1800 A1 for 36000 C1 for "surface is large enough as 36000 < 45238..." oe OR M1 for $\pi \times r^2 = 1800 \times 20$ M1 for 36000 ÷ π (=11459..) A1 for 107.0… C1 for "surface is large enough as 107…< 120" oe
Q9		Model provided for all candidates. Diagram enlarged and provided for MLP.	M1 for a method to find an area which is part of the cross section M1 (dep) for a complete method to find the total area of the cross section M1 (dep M2) for a complete method for the volume of the prism A1 cao OR M1 for a method to find the volume of cuboid, 8 × 8 × 12 M1 for a method to find the volume of the triangular prism M1 (depM2) for a complete method for the volume of the prism A1 cao

PMT

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q10	(a)	Diagram enlarged. Braille only: measurements given in the text.	M1 for $12.7^2 + 8.3^2$ or 161.29 + 68.89 or 230.18 M1 for $\sqrt{161.29 + 68.89}$ or $\sqrt{230.18}$ A1 for 15.2 or 15.17(168…)
Q10	(b)	Diagram enlarged. Braille only – measurements given in the text.	M1 for $\sin 40° = \frac{DE}{25}$ oe M1 for 25 sin 40° A1 for 16.1 or 16.06(969…)
Q13	(a)	Wording added 'There are four spaces to fill.'	B2 for all values correct (B1 for any one value correct)
Q13	(b)	Grid enlarged.	M1 ft for at least 6 points plotted correctly from their table A1 cao for correct curve drawn from (–4, –6) to (3, –6)
Q16		Diagram enlarged. Dot added at centre.	M1 for angle ABC = 180 – 67 (= 113) A1 cao OR M1 for obtuse angle AOC = 2 × 67 (= 134) A1 cao

PMT

PAPER: 5MB3H_01			
Question		**Modification**	**Notes**
Q18		Diagram enlarged.	M1 for correct use of sine rule to find *AB* or *BC*, eg $\frac{6}{\sin 25} = \frac{AB}{\sin 120}$ M1 for (*AB* =) $\frac{6}{\sin 25} \times \sin 120$ or (*BC* =) $\frac{6}{\sin 25} \times \sin 35$ M1(dep M2) for method to find area, eg 0.5×6×*AB*sin35 or 0.5×6×*BC* sin 120 A1 for answer in the range 21.1 – 21.2 OR (with perpendicular from *A* meeting *BC* extended at a point *X*) M1 for method to find *AX* M1 for method to find *BX* M1(dep M2) for 0.5 × "*AX*" × "*BC*" A1 for answer in the range 21.1 – 21.2
Q21		Diagram enlarged.	M1 for correct vector for $\overrightarrow{OY}$, eg $(\overrightarrow{OY}) = \overrightarrow{OA} + \overrightarrow{AY}$, may include terms in **a** and **b,** eg. 6**a** + $\overrightarrow{AY}$, 4**b** + 4**a** – **b** + $\overrightarrow{PY}$ M1 for $(\overrightarrow{AY} =) \frac{2}{3}\vec{A}P$ or $(\overrightarrow{PY} =) \frac{1}{3}\overrightarrow{PA}$ M1 for correct expression for $\overrightarrow{OY}$ in terms of **a** and **b,** eg $\frac{14}{3}$**a** + 2**b** C1 for $\frac{2}{3}$(7**a** + 3**b)** and "$\overrightarrow{OY}$ is parallel to the vector 7**a** +3**b" oe**

PMT

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London WC2R 0RL

Question 21

Relatively few students achieved any marks. Those that found a correct expression for *DQ* or *EQ* were usually able to write down a correct expression for *PQ* and simplify it to ½**b**. Mistakes were often made with the direction signs of the vectors and some students used ½**a** for the vector *EQ*. Those students obtaining all 4 marks generally made a very clear final statement. Some statements were insufficient with students mentioning "gradient" or "both have **b**" rather than stating that one vector was a multiple of the other.

Question 6

Many students struggled with this question, and for a number of reasons. Some clearly did not understand that the vectors needed to be multiplied by the values, and some of those who did were unable to. Many also struggled with the negative combined with a subtraction on the second vector whilst others thought the vector with the multiple was a mixed number and proceeded to 'convert' it to a top heavy fractions. There were also a number of students who lost the accuracy mark due to incorrect notation, giving their answer as a fraction.

Question 18

There were many students who gained some credit for their answers to this question on vectors and a significant number of answers were fully correct and concisely presented. Of those students who gained some marks but who were not successful in getting a correct expression for $\overrightarrow{DE}$, many were awarded 2 marks for a correct expression for either $\overrightarrow{OD}$, $\overrightarrow{DB}$ or $\overrightarrow{BE}$ in terms of **a** and **b**. Less successful attempts at answering the question often involved the incorrect use of the ratios given, for example quarters and thirds were often seen instead of fifths in expressions for $\overrightarrow{OD}$, $\overrightarrow{DB}$ and $\overrightarrow{BE}$.

Question 13

This question was successfully answered by just under a half of all students who took this paper. Nearly all students scored a mark for finding 3**a** as a column vector. However, students often failed to write their responses in a clear, logical manner. Few students manipulated the equation to get $-2\mathbf{b} = \begin{pmatrix} 8 \\ 17 \end{pmatrix} - 3\mathbf{a}$ or $2\mathbf{b} = 3\mathbf{a} - \begin{pmatrix} 8 \\ 17 \end{pmatrix}$ before substituting for **a**. This might have led to a clearer approach and the avoidance of errors with signs. As it was, examiners were often left not knowing whether the student was trying to find the value of $-2\mathbf{b}$ or $2\mathbf{b}$ with a consequent loss of marks for those students who did not get a fully correct final answer. Students who got one correct component of **b** were given the benefit of the doubt and rewarded with two of the three marks available.

Question 24

Many of the more high attaining candidates gained either partial or full credit for their answers to this question on vectors. The vector algebra seen was generally clearly presented with accurate simplification of expressions found. Most sudents who attempted the question gave a correct expression for either $\overrightarrow{CE}$ and/or $\overrightarrow{CF}$ in terms of $\boldsymbol{a}$ and $\boldsymbol{b}$. Of those candidates who gained marks but who were not successful in proving that *CF* is parallel to *DP*, it was often just the last stage which went awry. Candidates are advised that they should show examiners how their working leads to a proof that the required result is true. In this case, a statement that $\overrightarrow{DP} = 2\,\overrightarrow{CF}$ or equivalent and a conclusion that therefore *CF* and *DP* are parallel was expected.

Question 20

A slightly more straightforward vector problem allowed a good number of candidates to gain 1 or even 2 marks, for working with $\overrightarrow{RO}$ or $\overrightarrow{OR}$ and then applying the ratio to find $\overrightarrow{MO}$ or $\overrightarrow{MR}$. Unfortunately, many selected the incorrect vectors to add to gain the third mark, and it may be worth candidates writing out a clear vector expression such as $\overrightarrow{MT} = \frac{2}{5}\overrightarrow{RO} + \overrightarrow{OT}$ before working in terms of **a** and **b**. The other areas that caused candidates to lose marks were not clearly assigning their vectors, and much like in Question 5, not working correctly with brackets. At this grade the mark scheme is less lenient to incorrect algebra compared to earlier in the paper.

Question 22

Fully correct answers to this question were seen infrequently and there were many students who made no attempt to answer the question. However, a significant proportion of students were able to score at least one mark for writing down a correct vector equation and/or two simultaneous linear equations from the information given. The most success was achieved by students who wrote down and solved the equations $2a + 8b = 13$ and $6a + 2b = 6$ to get $b = \frac{3}{2}$ and $a = \frac{1}{2}$ then used this to write down a

correct expression for b in terms of a. Some students formed simultaneous equations but then used only one equation to find a relationship between a and b. Of those students who did solve the equations to get the value of a and the value of b, some failed to complete the question to get b in terms of a. Approaches using $a\binom{2}{6} + b\binom{8}{2} = k\binom{13}{6}$ where k was treated as a general constant or using $\frac{2a+8b}{6a+2b} = \frac{13}{6}$ as a start were rarely seen.

Question 18

The level of vector algebra was low and in was common to see blank responses for this question, with a majority having no idea how to start the proof. We did see a small number of candidates gaining some marks, normally one or two, typically for a correct expression for $\overrightarrow{EH}$ or $\overrightarrow{BC}$, but very few were able to complete the proof.

Question 19

The first mark was for finding $\overrightarrow{CB}$ or $\overrightarrow{BC}$ in terms of **a** and **b**. the second then required them to apply the ratio of 4 : 5 correctly to this. These marks were awarded relatively often, although some did apply the ratio incorrectly, normally when working from *N* to either *C* or *B*.

The last two marks were for an un-simplified and then a simplified expression for $\overrightarrow{MN}$. Again, a good number gained the third mark, and of these a significant number were able to simplify correctly. This is a place where using a calculator to simplify coefficients might have been a good idea. This step normally involves working with fractions and negatives, and most students attempted it manually.

In part (b) students had to recognise that $\overrightarrow{MN}$ was not a multiple of $\overrightarrow{OB}$. It was clear that many did actually understand that point, but lots struggled to articulate themselves accurately and so lost the communication mark.

Question 8

Many students were able to score at least one mark, and often two, in this question showing some understanding of vector arithmetic. Some students wrote out $2\begin{pmatrix}3\\2\end{pmatrix}+\begin{pmatrix}-1\\4\end{pmatrix}$ but were unable to proceed any further and pick up any marks. Some students tried to square the vector **a** rather than multiply it by 2 giving $\begin{pmatrix}9\\4\end{pmatrix}+\begin{pmatrix}-1\\4\end{pmatrix}$. Students that gained 2 marks often did so for writing the vector $\begin{pmatrix}5\\8\end{pmatrix}$ although there were differing levels of success at drawing it.

Unconventional notation was condoned for the method marks with many students missing out the brackets or even writing their answer as a fraction. The coordinate (5, 8) achieved M1 whereas just plotting the point (5, 8) without identification was M0. Most often the last mark was deducted for missing the arrow on the vector. When drawing vectors it is essential to indicate direction. Some students attempted to find the vector 2**a** + **b** by drawing but this method was often less successful. A number of these students had marks deducted as they only drew the vectors 2**a** and **b** without drawing the diagonal line of 2**a** + **b**, ie from start to finish, unfortunately without this diagonal line only one mark was awarded.