Chhetri Academy

chhetriacademy.com

GCSE MATHS

Statistics

Edexcel 1MA1 | Subtopic collection Topic 59

Histograms

Higher tier

Answers and examiner comments

52	2003-2025	Higher
question groups	available papers	tier

Using this booklet

1. Answer numbers match the companion question booklet.
2. Follow the original question number on marking tables.
3. Examiner comments are included where available.

Name _____ **Date** _____

Some entries use full source marking material in an appendix. Any unavailable mark schemes are identified beside the relevant answer.

Original question 19

The full source mark scheme begins on page 83. Use the original paper code, tier and question number shown above.

Original question 13

No matching mark scheme was available in the supplied archive for this question.

Original question 14

The full source mark scheme begins on page 137. Use the original paper code, tier and question number shown above.

Original question 17

No matching mark scheme was available in the supplied archive for this question.

Original question 19

No matching mark scheme was available in the supplied archive for this question.

22			$2c + 4r$	2	B2 for 2*c* +4*r* oe [B1 for 2*c* or 4*r* oe seen] Ignore any Left Hand Side = $2c + 4r$ {Note: ignore units or use of 'p'}

27	(a)		30 60	2	B1 cao B1 cao
	(b)		fd = 1.5 (ht 3cm) fd = 0.5 (ht 1cm)	2	M1 for at least one correct frequency density calculated for the last 2 bars (could be implied by one correct bar) or 1 sq = 5 cars A1 cao

Question	Working	Answer	Mark	Notes
25 (a)		12, 6	2	M1 for frequency density calculation (implied by one answer), or $1cm^2$=2 (trains), or fd=0.5 or $8cm^2$=16 A1 both 12 and 6
(b)	Bar of height 5cm (5-10) Bar of height 1cm (30-50)	Height 5cm Height 1cm	2	M1 for frequency density calculation (implied by one correct bar) or $1cm^2$=2 (trains) or fd=0.5 A1 for bar of height 5cm (5 to 10) AND for bar of height 1 cm (30 to 50) $8cm^2$=16

Question	Working	Answer	Mark	Notes
24 (a)	$\frac{8}{5}$ = 1.6	*Bar of height 3cm drawn*	2	M1 for 2cm² = 1 pupil oe **or** calculation of fd = 1.6 **or** bar of area 12 cm² but not correct shape A1 cao
(b)	6 + 8 + 6 + 5	25	2	B2 for 25 (B1 for frequency of 5 for number of students who watched between 20 and 30 hours)

Question	Working	Answer	Mark	Notes
10 (a)	$5\times16=80$ $12.5\times18=225$ $17.5\times10=175$ $27.5\times6=165$ $645\div50=12.9$ or $5.5\times16=88$ $13\times18=234$ $18\times10=180$ $28\times6=168$ $670\div50=13.4$	12.9	4	M1 for fx consistently within interval including ends (allow 1 error) M1 consistently using appropriate midpoints M1 (dep on first M1) for $\sum fx \div \sum f$ A1 for 12.9 or 13.4
(b)	$\frac{6}{50}\times\frac{5}{49}=\frac{30}{2450}$	$\frac{3}{245}$	2	M1 for $\frac{6}{50}\times\frac{5}{49}$ A1 for $\frac{3}{245}$ oe If M0A0, SC B1 for $\frac{9}{625}$ oe
(c)	$0\le d<10$ fd 1.6 $10\le d<15$ fd 3.6 $15\le d<20$ fd 2 $20\le d<35$ fd 0.4	Correct histogram	3	B2 for 4 correct histogram bars (± ½` square) (B1 for 2 or 3 histogram bars of different widths correct) B1 for frequency density label or key and consistent scaling SC if B0 then M1 for clear attempt to use frequency density or area

5MB1H_01
November 2010

5MB1H/01

14	(a)		5 15	2	B1 cao B1 cao
	(b)		fd = 2 (ht 4cm) fd = 2.5 (ht 5cm)	2	B2 for 2 correct bars B1 for 1 correct bar If B0 is scored then you can award M1 at least one correct frequency density calculated for the missing bars Or 1 sq = 2.5 plants oe

24	(a)		84, 60	2	B1 for 84 B1 for 60
	(b)			2	B1 for bar with width 160-180 and height 2cm (± 1 mm) B1 for bar with width 180-210 and height 6mm (± 1 mm)

1380_3H

Original question 11

The full source mark scheme begins on page 188. Use the original paper code, tier and question number shown above.

21	(a)		34, 12	2	M1 for frequency = fd × column width, can be implied by one frequency correct **or** fd correctly marked on vertical axis (1cm = 4 units) **or** identifying 1 cm^2 as frequency of 4 oe A1 34 and 12 both correct
	(b)		Bars of height 6 cm and 4.5 cm	2	B1 for bar of height 6 cm B1 for bar of height 4.5 cm

1380_4H

Question		Working	Answer	Mark	Notes
12	(a)		(20), (30), 45, 60, (48)	2	M1 for frequency = fd × column width, can be implied by 1 frequency correct OR fd correctly marked on vertical axis 2cm=1unit OR identifying 1 cm^2 as frequency of 5oe A1 45 and 60 both correct
	(b)		histogram bars	2	B2 for 2 correct histogram bars; heights at 6cm and 2.4cm (B1 1 correct bar)
	(c)	Area method: Total area 40.6 cm^2 For median: ÷2 = 20.3 0 to 40 is 19 cm^2 median lies 41–43 OR Proportionality method: Total 203 ÷ 2 = 101.5; 0 to 40 is 95 40–60: 6.5÷60×20=2.16 Median is 40+2.16 = 42.16 OR 204÷2=102; 0 to 40 is 95 40-60: 7÷60×20=2.3… Median is 40+2.33=42.33	41 – 43	2	Area method: M1 ft for calculation of total area and division by 2 (eg 40.6÷2 or 20.3) A1ft answer 41–43 OR Proportionality method: M1 ft for 203÷2=101.5 and 6.5÷60×20=2.16 or 204÷2=102 and 7÷60×20=2.33… A1 ft answer 41–43

5MB1H_01

Question		Working	Answer	Mark	Notes
12	(a)		Histogram	3	B3 for fully correct histogram (overlay) (B2 for 4 correct blocks B1 for 3 correct blocks) SC B1 for correct key, eg $1cm^2$=5(men) or correct freq÷ class interval for at least three frequencies
	(b)	$2x + 1.4x + x + 0.4x + 0.2x$	$5x$	2	M1 for finding total area of histogram, eg $20(cm^2)$, or fd for 10 to 20 ($0.1x$ oe) or area of one block in terms of x A1 for $5x$ oe

Original question 22

The full source mark scheme begins on page 202. Use the original paper code, tier and question number shown above.

Question		Working	Answer	Mark	Notes
13		$\frac{17.24}{57} \times 285 = 86.2$ $\frac{86.2}{285} \times 285 = 86.2$ $\frac{431}{1425} \times 285 = 86.2$	86 or 87 .	3	M1 for an attempt to relate total area to total frequency. Eg. using a fd scale of **1cm = 0.2** 285 ÷ [21+9.8+11.2+10+5 (=57)] (= 5) At least 3 of these 5 frequency values must be correct M1 for ('2.24'+10+5) × '5' A1 for 86 or 87 (accept 86.2) OR M1 for an attempt to relate total area to total frequency. Eg. using a fd scale of **1cm = 1** 285 ÷ [105+49+56+50+25 (=285)] (= 1) At least 3 of these 5 frequency values must be correct M1 for ('11.2'+50+25) × '1' A1 for 86 or 87 (accept 86.2) [SC: B1 can be awarded for correct labelling of fd axis using the scale of 1cm = 1 if M0 scored] OR M1 for an attempt to relate total area to total frequency. Eg. using a fd scale of **1cm = 5** 285 ÷ [525+245+280+250+125 (=1425)] (= 0.2) At least 3 of these 5 frequency values must be correct M1 for ('56'+250+125) × '0.2' A1 for 86 or 87 (accept 86.2)

5MB1H_01					
Question		Working	Answer	Mark	Notes
		(2.24+10+5) × 5 = for 17.24 × 5 = 86.2			OR M1 for 1 cm^2 = 5 farms M1 for ('2.24'+10+5) × 5 A1 for 86 or 87 (accept 86.2)

5MB1H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
14		$\frac{3}{6} \times \frac{2}{5} + \frac{2}{6} \times \frac{1}{5} = 8/30$ $1 - \frac{8}{30} = \frac{22}{30}$ OR $2 \times \frac{3}{6} \times \frac{2}{5} + 2 \times \frac{3}{6} \times \frac{1}{5} + 2 \times \frac{2}{6} \times \frac{1}{5}$ OR  	$\frac{22}{30} = \frac{11}{15}$	4	M1 for $\frac{3}{6} \times \frac{2}{5}$ or $\frac{2}{6} \times \frac{1}{5}$ M1 for $\frac{3}{6} \times \frac{2}{5} + \frac{2}{6} \times \frac{1}{5}$ M1 for $1 - (\frac{3}{6} \times \frac{2}{5} + \frac{2}{6} \times \frac{1}{5})$ A1 for $\frac{22}{30}$ oe OR M1 for $\frac{3}{6} \times \frac{2}{5}$ or $\frac{3}{6} \times \frac{1}{5}$ or $\frac{2}{6} \times \frac{1}{5}$ M1 for $\frac{3}{6} \times \frac{2}{5} + \frac{3}{6} \times \frac{1}{5} + \frac{2}{6} \times \frac{1}{5}$ M1 for $2 \times (\frac{3}{6} \times \frac{2}{5} + \frac{3}{6} \times \frac{1}{5} + \frac{2}{6} \times \frac{1}{5})$ A1 for $\frac{22}{30}$ oe OR M2 for $\frac{3}{6} \times \frac{3}{5}$ or $\frac{2}{6} \times \frac{4}{5}$ or $\frac{1}{6} \times \frac{5}{5}$ M1 for $\frac{3}{6} \times \frac{3}{5} + \frac{2}{6} \times \frac{4}{5} + \frac{1}{6} \times \frac{5}{5}$ A1 for $\frac{22}{30}$ oe

		OR A^1A^2 A^1A^3 $\mathbf{A^1B^1}$ $\mathbf{A^1B^2}$ $\mathbf{A^1C}$ A^2A^1 A^2A^3 $\mathbf{A^2B^1}$ $\mathbf{A^2B^2}$ $\mathbf{A^2C}$ A^3A^1 A^3A^{32} $\mathbf{A^3B^1}$ $\mathbf{A^3B^2}$ $\mathbf{A^3C}$ $\mathbf{B^1A^1}$ $\mathbf{B^1A^2}$ $\mathbf{B^1A^3}$ B^1B^2 $\mathbf{B^1C}$ $\mathbf{B^2A^1}$ $\mathbf{B^2A^2}$ $\mathbf{B^2A^3}$ B^2B^1 $\mathbf{B^2C}$ $\mathbf{CA^1}$ $\mathbf{CA^2}$ $\mathbf{CA^3}$ $\mathbf{CB^2}$ $\mathbf{CB^2}$			OR M2 for diagram showing all 30 possible outcomes (M1 for diagram showing 15 possible outcomes. ie without any repeats) M1 (dep) for identifying all outcomes in their diagram with different letters (underlining, circling, etc.) Or 30 – all outcomes where the letters are the same A1 for $\frac{22}{30}$ oe **Alternative scheme for replacement** M1 for $\frac{3}{6}\times\frac{3}{6}$ or $\frac{2}{6}\times\frac{2}{6}$ or $\frac{1}{6}\times\frac{1}{6}$ M1 for $1-(\frac{3}{6}\times\frac{3}{6}+\frac{2}{6}\times\frac{2}{6}+\frac{1}{6}\times\frac{1}{6})$ OR M1 for $\frac{3}{6}\times\frac{2}{6}$ or $\frac{3}{6}\times\frac{1}{6}$ or $\frac{2}{6}\times\frac{1}{6}$ M1 for $2\times(\frac{3}{6}\times\frac{2}{6}+\frac{3}{6}\times\frac{1}{6}+\frac{2}{6}\times\frac{1}{6})$

					OR M1 for a diagram showing all 36 'possible' outcomes A1 for $\frac{22}{36}$ (= 0.6111…) **Special cases:** B2 for $\frac{22}{36}$ (= 0.6111…) or $\frac{28}{36}$ (= 0.777..) or $\frac{16}{30}$ (= 0.533..) if M0 scored

Question		Working	Answer	Mark	Notes
24		<table><tr><th>Height h m</th><th>Freq</th><th>FD</th></tr><tr><td>$0 < h \leq 2$</td><td>7</td><td>3.5</td></tr><tr><td>$2 < h \leq 4$</td><td>14</td><td>7</td></tr><tr><td>$4 < h \leq 8$</td><td>18</td><td>4.5</td></tr><tr><td>$8 < h \leq 16$</td><td>24</td><td>3</td></tr><tr><td>$16 < h \leq 20$</td><td>10</td><td>2.5</td></tr></table>	3	3	B3 fully correct histogram with horizontal axis correctly scaled (B2 for 4 correct blocks **or** 5 correct blocks with incorrect or no scale) (B1 for 2 correct blocks of different widths **or** any 3 correct blocks) SC : B1 for key, eg. 1 cm^2 = 2 (trees) **or** correct values shown for (freq ÷ class interval) for at least 3 frequencies (3.5, 7, 4.5, 3, 2.5)

Question		Working	Answer	Mark	Notes
15	(a)	f.d. = 2, 12, 10, 3 e.g. each square = 4 people 1:15:10:1.5 e.g. each square = 2 people 2:30:20:3	histogram	3	B2 for all 4 bars fully correct with areas in the ratio 1:15:10:1.5 or 2:30:20:3 oe (B1 for 2 or 3 bars drawn in the correct ratio to each other) B1 for labelling the vertical axis f.d and consistent scaling or for a key
	(b)	Monday 40 + 6 = 46 more than 7 hours Tuesday One square = 12 ÷ 6 = 2 2 × 3.5 × 7 = 49 more than 7 hours	No with reason	2	M1 for a valid method to work out the frequency for more than 7 hours on Tuesday, (12 ÷ 6) × 3.5 × 7 or 2 × 3.5 × 7 or 7 × 7 or 2 × 24.5 or 49 A1 No with 46 and 49 seen

5MB1H_01

Question		Working	Answer	Mark	Notes
21		Total area = (0.12 × 40) + (0.36 × 20) + (0.7 × 20) + (0.56 × 20) + (0.18× 40) = 44.4 Area (140 < w < 200) = (0.36 × 20) + (0.7 × 20) + (0.56 × 20) = 32.4 32.4 ÷ 44.4	0.73	4	M1 for a method to find the frequency **or** the area of any one block M1 for a method (with correct values) to find total area of all blocks **or** 44.4 **or** 1110 **or** a correct method (with correct values) to find total area of middle 3 blocks **or** 32.4 **or** 810 M1 (dep on M2) for a correct method to find required proportion (could lead to a decimal or a percentage or a fraction) A1 for answer which rounds to 0.73 or 73% or $\frac{27}{37}$ or equivalent fraction

1MA0_2H

13	(a)		10 and 18	2	B2 Two correct values (B1 one correct value)
	(b)		Bars at heights 3cm and 2cm	2	B2 for two correct bars (B1 for one correct bar)

24		12 ÷ 10 = 1.2 15 ÷ 5 = 3 13 ÷ 5 = 2.6 18 ÷ 10 = 1.8 3 ÷ 15 = 0.2	Histogram	3	B3 for fully correct histogram (B2 for 4 correct blocks) (B1 for 3 correct blocks) (If B0, SC B1 for correct key eg $1cm^2$ = 2 (calls) Or frequency ÷ class interval for at least 3 frequencies) **NB** Apply the same mark scheme if a different frequency density is used.

PAPER: 1MA0_1H

Question		Working	Answer	Mark	Notes
14	(a)	0.016 × 500 = 8 0.03 × 500 = 15 0.08 × 250 = 20 0.096 × 250 = 24 0.026 × 1000 = 26	(8), 15, 20, 24, 26	2	M1 for correct calculation to find one frequency e.g. 0.03 × 500 or 0.08 × 250 or 0.096 × 250 or 0.026 × 1000 or for one frequency correct or establishing that 1 cm^2 = 2.5 fish A1 for all frequencies correct
	(b)	0.026 × 500 = 13 8 + 15 + 20 + 24 + 26 = 93	13 : 93	2	M1 ft for a complete correct method to find the number of fish over 2000 g ie 0.026 × 500 (=13) or '26'÷2 A1ft for 13 : 93 or '0.026 × 500' : 'total for all their fish' or '26'÷2 : 'total for all their fish' SCB1 for 93:13 given as the answer
	(c)	47th item needed	1292	2	M1 for a complete correct method to divide the area of the histogram into two equal parts **OR** for a complete correct method to interpolate for the 47th value A1 for answer in range 1290 to 1300

PAPER: 5MB1H_01

27	12×20 + 10.8×10 + 7×15 + 5×15 + 1.8×30 + 0.6×30 =240+108+105+75 +54+18 =528+72=600	12%	3	M1 for attempt to work out total area (eg =600) or area greater than 60 (eg =72) by using fd or counting squares M1 (dep) for $\frac{'72'}{'600'} \times 100$ oe (=12) A1 cao (must have % otherwise 2 marks)

PAPER: 1MA0_2H				
Question	**Working**	**Answer**	**Mark**	**Notes**
*28		Proof	3	M1 for one pair of equal angles or sides with reason M1 for second pair of equal angles or sides with reason C1 for proof completed correctly with full reasons and reason for congruence Acceptable reasons: *AD* common (oe eg both same) Angle *BAD* = angle *CDA* (angles in a semicircle are 90°) Angle *ABO* = angle *DCA* (angles in the same segment are equal) Triangle *ABD* and triangle *DCA* are congruent - ASA **OR** *BD* = *CA* (diameters of the circle) Angle *BAD* = angle *CDA* (angles in a semicircle are 90°.) *AD* common Triangle *ABD* and triangle *DCA* are congruent - RHS **OR** *BD* = *CA* (diameters of the circle) *AD* is common Angle *ADB* = angle *CAD* (base angles of an isosceles triangle are equal.) Triangle *ABD* and triangle *DCA* are congruent - SAS

Question 6:

	F	S	G	
W	12	55		96
M	**7**	**18**	9	**34**
	19	73		130

	F	S	G	
W	12	55	**29**	96
M			9	
	19	73	**38**	130

Modifications to the mark scheme for Modified Large Print (MLP) papers.

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 1MA0_2H			
Question		**Modification**	**Notes**
Q4		Model as well as diagram of prism provided. Question reversed.	Standard mark scheme
Q9		MLP – *a* changed to *x*	Standard mark scheme
Q11	(a) (c)	MLP – *x* changed to *y* MLP - *a* changed to *e*	 B2 for $3e^{-4}f^6$ **or** $\frac{3f^6}{e^4}$ (B1 for any two from 3, e^{-4} or $\frac{1}{e^4}$, f^6 in a product)
Q14	 (c)	Some frequencies changed: 6, 8, 14, 20, 2 Grid – *y* axis labelled in 4s up to 24. 2cm for 2. *x*- axis 2cm for 2	Standard mark scheme Standard mark scheme

PAPER: 1MA0_2H			
Question		**Modification**	**Notes**
Q15		Braille – information given about the diagram.	Standard mark scheme
Q17		Grid *x* 1.5 with 1.5 cm squares. For 36pt, squares will be close to 2cm.	Standard mark scheme
Q18		Braille – information given about the diagram.	Standard mark scheme
Q21	(b)	*a* changed to *f*	M1 for correct processes to isolate terms in *m* from other terms M1 for taking *m* out as a common factor A1 for $m = \frac{g-5}{f+3}$ **or** $m = \frac{5-g}{-f-3}$
Q22		MLP and Braille – diagram labelled clockwise from the top left. ABCD. *x* changed to *y*. Information given about the diagram.	Standard mark scheme
Q24		Model as well as diagram provided.	Standard mark scheme
Q26		MLP and Braille information given about the diagram.	Standard mark scheme
Q27		Diagram: 20-30 bar brought down to 9. 60-90 bar brought up to 2. 90-120 bar brought up to 1. 2cm grid.	Standard mark scheme

15			55 - 56	3	M1 for attempt to find 0.25×total area, eg 0.25×(3.6+14+5.6+7.2+1.4) (=7.95) condone one error M1 for ('7.95' – '3.6') ÷ 7 (=0.621…) A1 for 55 – 56 OR M1 for for attempt to find 0.25×total squares, eg 0.25×(90+350+140+180+35) (=198.75) condone one error M1 for ('198.75 – '90') ÷ 35 (=3.107…) A1 for 55 – 56

5MB1H_01

Paper_5MB1H_01

Question		Working	Answer	Mark	Notes
16	(a)	16 ÷ 20 = 0.8 27 ÷10 = 2.7 36 ÷ 15 = 2.4 6 ÷ 5 = 1.2	Correct histogram	3	M1 for recognising and showing evidence of using frequency density, eg at least 2 correct frequency densities or a key . A1 for all bars in correct proportions OR for one error with bars and correct labelling and scaling of axes A1 for fully correct histogram including axes labelled and scaled.
	(b)		$\frac{18}{85}$	3	M1 for using 40 cm on horizontal scale and using a correct method to convert area of bar to the right of 40 cm to frequency, eg 2.4 × 5. M1 (dep) for '12' + 6 A1 for $\frac{18}{85}$ oe OR M1 for $\frac{36}{3}$ M1 (dep) for '12' + 6 A1 for $\frac{18}{85}$ oe

Modifications to the mark scheme for Modified Large Print (MLP) papers.

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 5MB1H_01			
Question		**Modification**	**Notes**
Q02		2cm grid crosses, cha nged to solid circles. Right axis labeled	
Q07 Q09		Box for key put top left Box plot: median changed to 47.5, upper quartile changed to 52.5	
Q09	(C)	52kg changed to 52.5 kg	
Q10	(C)	Frequencies changed: 5, 20, 15, 15, 5 grid x axis – 2 cm for 10, y axis 2cm for 5	
Q13		2 cm grid, line moved parallel to original graph line to go through (0,20) (50, 80) (75, 110)	
Q16		Frequency column: 27 changed to 28 2 cm grid Number of weather stations changed from 85 to 86	

18	(a)	$5 \times 8 = 40$ $12.5 \times 15 = 187.5$ $17.5 \times 11 = 192.5$ $25 \times 10 = 250$ $40 \times 6 = 240$ $910 \div 50 = 18.2$	18.2	4	M1 for *fx* consistently within interval including ends (allow 1 error) M1 consistently using appropriate midpoints M1 (dep on first M1) for $\Sigma fx \div \Sigma f$ A1 for 18.2 cao
	(b)	$0 \leq t < 10$ fd 0.8 $10 \leq t < 15$ fd 3 $15 \leq t < 20$ fd 2.2 $20 \leq t < 30$ fd 1 $30 \leq t < 50$ fd 0.3	Correct histogram	3	B3 fully correct histogram with vertical axis correctly scaled. (B2 for 4 correct blocks or 5 correct blocks with incorrect or no scale) (B1 for 2 correct blocks of different widths or any 3 correct blocks or correct FD values for at least 3 frequencies) eg fd of 0.8, 3, 2.2, 1, 0.3

16		Table: 21 changed to 20 Top line 84 changed to 83 Grid: 2 cm	

PAPER: 1MA0_2H

Question		Working	Answer	Mark	Notes
26	(a)		(4), 9, 8, 10, 12	2	M1 for correct calculation to find one frequency e.g. 0.9×10 or 1.6×5 or 1×10 or 0.8×15 or for one frequency correct or shows that 1 $cm^2 = 1$ A1 for all frequencies correct
	(b)		$\frac{8}{43}$	2	M1 for 8 (people) or $\frac{2}{3}$ of "12" A1ft for 8 out of 43 stated as a percentage or fraction or decimal
	(c)		26000	2	M1 ft for finding the interval in which the "21.5th" or "22nd" value lies or 26 or 25.5 A1 for 26000 or 25500

13	F 90 126 144 120 60 54	345	5	M1 for use of F = FD × Int width A1 for any 3 Fs correct M1 for $\frac{60}{100}\times(90+126+144+120)$ $(=288)$ or $\frac{60}{100}\times 480$ (=288) M1 $\frac{1}{2}\times"(60+54)"(=57)$ or $\frac{1}{2}\times 114$ (=57) A1 cao

Modifications to the mark scheme for Modified Large Print (MLP) papers.

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 5MB1H_01			
Question		**Modification**	**Notes**
Q2	(a)	2 graphs provided, one for 'sandy soil' and one for 'clay soil'. Sandy soil graph line goes through (10, 25) and (20, 45) Clay soil graph line goes through (15, 20) and (25, 30) Points are moved to fit in with the new graph lines. Parts (b) and (c) have been reversed and relabelled.	B1 description eg Taller trees are older. Accept positive correlation.
Q2	(b)	2 graphs provided, one for 'sandy soil' and one for 'clay soil'. Sandy soil graph line goes through (10, 25) and (20, 45) Clay soil graph line goes through (15, 20) and (25, 30) Points are moved to fit in with the new graph lines. Parts (b) and (c) have been reversed and relabelled.	B1 19 – 21
Q2	(c)	2 graphs provided, one for 'sandy soil' and one for 'clay soil'. Sandy soil graph line goes through (10, 25) and (20, 45) Clay soil graph line goes through (15, 20) and (25, 30) Points are moved to fit in with the new graph lines. Parts (b) and (c) have been reversed and relabelled. 18 years has been changed to 15 years.	M1 for evidence of taking readings at two points from Sandy line, or increase excluding start eg 24÷10, 14÷5 A1 for answer 1.8 to 2.2 within given tolerance and adjusted appropriately.

PAPER: 5MB1H_01			
Question		**Modification**	**Notes**
Q2	(d)	2 graphs provided, one for 'sandy soil' and one for 'clay soil'. Sandy soil graph line goes through (10, 25) and (20, 45) Clay soil graph line goes through (15, 20) and (25, 30) Points are moved to fit in with the new graph lines. Parts (b) and (c) have been reversed and relabelled.	B2 for a complete explanation e.g.Trees grow at approximately twice the rate on sandy soil (B1 for a partial explanation e.g. Trees grow faster on sandy soil)
Q3	(a)	Question for questionnaire moved to the Diagram Book.	B1 e.g. people coming from the market are more likely to have used it.
Q3	(b)	Question for questionnaire moved to the Diagram Book.	B1 e.g. it's a leading question, "cheap" not defined, etc.
Q3	(c)	Question for questionnaire moved to the Diagram Book.	B1 for a suitable question e.g. How far have you travelled to get to the market? B1 for a set of at least 3 exhaustive, non-overlapping boxes with units, with no inequality symbols.
Q6	(a)	Bottom line has been drawn on the stem and leaf diagram.	B2 for a fully correct ordered stem and leaf diagram [B1 for a complete unordered diagram or for an ordered diagram with at most 2 errors] B1 (indep) key
Q6	(b)	Bottom line has been drawn on the stem and leaf diagram.	M1 for 36 ÷ (5 + 4) (=4) A1 cao

PAPER: 5MB1H_01			
Question		**Modification**	**Notes**
Q7	(a)	Frequency column on the table has been extended to allow room for working.	B1 cao
Q7	(b)	Frequency column on the table has been extended to allow room for working.	M1 for Σ(number of books × frequency) (=60) M1 for "60" ÷ "25" A1 cao SC B2 for an answer of 2.48
Q7	(c)	Frequency column on the table has been extended to allow room for working.	M1 for 15 × 4.4 (=66) M1 for a complete method eg ("60" + "66") ÷ (15 + "25") A1 cao

Original question 15

The full source mark scheme begins on page 222. Use the original paper code, tier and question number shown above.

23		$0<t\leq 5$ fd = 8 ÷ 5 = 1.6 $5<t\leq 15$ fd = 32 ÷ 10 = 3.2 $15<t\leq 30$ fd = 36 ÷ 15 = 2.4 $30<t\leq 40$ fd = 18 ÷ 10 = 1.8 $40<t\leq 60$ fd = 6 ÷ 20 = 0.3	Correct histogram	3	B3 for a fully correct histogram with vertical axis correctly scaled or with a key, eg. 2 cm^2 = 1 (B2 for at least 4 correct blocks with or without a scale or a key OR for all five fd correct) (B1 for 2 correct blocks of different widths or for at least three correct fd values)

PAPER: 1MA0_2H

PAPER: 5MB1H/01

Question		Working	Answer	Mark	Notes
15	(a)	12 ÷ 20 = 0.6 20 ÷ 10 = 2 17 ÷ 10 = 1.7 6 ÷ 15 = 0.4	Correct histogram	3	B3 for fully correct histogram (B2 for 3 correct blocks or all 4 frequency ÷ class interval, *y*-axis labelled and 1 correct block) (B1 for 2 correct blocks of different widths or for correct key eg 1cm^2 = 1 egg or for frequency ÷ class interval for at least 3 frequencies) Due to scale accept to within 1mm on plotting
	(b)		19	3	M1 for splitting one of relevant rectangles or for $\frac{7}{10}\times 20$ (=14) or $\frac{3}{10}\times 17$ (=5.1) M1 for (area of 53 – 63 interval) ÷ (total area) × 55 or for "14" + "5.1" A1 for 19

23	(i)		38 30	3	M1 for evidence of frequency density calculation, eg 6 on the frequency density axis for the height of the first column or 5 is 1 cm² can be implied by 30 as the second missing frequency A1 for 38 and 30
	(ii)		30–50 bar height 0.8 cm		B1 for 30 – 50 bar of height 0.8 cm

15	e.g. (0.8 + 0.8 + 0.8) + (3.6 + 3.6) + 7.2 + (4.4 + 4.4) = 25.6 cm^2 = 320 lizards So, 1 cm^2 = 12.5 lizards 7.2 + 4.4 + 4.4 = 16 16 × 12.5 = 200	200	3	M1 for equating area to frequency, eg 1 cm^2 = 12.5 lizards M1 for (7.2 + 4.4 + 4.4) × '12.5' A1 cao OR M2 for $\frac{16}{25.6} \times 320$ oe A1 cao

23		0.94 or 94% or $\frac{76}{81}$	3	M1 for method to work out total area eg 1.3×10 + 3.2×5 + 3.6×5 + 2.4×10 + 0.5×20 (=81) or area up to 100 grams eg 1.3×10 + 3.2×5 + 3.6×5 + 2.4×10 + 0.5×10 (=76) (In either case allow one error in reading a bar height) M1 for 1 – ((0.5×10)/"81") oe or for method to work out the total area and the area up to 100 grams (In both cases allow one error in reading a bar height) A1 for answer in range 0.938 to 0.94 or 93.8% to 94% or $\frac{76}{81}$ oe

Paper: 1MA0/2H

13		7	P1 P1 A1	for correct process to find any frequency, eg. “1.1” × 10 (= 11) or “2.8” × 10 (= 28) or “2.3” × 20 (= 46) or “1.4” × 20 (= 28) or “1.4” × 10 (= 14) or “0.7” × 30 (= 21) or for a correct process to find the total area and an area of any block, eg. using 1 cm^2 = 1 unit of area to get 53.6 and one of 4.4, 11.2, 18.4, 11.2, 5.6, 8.4 (dep P1) for complete process to find 20% of (“1.4” × 10 + “0.7” × 30), eg. $\frac{20}{100} \times "35"$ or $\frac{"5.6"+"8.4"}{"53.6"} \times 134 \times \frac{20}{100}$ cao

Question	Working	Answer	Mark	Notes
17 (a)	1.5, 6, 10.2, 7.2, 1.2	Histogram drawn	C1	for 2 correct bars of different widths or at least 3 correct frequency densities.
			C1	for all bars in correct proportions or 4 correct bars with axes scaled and labelled.
			C1	for fully correct histogram with axes scaled and labelled.
(b)		$\frac{123}{150}$	M1	for a method to find number of students in interval, eg $30 + 51 + 36 + \frac{1}{3} \times 18$ (= 123) or $150 - 15 - \frac{2}{3} \times 18$ (= 123)
			A1	for $\frac{123}{150}$ oe or 0.82 or 82%

Question	Answer	Mark	Mark scheme	Additional guidance
17 (a)	4, 6, 5, 4	M1	for a correct method to find at least 2 frequencies from bars of different widths, eg 10 × 0.4 (=4), 10 × 0.6 (=6), 5 × 1 (=5), 20 × 0.2 (=4)	
		A1	cao	
(b)	10	M1	for $\frac{23+1}{4}$ (=6) **or** $\frac{23}{4}$ (=5.75) could ft from their table in (a)	
		A1	for 10 or 9.375	Be aware of 10 coming from incorrect working ft does not apply to the A1

Paper: 1MA1/3H

Question	Answer	Mark	Mark scheme	Additional guidance
17 (a)	Histogram drawn	B3	for fully correct histogram eg relative heights 6, 3, 4, 2, 2	
		(B2	for 4 correct blocks **or** all 5 frequency ÷ class interval **and** 1 correct block)	
		(B1	for at least 2 correct blocks of different widths **or** for frequency ÷ class interval for at least 3 frequencies)	
(b)	66 to 71	M1	indication of the median in the third interval **or** proportional method shown	Just stating the interval is sufficient for this mark May be implied by the number on the answer line
		A1	ft answer between 66 and 71	Median is at (approx.) 68.75 by a proportional method

Question	Answer	Mark	Mark scheme	Additional guidance
21	210	M1	for method to find total frequency, 60 × 2 (= 120) + 30 × 5 (= 150) + 30 × 9 (= 270) + 15 × 6 (= 90) + 45 × 2 (= 90) or 720 **OR** for method to find the total area, 4 + 5 + 9 + 3 + 3 (= 24 cm^2)	Accept one error in total for the award of the method marks 24 must be from adding areas of bars not heights of bars
		M1	for finding the number of onions less than 60g or greater than 120 g = 120 + 90 + 90 (= 300), **OR** for finding the number of onions between 60g and 120g = 150 + 270 (= 420) **OR** for finding the area under the graph less than 60 or greater than 120 = 4 + 3 + 3 (= 10 cm^2) **OR** for finding the area under the graph between 60 and 120 = 5 + 9 (= 14 cm^2)	14 must be from adding areas of bars not heights of bars
		M1	(dep M2) for $1 - \frac{"300"}{"720"} (= \frac{7}{12})$ oe OR for $\frac{"420"}{"720"} (= \frac{7}{12})$ oe OR for $\frac{"14"}{"24"} (= \frac{7}{12})$ oe	Accept 58.3...%
		A1	cao	

Paper: 1MA1/3H

18	Bar of height 3.2	M1 M1 C1	method to find any frequency eg 1.2 × 2.5 (= 3) or 2 × 2.5 (= 5) or 2.8 × 5 (= 14) or 0.8 × 12.5 (= 10) **or** method to use areas eg 12 × 5 (=60) or 20 × 5 (=100) or 28 × 10 (=280) or 8 × 25 (=200) complete method to find total frequency for the four intervals eg "3" + "5" + "14" + "10" (=32) **or** "60" + "100" + "280" + "200" (=640) cao	Accept equivalent methods proportional to those shown

Paper: 1MA1/2H

17	7.645	P1	for process to use area to find at least one frequency, eg for first frequency $(7.2 - 6.4) \times 10$ (= 8) or $(7.2 - 6.4) \times 5$ (= 4) or $4 \times 5 \times 5$ (= 100)	Frequencies could be written on the graph
		P1	for process to find all frequencies, eg 8, 20, 40, 12 or multiples eg 4, 10, 20, 6 or 100, 250, 500, 150	Marks are for correct processes, one or more frequencies may be incorrect
		P1	(dep P2) for process to estimate mean, eg $((6.8 \times [8]) + (7.4 \times [20]) + (7.8 \times [40]) + (8.1 \times [12])) \div ([8] + [20] + [40] + [12])$	
		A1	for 7.645 (accept 7.65)	Award full marks if a correct answer is seen in working and is then incorrectly rounded.

Paper: 1MA1/3H

Question	Answer	Mark	Mark scheme	Additional guidance
17	30	P1	for process to find one correct frequency, eg. 0.8 × 5 (= 4) or 1.6 × 10 (= 16) or 2.2 × 10 (= 22) or 1.2 × 15 (= 18) **or** to find one correct area eg 5 × 8 (=40) or 10 × 16 (=160) or 10 × 22 (=220) or 15 × 12 (=180)	Accept equivalent methods proportional to those shown.
		P1	for process to find total number of people, eg. “4” + “16” + “22” + “18” (= 60) **or** for process to find total area eg “40” + “160” + “220” + “180” (= 600)	Condone 1 error in reading from the graph for 2nd and 3rd P marks
		P1	for process to find 20% of the total number of people, eg. “60” × 0.2 oe (= 12) **or** for process to find 20% of the total area “600” × 0.2 oe (=120)	
		A1	cao	NB: correct answer without supportive working gets 0 marks

Paper: 1MA1/3H

Question	Answer	Mark	Mark scheme	Additional guidance
17 (a)	histogram drawn	B3	for fully correct histogram, eg. relative heights 90, 96, 44, 8, 6	
		(B2	for 4 correct bars **or** for frequency ÷ class interval for all 5 frequencies and 2 correct bars of different widths)	
		(B1	for 2 correct bars of different widths or for frequency ÷ class interval for at least 3 frequencies)	
(b)	$0.4n$	M1	for finding ratio of heights or widths of bars, eg 5 : 1 or $\frac{1}{5}$, 1 : 2 or $\frac{n}{5}$ oe or $2n$ oe as answer or compares areas of bars, eg 6 and 2.4 or 3 and 1.2 or 150 and 60	Evidence for this mark may be seen on the diagram Any 2 numbers in the ratio 2.5 : 1 score M1
		A1	for $0.4n$ oe	

14	Histogram drawn	B3 (B2 (B1	for fully correct histogram, eg relative heights 1, 5, 6, 1.5 for 3 correct bars **or** for frequency ÷ class interval for at least 3 frequencies and 2 correct bars of different widths) for 2 correct bars of different widths **or** for frequency ÷ class interval for at least 3 frequencies)	Frequency densities are 1, 5, 6, 1.5

16 (a)	Bar of height 2.4 cm (fd = 1.2) drawn for interval 40 – 60 hours	M1	for using area to represent frequency, eg 11.2 (cm^2) represents 28 (students) or for showing correct scale on fd axis, 2 cm = 1 unit or correct process to find the height of bar, eg $\frac{28}{10}$ $(=2.8)$ or $\frac{24}{20}$ $(=1.2)$	May use any unit of area. Can be implied by frequency of 32 or 36.
		A1	cao	Can ignore scale if bar is correct
(b)	120	M1	ft their histogram for method to find total frequency, eg $(1.6\times 20)+(3.6\times 10)+28+24$ oe **or** for method to find total area under the histogram, eg $(4\times 3.2)+(2\times 7.2)+(2\times 5.6)+(4\times 2.4)$ $(=48)$ oe	Condone 1 error in reading fd values or measurements.
		A1	for 120 ft	

Paper: 1MA1/2H

Question	Answer	Mark	Mark scheme	Additional guidance
15 (a)	3	B1	for 3 correctly placed in the first row	
(b)	Correct bars	B2	for both bars correct, heights of 4.5 and 1.4	
		(B1	for one bar correct out of a maximum of 2 bars)	
(c)	18.5	M1	for finding the area of at least 2 bars eg, 2 × 2 (= 4) **or** 4 × 2 (= 8) **or** 4 × 4 (= 16) **or** 2 × 6.5 (= 13) **or** 4 × 0.5 (= 2)	Condone 1 error in reading fd values or measurements.
		M1	for ("43" + 1) ÷ 2 (= 22) **and** 10 ÷ 16 × 4 (= 2.5) **or** "43" ÷ 2 (= 21.5) **and** 9.5 ÷ 16 × 4 (= 2.375)	
		A1	For 18.5 or 18.375	

Question	Answer	Mark	Mark scheme	Additional guidance
13 (a)	Histogram drawn	B3	for fully correct histogram with axes scaled and labelled	Relative heights 2, 3.5, 4.4, 2, 0.8
		(B2	for 4 correct bars **or** for frequency ÷ class width for all 5 frequencies **and** 2 correct bars of different widths)	
		(B1	for 2 correct bars of different widths **or** for frequency ÷ class width for at least 3 frequencies)	
(b)	$\frac{67}{150}$	M1	for a method to find number of people in interval eg $\frac{1}{2} \times 70 + 22 + \frac{1}{3} \times 30$ (= 67) **or** $150 - 20 - 8 - \frac{1}{2} \times 70 - \frac{2}{3} \times 30$ (= 67) **or** 10 × [3.5] + 5 × [4.4] + 5 × [2] ft their histogram	Where [3.5], [4.4] and [2] are the heights of the bars of their histogram
		A1	for $\frac{67}{150}$ or ft their histogram	

16	Histogram completed	M1 M1 M1 C1	for method to use area to find at least one other frequency (not 11), eg 10×1.5 (= 15) **or** 20×0.3 (= 6) for method to find unknown frequencies, eg (60 – 11 – "15" – "6") ÷ 2 (= 14) ft for method to find an unknown frequency density, eg "14" ÷ 5 (= 2.8) **or** "14" ÷ 10 (= 1.4) **or** one correct bar drawn cao	 Condone one error in reading FD Ft one error from reading FD

Paper: 1MA1/2H

Paper: 1MA1/1H

Question	Answer	Mark	Mark scheme	Additional guidance
15	$\frac{130}{400}$ oe	M1	for a method to find the frequency (or area) of one relevant interval, eg 3×10 (= 30) or 5×20 (= 100) or 8×15 (= 120) or 10×15 (= 150) or 5×10 (= 50) or 8×10 (= 80) or, using $1\,cm^2 = 1$ unit, 3×2 (= 6) or 5×4 (= 20) or 8×3 (= 24) or 10×3 (= 30) or 5×2 (= 10) or 8×2 (= 16)	Evidence for this mark may be seen on the diagram Accept equivalent methods For M marks condone use of 2.5 for fd of 3 and/or 4.5 for fd of 5
		M1	for a method to find the total frequency (or area), eg "30" + "100" + "120" + "150" (= 400) or "6" + "20" + "24" + "30" (= 80)	
		M1	for a method to find the frequency (or area) between 20 g and 40 g, eg "50" + "80" (= 130) or "10" + "16" (= 26)	
		A1	for $\frac{130}{400}$ oe	Accept any equivalent fraction, decimal form, 0.32(5) or 0.33 or percentage form, 32.(5)% or 33%

Paper: 1MA1/1H

GCSE

Mark Scheme

Mathematics A (1387)

June 2003

Edexcel
Success through qualifications

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information please call our Customer Response Centre on 0870 240 9800, or visit our website at www.edexcel.org.uk.

June 2003

Publications Code UG014129

All the material in this publication is copyright
© London Qualifications Ltd 2003

CONTENTS

Notes on Marking Principles 2
Paper 5501 4
Paper 5502 11
Paper 5503 17
Paper 5504 24
Paper 5505 32
Paper 5506 40

NOTES ON MARKING PRINCIPLES

NOTES ON MARKING PRINCIPLES

1 Three types of mark are available

M marks - awarded for correct working seen or implied.
A marks - conditional accuracy marks which are awarded for accurate working following the award of M marks.
B marks - unconditional accuracy marks (independent of M).

2 Abbreviations

cao - correct answer only.
ft - follow through.
$\surd$ - Denotes a "follow through" answer.

SC - special case.
isw - ignore subsequent working.
oe - or equivalent (and appropriate).
NB: a candidate cannot benefit from both isw and ft.

3 If no working shown, then

correct answers normally score full marks
incorrect (even though nearly correct) answers score no marks.

4 Marking instructions

Misread loses A marks (and sometimes B marks) on that part but ft can be allowed on subsequent parts. M marks can still be earned. If in doubt contact your team leader.
If there is a wrong answer in the answer space DO CHECK the working in the body of the script.

- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.
- If there is a wrong answer in the answer space DO CHECK the working in the body of the script.
- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.

5 Style of marking

Answer correct: tick and write part mark in margin NEXT TO BRACKETED MARK.
Answer incorrect: cross, but show M, A or B marks if any earned in body of script and transfer the total of these to the margin next to the bracket mark.
Total for each double page at bottom right page (except for back if used), FINAL TOTAL IN RELEVANT BOX ON FRONT COVER.
Nought in margin for fully incorrect question or page AND FOR NO ATTEMPT.
Where no attempt has been made a line should be put in the answer space and zero in the margin next to the bracketed mark
CHOICE OF METHOD – No marks unless one answer is in answer space – then mark that.
CROSSED OUT WORK - if not replaced this should be marked (if legible).
There must always be a mark next to bracketed mark in the margin/

6 Follow Through Marks

Follow throughs are guided by two principles:

(a) Follow throughs which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous, do not award.

(b) Follow throughs which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

7 Probability

Probability answers must be given as fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least two decimal places (unless tenths).

Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.

8 Linear Equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

9 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

Paper 5501

No	Working	Answer	Mark	Notes
1 (a)		46	1	B1 cao
(b)		3.4	1	B1 oe
(c)		Arrow at 430	1	B1 allow ± half graduation
(d)		Arrow at 3.7	1	B1 allow ± half graduation Accept indications other than arrows as long as they are clear
2		1.60 2.05	2	B1) Condone B1) reversal
3 (a)	$\frac{18}{24}$ or $\frac{9}{12}$ or $\frac{6}{8}$	$\frac{3}{4}$	2	B2 for $\frac{3}{4}$ cao (B1 for $\frac{18}{24}$ or $\frac{9}{12}$ or $\frac{6}{8}$) SC B1 for $\frac{1}{4}$ only
(b)		16 squares shaded	1	B1 cao
4 (a)		line	1	B1 within overlay tolerance
(b)		midpoint	1	B1 within overlay tolerance ft from (a) ± 0.2cm
(c)		rectangle	1	B1 for rectangle 6 cm ± 0.2cm by 4 cm ± 0.2cm
5		kilograms, kg litre, *l* or cubic metres, m^3 inches, in	3	B1 B1 B1
6 (a)		parallel lines marked	1	B1
(b)		right angle marked	1	B1
(c)(i)		acute	1	B1
(ii)		reflex	1	B1

Paper 5501

No	Working	Answer	Mark	Notes
7 (i)		sphere	1	B1
(ii)		cylinder	1	B1 Accept circular prism
(iii)		pyramid	1	B1 Condone omission of "triangular" Accept tetrahedron
8 (a)(i)		40	2	B1 cao
(ii)		50		B1 cao
(b)		5 complete symbols	1	B1 cao
(c)		 and oe inc	1	B1
9 (i)		9, 37, 56, 59, 75	5	B1 cao
(ii)		0.067, 0.56, 0.6, 0.605, 0.65		B1 cao Ignore trailing zeros
(iii)		–10, –6, –4, 2, 5		B1 cao
(iv)		$\frac{2}{5}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}$		B2 for all 4 correct (B1 for any 3 in correct order) SC B1 for all 4 in reverse order (applies to(iv) only)
10 (a)			1	B1
(b)		Plotting (4, 24)	1	B1 ft from their matchsticks
(c)		60	1	B1 cao
(d)		$m = 6n$	2	B2 for $m = 6n$ oe (B1 for $6n$ oe or m = multiple of n except $m = n$)
11 (i)		6, 12	4	B1 cao
(ii)		4, 16		B1 cao
(iii)		3, 4, 6 or 3, 4, 6, 12		B1 Condone omission of 12
(iv)		8, 27		B1 cao

Paper 5501

No	Working	Answer	Mark	Notes
12		2.43	4	B1 for 36 or 0.36 B1 for 96 or 0.96 B1 for 125 or 1.25 If none of first 3 B1s awarded then SC B1 for four 24s and five 25s seen OR 4 × 24 and 5 × 25 seen B1 for 2.43 cao
13 (a)(i)		Edinburgh and Plymouth	3	B1 for Edinburgh or −7 B1 for Plymouth or 5
(ii)		12		B1ft from (i) if one positive and one negative
(b)		Cardiff and Belfast London and Plymouth	2	B1 for Cardiff and Belfast OR −6 and −4 B1 for London and Plymouth OR 3 and 5
14		5 9 8	3	B1 cao B1 cao B1 cao
15 (a)			2	✓ × 2 0 B2 1 0 B1 1 1 B1 2 1 B1
(b)	$2 \times 2 \times 2$	8	2	M1 for $2 \times 2 \times 2$ A1 for 8 cao
16 (a)		30	1	B1 cao
(b)		3	1	B1 cao
(c)	$2 \times 29 = 58$ $5 \times 30 = 150$ $2 \times 31 = 62$ $1 \times 32 = 32$ $\frac{302}{10} = 30.2$	30.2	3	M1 for freq × no pins M1 (dep on 1st M1) for totalling and ÷ 10 A1 for 30.2 cao

Paper 5501

No	Working	Answer	Mark	Notes
17		7 5 5 13 33 52 23	3	B3 all correct (B2 for 4, 5 or 6 correct B1 for 2 or 3 correct)
18 (a)(i) (ii) (iii) (iv) (b)		$4c$ p^4 $8g$ $10pr$ OR $10rp$ $10y - 15$	4 1	B1 oe B1 cao B1 oe B1 B1 cao Accept $10y + - 15$
19		$\frac{2}{3}$	3	M1 for 3 rows (9 squares) shaded M1 for 2 columns (10 squares) shaded A1 for $\frac{2}{3}$
	$\frac{3}{5} = \frac{9}{15}$ $\frac{2}{3} = \frac{10}{15}$	$\frac{2}{3}$	3	M1 for $\frac{3}{5} = \frac{9}{15}$ M1 for $\frac{2}{3} = \frac{10}{15}$ A1 for $\frac{2}{3}$
	$\frac{3}{5} = 0.6$ $\frac{2}{3} = 0.66$ or 0.67 or better	$\frac{2}{3}$	3	M1 for $\frac{3}{5} = 0.6$) Accept M1 for $\frac{2}{3} = 0.66$ or 0.67 or better) percentages A1 for $\frac{2}{3}$

Paper 5501

No	Working	Answer	Mark	Notes
20 (a)	955 × 48 7640 38200 45840 OR 48 × 955 240 2400 43200 45840	458.40	3	M1 for complete correct method (condone one computational error) A2 for 458.40 cao (A1 for digits 4584 OR ft if M1 awarded)
(b)	14. 5 48 | 696. 0 48 216 192 24 0 24 0	14.50	3	M1 for 1 as first digit in answer and remainder 21 M1 (dep) 4 as second digit in answer A1 for 14.50 (Accept 14.5)
21 (a)		$12x$	1	B1 oe
(b)		$12x + 10y$	2	B2 oe ft from (a) (B1 $12x$ + multiple of y or $10y$ seen) SC B1 for $x = 12x + 10y$ OR $y = 12x + 10y$

Paper 5501

No	Working	Answer	Mark	Notes
22	$\frac{1}{3}+\frac{1}{4}=\frac{4}{12}+\frac{3}{12}=\frac{7}{12}$ $1-\frac{7}{12}=\frac{5}{12}$	$\frac{5}{12}$	3	M1 for $\frac{4}{12}$ **and** $\frac{3}{12}$ oe A2 for $\frac{5}{12}$ oe (A1 for $\frac{7}{12}$) B1 for 1–"$\left(\frac{1}{3}+\frac{1}{4}\right)$" correctly evaluated
	$1-\frac{1}{3}=\frac{2}{3}$, $1-\frac{1}{4}=\frac{3}{4}$ $\frac{8}{12}-\frac{3}{12}=\frac{5}{12}$ or $\frac{9}{12}-\frac{4}{12}=\frac{5}{12}$	$\frac{5}{12}$	3	B1 for $\frac{2}{3}$ or $\frac{3}{4}$ seen M1 for $\frac{8}{12}-\frac{3}{12}=\frac{5}{12}$ or $\frac{9}{12}-\frac{4}{12}=\frac{5}{12}$ A1 for $\frac{5}{12}$ oe
		$\frac{5}{12}$	3	M1 for 0.25 and 0.33 or better A1 for 0.58 or better A1 for $0.41\dot{6}$ or recurring
23 (a) (b)(i) (ii)	 180 – (54 + 54)	54 72 Reason	1 3	B1 cao M1 for 180 – (54 + 54) A1 ft from (a) if $x < 90$ B1 for mentioning isosceles and equal or base angles or equal sides and equal or base angles

Paper 5501

No	Working	Answer	Mark	Notes
24 (a)		Bryani	2	M1 for 4×9 or $4 \times 3 \times 3$ or $4 \times x \times x$ or square x first or square 3 first A1 SC 4×3^2 with Bryani scores B2
(b)		64	1	B1 cao
25 (a)		 dotted line may be solid	3	B2 for rectangle base 3 squares and height 4 squares (B1 for rectangle with one correct dimension) B1 for horizontal line 1 cm from top) SC B2 for completely correct elevation on its side
(b)			2	B2 for perspective drawing showing slant face and cutout B1 for perspective drawing with either slant face cutout omitted
26 (a)	20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$	40	2	M1 for 20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$ A1 cao
(b)	$\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes	line from (45, 20) to (65, 0)	2	M1 for $\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes seen A1 for correct line SC If M0, B1 for line from (45, 20) to (t, 0) where $t > 45$ or B1 for a line of the correct gradient.

Paper 5502

No	Working	Answer	Mark	Notes
1 (a)		$\frac{7}{100}$	1	B1 cao accept 0.07
(b)		0.18	1	B1 cao
(c)	20 in 100 oe	40	2	M1 for sight of 20 in 100 or 20 × 2 A1 cao
2 (a)		6cm	2	B1 for 6 ± 0.2 or 60 ± 2 B1 indep for cm or mm consistent with 1st B1
(b)		At centre	1	B1 within overlay
(c)		Circle drawn	1	B1 all within overlay
3		See diagram	3	B3 all correct – see separate sheet (B2 for 3 correct B1 for 2 correct)
4 (a)		$\frac{1}{4}$ oe	1	B1 cao
(b)		0.75	1	B1 cao
(c)		75%	1	B1 cao
(d)(i)		9	2	B1 accept answer in range 9 - 9.2
(ii)		15 - 16		B1 accept answers in range15 - 16
5 (a)		9:30	1	B1 cao
(b)		2hrs 45 min	2	B2 for 2hr 45 min or 2 $\frac{3}{4}$ hr or 165 minutes B1 2:45 or 2.45 or 165 or 45min + 1hr + 1hr oe
(c)		17	1	B1 cao
6 (a)	96 × 4	3.84	2	M1 for 96 × 4 or digits 384 A1 cao
(b)	3 × 96 + 40 = 328	(0).56	2	M1 for 3 × 96 + 40 or digits 328 or digits 56 A1 cao accept 56p

Paper 5502

No	Working	Answer	Mark	Notes
7 (a)		54 000	1	B1 cao accept 54 thousand
(b)		50 000	1	B1 (accept ten thousand or 10 000) oe
8 (a)		14	1	B1 cao
(b)		6	1	B1 cao
(c)		Correct reflection	2	B2 fully correct (B1 correct reflection in a line parallel to the mirror line or condoning 1 block error in shape or position of shape)
9 (a)		Missing horiz label 1 (and 6) missing on vertical scale	2	B1 B1
(b)		Correct graph	2	B1 for bar up to 4 for yellow B1 for bar up to 2 for green
(c)		Blue	1	B1 cao
(d)	3 + 5 + 4 + 2	14	1	B1 ft from (b)
(e)		$\frac{3}{14}$	1	B1 ft on '14'
10	Barry (8) because you double Kath (7) because you add, 1,2,3		2	B1 oe B1 oe SC: B1 for correct rules only
11 (a)		$2n$	1	B1 for $2n$ or $n + n$ OR $2 \times n$ OR $n \times 2$ OR $n2$
(b)		$2n + 15$	1	B1 for "$2n$" + 15 oe
(c)		20q	1	B1 cao
12 (a)	1 + 3 + 5 + 8 + 5	22	2	M1 add frequencies A1 cao
(b)		No, is>no of cups of coffee in the table	1	B1 'average cannot be bigger than 6' oe OR 'Average must be less than 6 oe'
13 (a)		Trapezium	1	B1 cao ignore spelling
(b)		(2, 3)	1	B1 cao
(c)		Isosceles	1	B1 cao ignore spelling
(d)		Q correct	1	B1 cao

Paper 5502

No	Working	Answer	Mark	Notes
14 (a)		250 000	1	B1 cao
(b)	$\frac{28}{4}$	7	2	M1 for $\frac{28}{4}$ oe or "250000" × 28 A1 cao **SC** B1 for 7 000 000
15 (a)		10	1	B1 cao
(b)		5.5	1	B1 ±0.3 pounds
(c)	$\frac{110}{22}$	50	3	M1 for use of graph at 11 or $\frac{110}{22}$ A1 for 5 A1 cao **SC** B2 for 49.5 – 50.6

Paper 5502

No	Working	Answer	Mark	Notes
16 (a)	269.30 – 56.80 = 212.50 $\frac{212.50}{42.50}$	6	2	M1 for $\frac{269.30-56.80}{42.50}$ or 5 seen A1 cao
(b)	5% of £269.30 £269.30 – "£13.465" OR $\frac{95}{100}\times 269.30$	255.83 or 255.84	3	M1 for (5 ÷ 100) × 269.30 M1 for 269.30 – "13.465" A1 cao OR M2 for $\frac{95}{100}\times 269.30$ A1 cao **Alternative Method:** M1 for $\frac{5}{100}\times 56.80$ (= 2.84) and $\frac{5}{100}\times 42.50$ (=2.12(5)) (OR 53.96 AND 40.38 (40.375) seen M1 for 56.80 – "2.84" (= 53.96) 42.50 – "2.12(5)" (= 40.375 or 40.38) "5" × "40.375" + "53.96" A1 cao
17	$4.1^2\times 1.07 = 16.81\times 1.07$	17.9867	2	M1 for ("4.1") followed by squaring, or sight of 16.81 A1 cao SC: *B1 for 18 or better with no working*
18 (a)		60	1	B1 cao
(b)	360 – 60 – 90 – 90	120	2	M1 for 360 – " 60 " – 90 – 90 or 180 – "60" A1 cao
(c)	6 × 2	12	2	M1 for 6 × 2 A1 cao for 12

Paper 5502

No	Working	Answer	Mark	Notes
19 (a)(i)	$240 \times 5 = 1200$	1250	3	B1 cao 1250
(ii)	$\frac{50}{'1250'}$	$\frac{1}{25}$		M1 cao $\frac{50}{'1250'}$ A1 for $\frac{1}{25}$ in its simplest form
(b)	$\frac{60}{100} \times 1000 = 600$	12:5	3	M1 for $\frac{60}{100} \times 1000$ oe A1 for 600 A1 cao
20 (a)		$x + 2$	1	B1 accept $2 + x$ but not $x = x + 2$
(b)	$x + 5 + x + 5 + x + 2 + x + 2$	$4x + 14$	2	M1 adding 4 sides, two of which are '$x + 2$' (all sides to be linear expressions in x) **SC** $x + 5 + x + 2 \times 2$ gets M1 A1 for correct simplified answer or $(20 - 14) \div 4$ oe gets M1
(c)	'$4x + 14$' = 20	1.5 oe	2	M1 for equation A1 cao
21	$\sum f = 90$	Angles drawn, labelled	3	M1 for 1 person = 4°or one angle correct in table or pie chart A1 any 2 correctly drawn angles in pie chart A1 fully correct chart labelled
22 (a)		$2p - q$	2	B1 cao for $2p$ B1 cao for $-q$ accept $(-q + 2p)$, $2p - 1q$ and $2p + -q$
(b)	$5x = 3 + 4$	1.4	2	M1 for either (+3 or sight of 7) or (÷ 5 or sight of 0.8 and 0.6) A1 cao accept $\frac{7}{5}$ or $1\frac{2}{5}$

Paper 5502

No	Working	Answer	Mark	Notes
23 (a)		$\frac{4\times5}{2}$	1	B1 cao
(b)	$1+2+3+4+5+6+7+8$	$\frac{8\times9}{2}$	1	B1 cao
(c)	$\frac{100\times101}{2}$	5050	1	B1 cao
24	0 | 5 7 8 8 1 | 0 0 0 0 2 5 5 5 6 2 | 0 0 0 4 5 3 | 3 5 Key 1 | 2 = 12 (min)	See working	3	B1 for stem 0, 1, 2, 3 or 0, 10, 20, 30 B1 for accurate unordered leaves condone 1 error or omission B1 for key and ordered leaves all correct
25	$3.2\times2.8=8.96$ $2\times4.5\times2.8=25.2$ $2\times4.5\times28.8=28.8$ $\frac{62.96}{2.5}\times2.99$	£75.30	5	M1 for area of any face found correctly M1 for 2 areas found correctly A1 for 62.96 or 54 M1 for $\frac{'62.96'}{2.5}\times2.99$ Al cao Alternate method for candidates who round up $\frac{"62.92"}{2.5}$ M1 for "26" × 2.99 A1 for £77.74 cao **SC**: *for top included B2 for 71.92 m^2 seen or* B3 for £86.02 seen **SC** B4 for £64.58 or £65.78 seen
26	$2.5\times10\,000$	25 000	2	M1 for $2.5\times100\times100$ A1 cao

Paper 5503

No	Working	Answer	Mark	Notes
1 (a)(i)		$8g$	2	B1 oe
(ii)		$10rp$		B1 for $10pr$ or $10\ rp$
(b)		$10y - 15$	1	B1 cao accept $10y$ - + 15
(c)	$6x + 8 - 12x + 15$	$-6x + 23$	2	M1 for 3 correct terms out of 4 A1 cao
2		$m = 6n$	2	B2 for $m = 6n$ oe accept $6 \times n$, n 6 (B1 for $6n$ alone, or $6n$ +1 oe OR m = multiple of n except $m = n$)
3 (i)		0.067, 0.56, 0.6, 0.605, 0.65	1	B1 cao Ignore trailing zeros
(ii)		−10, −6, −4, 2, 5	1	B1 cao
(iii)		$\frac{2}{5}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}$	2	B2 all four correct (B1 any three in correct order) SC: B1 all 4 in reverse order
4		7 5 5 13 33 52 23	3	B3 all correct (B2 for 4, 5 or 6 correct entries) (B1 for 2, 3 correct entries)

Paper 5503

No	Working	Answer	Mark	Notes
5		$\frac{2}{3}$	3	M1 for 3 rows (9 squares) shaded M1 for 2 columns (10 squares) shaded A1
	OR $\frac{3}{5}=\frac{9}{15}$ $\frac{2}{3}=\frac{10}{15}$ Therefore $\frac{2}{3}>\frac{3}{5}$	$\frac{2}{3}$		M1 for $\frac{3}{5}=\frac{9}{15}$ M1 for $\frac{2}{3}=\frac{10}{15}$ A1
	OR $\frac{3}{5}=0.6$ or percent $\frac{2}{3}=0.66$ or 0.67 or better Therefore $\frac{2}{3}>\frac{3}{5}$	$\frac{2}{3}$		M1 for $\frac{3}{5}=0.6$ or percent M1 for $\frac{2}{3}=0.66$ or 0.67 or better A1
6	$2\times 29=58$ $5\times 30=150$ $2\times 31=62$ $1\times 32=32$ $\frac{302}{10}=30.2$	30.2	3	M1 for freq × no. pins (at least 3) M1 for totalling and for ÷ 10 (dep on 1st M1) A1 cao
7 (a)			2	B1 for 180° rotation (wrong centre) B1 cao
(b)			3	B1 for any enlargement sf other than 1 B1 for all sides halved B1 for position

Paper 5503

No	Working	Answer	Mark	Notes
8 (a)		$12x$	1	B1 oe
(b)		$12x + 10y$	2	B2 oe or ft from (a) (B1 for $12x$ + a multiple of y or $10y$ seen) **SC** B1 for $x = 12x + 10y$ or $y = 12x + 10y$
9	$\frac{1}{3}+\frac{1}{4}=\frac{4}{12}+\frac{3}{12}=\frac{7}{12}$ $1-\frac{7}{12}=\frac{5}{12}$ OR $1-\frac{1}{3}=\frac{2}{3}, 1-\frac{1}{4}=\frac{3}{4}$ $\frac{9}{12}-\frac{4}{12}=\frac{5}{12}, \frac{8}{12}-\frac{3}{12}=\frac{5}{12}$	$\frac{5}{12}$	3	M1 for $\frac{4}{12}$ and $\frac{3}{12}$ oe A2 for $\frac{5}{12}$ oe (A1 for getting $\frac{7}{12}$) (B1 for $1 - \frac{"1"}{3} + \frac{1"}{4}$ correctly evaluated) OR B1 for $\frac{2}{3}$ or $\frac{3}{4}$ seen M1 for $\frac{3}{4}-\frac{1}{3}=\frac{9}{12}-\frac{4}{12}$ or $\frac{2}{3}-\frac{1}{4}=\frac{8}{12}-\frac{3}{12}$ A1 for $\frac{5}{12}$ oe OR M1 0.25 and 0.33 or better A1 for 0.58 or better A1 for 0.416 *recurring*
10 (a)		54	1	B1 cao
(b)(i)	180 – (54 +54)	72	3	M1 for 180 – (“54” + “54”) A1 for 72° ft from (a) if $x < 90°$
(ii)				B1 for mentioning isosceles and equal or base angles or equal sides and equal or base angles

Paper 5503

No	Working	Answer	Mark	Notes
11 (a)	Bryani was correct $4 \times 3^2 = 4 \times 9 = 36$	Bryani	2	M1 for 4×9 or $4 \times 3 \times 3$ or "square the three/x then multiply by four" A1 Bryani **SC** 4×3^2 with Bryani gets B2
(b)			1	B1 cao
12 (a)			3	B2 for rectangle height 4 squares, base 3 squares (B1 for rectangle with one correct dimension) B1for line 1 square from the top **SC** B2 for completely correct elevation on its side
(b)			2	B2 for perspective drawing showing slant face and cut out (B1 for perspective drawing with either slant face or cut out omitted or one aspect incorrect)
13 (a)	20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$	40	2	M1 for 20×2 or $\frac{20}{30} \times 60$ or $20 \div \frac{1}{2}$ A1 cao
(b)	$\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes seen	Line from (45, 20) to (65, 0)	2	M1 for $\frac{20}{60}$ or $\frac{1}{3}$ or 20 minutes seen A1 for correct line **SC** If M0, B1 for line from (45, 20) to (t, 0) where $t > 45$ or a line of the correct gradient
14 (i)		119.31	3	B1 cao
(ii)		119 310		B1 cao
(iii)		1.23		B1 cao

Paper 5503

No	Working	Answer	Mark	Notes
15	$\frac{10}{100}\times 12000$ 12 000 – 1200 = 10 800 $\frac{10800}{10}=1080$ 10 800 – 1080 = £9720	£9720	3	M1 for $\frac{10}{100}\times 12000$ or sight of 1200 or 2400 or 10 800 or 9600 M1 (dep) for $\frac{10}{100}\times(12\,000-\frac{10}{100}\times 12000)$ or sight of 1080 A1 cao Alternative mark scheme M2 for $12000\times(1-\frac{10}{100})^2$ (M1 for $12000\times(1-\frac{10}{100})$ A1 cao
16 (a)	$2p = 6$	$p = 3$	2	M1 for $7p - 5p = 8 - 2$ or $2p$ or 6 A1 cao
(b)	$7r - 5r = -20 - 2$	–11	2	M1 for $7r + 2 = 5r - 20$ or $\frac{7r}{5}+\frac{2}{5}=r-4$ or $7r - 5r = 20 - 2$ or $\frac{7r}{5}-r=-4-\frac{2}{5}$ A1 cao
17		$5n + 1$	2	B2 oe (B1 for $5n$ seen) NB: n = gets B1 max

Paper 5503

No	Working	Answer	Mark	Notes
18 (a)	–1, 0, 1		2	B2 for –1, 0, 1 (B1 for –1, 0 or 0,1 or –1,1 or –2, -1, 0, 1 only)
(b)	(–1, –1), (0, –1), (1, –1), (0, 0), (1, 0), (1, 1)		3	B3 for 6 points correct B2 for 3 points correct B1 for 1 point correct **NB** –B1 each additional point over six
19 (a)	Triangle with vertices at (0,0) (0,–2) and (3,0)		2	M1 for correct orientation A1 cao
(b)	Rotation, 180°, centre (0,1) Enlargement sf - 1 centre (0,1)		2	B2 for 180° 'rotation' centre (0, 1) B2 for Enlargement sf - 1 centre (0,1) (B1 for any two of the three parts) NB: B0 if additional transformation is included
20	Bisector of $\angle BAC$ Arc around A Region		3	B3 cao (B2 for either two correct boundaries, no shading/ wrong shading or one correct boundary, one incorrect boundary with valid shading) (B1 for either two incorrect boundaries but one drawn from A and one intersection, with valid shading or one correct boundary) Ignore shading outside the triangle
21		Length Volume Area	3	B1 for Length B1 for Volume B1 for Area
22 (a)	Unbiased question with choices		2	B1 for unbiased question B1 for at least 2 choices
(b)			2	Classification 1: A biased question Classification 2: A restricted sample of people Classification 3: Not specifying a range of foods Classification 4: Nothing to do with eating habits B2 reasons which satisfy 2 different classifications (B1 a reason which satisfies one classification)



Paper 5503

No	Working	Answer	Mark	Notes
23 (a)	$6 \times 10^2 \times 8 \times 10^4$ 48×10^6	4.8×10^7	3	M1 for $6 \times 10^a \times 8 \times 10^b$ oe, a and b integers including 0 A1 for 48×10^6 oe A1 cao
(b)	200 000 + 30 000	230000	2	B2 cao (B1 for sight of 200 000 or 30 000 or 2.3×10^5 or 23×10^4)
24 (i)		64	4	B1 cao
(ii)		3		B1 cao
(iii)	$\sqrt{16 \times 9} = \sqrt{144}$	12		B2 cao (B1 for sight of $\sqrt{2^4} \times \sqrt{9}$ or better, or 144 seen)
25 (i)	Tangent 90° to diameter /radius/ line from (through) centre	27°	4	B1 for 27° cao B1 for reason
(ii)	180 –(90+ "27") angle in semicircle (is 90°)/Alternate segments /angle at centre twice at circumference	63°		B1 ft for 90–"27" if not 63° B1 for reason
26 (a)(i)		152	2	B1 cao
(ii)		177		B1 cao
(b)			3	B1 for median marked at 167 B1 ft for postion of box with its ends at "152" and "177" B1 for position of whiskers with ends at 132 and 182 NB: For any points plotted between 141 and 149 give a tolerance of an extra ± 1 square
27	$x^2 + xy + xy + y^2$	$x^2 + 2xy + y^2$	2	M1 for at least 3 of the 4 terms correct A1 cao
		25	2	M1 for recognising 3.47 + 1.53 (= 5) A1 cao

Paper 5504

No	Working	Answer	Mark	Notes
1 (a)	$4.1^2 \times 1.07 = 16.81 \times 1.07$	17.9867	2	M1 for (4.1) followed by squaring, or sight of 16.81 A1 cao SC: B1 for 18 or better with no working
(b)		$(1.6 + 3.8 \times 2.4) \times 4.2$	1	B1 cao Allow additional brackets if they give an expression with value 45.024
2 (a)	269.30 – 56.80 = 212.50	6	2	M1 for $\frac{269.30 - 56.80}{42.50}$ or 5 seen A1 cao
(b)	5% of £269.30 £269.30 – "£13.465" OR $\frac{95}{100} \times 269.30$	255.83 or 255.84	3	M1 for (5 ÷ 100) × 269.30 M1 (dep) for 269.30 – "13.465" A1 cao OR M2 for $\frac{95}{100} \times 269.30$ A1 cao **Alternative Method:** M1 for $\frac{5}{100} \times 56.80$ (= 2.84) and $\frac{5}{100} \times 42.50$ (=2.12(5)) (OR 53.96 AND 40.38 (40.375) seen M1 for 56.80 – "2.84" (= 53.96) 42.50 – "2.12(5)" (= 40.375 or 40.38) "5" × "40.375" + "53.96" A1 cao

Paper 5504

No	Working	Answer	Mark	Notes
3 (a)		60	1	B1 cao
(b)	360 – 60 – 90 – 90	120	2	M1 for 360 – "60" – 90 – 90 or 180 – "60" A1 cao
(c)	6 × 2	12	2	M1 for 6 × 2 A1 cao 12
(d)		Correct drawing	2	B2 for triangle and construction lines (see overlay) (B1 for 1 line of length 4cm and correct arcs crossing OR for correct triangle with either no arcs or incorrect arcs) SC: B1 similar triangle drawn with construction lines
4 a(i)	240 × 5 = 1200	1250	3	B1 cao 1250
(ii)	$\frac{50}{1250}$	$\frac{1}{25}$		M1 $\frac{50}{"1250"}$ A1 oe in its simplest form
(b)	$\frac{60}{100} \times 1000 = 600$ 600: 250	12:5	3	M1 for $\frac{60}{100} \times 1000$ oe A1 for 600 A1 cao
5 (a)		$x + 2$	1	B1
(b)	$x + 5 + x + 5 + x + 2 + x + 2$	$4x + 14$	2	M1 adding 4 sides, two of which must be '$x + 2$' (all sides to be linear expressions in x) A1 for correct simplified answer
(c)	"$4x + 14$" = 20	1.5	2	M1 for equation "$4x + 14$" = 20 OR $\frac{20-14}{4}$ oe A1 cao

Paper 5504

No	Working	Answer	Mark	Notes
6 (a)		$2p - q$	2	B1 cao for $2p$ B1 cao for $-q$ accept $-q + 2p$ and $2p - 1q$
(b)	$5x = 3 + 4$	1.4	2	M1 for either (+3 or sight of 7) or (÷ 5 or sight of 0.8 and 0.6) A1 cao accept $\frac{7}{5}$ or $1\frac{2}{5}$
7 (a)		$\frac{4\times5}{2}$	1	B1 cao
(b)	$1 + 2 + 3 + 4 + 5 + 6 + 7 + 8$	$\frac{8\times9}{2}$	1	B1 cao
(c)	$\frac{100\times101}{2}$	5050	1	B1 cao
(d)		$\frac{n(n+1)}{2}$	2	B2 cao (B1 for any quadratic in n)
8	$3.2 \times 2.8 = 8.96$ $2 \times 4.5 \times 2.8 = 25.2$ $2 \times 4.5 \times 3.2 = 28.8$ $\frac{62.96}{2.5} \times 2.99$	75.30	5	M1 for area of any face found correctly M1 for 2 areas seen A1 for 62.96 or 54 M1 for $\frac{'62.96'}{2.5} \times 2.99$ A1 cao Alternate method for candidates who round up $\frac{"62.96"}{2.5}$ M1 for "26" x 2.99 A1 for £77.74 cao SC: for top included B2 for 71.92 m^2 seen or B3 for £86.02 or £86.71 SC: B4 for £64.58 or £65.78 seen

Paper 5504

No	Working	Answer	Mark	Notes
9	2.5×10000	25000	2	M1 for 2.5×10000 or $2.5 \times 100 \times 100$ A1 cao
10 (a)	$\sum f = 90$ (88), 144, 32, 96	Angles drawn, labelled	3	M1 for 1 person = 4° or one angle correct in table or pie chart A1 any 2 angles correctly drawn in pie chart A1 fully correct chart labelled
(b)	$0.38 + 0.27 + 0.15$	0.20	2	M1 1 – sum A1 cao
11	0 | 5 7 8 8 1 | 0 0 0 0 2 5 5 5 6 2 | 0 0 0 4 5 3 | 3 5 Key 1 | 3 = 13 (min)	See working column	3	B1 for stem 0, 1, 2, 3 or 0, 10, 20, 30 B1 for accurate unordered leaves condone 1 error or omission B1 for key and ordered leaves all correct
12 (a)	$V = \pi \times 4^2 \times 10$	502 – 503	2	M1 for $\pi \times 4^2 \times 10$ A1 502 – 503
(b)	$P^2 = 10^2 + 8^2$ $P = \sqrt{164}$	$\sqrt{164} < 13$	3	M1 for sight of a correct right-angled triangle M1 for $10^2 + 8^2$ A1 for conclusion based on a correct calculation or 12.8 seen
13 (a)	2×30	$2 \times 2 \times 3 \times 5$	4	M1 for systematic method, eg division, factor trees (at least one prime) A1 cao
	2×48	$2^5 \times 3$		M1 for systematic method, eg division, factor trees (at least one prime) A1 cao
(b)		12	1	B1 cao
(c)	$2^5 \times 3 \times 5$	480	2	B2 cao B1 for $2^5 \times 3 \times 5$ or any correct common multiple

Paper 5504

No	Working	Answer	Mark	Notes
14 (a)		$150 < C \leq 200$	2	M1 use of cum freq to find the cost of the 20th or 20.5^{th} car OR $\frac{1}{2}\Sigma f$ or $\frac{1}{2}(\Sigma f + 1)$ A1 eg 150 to 200
(b)		No, because the 21st value is in the same interval	1	B1 20.5^{th} or 21st in same interval or an alternative correct explanation
(c)	80% = 5200 $\frac{5200}{80} \times 100$	6500	3	M1 for (100 – 20) % = 5200 M1 for $\frac{5200}{"80"} \times 100$ A1 cao
15 (a)	$x^2(x + 1) = 230$	AG	2	M1 for $x \times x \times (x+1)$ or $x \times x \times x + 1$ oe A1 cao from $x \times x \times (x+1)$
(b)	‘5 – 150 5.1 – 158.7 5.2 – 167.6 5.3 – 177.0 5.4 – 186.6 5.5 – 196.6 5.6 – 207.0 5.7 – 217.7 5.8 – 228.8 5.9 –240.2 5.85 – 234.4	5.8	4	B 2 for trial between 5.8 and 5.9 inclusive (B1 for different trial between 5 and 6 inclusive) B1 for different trial between 5.8 and 5.85 (not including 5.8) B1 (dep on at least one previous B1) cao for 5.8, 5.81, 5.811

Paper 5504

No	Working	Answer	Mark	Notes
16	$\pi \times \left(\frac{15}{2}\right)^2 = 176.715$	88.4cm^2	3	M1 for $\pi \times \left(\frac{15}{2}\right)^2$ A1 88.3 – 88.4 B1(ind) for cm²
17 (a)	$5 = 0.5x + 1$	8	2	M1 for $5 = 0.5x + 1$ A1 cao
(b)		$y = \frac{1}{2}x + c$	1	B1 $y = \frac{1}{2}x + c$, c ≠ 1, oe
(c)		$x = 2y - 2$ OR $x = 2(y - 1)$	2	M1 for correctly multiplying both sides by 2 or correctly isolating $\frac{x}{2}$ A1 for $x = 2(y - 1)$, $x = \frac{y-1}{0.5}$, $x = \frac{y-1}{\frac{1}{2}}$ oe SC: B1 for $x = 2y - 1$
18	$4x - 6y = 22$ $15x + 6y = 54$ $19x = 76$	$x = 4,\ y = -1$	4	M1 for coefficients of x or y the same followed by correct operation , one arithmetical error A1 cao M1(dep on previous M mark) for sub for other variable A1 cao Trial and improvement –0 unless both x and y correct values found

Paper 5504

No	Working	Answer	Mark	Notes
19 (a)	$SF = \frac{10}{6}$ $\frac{10}{6} \times 4.8 = 8$	8	2	M1 for sight of $\frac{10}{6}$ or $\frac{6}{10}$ or 1.67 or better or $\frac{CD}{10} = \frac{4.8}{6}$ A1 cao
(b)	$\frac{10}{6} \times 4.5 - 4.5 = 3$	19.8	2	M1 for use of SF from (a) to find *BC* or *AC* and adding 4 sides A1 cao
20	$\frac{6 \times 10^{15}}{3.2 \times 10^{8}}$ 1.875×10^{7}	4.3×10^{3}	3	B3 for 4.3×10^{3} to 4.34×10^{3} (B2 for 1.875×10^{7} oe or 4300 to 4340 or final answer of 1.9×10^{7}) (B1 for sight of 6×10^{15} oe or 3.2×10^{8} oe)
21	$8.5 \times \tan 38 = 8.5 \times 0.7813$	6.64	3	M1 for correct use of trig, eg $\tan 38 = \frac{opp}{8.5}$ M1 for 8.5 × tan38 A1 6.64 – 6.641

Paper 5504

No	Working	Answer	Mark	Notes
22 (a)		No, as you would expect about 100. Yes, as it is possible to get 200 sixes with a fair dice	1	B1 for a consistent answer
(b)	$\frac{1}{6}, \frac{5}{6}$ + labels		3	B1 for $\frac{5}{6}$ on the red dice, *not six* branch B1 for a fully complete tree diagram with all branches labelled B1 for $\frac{1}{6}$, $\frac{5}{6}$ on all remaining branches as appropriate

Paper 5505

No	Working	Answer	Mark	Notes
1 (i) (ii) (iii)		119.31 119310 1.23	3	B1 cao B1 cao B1 cao
2	$\frac{10}{100}\times 12000 = 1200$ 12 000 – 1200 = 10 800 10 800 ÷ 10 =1080 10 800 – 1080 = £9720	9720	3	M1 for $\frac{10}{100}\times 12\,000$ or sight of 1200 or 2400 or 10 800 or 9600 M1 (dep) for $\frac{10}{100}\times(12\,000 - \frac{10}{100}\times 12\,000)$ or sight of 1080 A1 cao Alternative markscheme M2 for $12000\times\left(1-\frac{10}{100}\right)^2$ (M1 for $12000\times\left(1-\frac{10}{100}\right)$) A1 cao
3	$7r - 5r = -20 - 2$	–11	2	M1 for $7r + 2 = 5r - 20$ or $\frac{7r}{5} + \frac{2}{5} = r - 4$ or $7r - 5r = -20 - 2$ or $\frac{7r}{5} - r = -4 - \frac{2}{5}$ A1 cao

Paper 5505

No	Working	Answer	Mark	Notes
4 (a)		–1, 0, 1	2	B2 for –1, 0, 1 only (B1 for –1, 0 or 0,1 or –1, 1 or –2, -1, 0, 1 only)
(b)		(–1,–1)(0,–1) (1,–1) (0, 0) (1, 0) (1, 1)	3	B3 for 6 points correct (B2 for 3 points correct) (B1 for 1 point correct) *NB: –B1 for each additional point over six*
5		$5n + 1$	2	B2 oe (B1 for $5n$ seen) NB: n = gets B1 max
6 (a)	Triangle with vertices at (0,0) (0,–2) and (3,0)		2	M1 for correct orientation A1 cao
(b)	Rotation, 180°, centre (0,1) Enlargement sf –1 centre (0,1)		2	B2 for 180° 'rotation' centre (0, 1) or for Enlargement sf - 1 centre (0,1) (B1 for any two of the three parts) NB: B0 if additional transformation is included
7	Bisector of $\angle BAC$ Arc around A Region	See overlay	3	B3 cao (B2 for either two correct boundaries, no shading/ wrong shading or one correct boundary, one incorrect boundary with valid shading) (B1 for either two incorrect boundaries but one drawn from A and one intersection, with valid shading or one correct boundary) Ignore shading outside the triangle
8		Length Volume Area	3	B1 for Length B1 for Volume B1 for Area



Paper 5505

No	Working	Answer	Mark	Notes
9 (a)	Unbiased question with choices		2	B1 for unbiased question B1 for at least 2 choices
(b)	Leading question and a restricted sample		2	Classification 1: A biased question Classification 2: A restricted sample of people Classification 3: Not specifying a range of foods Classification 4: Nothing to do with eating habits B2 reasons which satisfy 2 different classifications (B1 a reason which satisfies one classification)
10 (a)	$6 \times 10^2 \times 8 \times 10^4$ $48 \times 10^6 = 4.8 \times 10^7$	4.8×10^7	3	M1 for $6 \times 10^a \times 8 \times 10^b$ oe, a and b integers including 0 A1 for 48×10^6 oe A1 cao
(b)	200 000 + 30 000 = 230 000	230 000	2	B2 cao (B1 for sight of 200 000 or 30 000 or 2.3×10^5 or 23×10^4)
11 (a)	$x^2 + xy + xy + y^2$	$x^2 + 2xy + y^2$	2	M1 for at least 3 of the 4 terms correct A1 cao
(b)		25	2	M1 for recognising 3.47 + 1.53 (= 5) A1 cao
12 (i)	Tangent 90° to diameter /radius/ line from (through) centre	27°	4	B1 for 27° cao B1 for reason
(ii)	180 –(90+ "27") angle in semicircle (is 90°)/Alternate segments /angle at centre twice at circumference	63°		B1 ft for 90–"27" if not 63° B1 for reason
13 (i)		p^9	1	B1 cao
(ii)		$6q^6$	2	B2 for $6q^6$ (B1 for sight of $\frac{6q^9}{q^3}$ or $3q \times 2q^5$ or $3q^4 \times 2q^2$ or $6 \times q \times q \times q \times q \times q \times q$ or final answer of the form kq^6, $k > 0$)

Paper 5505

No	Working	Answer	Mark	Notes
14 (a)(i)		152	2	B1 cao
(ii)		177		B1 cao
(b)			3	B1 for median marked at 167 B1 ft for postion of box with its ends at "152" and "177" B1 for position of whiskers with ends at 132 and 182 NB: For any points plotted between 141 and 149 give a tolerance of an extra ± 1 square
15	(arc =) $\frac{40}{360} \times 2\pi \times 9$ $= 2\pi$	$2\pi + 18$	4	M1 for $\frac{40}{360} \times$ M1 for $2\pi \times 9$ M1 (dep) for $\frac{40}{360} \times 2\pi \times 9$ oe A1 for $\frac{18 \times \pi}{9} + 18$ oe exact form
16 (i)		1	1	B1 cao
(ii)		$\frac{1}{16}$	1	B1 cao accept 0.0625
(iii)		64	1	B1 cao condone ± 64

Paper 5505

No	Working	Answer	Mark	Notes
17 (a)	$F \propto 1/x^2$ $F = k/x^2$ $4 = k / 3^2$ $F = 36/x^2$	$F = \frac{36}{x^2}$	3	M1 for $F = k/x^2$ seen or implied. ($k \neq 1$) M1 (dep) for subst. or sight of $k = 36$ A1 for $F = 36/x^2$
(b)		9	1	B1 ft for 9 (ft on $F = kx^n$, $n \neq 0$)
(c)	$64 = \frac{36}{x^2}$ $x^2 = \frac{36}{64}$ $x = \pm\frac{3}{4}$	$\frac{6}{8}$	2	M1 for $x^2 = \frac{"36"}{64}$ A1 for $\frac{6}{8}$ oe (condone $\pm$) *SC: Use of $F = kx^2$ max M1 M1 A0 B1 ft M0 A0* *SC: Use of* $F = \frac{k}{\sqrt{x}}$ *max M1 M1 A0 B0 M1* ($\sqrt{x} = \frac{4\sqrt{3}}{64}$) *A0*
18	$(5+\sqrt{3})(5-\sqrt{3}) = 5\times5 - 5\sqrt{3} + 5\sqrt{3} - \sqrt{3}\sqrt{3}$ $= 5\times5 - 3$ $\frac{22}{\sqrt{22}} = \frac{22\sqrt{22}}{22}$	$\sqrt{22}$	3	B1 for correct expansion 25 - $5\sqrt{3}$ + $5\sqrt{3}$ - $\sqrt{3}$ $\sqrt{3}$ with 1st three terms reducing to 25 **without** any errors seen B1 (indep) for $\sqrt{3}$ $\sqrt{3}$ = 3 B1 for $\sqrt{22}$ coming from $\frac{22}{\sqrt{22}}$ (S.C $\frac{(5+\sqrt{3})(5-\sqrt{3})\sqrt{22}}{22}$ gets B1)

Paper 5505

No	Working	Answer	Mark	Notes
19 (a)		60 40	2	B1 cao B1 cao
(b)		correct bars	2	B1 for $30 < x \leq 40$ with an area of 2½ squares B1 for $40 < x \leq 70$ with an area of 3 squares *SC:* $\frac{0}{4}$ *give M1 if clearly using area or frequency density*
20 (a)	$6x + 8 - 12x + 15$	$-6x + 23$	2	M1 for 3 of the 4 terms $6x$, $+8$, $-12x$, $+15$ correct A1 cao
(b)		$32x^5y^{15}$	2	B2 cao (B1 for two of 32, x^5, y^{15})
(c)	$\frac{(n+1)(n-1)}{n+1} \times \frac{2}{n-2}$ $\frac{2(n-1)}{n-2}$	$\frac{2(n-1)}{n-2}$	3	M1 for $k(n + 1)(n - 1)$ M1 dep for $\frac{(n+1)(n-1)}{(n+1)} = n - 1$ A1 for $\frac{2(n-1)}{n-2}$
21	Vertices at $(-3, -1\frac{1}{2}), (-4\frac{1}{2}, -1\frac{1}{2}), (-3, -4\frac{1}{2})$		3	B1 for all sides $\times 1\frac{1}{2}$ B1 for correct orientation with 2 vertices almost correct B1 cao

Paper 5505

No	Working	Answer	Mark	Notes
22	Total = 3 + 5 + 2 (=10) $\frac{3}{10}\times\frac{3}{10}\times\frac{5}{10}\left(=\frac{45}{1000}\right), \frac{3}{10}\times\frac{3}{10}\times\frac{2}{10}\left(=\frac{18}{1000}\right)$ $\frac{5}{10}\times\frac{5}{10}\times\frac{3}{10}\left(=\frac{75}{1000}\right), \frac{5}{10}\times\frac{5}{10}\times\frac{2}{10}\left(=\frac{50}{1000}\right)$ $\frac{2}{10}\times\frac{2}{10}\times\frac{3}{10}\left(=\frac{12}{1000}\right), \frac{2}{10}\times\frac{2}{10}\times\frac{5}{10}\left(=\frac{20}{1000}\right)$ $3\times\left(\frac{"45"}{1000}+\frac{"18"}{1000}+\frac{"75"}{1000}+\frac{"50"}{1000}+\frac{"12"}{1000}+\frac{"20"}{1000}\right)$ $\frac{660}{1000}$	$\frac{660}{1000}$ oe	5	M3 for all six expressions seen OR their combined equivalents (M2 for four expressions seen OR their combined equivalents) (M1 for two expressions seen OR their combined equivalents) M1 sum of 18 relevant products condone 1 slip A1 for $\frac{660}{1000}$ oe *SC: without replacement maximum M4 A0* SC: Just 2 beads: Answer either $\frac{38}{100}$ oe OR $\frac{28}{90}$ oe B1

Paper 5505

No	Working	Answer	Mark	Notes
23 (a)(i)		$6\mathbf{b} - 6\mathbf{a}$	2	B1 for $6\mathbf{b} - 6\mathbf{a}$ oe
(ii)		$6\mathbf{a}$		B1 for $6\mathbf{a}$ oe
(b)	$\overrightarrow{EX} = \overrightarrow{EB} + \overrightarrow{BX}$ $= 12\mathbf{b} + \frac{1}{2}\overrightarrow{BC}$	$12\mathbf{b} - 3\mathbf{a}$	2	M1 for $\overrightarrow{EX} = \overrightarrow{EB} + \overrightarrow{BX}$ oe vector journey in a form ready for straightforward substitution A1 for $12\mathbf{b} - 3\mathbf{a}$ oe
(c)	$\overrightarrow{AY} = \frac{5}{3}\overrightarrow{AB}$ or $\overrightarrow{BY} = \frac{2}{3}\overrightarrow{AB}$ $\overrightarrow{EY} = 16\mathbf{b} - 4\mathbf{a}$ or $\overrightarrow{XY} = 4\mathbf{b} - \mathbf{a}$ $\overrightarrow{EY} = 4\overrightarrow{XY}$ or $\overrightarrow{EX} = 3\overrightarrow{XY}$ or $\overrightarrow{EY} = \frac{4}{3}\overrightarrow{EX}$	Printer Answer	3	B1 for either $\overrightarrow{AY} = \frac{5}{3}\overrightarrow{AB}$ or $\overrightarrow{BY} = \frac{2}{3}\overrightarrow{AB}$ oe B1 ft for either $\overrightarrow{EY} = 16\mathbf{b} - 4\mathbf{a}$ or $\overrightarrow{XY} = 4\mathbf{b} - \mathbf{a}$ ft only on parts (*a*) and (*b*) B1 for either $\overrightarrow{EY} = 4\overrightarrow{XY}$ or $\overrightarrow{EX} = 3\overrightarrow{XY}$ or $\overrightarrow{EY} = \frac{4}{3}\overrightarrow{EX}$ oe **plus** conclusion of *E*, *X*, *Y* on the same straight line
24 (a)(i)		$(5, -4)$	4	B1 cao
(ii)		$(2, -9)$		B1 cao
(iii)		$(2, 4)$		B1 cao
(iv)		$(1, -4)$		B1 cao
(b)		$(x-2)^2 - 4$	4	B4 for $(x-2)^2 - 4$ oe eg. $x^2 - 4x$ (B3 for $(x+2)^2 - 4$ or $(x-2)^2 + 4$) (B2 for $x^2 - 4$ or $(x-2)^2$ OR $x^2 + bx$, $b \neq 0$ OR $(x+2)^2 + 4$ OR $f(x-2) - 4$) (B1 for $x^2 + 4$ or $(x+2)^2$ or $ax^2 + bx$ or $x^2 + bx + c$ OR $x - 2 - 4$ or $x^2 - 2 - 4$, $a, b, c \neq 0$)

Paper 5506

No	Working	Answer	Mark	Notes
1 (a)	$V = \pi \times 4^2 \times 10$	502 – 503 cm^3	2	M1 for $\pi \times 4^2 \times 10$ A1 502 – 503
(b)	$P^2 = 10^2 + 8^2$ $P = \sqrt{164}$	$\sqrt{164} < 13$	3	M1 for sight of correct right angled triangle M1 for $10^2 + 8^2$ A1 for conclusion based on a correct calculation Or 12.8 seen
2 (a)(i)	2×30	$2 \times 2 \times 3 \times 5$	4	M1 for systematic method, eg division, factor trees (at least one prime) A1 cao
(ii)	2×48	$2^5 \times 3$		M1 for systematic method, division, factor trees (at least one prime) A1 cao
(b)		12	1	B1 cao
(c)	$2^5 \times 3 \times 5$	480	2	B2 cao B1 for $2^5 \times 3 \times 5$ or any correct common multiple

Paper 5506

No	Working	Answer	Mark	Notes
3 (a)		$150 < C \leq 200$	2	M1 use of cum freq to find the cost of the 20th or 20.5^{th} car OR $\frac{1}{2}\Sigma f$ th or $\frac{1}{2}(\Sigma f + 1)$ th car. A1 eg 150 to 200, 150 – 200
(b)		No, because the 21st value is in the same interval	1	B1 for 20.5^{th} or 21st value in the same internal consistent with 'a' OR Refers to the median value being low in the interval (statement to be mathematically correct) See additional sheet
(c)	80% = 5200 $\frac{5200}{80} \times 100$	6500	3	M1 for (100 – 20)% = 5200 M1 for $\frac{5200}{"80"} \times 100$ A1 cao

Paper 5506

No	Working	Answer	Mark	Notes
4 (a)	$x^2(x+1) = 230$	AG	2	M1 for $x \times x \times (x+1)$ or $x \times x \times x + 1$ oe, $x^2(x+1)$, $x^2 \times x + 1$ A1 cao from $x \times x \times (x+1)$, no need to see 230
(b)	5 – 150 6 – 252 5.1 – 158.7 5.2 – 167.6 5.3 – 177.0 5.4 – 186.6 5.5 – 196.6 5.6 – 207.0 5.7 – 217.7 5.8 – 228.8 5.9 –240.2 5.85 – 234.4	5.8	4	B2 for trial between 5.8 and 5.9 inclusive evaluated (B1 for trial between 5 and 6 inclusive evaluated) B1 for different trial between 5.8 and 5.85 (not including 5.8) B1 dep on at least are previous B1 5.8, 5.81, 5.811
5	$\pi \times \left(\frac{15}{2}\right)^2 = 176.715$	88.4 cm^2	3	M1 for $\pi \times \left(\frac{15}{2}\right)^2$ seen A1 88.3 – 88.4 B1 for cm² (independent)

Paper 5506

No	Working	Answer	Mark	Notes
6 (a)	$5 = 0.5x + 1$	8	2	M1 for $5 = 0.5x + 1$ A1 cao
(b)		$y = \frac{1}{2}x + c$	1	B1for $y = \frac{1}{2}x + c$, $c \neq 1$, oe
(c)		$x = 2y - 2$ OR $x = 2(y - 1)$	2	M1for correctly multiplying both sides by 2 or correctly isolating $\frac{x}{2}$ A1 for $x = 2(y - 1)$, $x = \frac{y-1}{0.5}$, $\frac{y-1}{\frac{1}{2}}$ oe SC B1 for x= 2y -1
7	$4x - 6y = 22$ $15x + 6y = 54$ $19x = 76$	$x = 4$, $y = -1$	4	M1 for coefficients of x or y the same followed by correct operation, allow one arithmetical error A1 cao M1 (dep) for correct sub for other variable A1 cao Trial and improvement 0 marks unless both correct values of x and y found
8 (a)	SF = $\frac{10}{6}$ $\frac{10}{6} \times 4.8 = 8$	8	2	M1 for sight of $\frac{10}{6}$ or $\frac{6}{10}$ or 1.67 or better or $\frac{CD}{10} = \frac{4.8}{6}$ A1 cao
(b)	$\frac{10}{6} \times 4.5 - 4.5 = 3$	19.8	2	M1 for use of SF from "a" to find AC or BC or $\frac{BC}{4.5} = \frac{4}{6}$ and adding 4 sides A1 cao

Paper 5506

No	Working	Answer	Mark	Notes
9	$\frac{6\times10^{15}}{3.2\times10^{8}}$ 1.875×10^{7}	4.3×10^{3}	3	B3 for 4.3×10^{3} to 4.34×10^{3} (B2 for 1.875×10^{7} oe or 4300 to 4340, final answer of 1.9×10^{7} B1 for sight of 6×10^{15} oe or 3.2×10^{8} oe)
10	$8.5\times\tan38$ $=8.5\times0.7813$ $\frac{8.5}{\sin(90-38)}=\frac{AB}{\sin38}$ $AB=\frac{8.5\times\sin38}{\sin(90-38)}$ $=\frac{5.2331}{0.788}=6.64$	6.64	3	M1 for correct use of trig, eg $\tan 38=\frac{opp}{8.5}$ M1 for $8.5\times\tan38$ A1 6.64 – 6.641 OR M1 for correct substitution into the sine rule M1 (dep) for correct rearrangement for AB= A1 6.64 – 6.641
11 (a)		No, as you would expect about 100. Yes, as it is possible to get 200 sixes with a fair dice	1	B1 for a consistent answer See additional sheet

Paper 5506

No	Working	Answer	Mark	Notes
(b)	$\frac{1}{6}, \frac{5}{6}$ + labels		3	B1 for $\frac{5}{6}$ on the red dice, *not six* branch B1 for a fully complete tree diagram with all branches labelled B1 for $\frac{1}{6}$ and $\frac{5}{6}$ on all remaining branches as appropriate
(c)(i)	$\left(\frac{1}{6}\right)^2$	$\frac{1}{36}$	2	M1 $\left(\frac{1}{6}\right)^2$ or $\frac{1}{6}\times\frac{1}{6}$ only or 0.28 A1 $\frac{1}{36}$ or 0.03 or better
(ii)	$1-\left(\frac{5}{6}\right)^2$ OR $\frac{1}{6}\times\frac{5}{6}+\frac{5}{6}\times\frac{1}{6}+\frac{1}{6}\times\frac{1}{6}$	$\frac{11}{36}$	3	M2 for $1-\left(\frac{5}{6}\right)^2$ or $1-\frac{5}{6}\times\frac{5}{6}$ A1 cao OR M1 for $\frac{1}{6}\times\frac{5}{6}$ oe M1 for 2 or 3 only of $\frac{1}{6}\times\frac{5}{6}, \frac{5}{6}\times\frac{1}{6}$, "a" A1 for $\frac{11}{36}$ or 0.31 or better

Paper 5506

No	Working	Answer	Mark	Notes
12 (a)	$\pi \times 30 \times \frac{7.5^2}{3} - \pi \times 10 \times \frac{2.5^2}{3} = 1767 - 65$	1700	3	M1 for either $\pi \times 30 \times \frac{7.5^2}{3}$ or $\pi \times 10 \times \frac{2.5^2}{3}$ M1 (dep) for difference A1 1700 – 1702 **SC** B1 Using d instead of r, 6800 – 6808
(b)	$\frac{S}{2\pi d} = \sqrt{h^2 + d^2}$ $\left(\frac{S}{2\pi d}\right)^2 = h^2 + d^2$	$h = \sqrt{\frac{S^2 - 4\pi^2 d}{4\pi^2 d^2}}$	3	M1 for correctly isolating $\sqrt{h^2 + d^2}$ or $h^2 + d^2$ or $h + d$ or kh^2 or kh M1(indep) squaring both sides A1 $h = \sqrt{\frac{S^2 - 4\pi^2 d^4}{4\pi^2 d^2}}$, $h = \frac{\sqrt{S^2 - 4\pi^2 d^4}}{2\pi d}$ $h = \sqrt{\left(\frac{S}{2\pi d}\right)^2 - d^2}$
(c)	$\left(\frac{30}{20}\right)^2 \times 450$ or $450 \div \left(\frac{20}{30}\right)^2$	1012.5	2	M1 for sight of correct SF^2 including 4:9 A1 1010 to 1013

Paper 5506

No	Working	Answer	Mark	Notes
13 (a)	$\frac{2x(x+20)}{2} = 400$	As given	2	M1 $\frac{2x(x+20)}{2}$ or $\frac{2x \times x + 20}{2}$ or $2x(x + 20) = 800$ A1 cao following correct working, no need for = 400 *SC B1* $2x \times x + \frac{1}{2} \times 2x(10 - \frac{x}{2}) \times 2$
(b)	$\frac{-20 \pm \sqrt{20^2 - 4 \times 1 \times (-400)}}{2}$ $= \frac{-20 \pm 44.721}{2}$	12.361	3	M1 for correct sub, up to signs, in the quad formula A1 for 44.7 or $\sqrt{2000}$ A1 for 12.3606 – 12.361, ignore negative solution T.I B3 for 12.361 OR Completing the square M1 for $(x+10)^2$ seen A1 for $-10 \pm \sqrt{500}$ A1 for 12.3606 – 12.361 ignore negative solution

Paper 5506

No	Working	Answer	Mark	Notes
14 (a)	$0.5\times 8\times 15\times \sin 70^0$	56.4	2	M1 for correct sub into area formula A1 56.38 – 56.4
(b)	$AB^2 = 8^2 + 15^2 - 2\times 8\times 15\times \cos 70^0 = 206.9$ EITHER $0.5\times AB\times CX$ ='56.38' OR $\frac{\sin B}{8} = \frac{\sin 70}{\sqrt{'206.9'}}$ $B = 31.5$ 15 sin '31.5'	7.84	4	M1 for correct sub into cos rule A1 for 206.9 - 207 or 14.38 – 14.4 EITHER M1 for use of area rule to find CX A1 7.83 – 7.84 OR M1 for correct use of sine rule to find sin B or sin A and then sight of 15sin B *or* 8 sin A A1 7.83 – 7.84

Paper 5506

No	Working	Answer	Mark	Notes
15 (a)	$4a^2 - 4a + 1 - (4b^2 - 4b + 1) =$	AG	3	Expansion Method M1 for a correct expansion of any one of the three terms M1(dep) on an attempt to expand all 3 terms and show LHS = RHS A1 fully correct algebra RHS exp is $4\left(a^2 + ab - a - ba - b^2 + b\right)$
	$4(a^2 - b^2) - 4(a - b)$ $4(a - b)(a + b - 1)$			
	OR $((2a - 1) - (2b - 1))((2a - 1) + (2b - 1))$ $(2a - 2b)(2a + 2b - 2)$			OR Factorisation Method M1 for attempt to use difference of 2 squares on LHS M1 for one bracket correctly simplified A1 fully correct
(b)	Any 2 odd square numbers have the above form If a and b are both even or odd then $a - b$ is even, so $4(a - b)$ is a multiple of 8 If one of a,b is odd, then $a + b - 1$ is even, so $4(a + b - 1)$ is a multiple of 8		3	B1 'any 2 square nos have the above form' (may be implied by sight of $(2a - 1)^2 - (2b - 1)^2$ in part (b)) B1 first reason B1 second reason SC B1 for $(2r + 1)^2 - (2r - 1)^2$ B1 for 8r
16 (a)	$g_L = \dfrac{2 \times 4.495}{1.35^2 \times \sin 30.5}$	9.719	4	B2 for any 4 of 4.505, 1.25, 29.5, 4.495, 1.35, 30.5 seen (B1 for any two or three seen) B1 for 11.710 – 11.7103
	$g_u = \dfrac{2 \times 4.505}{1.25^2 \times \sin 29.5}$	11.710		B1 cao 9.719 – 9.71904
(b)	Round, until lower and upper bounds agree	10	1	B1 for 10 + reason " they agree to this level of accuracy"

Paper 5506

No	Working	Answer	Mark	Notes
17 (a)(i)		xy	3	B1 cao
(ii)		y^2		B1 for y^2 or $y \times y$
(iii)		$\frac{x}{2}$		B1 for $\frac{x}{2}$ or $0.5x$ or $2^{-1}x$
(b)	Divide to get $2y = 1$	$q = -1$	2	M1 for $2y = 1$ or $\frac{x}{2} = 32$ or $p + q = 5$ or $1 + p + 2q = 5$
		$p = 6$		A1 cao
18 (a)	$x^2 - 2mx + m^2 - k$	$k = m^2$	2	M1 for correct exp of $(x - m)^2$ or correct completion of the square eg $\left(x - \frac{2m}{2}\right)^2 - \left(\frac{2m}{2}\right)^2$
				A1 cao
				SC B1 for $k = -m^2$
(b)(i)	Min value is $-m^2$	$-m^2$	3	M1 for recognition that min value occurs when $(x - m)^2 = 0$ (either (b)(i) or (b)(ii) correct implies this M1)
(ii)	$x = m$	m		A1 ft on 'k', " $-k$" gets M1 A0
				A1 cao
19	$0.06 \times 0.05 = 0.003$	No	2	M1 for 0.06×0.05
				A1 correct conclusion based on 0.003 or 0.06 x 0.05 stated as $\neq 0.0011$
				OR M1 for statement that for the two events to be independent P (BL and CL) = P(BL) × P(CL)

Paper 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
20		50 50 4	3	B1 50 or $\frac{100}{2}$ B1 for 50 or "a" B1 4 or $\frac{360}{90}$ oe

UG014129
51

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4LN

Telephone 01623 467467
Fax 01623 450481

Order Code UG014129 June 2003

For more information on Edexcel qualifications please contact our
Customer Response Centre on 0870 240 9800
or email: enquiries@edexcel.org.uk
or visit our website: www.edexcel.org.uk

London Qualifications Limited. Registered in England and Wales no.4496750
Registered Office: Stewart House, 32 Russell Square, London WC1B 5DN



edexcel

GCSE
Edexcel GCSE
Mathematics A 1387

July 2004

advancing learning, changing lives

Mark Scheme

Edexcel GCSE
Mathematics A 1387

London Qualifications is one of the leading examining and awarding bodies in the UK and throughout the world. It incorporates all the qualifications previously awarded under the Edexcel and BTEC brand. We provide a wide range of qualifications including general (academic), vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, our centres receive the support they need to help them deliver their education and training programmes to learners.

For further information please call Customer Services on 0870 240 9800, or visit our website at www.edexcel.org.uk

July 2004

Publications Code UG015616

All the material in this publication is copyright
© London Qualifications Ltd 2004

CONTENTS

Notes on Marking Principles....2
Paper 5514....4
Paper 5515....8
Paper 5516.... 11
Paper 5517.... 15
Paper 5518.... 19
Paper 5519.... 23

NOTES ON MARKING PRINCIPLES

NOTES ON MARKING PRINCIPLES

1 Three types of mark are available

M marks - awarded for correct working seen or implied.
A marks - conditional accuracy marks which are awarded for accurate working following the award of M marks.
B marks - unconditional accuracy marks (independent of M).

2 Abbreviations

cao - correct answer only.
ft - follow through.
$\surd$ - Denotes a "follow through" answer.

SC - special case.
isw - ignore subsequent working.
oe - or equivalent (and appropriate).
NB: a candidate cannot benefit from both isw and ft.

3 If no working shown, then

correct answers normally score full marks
incorrect (even though nearly correct) answers score no marks.

4 Marking instructions

Misread loses A marks (and sometimes B marks) on that part but ft can be allowed on subsequent parts. M marks can still be earned. If in doubt contact your team leader.
If there is a wrong answer in the answer space DO CHECK the working in the body of the script.

- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.
- If there is a wrong answer in the answer space DO CHECK the working in the body of the script.
- If it is clear from working that the "correct" answer has been obtained from incorrect working, award no marks. If in doubt contact your team leader.

5 Style of marking

Answer correct: tick and write part mark in margin NEXT TO BRACKETED MARK.
Answer incorrect: cross, but show M, A or B marks if any earned in body of script and transfer the total of these to the margin next to the bracket mark.
Total for each double page at bottom right page (except for back if used), FINAL TOTAL IN RELEVANT BOX ON FRONT COVER.
Nought in margin for fully incorrect question or page AND FOR NO ATTEMPT.
Where no attempt has been made a line should be put in the answer space and zero in the margin next to the bracketed mark
CHOICE OF METHOD – No marks unless one answer is in answer space – then mark that.
CROSSED OUT WORK - if not replaced this should be marked (if legible).
There must always be a mark next to bracketed mark in the margin/

6 Follow Through Marks

Follow throughs are guided by two principles:

(a) Follow throughs which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous, do not award.
(b) Follow throughs which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

7 Probability

Probability answers must be given as fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least two decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.

6 Linear Equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

8 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

PAPER 5501					
No		**Working**	**Answer**	**Mark**	**Notes**
1			K: 1020 L: 8.06	2	B1 for 1020 or 1,020 B1 for 8.06 Accept £8.06p and £8,06
2	(a)		28, 33	2	B1, B1 (B1 ft for "28" + 5 if both numbers >25)
	(b)			1	B1 for add 5, +5, for going up in 5's the difference is 5 oe
	(c)			1	B1 for they end in 3 and 8, it ends in 7; or refers to 383 and 388 are in sequence
3	(a)(i)		$\frac{1}{2}$	2	B2 for ½ accept half (B1 for an equivalent unsimplified fraction eg 4/8 or 50% or 0.5)
	(ii)		2 rectangles shaded	1	B1 for correct shading (any 2 rectangles)
	(b)(i)		40	4	B1 for 40 cao
	(ii)		150 000		B1 for 150 000 cao (accept 150,000 not 150.000)
	(iii)		6.55		B1 for 6.55 cao not $6.5^{½}$
	(iv)		$\frac{3}{8}$		B1 for $\frac{3}{8}$ oe accept 0.375
	(c)		Cross 3cm from *A*	1	B1 mark a cross 3 cm (±2 mm) from *A*
4	(a)(i)		metres	2	B1 for metres (m)
	(ii)		grams		B1 for grams (g)
	(b)		miles	1	B1 for miles
5	(a)		(3, 2)	1	B1 for (3, 2)
	(b)(i)		*Q* at (0, 3)	1	B1 for *Q* plotted correctly on *y*-axis at (0, 3) ±2mm
	(ii)		*R* at (−2, −3)	1	B1 for *R* plotted correctly at (−2, −3) ±2mm
6	(a)(i)		30	2	B1 for 30
	(ii)		25		B1 for 25
	(b)(i)		4 shapes	2	B1 for drawing 4 shapes
	(ii)		1½ shapes		B1 for drawing 1½ shapes

PAPER 5501					
	No	Working	Answer	Mark	Notes
7	(a)		30	1	B1 for 30
	(b)	$20 \times 3 + 5$	65	2	M1 for $20 \times 3 + 5$ oe A1 for 65
8	(a)		Draws perp.	1	B1 for correctly drawing perp must touch line or cut line *AB* ± 2mm
	(b)		Sketches a cylinder	1	B1 for sketching cylinder
9	(a)	1974 6580 8554	855.4	3	M1 for complete method with relative place value correct, condone 1 error in multiplication A2 cao (A1 for digits 8554 seen or A1 for "855.4" dependent on 1 arithmetic error)
	(b)	990 ÷ 69 = 14.3…. or 14 rem 24	14	4	Method 1 - Everything excluding long division M2 for a valid method with no errors …need to see 966 or 1035 (M1 for a valid method, including estimation, that reaches an answer between 901 and 1099 inclusive. Ignore errors.) A2 for 14 with a fully correct method (A1 (dep on 1st M1) for an answer between 14 and 15 inclusive. Ignore errors. Cannot have used estimation.) Note: Estimation can only score a maximum of M1 unless accompanied by another method or 966 or 1035 see (award M2) Method 2 – long division method M2 for 990 ÷ 69 = integer answer between 12 and 19 inclusive and rem 30 seen (M1 for 990 ÷ 69 with 1 in tens column.) A2 for 14 with fully correct method (A1 (dep on 1st M1) for an answer between 14 and 15 inclusive.)

PAPER 5501					
	No	Working	Answer	Mark	Notes
10	(a)(i)		7	2	B1 for 7
	(ii)		−10		B1 for −10
	(b)(i)		6	2	B1 for 6 (accept −6)
	(ii)		8		B1 for 8 (accept −8)
	(c)		−7	1	B1 for −7 cao
11	(a)		B and D	2	B1, B1 (–1 each extra)
	(b)(i)		A	2	B1 for A
	(ii)		3		B1 for 3
12	(a)	800 − 144	656	2	M1 for at least 1 digit correct and in correct position needs to be 3 digit number A1 for 656
	(b)	144 is less than 200 so Trudy is wrong		2	B1 for 200 B1ft for 'correct' explanation based on cand's "200"
	(c)	$\frac{45\times800}{100}$	360	2	M1 for 45 × 800 ÷ 100 oe A1 for 360
	(d)	$\frac{176}{800}\times100$	22	2	M1 for $\frac{176}{800}\times100$ oe A1 for 22
13	(a)(i)		25	4	B1 for 25
	(ii)		28		B1 for 28
	(iii)		5 and 20		B1 for 5 and 20
	(iv)		26 and 33		B1 for 26 and 33
	(b)	$2^3 = 8$ or $2^3 = 2 \times 2 \times 2$ (which is not 6)		1	B1 for valid 'explanation'
14			10 m^2	2	B1 for 10 B1 indep for m^2.

PAPER 5501					
No		**Working**	**Answer**	**Mark**	**Notes**
15	(a)	$\frac{11}{12}-\frac{10}{12}$	$\frac{1}{12}$	2	M1 for correctly writing both fractions to a common denominator A1 for $\frac{1}{12}$ oe
	(b)	$\frac{70\times400}{200}$	140	2	B2 for 140 (accept 136) (B1 for sight of any two of 70, 400 or 200)
16	(a)		$2y$	1	B1 for $2y$ or $2\times y$, $y2$, $y\times2$
	(b)		$3p^2$.	1	B1 for $3p^2$ or $3\times p^2$ or $p^2\times3$ or $p^2 3$
	(c)		$x(x-3)$	2	B2 for $x(x-3)$ (B1 for x (x…….)) $\frac{25}{60}$ wrongly cancelled gets B1 ISW
17	(a)		2 23 9 34 15 2 9 26 17 25 18 60	3	B3 for all correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
	(b)		$\frac{25}{60}$	1	B1 for $\frac{25}{60}$ or $\frac{5}{12}$ oe $\frac{25}{60}$ wrongly cancelled gets B1 ISW

PAPER 5501					
	No	**Working**	**Answer**	**Mark**	**Notes**
18	(a)(i)		60	2	B1 for 60
	(ii)				B1 for all **angles equal** so equilateral triangle oe
	(b)(i)		130	3	B1 for 130
	(ii)				B1 for isosceles triangle oe or 2 angles equal accept $\angle Q = \angle R$ B1 for angles on a straight **line** add up to **180°**oe (180° could be in working)
	(c)		64	1	B1 for 64
19	(a)		1	1	B1 for 1
	(b)		3	1	B1 for 3
	(c)	$9\times1+3\times2+5\times3+3\times4$ (= 42) "42" ÷ 20	2.1	3	M1 for 9×1, 3×2, 5×3, 3×4 or for 42 seen M1 (dep) for "42" ÷ 20 A1 for 2.1 or $2\frac{1}{10}$ or $2\frac{2}{20}$
20	(a)		09 05	1	B1 for 09 05 oe
	(b)		7	1	B1 for 7
	(c)		10	1	B1 for 10
	(d)	7km in 20 mins	21	3	B1 for 20 (minutes) oe or $\times3$ seen M1 for $\frac{"7"}{"20"}$ A1 cao [SC: "7" × 3 seen gets B1 M1]

PAPER 5501				
No	**Working**	**Answer**	**Mark**	**Notes**
21	Vertices of new triangle at (2, −2) (2, 6) and (6, −2)	Correct triangle drawn	3	B3 cao (B2 for either correct triangle in wrong translated position or for 2 vertices correct) (B1 a right angled triangle with horizontal length 4 or vertical length 8 in the same orientation as the shaded triangle) SC B2 for s + 3 totally correct
22 (a) (b) (c)	 8560 ÷ (10 × 10)	8 Pentagon 85.6	1 1 2	B1 for 8 B1 for pentagon M1 for 8560 ÷ (10 × 10) oe A1 for 85.6
23	$2x + x + 100 + 47 = 360$ $2x + x = 360 - 100 - 47$ $x = 71$ Largest angle = $2x$ =	142	4	M1 for $2x + x + 100 + 47 = 360$ or 360 – 147 or 213 seen M1 dep for correctly separating x-terms and non x terms or "360 – 147" ÷ 3 A1 for $x = 71$provided M2 awarded A1 ft for 142

PAPER 5502					
	No	**Working**	**Answer**	**Mark**	**Notes**
1	(a)		Diagram	1	B1 cao
	(b)		22, 26	1	B1 cao
	(c)		46	1	B1 cao
2	(a)	£5.95 + £1.62 + 25p + 2 × 48p = £8.78	1.22	3	M1 for 3 of £5.95, £1.62, 25p, 48p seen with intention to add or 170 seen A1 for £8.78 A1 ft from " 8.78"
	(b)	160 ÷ 12 = 13.(333…)	14	2	M1 for 160 ÷ 12, or 13 or better or 12 × 13 or 12 × 14 A1 14 cao
	(c)	"14" × £12.20 = £170.8**0**	170.80	2	M1 for "14" × 12.20 A1 for 170.8**0** or 158.6**0** if 13 in (b) NB: 170.8 or 158.6 gets M1 A0
3	(a)(i)		1459	1	B1 cao
	(ii)		9541	1	B1 cao
	(b)		9 + 5 = 14	1	B1 cao
	(c)		0	1	B1 cao
4	(a)		1123	1	B1 cao
	(b)		1125	1	B1 cao
	(c)		43	1	B1 cao
	(d)	67 – "43"	24	2	M1 for 67 – "43" A1 ft from "43"
	(e)(i)		80	2	B1 cao
	(ii)		56 – 58		B1 answer in range 56 to 58 inc
	(f)		Completes graph	2	B2 for (Apr , 70) and (May , 60 < p < 65) plotted and joined (B1 for one point plotted)

PAPER 5502					
	No	Working	Answer	Mark	Notes
5	(a)(i)		4	1	B1 cao
	(ii)		10	1	B1 cao
	(b)		28	2	B2 for 28 (B1 for 29 or 14×2)
6	(a)		$\frac{3}{4}$	1	B1 accept 0.75, 75%
	(b)	$\frac{3}{4} \times 28 = 21$	15.54	3	M1 for $\frac{3}{4} \times 28$ or 21 seen M1 for "21" $\times$ 74 or 28×74 or 7×74 A1 cao
	(c)(i)		44 – 46	2	B1 44 – 46 inc
	(ii)		5.2 – 5.4		B1 5.2 – 5.4 inc
7	(a)		Correct shape	1	B1
	(b)		Correct shape	1	B1
8	(a)		Hexagon	1	B1
	(b)		Right	1	B1 cao
	(c)		Drawing	2	B2 for 7 additional hexagons with at least 2 points at which 3 hexagons meet (B1 for one point at which 3 hexagons meet)
9	(a)		$\frac{3}{4}$	2	M1 $\frac{80-20}{80}$ or $\frac{60}{80}$ or 0.75 (oe) A1 cao Do not accept equivalents
	(b)		15	1	B1 cao
	(c)		Iron and Cook set	1	B1 for both Accept mixer and fryer
	(d)		Microwave	1	B1 cao
	(e)		Bar chart	2	B1 one column (mixer): 70, 90 B1 second column (fryer): 70, 85 [SC: B1 columns correct but reversed]

PAPER 5502					
No		**Working**	**Answer**	**Mark**	**Notes**
10	(a)		Acute	1	B1
	(b)		Reflex	1	B1
	(c)		reason	1	B1 reason eg 120 + 230 = 350 ≠ 360
11	(a)		$\frac{1}{2}$	1	B1 for $\frac{1}{2}$ oe
	(b)		reason	1	B1 for reason e.g could get 30 heads
12	(a)	7.5×4	30	2	M1 for 7.5×4 or $7\frac{1}{2} \times 4$ A1 cao
	(b)	$\sqrt{(42.25)}$	6.5	2	B2 for 6.5 (B1 for 42.25)
13	(a)		$6x$	1	B1 accept $6 \times x$, $x6$ oe
	(b)		$x - 4$	1	B1 cao
	(c)		$12(x - 4)$	1	B1 for $12(x - 4)$ or $12x - 48$ or $12 \times x - 48$ oe
	(d)		$18x - 48$	2	M1 for "$6x$" + "$12(x-4)$" or "$6x$" + "$12x - 48$" oe A1 ft $18x - 48$, $2(9x - 24)$, $3(6x - 16)$, $6(3x - 8)$
14	(a)		90	2	B2 for 90 ± 2 (B1 for 9 ± 0.2)
	(b)		317	1	B1 for 317 ± 4
15	(a)	Accurate drawing of triangle		2	B1 cao for 38° (±2°) B1 cao (indep) for BC drawn 7.3 cm (±2mm) and completing the triangle.
	(b)	Measure angle A	63°	1	B1 for 63° (±2°) or ft their diagram (±2°)

PAPER 5502					
	No	Working	Answer	Mark	Notes
16	(a)	0 × 1 = 0 1 × 1 = 1 2 × 4 = 8 3 × 8 = 24 4 × 8 = 32 5 × 7 = 35	100	3	M1 col 1 × col 2 (at least 3 shown) M1 (dep) sum of totals A1 cao SC: M1 M1 A0 for 101
	(b)	Either 35.50 ÷ 1.42 = 25; 26.99 – 25 = 1.99 Cheaper in the USA Or 26.99 × 1.42 = 38.33; 38.33 – 35.50 = 2.83 Cheaper in the USA		3	M1 35.50 ÷ 1.42 A1 25 seen B1 ft (dep on M1) £1.99 with conclusion (must have units) **OR** M1 26.99 × 1.42 A1 38.32 or 38.33… seen B1 ft (dep on M1) $2.83 or $2.82 with conclusion (must have units) must be to 2dp
17	(a)		-1, (1), (3), 5, 7, 9	2	B2 cao (B1 for 2 values)
	(b)		Graph	2	B1 ft for plotting points ±1/2 square B1 cao for line between $x = -2$ and $x = 3$
	(c)(i)		0.4	2	B1 for 0.4 or ft from single straight line with positive gradient
	(ii)		1.2		B1 for 1.2 or ft from single straight line with positive gradient

PAPER 5502					
	No	Working	Answer	Mark	Notes
18	(a)		Plots	1	B1 cao
	(b)		description	1	B1 dynamic relationship or "positive" (correlation)
	(c)		line of best fit	1	Line within overlay region, and to the extent of.
	(d)(i)	280 g →	reading	2	B1 ft from single straight line of positive gradient (±1/2 square)
	(ii)	120 pages →	reading		B1 ft from single straight line of positive gradient (±1/2 square)
19			200 150 225 150 10	3	B3 cao (B2 for three correct, or B1 for one correct)
20	(a)			2	B1 B1 for correct sketch – ignore additional internal lines B1 B0 for rectangle or right-angled triangle or correct shape without line NB 3-D sketch gets B0 B0
	(b)			2	B1 B1 for correct plan – ignore internal lines – accept a rotated plan, not reflected B1 B0 for a single rectangle drawn NB 3-D sketch gets B0 B0
21	(a)	163.25 – 35.50 = 127.75 127.75 ÷ 18.25 = 7	8	3	M1 163.25 – 35.50 (or sight of 127.75) M1 (dep) "127.75" ÷ 18.25 A1 cao SC: M2 for 7 days
	(b)	1.175 × £64 or $\frac{17.5}{100}$ × 64 or £11.20 seen	75.20	2	M1 1.175 × 64, 0.175 × 64 oe or 11.20 or 75.2 seen or 6.40, 3.20 and 1.60 seen A1 cao

PAPER 5502					
	No	Working	Answer	Mark	Notes
22	(a)		6	1	B1 cao
	(b)		$t^2 - 2t$	1	B1 oe
	(c)		$3(y - 4)$	1	B1 Accept $3 \times (y - 4)$ or $3(y - 4$
23	(a)		Reason	1	B1 Makes some mention of bias either directly or making reference to an insufficient or biased range of responses
	(b)		Reason	1	B1 (a) an insufficient range of responses (b) No mention of money (c) No time frame in the question (d) Misunderstanding of "A lot" and "Not much"

PAPER 5503					
No		**Working**	**Answer**	**Mark**	**Notes**
1			10.	1	B1 for 10
2	(a)(i)		$3e + 2f$	2	B1 for $3e + 2f$
	(ii)		$3p^2$		B1 for $3p^2$
	(b)	$5 \times -3 + 1$	-14	2	M1 for $5 \times -3 + 1$ A1 for -14
3		1974 $\frac{6580}{8554}$	855.4	3	M1 for complete correct method with relative place value correct, condone 1 error in multiplication. A2 cao (A1 for digits 8554 seen or A1 ft for "855.4" dependent on one arithmetic error only)
4	(a)		2 23 9 34 15 2 9 26 17 25 18 60	3	B3 all correct (B2 for 4 or 5 entries correct) (B1 for 2 or 3 entries correct)
	(b)		$\frac{25}{60}$	1	B1 for $\frac{25}{60}$ or $\frac{5}{12}$ oe
5	(a)	$\frac{45}{100} \times 800$	360	2	M1 for $\frac{45}{100} \times 800$ oe A1 for 360
	(b)	$\frac{176}{800} \times 100$	22	2	M1 for $\frac{176}{800} \times 100$ oe A1 for 22

PAPER 5503					
	No	Working	Answer	Mark	Notes
6	(a)	$\frac{11}{12}-\frac{10}{12}$	$\frac{1}{12}$	2	M1 for correctly writing both fractions to a common denominator A1 for $\frac{1}{12}$ oe
	(b)	$\frac{70\times400}{200}$	140	2	B2 for 140 (accept 136) (B1 for sight of any two of 70, 400 or 200)
7	(a)(i)		60	2	B1 for 60
	(ii)				B1 for all angles equal so equilateral triangle oe
	(b)(i)		130	3	B1 for 130
	(ii)				B1 for isosceles triangle oe B1 for angles on a straight line add up to 180°oe
	(c)		64	1	B1 for 64
8		$9\times1+3\times2+5\times3+3\times4$ (=42) "42" ÷ 20	2.1	3	M1 for 9×1, 3×2, 5×3, 3×4 or for 42 seen M1 (dep) "42" ÷20 A1 for 2.1 or $2\frac{1}{10}$ or $2\frac{2}{20}$
9	(a)		09 05	1	B1 for 09 05 oe
	(b)		7	1	B1 for 7
	(c)		10	1	B1 for 10
	(d)	7 km in 20 minutes	21	3	B1 for 20 (minutes) oe or $\times3$ seen M1 for $\frac{"7"}{"20"}$ A1 cao SC: "7" $\times3$ gets B1 M1

PAPER 5503					
No		**Working**	**Answer**	**Mark**	**Notes**
10		Vertices of new triangle at (2, −2) (2, 6) and (6, −2)	Correct triangle drawn	3	B3 cao (B2 for either correct triangle in wrong translated position or for 2 vertices correct) (B1 for a right-angled triangle with horizontal length 4 or vertical length 8 in the same orientation as shaded triangle) SC B2 for scale factor 3, totally correct
11		$2^2 + 3 =$	7 which is not even	2	B2 (B1 for correctly evaluating $n^2 + 3$ with a prime number value for n.)
12	(a)	360 ÷ 5	72	2	M1 for 360 ÷ 5 oe A1 for 72
	(b)	8560 ÷ (10 × 10)	85.6	2	M1 for 8560 ÷ (10 × 10) oe A1 for 85.6
	(c)(i)	Least length = 100.5	100.5	2	B1 for 100.5
	(ii)	Greatest length = 101.5	101.5		B1 for 101.5 or 101.499 or better
13		$2x + x + 100 + 47 = 360$ $2x + x = 360 - 100 - 47$ $x = 71$ Largest angle = $2x$ =	142	4	M1 for $2x + x + 100 + 47 = 360$ or 360 – 147 or 213 seen M1 (dep) for correctly separating x-terms and non x terms or $\frac{"360-147"}{3}$ A1 for $x = 71$ A1 ft (provided M2 awarded) for "142"
14	(a)(i)		2.21	2	B1 for 2.21
	(ii)		0.013		B1 for 0.013
	(b)	LCM = 3 × 13 × 17 = 3 × 221	663	2	M1 for 3 × 13 × 17 oe A1 for 663

PAPER 5503					
No		**Working**	**Answer**	**Mark**	**Notes**
15			Tick boxes 1,3 & 6	3	B1 for $\frac{\pi abc}{2d}$; B1 for $2a^2$; B1 for $2\left(c^2+d^2\right)$ (−B1 for each additional expression ticked (>3) to a minimum of 0)
16		200×0.2	40	2	M1 for 200×0.2 or $\frac{40}{200}$ seen A1 for 40
17	(a)	eg $2 \mid \underline{108}$ $2 \mid \underline{54}$ $3 \mid \underline{27}$ $3 \mid \underline{9}$ 3	$2^2 \times 3^3$	3	M2 for full systematic method of at least 4 divisions by prime numbers oe factor trees; condone 1 calculation error (M1 for 108 written as a correct product(s) with only one non-prime or equivalent division or a full process with 2 calculation errors.) A1 for $2^2 \times 3^3$ (accept $2 \times 2 \times 3 \times 3 \times 3$)
	(b)		12	1	B1 for 12
18		Perpendicular from P to intersecting arcs (within tramlines); perpendicular at least 2cm long		2	M1 relevant pair of arcs crossing within tramlines A1 SC M1 AO for a full construction of a perpendicular to AB not passing through P.
19		15×10	150 cm^3.	3	M1 for 15×10 A1 for 150 B1 for cm^3.

PAPER 5503					
No		**Working**	**Answer**	**Mark**	**Notes**
20			$y = 2x + 6$	2	B2 for $y = 2x + 6$ (B1 for $y = 2x + k$, $k \neq 1$ or for $y = mx + 6$, m $\neq 0$ or for $2x + 6$)
21		$6\frac{2}{5} = \frac{32}{5}$ Area of triangle = $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ (=2) Length of a side of sq. = $\sqrt{18 \times "2"}$ (=6) Perimeter of square = 4 × 6	24	5	B1 for $6\frac{2}{5} = \frac{32}{5}$ oe or $3\frac{1}{5} = \frac{16}{5}$ or $\frac{30}{8} + \frac{2}{8}$ oe (or implied by area of triangle=2) M1 for $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ oe M1 for area of square = 18 × product of two lengths A1 for $\sqrt{18 \times "2"}$ A1 for 24
22	(a)		$x(x-3)$	2	B2 for $x(x-3)$ (B1 for x (…….))
	(b)		k^3.	1	B1 for k^3.
	(c)(i)	$4x + 20 + 3x - 21$	$7x - 1$	4	M1 for **three** of 4 terms $4x + 20 + 3x - 21$ (or better) A1 for $7x - 1$
	(ii)	$x^2 + 3yx + 2xy + 6y^2$	$x^2 + 5xy + 6y^2$		M1 for **three** of 4 terms $x^2 + 3yx + 2xy + 6y^2$ A1 for $x^2 + 5xy + 6y^2$
	(d)		$(p+q)(p+q+5)$	1	B1 for $(p+q)(p+q+5)$

PAPER 5503					
	No	Working	Answer	Mark	Notes
23	(a)		32	1	B1 for 32 (accept 31.5 to 33.5 inclusive)
	(b)			3	B1 for ends of whiskers at 9 and 57, (with a box) B1 for ends of box at 16 and 45 (± 0.5) B1 for median marked at 32 or a complete box and whisker diagram drawn with a median inside the box
	(c)	Median(B)>Median(G); on average boys take longer IQR(B)>IQR(G); times for boys have a greater spread		2	B1 eg for comparison of medians B1 eg for comparison of (interquartile) ranges
24	(a)	$6y = 15 - 5x$	$y = \frac{15-5x}{6}$	2	M1 for either $6y = 15 - 5x$ or -6 $y = 5x - 15$ or for $\frac{5x}{6} + y = \frac{15}{6}$ or a correct ft on sign error to $y=$ A1 for $y = \frac{15-5x}{6}$ oe
	(b)	$6k + 5(-21) = 15$	20	2	M1 for substitution of $x = -21$ (or $x = 21$) into given equation or candidate's answer to (a) A1 for $k = 20$
	(c)(i)		Region R indicated	3	B2 correct region shaded (accept unshaded if R clear) (B1 shaded or R region satisfies 3 of the 4 given inequalities with same boundaries)
	(ii)		(1,1)		B1 for (1,1)

PAPER 5503					
	No	Working	Answer	Mark	Notes
25	(a)		60	1	B1 for 60 B1 for *BDC* = 25
	(b)	eg Angle *BDC* = 25 *ADB* = 60 − 25	35	2	[Award the mark for equivalent approaches] B1 ft for (a) − 25
	(c)	Ben is correct; angle *DAB* = 65 + 25 = 90 and since angle in a semi-circle is 90°, *BD* must be a diameter		1	B1 for full valid justification

PAPER 5504					
	No	Working	Answer	Mark	Notes
1	(a)	Accurate drawing of triangle		2	B1 cao for 38º (±2º) B1 cao (indep) for BC drawn 7.3 cm (±2mm) and completing the triangle.
	(b)	Measure angle A	63º	1	B1 for 63º (±2º) or ft their diagram (±2º)
2	(a)	0 × 1 = 0 1 × 1 = 1 2 × 4 = 8 3 × 8 = 24 4 × 8 = 32 5 × 7 = 35	100	3	M1 col 1 × col 2 (at least 3 shown) could be implied by answers. M1 (dep) sum of totals A1 cao SC: M1 M1 A0 for 101
	(b)	Either $35.50 ÷ 1.42 = £25; £26.99 - £25 = £1.99 Cheaper in the USA Or £26.99 × 1.42= $38.33; $38.33 - $35.50 = $2.83 Cheaper in the USA		3	M1 35.50 ÷ 1.42 A1 25 seen B1 ft (dep on M1) £1.99 with conclusion; units needed OR: M1 26.99 × 1.42 A1 38.33 or 38.32…seen B1 ft (dep on M1) $2.83 or $2.82 with conclusion; units needed
3	(a)		-1, (1), (3), 5, 7, 9	2	B2 cao (B1 for 2 values)
	(b)		Graph	2	B1 ft for plotting points ±1/2 square B1 cao for line between $x = -2$ and $x = 3$
	(c)(i)		0.4	2	B1 0.4 or ft single straight line with positive gradient
	(ii)		1.2		B1 1.2 or ft single straight line with positive gradient

PAPER 5504					
	No	Working	Answer	Mark	Notes
4	(a)			2	B1 Rectangle drawn B1 Triangle attached Ignore additional internal lines; 3D sketch gets B0
	(b)			2	B1 single Rectangle drawn B1 Attached smaller rectangle drawn Accept rotated plan, Ignore internal lines down; Nets get B0
5	(a)	£163.25 - £35.50 = £127.75 £127.75 ÷ £18.25 = 7	8	3	M1 163.25 – £35.50 (or sight of £127.75) M1 (dep) "127.75" ÷ 18.25 A1 cao SC: M2 for 7 days
	(b)	1.175 × £64 or $\frac{17.5}{100}$ × 64 or £11.20 seen	75.20	2	M1 1.175 × 64.00 0.175 × 64 oe or 11.20 or 75.2 seen OR 6.40, 3.20, and 1.60 seen A1 cao
	(c)	54 × £2.38 =	128.52	2	M1 54 × £2.38 A1 cao
6		Tessellation	Drawing	2	B2 seven additional hexagons, with at least 2 points at which 3 hexagons meet (B1 one point at which 3 hexagons meet).
7			200 150 225 150 10	3	B3 cao (B2 for three correct, or B1 for one correct)
8	(a)(i)	Bearing measured	70	2	B1 70 ± 4
	(ii)	Bearing measured	317		B1 317 ± 4
	(b)	Locus (circle) drawn & shaded		2	B1 circle ±2 mm centre Manchester B1 shading with accurate or approximate circle within tolerance

PAPER 5504					
	No	**Working**	**Answer**	**Mark**	**Notes**
9	(a)		Plots	1	B1 cao
	(b)		Description	1	B1 dynamic relationship or "positive" (correlation)
	(c)		Line of best fit	1	Line within overlay region, and to the extent of.
	(d)(i)	280 g →	Reading	2	B1 ft from single straight line of positive gradient (±1/2 square) B1 ft from single straight line of positive gradient (±1/2 square)
	(ii)	120 pages →	Reading		
10	(a)	$\frac{1}{6}, \frac{3}{8}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}$		2	B2 for all correct (B1 for 1 error or all correct but wrong order, or use of a common denominator decimals).
	(b)	3/5, 65%, 2/3, 0.72, 3/4		2	B2 for all correct (B1 for 1 error or all correct but wrong order or conversions to decimals oe)
11	(a)		182, 178, 180, 184	2	M1 mean of any three consecutive months, eg (147 + 161 + 238) ÷ 3 oe A1 cao
	(b)	Sale price = 80% Fun Friday price = 70% of 80% = 56% (oe)		2	B1 B1 for a fully correct explanation involving a worked example (oe) B1 a partially complete explanation

PAPER 5504					
	No	Working	Answer	Mark	Notes
12	(a)		Reason	1	B1 makes a mention of bias, either directly, or making reference to an insufficient or biased range of responses.
	(b)(i)		Reason	1	B1 make some mention of any one of (a) an insufficient range of responses (b) no mention of money (c) no time frame in the question (d) misunderstanding of "a lot" or "not much"
	(ii)		Question	2	B2 B1 for each of the following upto B2 (a) an improved question eg time frame made clear (b) response boxes (imperfect) (c) response boxes no errors OR For suggesting a generally improved question (a) a question clearly in the context of changes to the canteen (b) at least 3 boxes showing a full range of responses
13	(a)	$20y - 18y = 16 - 9$ oe $2y = 7$	$3\frac{1}{2}$ oe	3	M1 $20y - 18y = 16 - 9$ oe M1 $2y = 7$ A1 cao
	(b)	$40 - x = 3(4 + x)$ $40 - x = 12 + 3x$ $40 - 12 = x + 3x$ $4x = 28$	7	3	M1 multiplying through by 3: $3 \times \frac{40 - x}{3} = 3 \times 4 + 3 \times x$ A1 $40 - 12 = x + 3x$ A1 cao
14	(a)		$12(x\text{-}4)$	1	B1 for $12(x - 4)$ or $12x - 48$ or $12 \times x - 48$ oe $x = 12(x - 4)$ gets B0
	(b)		$18x\text{-}48$	2	M1 $6x +$ "$12(x - 4)$" or $6x +$ "$12x - 48$" A1 ft a linear expression $18x - 48$, $2(9x - 24)$, $3(6x - 16)$, $6(3x - 8)$

PAPER 5504					
No		**Working**	**Answer**	**Mark**	**Notes**
15		x $x^3 - 2x$ 4.1 60.7(21) 4.2 65.6(88) 4.3 70.9(07) Accept answer to 1 dp 4.22 66.7(114) or more 4.23 67.2(269) 4.24 67.7(45) 4.25 68.2(656)	4.2	4	B2 for trial $4.1 \leq x \leq 4.3$ evaluated (B1 for trial $4 < x < 5$ evaluated) B1 for different trial $4.225 \leq x \leq 4.25$ evaluated B1 (dep on at least one previous B1) for 4.2 cao
16	(a)		1.0×10^{-9}	1	B1 accept 1×10^{-9} or just 10^{-9}
	(b)	$1 \div 5 \times 10^{-9}$	2×10^{8}	2	M1 for $1 \div$ ("5×10^{-9}") or digit 2 with zeros only seen Condone omission of bracket for M1. A1 cao
17	(a)	$\frac{6.27 \times 4.52}{4.81 + 9.63} = \frac{28.3404}{14.44} = 1.962631579$	1.9626…	2	B2 for 1.9626… (B1 for 28.34 or 14.44)
	(b)		1.96	1	B1 ft from (a) as rounded to 1dp or 2dp. Do not accept 2, 2.00, but accept 2.0
18			$d = 4n + 6$	2	B2 $d = 4n + 6$ oe (B1 $d = 4n + k$, k an integer or 0, $4n + 6$, $n = 4n + 6$)
19		Vol = $\pi \times 3.8^2 \times 2.5 = \pi \times 14.44 \times 2.5$ = 45.36… × 2.5 = 113.411 Mass = "113" × 1.5 = 170.1165	170	4	M1 for $\pi \times r^2 \times 2.5$ where r = is 3.8 or 7.6 A1 if $r = 3.8$ M1 for "113" × 1.5 A1 for 169.5 – 170.3 cao

PAPER 5504					
	No	Working	Answer	Mark	Notes
20	(a)	$10^2 + 6^2$ or 136 $\sqrt{(100+36)}$ or $\sqrt{136} = 11.66\ldots$	11.7	3	M1 $10^2 + 6^2$ or 136 seen M1 (dep) $\sqrt{100+36}$ or $\sqrt{136}$ A1 11.66 – 11.7
	(b)	$\cos x = \frac{8}{10}$ or 0.8 $x = \cos^{-1} 0.8 = 36.869^\circ$	36.9	3	M1 for $\cos = \frac{8}{10}$, cos = 0.8 (oe) M1 (dep) for $\cos^{-1}$ (oe) A1 for 36.86 – 36.9
21		$18x - 6y = 99$ $\quad 12x - 4y = 66$ $8x + 6y = 18$ $\quad 12x + 9y = 27$ $26x = 117$ $\quad -13y = 39$	$x = 4\frac{1}{2}$ $y = -3$	4	M1 correct process to eliminate either x or y (condone one error) A1 cao for non – eliminated one. M1 (dep on 1st M1) for correct substitution of their found value. A1 cao (need both)
22		$\frac{133}{72} = 1.8472, \frac{160}{82} = 1.9512$ OR $\frac{72}{133} = 0.54135, \frac{82}{160} = 0.5125$ OR $\frac{160}{133} = 1.203...., \frac{82}{72} = 1.1388....$ OR $\frac{133}{160} = 0.83125...., \frac{72}{82} = 0.878$	1.84..≠ 1.95.. 1.20..≠ 1.13..	3	M1 for $\frac{133}{72}$ (= 1.8472…) oe Accept 1.8, 1.85 M1 for $\frac{160}{82}$ (=1.9512…) oe consistent pairing Accept 2.0, 1.9 **OR** M1 for $\frac{160}{133}$ (= 1.203…) oe M1 for $\frac{82}{72}$ (=1.1388) oe A1 for enough decimal places to show that the ratios are not equal; since the scale factors are different the shapes cannot be similar. NB Do Not need conclusion

PAPER 5504					
	No	Working	Answer	Mark	Notes
23	(a)		0.4, 0.6 0.4, 0.6, 0.4	2	B1 for LHS: (0.6), 0.4 B1 for RHS: 0.6, 0.4, 0.6, 0.4
	(b)	(30 × 42) – (25 × 42.8) = 1260 – 1070 = 190 190 ÷ 5 =	38	3	M1 for (30 × 42) – (25 × 42.8) or 1260 – 1070 or 190 seen M1 (dep) for "190" ÷ 5 A1 cao
24	(a)	£12000 × 0.25 = £3000; £12000 – £3000 = £9000 £9000 × 0.25 = £2250; £9000 – £2250 = £6750 £6750 × 0.25 = £1687.50; £6750 – £1687.50 =	£5062.50	3	M1 for 12000 × 0.75 (=9000) oe or £3000 or £23437.50 seen M1 (dep) for at least two further depreciation calculations (complete steps) A1 cao **OR** M2 for $12000 \times (0.75)^3$ or 5062.50 seen (M1 for $12000 \times (0.75)^n$, $n = 2$ or 4)
	(b)	0.8 × 0.8 × 0.8 × 0.8 (oe)	0.4096	2	M1 0.8^4 (oe) A1 cao

PAPER 5505					
	No	Working	Answer	Mark	Notes
1	(a)(i) (ii) (b)	 LCM = 3 × 13 × 17 = 3 × 221	2.21 0.013 663	2 2	B1 for 2.21 B1 for 0.013 M1 for 3 × 13 × 17 oe A1 for 663
2			Tick boxes 1,3 & 6	3	B1 for $\frac{\pi abc}{2d}$; B1 for $2a^2$; B1 for $2(c^2 + d^2)$ (−B1 for each additional expression ticked (>3) to a minimum of 0)
3	(a) (b)	200 × 0.2 0.2 + 0.4	40 0.6	2 2	M1 for 200 × 0.2 or $\frac{40}{200}$ seen A1 for 40 M1 for 0.2 + 0.4 A1 for 0.6
4	(a) (b)	eg 2 | 108 2 | 54 3 | 27 3 | 9 3 12	$2^2 \times 3^3$ 12	3 1	M2 for full systematic method of at least 4 divisions by prime numbers oe factor trees; condone 1 calculation error (M1 for 108 written as a correct product with only one non-prime or equivalent division or a full process with 2 calculation errors.) A1 for $2^2 \times 3^3$ (accept 2 × 2 × 3 × 3 × 3) B1 for 12
5		Perpendicular from P to intersecting arcs (within tramlines); perpendicular at least 2 cm long		2	M1 relevant pair of arcs crossing within tramlines A1 SC M1A0 for full construction of a line perpendicular to AB not through P
6		15 × 10	150 cm^3.	3	M1 for 15 × 10 A1 for 150 B1 for cm^3

PAPER 5505					
	No	Working	Answer	Mark	Notes
7	(a)		k^3.	1	B1 for k^3.
	(b)(i)	$4x + 20 + 3x - 21$	$7x - 1$	4	M1 for **three** of 4 terms $4x + 20 + 3x - 21$ (or better) A1 for $7x - 1$
	(ii)	$x^2 + 3yx + 2xy + 6y^2$	$x^2 + 5xy + 6y^2$		M1 for **three** of 4 terms $x^2 + 3yx + 2xy + 6y^2$ A1 for $x^2 + 5xy + 6y^2$
	(c)		$(p+q)(p+q+5)$	1	B1 for $(p+q)(p+q+5)$
	(d)		m^8	1	B1 for m^8.
	(e)		$6r^3t^6$	2	B2 for $6r^3t^6$ (B1 for $.....r^3t^6$ or for $6....t^6$)
8	(i)	Least length = 100.5	100.5	2	B1 for 100.5
	(ii)	Greatest length = 101.5	101.5		B1 for 101.5; ACCEPT 101.499 or better
9		$6\frac{2}{5} = \frac{32}{5}$ Area of triangle = $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ (= 2) Length of a side of sq. = $\sqrt{18 \times "2"}$ (=6) Perimeter of square = 4 × 6	24	5	B1 for $6\frac{2}{5} = \frac{32}{5}$ oe or $3\frac{1}{5} = \frac{16}{5}$ oe or $\frac{30}{8} + \frac{2}{8}$ oe (or implied by area of triangle=2) M1 for $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ oe M1 for (area of square) = 18 × product of two lengths A1 = $\sqrt{18 \times "2"}$ A1 for 24

PAPER 5505					
	No	Working	Answer	Mark	Notes
10	(a)	$6y = 15 - 5x$	$y = \frac{15-5x}{6}$	2	M1 for either $6y = 15 - 5x$ or for $\frac{5x}{6} + y = \frac{15}{6}$ or $-6y = 5x - 15$ or a correct ft on sign error to $y =$ A1 for $y = \frac{15-5x}{6}$ oe
	(b)	$6k + 5(-21) = 15$	20	2	M1 for subst. of $x = -21$ (or $x = 21$) into given eq^n or answer to (a) A1 for $k = 20$
	(c)(i)		Region R indicated	3	B2 correct region shaded (accept unshaded if R clear) (B1 shaded (R) region satisfies 3 of the 4 given inequalities with same boundaries)
	(ii)		(1,1)		B1 for (1,1)
11	(a)		$y = 2x + 6$	2	B2 for $y = 2x + 6$ (B1 for $y = 2x + k$, $k \neq 1$ or for $y = mx + 6$, $m \neq 0$ or for $2x+6$)
	(b)	Grad of $AB = 2$; Grad of $BC = \frac{-1}{"2"}$	$y = -\frac{1}{2}x + 6$	2	M1 for Grad of $BC = \frac{-1}{"2"}, -\frac{1}{2}$ or grad of $BC = -\frac{1}{\text{grad of AB}}$ A1 ft for $y = -\frac{1}{2}x + 6$ oe ft on wrong coeff. of x in (a)
	(c)	Eg A rectangle is always a cyclic quadrilateral because the opposite angles of a rectangle always add up to 180° (90+90)		1	B1 for valid explanation. (eg lines from the pt of int. of diagonals of rect to all 4 vertices are equal(radii))
12		Vertices of new triangle at $(-1, -4)$ $(-4, 2)$ and $(2, 2)$	Correct triangle drawn	3	B3 cao (B2 for 3 correct vertices no triangle or triangle with 2 correct vertices (B1 for triangle with two of six co-ordinates correct from using P as centre or any isosceles triangle with base 6 and height 6)

PAPER 5505					
	No	Working	Answer	Mark	Notes
13	(a)		32	1	B1 for 32 (accept 31.5 to 33.5 inclusive)
	(b)			3	B1 for ends of whiskers at 9 and 57 (with a box) B1 for ends of box at 16 and 45/46 ($\pm$ 0.5) B1for median marked at "32"or complete box and whisker diagram drawn with a median inside the box
	(c)	Median(B)>Median(G); on average boys take longer IQR(B)>IQR(G); times for boys have a greater spread		2	B1 eg for comparison of medians (ft on diagrams) B1 eg for comparison of (interquartile) ranges (ft on diagram)
14		Reading top to bottom frequencies are 20; 18; 45; 52		2	B2 all correct (B1 for one frequency correct)
15	(a)	$1600 = p\,q^0$.; $400 = pq^2$ $q^2 = 0.25$	$p = 1600$ $q = 0.5$	3	M1 for either $400 = pq^2$ or $1600 = pq^0$ B1 for $p = 1600$ A1 for $q = 0.5$
	(b)	$V = pq^{-2}$	6400	2	M1 for recognition that t = –2 A1 for 6 400

PAPER 5505					
No		**Working**	**Answer**	**Mark**	**Notes**
16	(a)	(I) $AB = AC$ (triangle ABC is isosceles) (II) PB = PC tangents (from a point to a circle are) equal (III) AP = AP (common side) so the 2 triangles are congruent ,SSS.		3	B3 for I, II, III with congruency reason (B2 for any two of I, II or III) (B1 for any one of the I, II or III)
	(b)	$BPC = 20°$ PBC (or PCB) = 90-1/2 "20" (= 80°) $BAC = PBC$ = "80"	50°	4	B4 for 50° (B3 for $BAC = 80°$) (B2 for $PBC = 80°$ or $PCB = 80°$) (B1 for $APC = 10°$ or $BPC = 20°$ or a middle angle = 90°) SC if clear numerical slip seen eg "PBC = 180 – 90 – 10 = 70" then goes on to get correct ft angle $ABC = 55$ deduct 1 from total so this cand would get B4–1 = B3
17	(a)	$\overrightarrow{OP} = \overrightarrow{OA} + \overrightarrow{AP}$ $= \overrightarrow{OA} + \frac{2}{3}(6\mathbf{c} - 6\mathbf{a})$ $= 6\mathbf{a} + 4\mathbf{c} - 4\mathbf{a}$	$2\mathbf{a} + 4\mathbf{c}$	3	M1 for $\overrightarrow{OP} = \overrightarrow{OA} + \overrightarrow{AP}$ or any correct vector journey involving $\overrightarrow{OP}$ M1 for $\overrightarrow{AP} = \frac{2}{3}(6\mathbf{c} - 6\mathbf{a})$ oe or $\overrightarrow{CP} = \frac{1}{3}(-6\mathbf{c} + 6\mathbf{a})$ oe or reverse vectors A1 for $2\mathbf{a} + 4\mathbf{c}$ oe (accept unsimplified)
	(b)	Eg $\overrightarrow{OM} = \overrightarrow{OC} + \overrightarrow{CM} = 6\mathbf{c} + 3\mathbf{a}$ $\overrightarrow{OM} = 1.5\overrightarrow{OP}$	$\overrightarrow{OM} = 1.5\overrightarrow{OP}$ so OPM is a straight line	2	B1 for $\overrightarrow{OM} = 6\mathbf{c} + \frac{1}{2}(6\mathbf{a})$ or $\overrightarrow{PM} = 2\mathbf{c} + \mathbf{a}$ unsimplified or reverse vectors B1 for a fully correct proof.

PAPER 5505					
	No	Working	Answer	Mark	Notes
18	(a)		4	1	B1 for 4 condone $\pm$ 4
	(b)		2	1	B1 for 2 condone ± 2
	(c)	$\sqrt{160} = 4\sqrt{10}$; $\left[\frac{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)-\sqrt{2}\times\sqrt{5}}{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)}\right]\times 100$ $\left[\frac{6\sqrt{10}-\sqrt{10}}{6\sqrt{10}}\right]\times 100$	$\frac{500}{6}$	4	B1 for **either** $\sqrt{160} = 4\sqrt{10}$ or $\sqrt{8} = 2\sqrt{2}$ **or** $\sqrt{20} = 2\sqrt{5}$ M1 for $\left[\frac{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)-\sqrt{2}\times\sqrt{5}}{\sqrt{8}\left(\sqrt{5}+\sqrt{20}\right)}\right]$ oe $(\times 100)$ B1 for either $6\sqrt{10} - \sqrt{10}$ or $6\sqrt{10}$ A1 for $\frac{500}{6}$ (accept 83.3 if no obvious earlier error)

PAPER 5505					
	No	Working	Answer	Mark	Notes
19	(a)(i)		$(2x-7)(x-14)$	3	M1 x^2 term and constant term (± 98) obtained or $2x(x-14) - 7(x-14)$ or $x(2x-7) - 14(2x-7)$ A1 for $(2x-7)(x-14)$
	(ii)		$x = \frac{7}{2}$; $x = 14$		B1ft ft (i) provided of form $(2x\pm a)(x\pm b)$
	(b)(i)		$\frac{7}{n+7}$	3	B1 for $\frac{7}{n+7}$ oe
	(ii)	$\frac{7}{n+7} = \frac{2}{5} \Rightarrow 2(n+7) = 5 \times 7$ $2n = 21$	n=10.5 is not possible since n has to be an integer		M1 for $2(n+7) = 5\times 7$ or $n+7=5\times 3.5$ (can be implied) ft (b)(i) fractional in terms of n and < 1 A1 ft for n=10.5 not possible (since n not integer) oe
	(c)	$2\times\left(\frac{n}{n+7}\right)\times\left(\frac{7}{n+7}\right) = \frac{4}{9}$ $14n\times 9 = 4(n+7)^2$ $14n\times 9 = 4(n^2+14n+49)$ $4n^2+56n+196-126n = 0$	$2n^2 - 35n + 98 = 0$	5	M1 for $\left(\frac{n}{n+7}\right)\times\left(\frac{7}{n+7}\right)$ seen M1 for $2\times\left(\frac{n}{n+7}\right)\times\left(\frac{7}{n+7}\right)$ oe $(=\frac{4}{9})$ M1(dep on 1st M) elimination of fractions within an equation B1 3 terms correct in expansion of $(n+7)^2 = n^2 + 7n + 7n + 49$ A1 full valid completion to printed answer

PAPER 5505					
	No	Working	Answer	Mark	Notes
	(d)	$\frac{7}{n+7}\times\frac{7}{n+7}=\frac{7}{21}\times\frac{7}{21}=$	$\frac{1}{9}$	2	M1 for $\frac{7}{n+7}\times\frac{7}{n+7}$ or better or ft [answer (b)(i)]^2 or $1-\frac{4}{9}-(\frac{n}{n+7})^2$ A1 for $\frac{1}{9}$ oe cao
20	(a)(i)		$y=1+\sin x$	2	B1 for $y=1+\sin x$
	(ii)		$y=2\sin x$		B1 for $y=2\sin x$ SC both (i) f(x) + 1, (ii) 2f(x) B1
	(b)	Stretch parallel to y-axis scale factor 3 Stretch parallel to x-axis scale factor $\frac{1}{2}$		3	M1 for 'stretch' A1 for Stretch parallel to y-axis scale factor 3 oe A1 for Stretch parallel to x-axis scale factor $\frac{1}{2}$ oe SC if M0 award BI for "sf 3 vertically" **and** "sf $\frac{1}{2}$ horizon."

PAPER 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
1		Question	2	B1 for each of the following upto B2 (a) an improved question eg time frame made clear (b) response boxes (imperfect) (c) response boxes no errors OR For suggesting a generally improved question (a) a question clearly in the context of changes to the canteen (b) at least 3 boxes showing a full range of responses
2	Locus (circle) drawn & shaded		2	B1 circle centre Manchester ±2mm B1 shading with accurate or approximate circle within tolerance
3 (a)		182, 178, 180, 184	2	M1 mean of any three consecutive months, eg (147 + 161 + 238) ÷ 3 oe A1 cao
(b)	Sale price = 80% Fun Friday price = 70% of 80% = 56% (oe)		2	B1 B1 for a fully correct explanation involving a worked example (oe) B1 a partially complete explanation
4	x $x^3 - 2x$ 4.1 60.7(21) 4.2 65.6(88) 4.3 70.9(07) 4.22 66.7(114) 4.23 67.2(269) 4.24 67.7(45) 4.25 68.2(656)	4.2	4	B2 for trial $4.1 \leq x \leq 4.3$ evaluated (B1 for trial $4 < x < 5$ evaluated) B1 for different trial $4.225 \leq x \leq 4.25$ evaluated B1 (dep on at least one previous B1) for 4.2 cao

PAPER 5506					
	No	Working	Answer	Mark	Notes
5	(a)		1.0×10^{-9}	1	B1 accept 1×10^{-9} or just 10^{-9}
	(b)	$1 \div (5 \times 10^{-9})$	2×10^{8}	2	M1 for $1 \div$ ("5×10^{-9}") or digit 2 with zeros only seen Condone omission of bracket for M1. A1 cao
6	(a)	$\frac{6.27 \times 4.52}{4.81 + 9.63} = \frac{28.3404}{14.44} = 1.962631579$	1.9626…	2	B2 for 1.9626… (B1 for 28.34… or 14.44)
	(b)		1.96	1	B1 ft from (a) as rounded to 1dp or 2dp. Do not accept 2, 2.00, but accept 2.0
7			$d = 4n + 6$	2	B2 $d = 4n + 6$ oe (B1 $d = 4n + k$, k an integer $\neq 6$, $4n + 6$, $n = 4n + 6$)
8		Vol = $\pi \times 3.8^2 \times 2.5 = \pi \times 14.44 \times 2.5$ $= 45.36\ldots \times 2.5 = 113.411$ Mass = "113" × 1.5 = 170.1165	170	4	M1 for $\pi \times r^2 \times 2.5$ where r = is 3.8 or 7.6 A1 if $r = 3.8$ M1 for "113" × 1.5 A1 for 169.5 – 170.3 cao

PAPER 5506					
	No	Working	Answer	Mark	Notes
9	(a)	$10^2 + 6^2$ or 136 $\sqrt{}(100 + 36)$ or $\sqrt{}136 = 11.66\ldots$	11.7	3	M1 for $10^2 + 6^2$ or 136 seen M1 (dep) $\sqrt{100+36}$ or $\sqrt{136}$ A1 11.66 – 11.7
	(b)	Cos x = 8/10 or 0.8 $x = \cos^{-1} 0.8 = 36.869^o$	36.9	3	M1 for cos = $\frac{8}{10}$, cos = 0.8(oe) M1 (dep) for $\cos^{-1}$ (oe) A1 for 36.86 – 36.9 M1 Use of sine rule and x found M1 for $x = 90 - \sin^{-1}$("0.8") A1 for 36.86 – 36.9
10		$18x - 6y = 99$ $\quad 12x - 4y = 66$ $8x + 6y = 18$ $\quad 12x + 9y = 27$ $26x = 117$ $\quad -13y = 39$	$x = 4\ ½$ $y = -3$	4	M1 correct process to eliminate either x or y (condone one error) A1 cao for non – eliminated one. M1 (dep on 1st M1) for correct substitution of their found value. A1 cao (need both)

PAPER 5506					
	No	Working	Answer	Mark	Notes
11		$\frac{133}{72} = 1.8472\,,\ \frac{160}{82} = 1.9512$ OR $\frac{72}{133} = 0.54135, \frac{82}{160} = 0.5125$ OR $\frac{160}{133} = 1.203...., \frac{82}{72} = 1.1388....$ OR $\frac{133}{160} = 0.83125...., \frac{72}{82} = 0.878$	1.84..≠ 1.95.. 1.20..≠ 1.13..	3	M1 for $\frac{133}{72}$ (= 1.8472…) oe Accept 1.8, 1.85 M1 for $\frac{160}{82}$ (=1.9512…) oe consistent pairing Accept 2.0, 1.9 **OR** M1 for $\frac{160}{133}$ (= 1.203…) oe M1 for $\frac{82}{72}$ (=1.1388) oe A1 for enough decimal places to show that the ratios are not equal; since the scale factors are different the shapes cannot be similar. NB Do Not need conclusion
12	(a)	£12000 × 0.25 = £3000; £12000 – £3000 = £9000 £9000 × 0.25 = £2250; £9000 – £2250 = £6750 £6750 × 0.25 = £1687.50; £6750 – £1687.50 =	£5062.50	3	M1 for 12000 × 0.75 (=9000) oe or £3000 or £23437.50 seen M1 (dep) for at least two further depreciation calculations (complete steps) A1 cao **OR** M2 for 12000 × $(0.75)^3$ or 5062.50 seen (M1 for 12000 × $(0.75)^n$, $n = 2$ or 4)
	(b)	0.8 × 0.8 × 0.8 × 0.8 (oe)	0.4096	2	M1 0.8^4 (oe) A1 cao

PAPER 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
13	$r^2 = \frac{3\times10}{\pi h} = \frac{3\times10}{\pi 1.5} = \frac{3\times10}{4.712..} = 6.36, 6.37$ $r = \sqrt{6.366\ldots}$	2.52-2.54	3	M1 for correct rearrangement to give $r^2 = \frac{3V}{\pi h}$ or $30 \div 4.712..$ or $6.36 - 6.37$ *Allow 0.3, 0.33 for* $\frac{1}{3}$ M1 (dep) for $\sqrt{}$ A1 cao 2.52 – 2.54
14 (a)		0.4 0.6,0.4, 0.6,0.4	2	B1 for LHS: (0.6), 0.4 B1 for RHS: 0.6, 0.4, 0.6, 0.4
(b)	0.6×0.6	0.36	2	M1 $0.6 \times$ "0.6" [0 < "0.6" < 1] A1 cao
(c)	$(30 \times 42) - (25 \times 42.8) = (1260 - 1070 = 190$ $190 \div 5 =$	38	3	M1 for $(30 \times 42) - (25 \times 42.8)$ or $1260 - 1070$ or 190 seen M1(dep) for "190" ÷ 5 A1 cao 38

PAPER 5506					
	No	Working	Answer	Mark	Notes
15	(a)	$S = \frac{k}{f^2};\ 125 = \frac{k}{8^2};\ k = 8000$	$S = \frac{8000}{f^2}$	3	M1 for $S = \frac{k}{f^2}$ M1 for $125 = \frac{k}{8^2}$ A1 cao These marks can be awarded if the full formula appears in part (b), rather than in part (a). [SC: $S = \frac{125}{64} f^2$ M1 M0 A0, $S = 1.95(3125) f^2$ M1 M0 A0 $S = \frac{1000}{f}$ M1 M0 A0]
	(b)		500	1	B1 cao

PAPER 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
16	$0.5 \times 3.2 \times 8.4 \times \text{Sin } B = 10$ sin $B = 0.74404\ldots$ 48.077 $AC^2 = 3.2^2 + 8.4^2 - 2 \times 3.2 \times 8.4 \times \cos B$ $AC^2 = 44.8815\ldots..$ $AC = 6.69\ (936\ldots)$ Perimeter = 18.3 Use the altitude AD, $\frac{h \times 8.4}{2} = 10 \Rightarrow h = (2.381)$ $BD = \sqrt{3.2^2 - h^2} = 2.139$ $DC = 6.261$ $AC = \sqrt{2.38^2 + '6.261'^2} = 6.69(936)$ Perimeter = 18.3	18.3	6	M1 for $0.5 \times 3.2 \times 8.4 \times \sin B$ (= 10) A1 for sin B = 0.74(404…) or B = 47.7 – 48.1 M1 for $3.2^2 + 8.4^2 - 2 \times 3.2 \times 8.4 \times \cos$ "48.077" M1(dep) for AC^2 = "44.8(815)"….. with correct order of evaluation A1 $AC = 6.69 - 6.7$ A1 18.29 – 18.3 **for** $\frac{h \times 8.4}{2} = 10 \Rightarrow h = (2.381)$ M1 for $BD^2 = 3.2^2 - "2.381"^2$ A1 $BD = 2.1 - 2.2$ M1 (dep) $AC^2 = "2.381"^2 + "6.261"^2$ A1 $AC = 6.69 - 6.7$ A1 18.29 – 18.3

PAPER 5506					
	No	Working	Answer	Mark	Notes
17	(a)	$(x + 8)^2 = x^2 + (x + 5)^2$ $x^2 + 16x + 64 = 2x^2 + 10x + 25$		4	B1 for angle $OTA = 90°$ (implied by use of Pythagoras with OA as hypotenuse) M1 $(x + 8)^2 = x^2 + (x + 5)^2$ oe M1 for correct squaring of x+8 or x+5 A1 for completion following correct working
	(b)	$(+6 + \sqrt{(6^2 - - 4\times39)})/2$ $= (6 \pm \sqrt{(36 + 156)})/2$ $= (6 \pm \sqrt{192})/2 = (6 + 13.856)/2$	9.93	3	M1 for substitution into quadratic formula, allow sign errors in b and c M1 for $x = (6 \pm \sqrt{192})/2$ [+ alone will do] A1 for 9.92 – 9.93
					M1 for $(x-3)^2-9-39$ M1 for $x = 3 \pm \sqrt{48}$ [+ alone will do it] A1 for 9.92 – 9.93 T&I= 9.93 gets M1,M0,A0
18		$EB = 60 \times \tan 30°$ $BD = \sqrt{(60^2 + 60^2)}$ $\tan BDE = 34.64 \div 84.85$	22.2°	4	M1 for $EB = 60\times\tan30$ M1 for $BD = \sqrt{(60^2 + 60^2)}$ M1 for $\tan BDE =$ "34.64" ÷ "84.85" A1 22.17 – 22.21
		$EB = 60 \times \tan 30°$ (= 34.64) $ED^2 = 60^2 + \left(\frac{60}{\cos 30}\right)^2$ $ED = \sqrt{8400} = (91.65)$ $\text{Angle} = \sin^{-1}\left(\frac{EB}{\sqrt{8400}}\right) = 22.2$			M1 for $EB = 60 \times \tan 30°$ oe M1 for fully correct method for ED M1 for $\sin BDE = \left(\frac{'34.84'}{'\sqrt{8400}'}\right)$ (oe) A1 22.17 – 22.21

PAPER 5506					
	No	Working	Answer	Mark	Notes
19	(a)	$x^2 + 6^2 = 25$ $x^2 = -11$		2	M1 substitutes y=6 to get x^2+36 = 25 A1 deduces x^2=25-36<0; impossible, (so line does not intersect curve)
	(b)	$y^2 = (2x - 2)^2$ $x^2 + (2x - 2)^2 = 25$ $x^2 + 4x^2 - 8x + 4 = 25$ $5x^2 - 8x + 4 - 25 = 0$ $5x^2 - 8x - 21 = 0$ $(5x + 7)(x - 3) = 0$ or $x = \frac{--8 \pm \sqrt{(-8)^2 - 4 \times 5 \times (-21)}}{10} = \frac{8 \pm \sqrt{484}}{10}$ So $x = 3$, or $x = -1.4$ Subs into either equation gives: $y = 4$, or $y = -4.8$	$x = 3$ $y = 4$ $x = -1.4$ $y = -4.8$	6	M1 $y^2=(2x-2)^2$ seen or implied A1 $x^2 + 4x^2 - 8x + 4 = 25$ (oe expanded form) M1 correct attempt to solve 3 term quadratic equation A1 $x = 3$, $x = -1.4$ M1 (dep on previous Ms) Sub one value of x into either equation gives: A1 $y = 4$, $y = -4.8$ NB: incorrectly matched pairs loses the final A1.

PAPER 5506					
	No	Working	Answer	Mark	Notes
20	(a)	400.005 ÷ 59.95 = 6.672310 6.672310 ÷ 1000 × 3600 = 24.02032 399.995 ÷ 60.05 = 6.661032 6.661032 ÷ 1000 × 3600 = 23.97972	24.020 23.980	5	B1 for 400.005 or 59.95 seen oe M1 for "400.005" ÷ "59.95" where "400.005" ∈ [400.005,400.5] and "59.95" ∈ [59.5,59.95] oe B1 for 399.995 or 60.05 seen oe M1 for "399.995" ÷ "60.05" where "399.995" ∈ [399.5,399.95] and "60.05" ∈ [60.05,60.5] oe A1 23.979-23.980 and 24.020-24.0204
	(b)	24.0 because to 1dp the answers are the same		1	B1cao for 24.0 with reason
	(c)	177 × 50/477 = 18.553 → 19 → 18 111 × 50/477 = 11.635 → 12 86 × 50/477 = 9.0147 → 9 82 × 50/477 = 8.595 → 9 21 × 50/477 = 2.201 → 2	18 12 9 9 2	3	M1 for $\frac{50}{"477"} \times 177$ or 111 or 86 or 82 or 21 A1 for all integers or better answers, at least 3 correct A1 cao
21	(a)	$40 - x = 3(4 + x)$ $40 - x = 12 + 3x$ $40 - 12 = x + 3x$ $4x = 28$	7	3	M1 multiplying through by 3: $3 \times \frac{40 - x}{3} = 3 \times 4 + 3 \times x$ A1 $40 - 12 = x + 3x$ A1 cao
	(b)	$\frac{2x(2x - 3)}{(2x - 3)(2x + 3)} = \frac{2x}{2x + 3}$	$\frac{2x}{2x + 3}$	3	B1 for $(2x-3)(2x+3)$ B1 for $2x(2x-3)$ or $(2x + 0)(2x + 3)$ B1 cao

PAPER 5506				
No	**Working**	**Answer**	**Mark**	**Notes**
22	Distance from x axis is y. Distance from (0, 2) is $\sqrt{(x^2+(y-2)^2)}$ $y^2 = x^2 + (y-2)^2$ $y^2 = x^2 + y^2 - 4y + 4$ $0 = x^2 - 4y + 4$ $4y = x^2 + 4$ and finish		4	B1 for $(x-0)^2 + (y-2)^2$ or $\sqrt{((x-0)^2+(y-2)^2)}$ oe seen B1 for $y = \sqrt{(x-0)^2+(y-2)^2}$ or $y^2 = (x-0)^2 + (y-2)^2$ oe B1 $(y-2)^2 = y^2 - 4y + 4$ seen B1 for $4y = x^2 + 4$ and finish

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone: 01623 467467
Fax: 01623 450481
Email: publications@linneydirect.com

Order Code UG015616 July 2004

For more information on Edexcel and BTEC qualifications please contact
Customer Services on 0870 240 9800
or http://enquiries.edexcel.org.uk
or visit our website: www.edexcel.org.uk

London Qualifications Limited, trading as Edexcel. Registered in England and Wales No. 4496750
Registered Office: Stewart House, 32 Russell Square, London WC1B 5DN



PMT

Mark Scheme (Results)

June 2011

Modular Mathematics (GCSE)
Unit 1: 5MB1H_01 (Higher)

PMT

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at www.edexcel.com.

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:
http://www.edexcel.com/Aboutus/contact-us/

June 2011
Publications Code UG028422
All the material in this publication is copyright
© Edexcel Ltd 2011

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labeling conventions.

ii*) select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect canceling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations
Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

12 Parts of questions
Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers
Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

Guidance on the use of codes within this mark scheme
M1 – method mark A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

5MB1H_01				
Question	**Working**	**Answer**	**Mark**	**Notes**
1 (i)	 OR 96 – 37 = 59 children can swim 18 + 21 = 39 children in Y5 or Y6 can swim 59 – 39	20	4	M1 for including 4 of the 6 pieces of information given in a clearly labeled two-way table A1 for 20 or 20 out of 96 or 20/96 OR M1 for a correct method that leads to the number of children in year 4 that can swim eg 96 – 37 (= 59) children can swim 18 + 21 = (39) children in Y5 or Y6 can swim '59' – '39' A1 for 20 or 20 out of 96 or 20/96
(ii)	 OR 20 + 11 = 31children in Y4 30 + 31 = 61 children in Y4 or Y6 96 – 61	35		M1 for one correct calculation leading to a 'new', piece of information in a clearly labeled two-way table A1 for 35 or 35 out of 96 or $\frac{35}{96}$ OR M1 for a correct method that leads to the total number of children in year 5 eg '20' + 11 (= 31) children in Y4 30 + '31' (=61) children in Y4 or Y6 96 – '61' children in Y5 A1 for 35 or 35 out of 96 or $\frac{35}{96}$ SC If M1 not earned then award B1 if ans(ii) = 55 – ans (i)

PMT

5MB1H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
2	(a)	See scatter graph	(12,14) and(17,20) plotted	1	B1 for correct plotting of both points
	(b)			1	B1 Positive (accept a correct relationship)
	(c)		16 to 18	2	B2 16 – 18 (M1 for a single line segment from m=11 to m = 16 within overlay or a vertical line drawn from m = 15 A1 for an answer in the range 16-18 or ft their line of best fit ± 2mm)

5MB1H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
3	(a)	$\frac{15}{100} \times 2400 = 360$ 2400 – 360 = 2040 2040 ÷ (1 + 2 + 5) = 255 255 × 1, 255 × 2, 255 × 5	Cement = 255 Sand = 510 Stone = 1275	4	M1 for $\frac{15}{100} \times 2400$ or $\frac{85}{100} \times 2400$ M1 for '2040' ÷ (1 + 2 + 5) M1 (dep on previous M1) for '255' × 1 or '255' × 2 or '255' × 5 either explicitly or with answer(s) on the answer line. A1 for all 3 correct masses
	(b)*	255 × 30 – 6500 = 1150 so not enough OR 6500 ÷ 255 = 25.5 m^3 worth of cement Which is less than 30 so not enough OR 6.5 tonnes of concrete gives 6.5 × 8 = 52 tonnes of dry mixture 30 m^3 of concrete requires 30 × 2400 × 0.85 kg of dry mixture = 61200kg of dry mixture (=61.2 tonnes)	No, with justification	3	B1 for a correct conversion between kg and tonnes e.g. 6.5 tonnes = 6500 kg (but not 1 tonne = 1000 kg) M1 for '255' × 30 C1(dep on M1) for an answer of No (Yes), supported by (ft) calculations. OR B1 for a correct conversion between kg and tonnes e.g. 6.5 tonnes = 6500 kg (but not 1 tonne = 1000 kg) M1 for 6500 ÷ '255' C1(dep on M1) for an answer of No (Yes), supported by (ft) calculations OR M1 for 6.5 × 8 and 30 × 2400 × 0.85 B1 for a correct between kg and tonnes e.g. 61200 kg = 61.2 kg (but not 1 tonne = 1000 kg) C1 (dep on M1) for an answer of No (Yes), supported by (ft) calculations.

PMT

5MB1H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
4	(a)		14 | 8 15 | 2 4 16 | 0 2 4 4 8 17 | 0 3 4 9 14 | 8 = 148 cm	3	B2 for a fully correct ordered diagram (B1 for correct unordered diagram or ordered with at most two errors) B1 for a correct key eg 14 | 8 = 148 cm (cm not required)
	(b) *	Boy's Median = 170 Girl's Media = 164 Boy's Mean = 170(.38) Girl's Mean= 164 Boy's Range = 27 Girl's Range = 31 Boy's IQR = 14 Girl's IQR = 14.5 – 15.25 15 | 7 9 16 | 2 6 8 9 17 | 0 3 4 6 6 18 | 1 4	Compares: medians/means + Range + Spread	3	A maximum 2B marks from: B1 for a correct mean or median for either the boys or the girls. B1 for a correct measure of spread for either the boys or the girls. B1 for a correct stem and leaf diagram drawn for the boys (no need for a key) C1 for any correct comparison based on 2 medians(ft) or 2 means(ft) or 2 ranges(ft) or 2 IQRs (ft) or a correct statement following from comparing the correct stem and leaf diagrams. This comparison must be a general statement which compares the boys' heights with the girls' heights.
5	(a)	$1 – 0.3 – 0.1 – x$	$0.6 – x$	2	M1 for $0.3 + 0.1 + x + P(2) = 1$ oe A1 $0.6– x$ or $\frac{6}{10} - x$ or $1 - (0.4 + x)$ or $1 - 0.3 - 0.1 - x$
	(b)	$0.3 + 0.1$	0.4	1	B1 cao
	(c)		$300x$	1	B1 for $300x$, $x \times 300$ oe

PMT

5MB1H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
6	(a)		Overlapping response boxes Leading/biased question Age too personal Missing units	2	B2 for any two of: overlapping response boxes, too personal to ask person's age, a leading or biased question, no units (B1 – just one of the above).
	(b)		How many pieces of fruit do you eat each day? 0 to2, 3 to 4, over 4	2	B1 for a sensible question including a time period B1 for at least 3 response boxes. Any pairs of response boxes must not overlap
	(c)	$\frac{73}{536+384+48+73} \times 100$	7	2	M1 for $\frac{73}{536+384+48+73} \times 100$ Or $100 \div \left\langle \frac{536+384+48+73}{73} \right\rangle$ Or 7.01… A1 for 7
7		1200×1.035^3 Or 1200 × 1.035 = 1242 1242 × 1.035 = 1285.47 1285.47× 1.035 = 1330.46	1330.46	3	M2 for 1200×1.035^3 A1 1330.46 – 1330.47 Or M1 1200 × 1.035 M1(dep) for ' 1242'× 1.035 and '1285.47'× 1.035 A1 1330.46 – 1330.47 [SC: B1 for 42 or 84 or 126 or 1242 or 1284 or 1326 seen, if M0 scored]

5MB1H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
8	(a)		$25 \leq t < 30$	1	B1 for the interval $25 \leq t < 30$ described unambiguously
	(b)	12.5×6 + 17.5×4 + 22.5×24 + 27.5×44 + 32.5×10 + 37.5×4 =75+70+540+1210+325+150 = 2370 2370 ÷ 92	25.8	4	M1 for finding at least 4 products fx consistently within interval (including end points) M1 (dep) for use of at least 4 correct midpoints M1 (dep on first M) for $\frac{\Sigma fx}{\Sigma f}$ A1 25.76 – 25.8
	(c)	Cf table: 6, 10, 34, 78, 88, 92 Cf graph	Correct CF graph	3	B1 Correct cumulative frequencies (may be implied by correct heights on the grid) M1 for at least 5 of "6 points" plotted consistently within each interval A1 for a fully correct cf graph.
	(d)	Median at the (92+1)÷2 = 46.5 day temp IQR = UQ – LQ = 69.75th day temp – 23.25th day temp	Median = 25.4 IQR = 6.7	3	B1 for a median in the range 24.5−26.5 or B1 ft read off at CF = 46(.5) from a CF graph tol $\frac{1}{2}$ square M1 for reading off temps at CF= 69 or 69.75 (±0.5) and 23 or 23.25(±0.5) from a CF graph and subtracting A1 ft

PMT

5MB1H_01					
Question		**Working**	**Answer**	**Mark**	**Notes**
9		Gradient $= \frac{68-32}{20}$	$m = 1.8$, $k = 32$	3	M1 for attempting to find the gradient of the line eg from height ÷ base of a triangle with the line as hypotenuse A1 for $m = 1.7 - 1.9$ B1 for $k = 31$ - 33
10	(a)		0.8 0.6 0.2 0.4 0.5 0.5	2	B2 for all 6 correct probabilities (B1 for two correct probabilities)
	(b)	0.6 × 0.8	0.48	2	M1 for '0.6' × '0.8' ft probability tree diagram A1 cao
	(c)	1 – (0.4 × 0.5) 1 – 0.2 OR 0.6 × 0.8 + 0.6 × 0.2 + 0.4 × 0.5	0.8	3	M2 for 1 – '0.4' × '0.5' ft probability tree diagram A1 cao OR M2 for '0.6' × '0.8' + '0.6' × '0.2' + '0.4' × '0.5' (M1 for any two of '0.6' × '0.8', '0.6' × '0.2', '0.4' × '0.5'added) A1 cao

PMT

5MB1H_01				
Question	**Working**	**Answer**	**Mark**	**Notes**
11 (a)		Histogram Fd axis labelled say 0 to 8 5 – 10, fd = 1.6 10 – 12, fd = 7.5 12 – 16, fd = 6 16 – 20, fd = 4 20 – 25, fd = 2	3	M1 for fd axis labeled and 1 correct block or for one correct division eg 8 ÷ 5 or 1.6 A1 for at least 2 additional correct blocks or 2 additional frequency densities calculated A1 for completely correct histogram SC B2 for a fully correct but unlabeled histogram OR M1 for area key and 1 correct block A1 for at least 2 additional correct blocks A1 for completely correct histogram SC B2 for a fully correct but unlabeled histogram
(b)		32	2	M1 for 6 × 1 + 16 + 10 A1 cao

PMT

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481
Email publication.orders@edexcel.com

Order Code UG028422 June 2011

For more information on Edexcel qualifications, please visit
www.edexcel.com/quals

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual

Llywodraeth Cynulliad Cymru
Welsh Assembly Government

Rewarding Learning

PMT

edexcel

Mark Scheme (Results)

Summer 2012

GCSE Mathematics (Linear) 1MA0
Foundation (Non-Calculator) Paper 1F

PMT

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at www.edexcel.com.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2012
Publications Code UG032619
All the material in this publication is copyright
© Pearson Education Ltd 2012

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labeling conventions.

ii) *select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate*.
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect canceling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.
Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

12 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers

Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

Guidance on the use of codes within this mark scheme
M1 – method mark A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
1	(a)		380	1	B1 cao
	(b)		6.2	1	B1 cao
	(c)		Arrow at 34	1	B1 cao
2	(a)		8	1	B1 for 8 ± 0.2
	(b)		35	1	B1 for 35 ± 2°
	(c)		Circle drawn	1	B1 for all parts within ± 2mm, (use overlay)
3	(a)		4, 7, 4, 3, 2	2	M1 for at least 3 correct tallies **or** at least 3 correct frequencies A1 for all frequencies correct
	(b)		7	1	B1 for 7 **or** ft from frequencies in (a) or tallies if no frequencies
	(c)		Diagram drawn	3	M1 for bar chart or other suitable chart with at least 3 correct heights for their scale (can f.t.) A1 for all 5 bars correctly labelled and vertical axis correctly scaled A1 for fully correct **or** ft frequencies in (a) **OR** M1 for pictogram with at least 3 correct rows (can f.t.) A1 for correct labels on all 5 rows and correctly key A1 for fully correct **or** ft frequencies in (a) **OR** M1 for pie chart with at least 3 correct sectors ± 2° (can f.t.) A1 for all 5 sectors correctly labelled A1 for fully correct **or** ft frequencies in (a)

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
4		£1.18 + 94p = £2.12 £5 – £2.12 – 30p = £2.58 £2.58 ÷ 2 =	1.29	3	M2 for (5 – 1.18 – 0.94 – 0.30) ÷ 2 oe **or** digits 129 (M1 for 1.18 + 0.94 **or** 2.12 seen **or** 1.18 + 0.94 + 0.30 oe **or** 2.42 seen **or** 5 – 1.18 – 0.94 oe **or** 2.88 seen **or** (5 – 1.18 – 0.94) ÷ 2 **or** 1.44 seen **or** 5 – 1.18 – 0.94 – 0.30 oe **or** 2.58 seen) A1 cao **NOTE**: Accept working in £ or pence
5	(a)(i)		(2, 3)	2	B1 cao
	(ii)		(–3, 1)		B1 cao
	(b)		Point plotted at (3, –4)	1	B1 cao
6	(a)		– 5	1	B1 cao
	(b)		6	1	B1 for 6 or – 6
	(c)		3	1	B1 cao
7			(P, B), (P, S), (P, L) (M, B), (M, S), (M, L) (H, B), (H, S), (H, L)	2	M1 for any 3 combinations with no incorrect combinations A1 for all 9 combinations with no duplicates or extras
8	(a)		Walk	1	B1 cao
	(b)	24 ÷ 4 =	6	2	M1 for 24 ÷ 4 oe or ¼ oe seen A1 cao

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
9	(a)		Isosceles triangle	1	B1 for isosceles triangle
	(b)		Rectangle with area 12 cm^2	2	M1 for rectangle drawn A1 cao
10	(a)		A marked at 0	1	B1 for A marked at 0 (within overlay)
	(b)		B marked at 1/4	1	B1 for B marked at 1/4 (within overlay)
11	(a)		9	1	B1 cao
	(b)		33	2	M1 for 5 × 5 **or** 25 seen in the working **or** 2 × 2 × 2 **or** 8 seen in the working A1 cao
12	(a)		20	2	M1 3 × 3 × 3 oe seen or drawn or 27 seen or use of 3 layers A1 cao
	(b)			2	B2 for correct view (B1 for  or )
13	(a)(i)		07 29	2	B1 for 07 29
	(ii)		36		B1 for 36 or ft difference between (i) and 06 53
	(b)		07 51	1	B1 cao
	(c)		09 55	1	B1 for 09 55 or 9 55 or five to ten

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
14		2 + 8 + 2 + 8 = 20 20 ÷ 4 =	5	4	M2 for 2 + 8 + 2 + 8 oe **or** 20 seen **or** (2 + 8) ÷ 2 oe (M1 for the sum of 3 sides of the rectangle) M1 (dep) for the sum of 3 or 4 sides of the rectangle ÷ 4 **or** an attempt to evaluate (2 + 8) ÷ 2 oe to get the length of one side A1 cao **SC**: B1 for an answer of 4 coming from $\sqrt{2\times 8}$ oe
15	(a)		4	1	B1 cao
	(b)	9.5 – 4.75 = **OR** 9.5 ÷ 2 =	4.75	2	M1 for 9.5 – 4.75 **or** 9.5 ÷ 2 **or** 4.75 – 9.5 A1 cao
	(c)		6	1	B1 cao
	(d)	12 × 4 =	48	2	M1 for × 4 seen **or** identifying + 0.5 for every 2 inches **or** 12 + 12 + 12 + 12 oe **or** build up method eg 12, 24, 36, 48 allow one error A1 cao
16	(a)		trapezium	1	B1 for trapezium or isosceles trapezium
	(b)			2	B2 for correct tessellation (at least 5 more shapes) (B1 for at least 4 shapes (including initial shape) correctly tessellating)

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
17*		S: 35 ÷ 100 × 40 = 14 W: 40 ÷ 8 × 3 = 15 **OR** D: 16 ÷ 40 (× 100) = 0.4 (40%) W: 3 ÷ 8 (× 100) = 0.375 (37.5%) **OR** D: $\frac{16}{40} = \frac{80}{200}$ S: $\frac{35}{100} = \frac{70}{200}$ W: $\frac{3}{8} = \frac{75}{200}$	Debbie and correct calculations	4	**Compares Marks out of 40 or fractions with denominator of 40** M1 for 35 ÷ 100 × 40 oe **or** 14 seen (or 14/40 seen) M1 for 40 ÷ 8 × 3 **or** 15 seen (or 15/40 seen) A1 for 14 and 15 **or** $\frac{14}{40}$ and $\frac{15}{40}$ C1 (dep on M1) for correct conclusion for their working QWC with 3 comparable marks: Decision and justification should be clear with working clearly presented and attributable. **OR Decimals (or Percentages)** M1 for 16 ÷ 40 (× 100) oe **or** 0.4 (or 40) seen M1 for 3 ÷ 8 (× 100) oe **or** 0.375 (or 37.5) seen A1 for 0.4 and 0.375 (**or** 40 and 37.5) C1 (dep on M1) for correct conclusion for their working QWC: with 3 comparable decimals (or percentages: Decision and justification should be clear with working clearly presented and attributable. **OR Compares Fractions with denominator other than 40** M1 for attempt to convert all to fractions with a common denominator other than 40 M1 for at least 1 correct A1 for $\frac{80}{200}$ and $\frac{70}{200}$ and $\frac{75}{200}$ oe C1 (dep on M1) for correct conclusion for their working QWC with 3 comparable fractions: Decision and justification should be clear with working clearly presented and attributable.

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
18*	(a)		10	1	B1 cao
	(b)	 **Miles** 0 10 20 30 40 50 **Ed** 0 15 30 45 60 75 **Bill** 10 20 30 40 50 60	Ed is cheaper up to 20 miles, Bill is cheaper for more than 20 miles	3	M1 for correct line for Ed intersecting at (20,30) ±1 sq tolerance **or** $10 + x = 1.5x$ oe C2 (dep on M1) for a correct full statement ft from graph eg. Ed cheaper up to 20 miles **and** Bill cheaper for more than 20 miles (C1 (dep on M1) for a correct conclusion ft from graph eg. cheaper at 10 miles with Ed ; eg. cheaper at 50 miles with Bill eg. same cost at 20 miles; eg for £5 go further with Bill **or** A general statement covering short **and** long distances eg. Ed is cheaper for shorter distances **and** Bill is cheaper for long distances) **OR** (continued on next page)

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
18 (contd)					**OR** M1 for correct method to work out Ed's delivery cost for at least 2 values of n miles where $0 < n \leq 50$ **or** for correct method to work out Ed and Bill's delivery cost for n miles where $0 < n \leq 50$ C2 (dep on M1) for 20 miles linked with £30 for Ed and Bill with correct full statement eg. Ed cheaper up to 20 miles **and** Bill cheaper for more than 20 miles (C1 (dep on M1) for a correct conclusion eg. cheaper at 10 miles with Ed; eg. cheaper at 50 miles with Bill eg. same cost at 20 miles; eg for £5 go further with Bill **or** A general statement covering short **and** long distances eg. Ed is cheaper for shorter distances **and** Bill is cheaper for long distances) SC: B1 for correct full statement seen with no working eg. Ed cheaper up to 20 miles **and** Bill cheaper for more than 20 miles QWC Decision and justification should be clear with working clearly presented and attributable

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
19		1,96 × 2.25 = 4.41 **OR** 4.23 ÷ 9 = 0.47 1.96 ÷ 4 = 0.49 **OR** 4.23 × 4 = 16.92 1.96 × 9 = 17.64 **OR** 4.23 ÷ 9 = 0.47 0.47 × 4 = 1.88 **OR** 1.96 ÷ 4 = 0.49 0.49 × 9 = 4.41 **OR** 9 ÷ 4.23 = 2.12 4 ÷ 1.96 = 2.04	Pack of 9	3	M2 for a fully correct method to enable a conclusion eg 1.96 × 2¼ **OR** M1 for 4.23 ÷ 9 or 423 ÷ 9 **or** 0.47 seen **or** 47 seen M1 for 1.96 ÷ 4 or 196 ÷ 4 or 0.49 seen or 49 seen **OR** M1 for 4.23 × 4 **or** 423 × 4 **or** 16.92 seen **or** 1692 seen M1 for 1.96 × 9 **or** 196 × 9 **or** 17.64 seen **or** 1764 seen **OR** M1 for 4.23 ÷ 9 **or** 423 ÷ 9 **or** 0.47 seen **or** 47 seen M1 for 0.47 × 4 **or** 47 × 4 **or** 1.88 seen **or** 188 seen **OR** M1 for 1.96 ÷ 4 **or** 196 ÷ 4 **or** 0.49 seen **or** 49 seen M1 for 0.49 × 9 **or** 49 × 9 **or** 4.41 seen **or** 441 seen **OR** M1 for 9 ÷ 4.23 **or** 2.12(...) seen **or** 2.13 seen M1 for 4 ÷ 1.96 **or** 2.04(...) seen A1 for Pack of 9 and fully correct calculations **NOTE:** B0 for an answer of 9 not supported by working.
20	(a)		6	1	B1 cao
	(b)		44	1	B1 cao
	(c)		31	2	M1 for 60 – 29 **or** 29 – 60 **or** any correct method that is attempting to find the difference between 29 and 60 (allow 1 arithmetic error) A1 cao

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
21*		Angle *DBC* = (180 – 50) ÷ 2 Base angles of isosceles triangle are equal Angle *ABD* = 180 – 65 Angles on a straight line add up to 180 *x* = 180 – 20 – 115 Angles in a triangle add up to 180 **OR** Angle *DBC* = (180 – 50) ÷ 2 Base angles of isosceles triangle are equal *x* = 65 – 20 Exterior angle of triangle is equal to sum of interior opposite angles **OR** Angle *DCB* = (180 – 50) ÷ 2 Base angles of isosceles triangle are equal *x* = 180 – 50 – 20 – 65 Angles in a triangle add up to 180	45 with reasons	4	M1 for (180 – 50) ÷ 2 oe **or** 65 seen M1 for 180 – 20 – (180 – "65") **or** "65" – 20 or 180 – 50 – 20 –'65' oe C2 for *x* identified as 45 with full reasons QWC: Reasons clearly laid out with correct geometrical language used (C1 (dep on M1) for one reason QWC: Reasons clearly laid out with correct geometrical language used) NOTE: *x* = 45 with no working or without any correct angles marked on the diagram cannot score.

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
22	(a)	360 ÷ 60 = 6 300 ÷ 60 = 5 6 × 5 =	Yes and 30	3	M1 for dividing side of patio by side of paving slab eg 360 ÷ 60 **or** 300 ÷ 60 **or** 3.6 ÷ 0.6 **or** 3 ÷ 0.6 **or** 6 and 5 seen **or** 6 divisions seen on length of diagram **or** 5 divisions seen on width of diagram M1 for correct method to find number of paving slabs eg (360 ÷ 60) × (300 ÷ 60) oe **or** 6 × 5 **or** 30 squares seen on diagram (units may not be consistent) A1 for Yes **and** 30 (or 2 extra) with correct calculations **OR** M1 for correct method to find area of patio **or** paving slab eg 360 × 300 **or** 108000 seen **or** 60 × 60 **or** 3600 seen **or** 3.6 × 3 **or** 10.8 seen **or** 0.6 × 0.6 **or** 0.36 seen M1 for dividing area of patio by area of a paving slab eg. (3.6 × 3) ÷ (0.6 × 0.6) oe (units may not be consistent) A1 for Yes **and** 30 (or 2 extra) with correct calculations **OR** M1 for method to find area of patio and area of 32 slabs eg. 60 × 60 × 32 **or** 360 × 300 M1 for method to find both areas eg. 60 × 60 × 32 **and** 360 × 300 (units may not be consistent) A1 for Yes **and** 115200 **and** 108000 **OR** Yes **and** 11.52 **and** 10.8 **NB :** Throughout the question, candidates could be working in metres or centimetres

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
22	(b)	1726 25890 27616 **8 6 3** **2** 2 4 1 8 9 **3** **7** 1 6 1 2 6 **2** 6 1 6 800 60 3 30 24000 1800 90 2 1600 120 6 24000 + 1800 + 90 + 1600 + 120 + 6 = 27616	276.16	3	M1 for complete correct method with relative place value correct. Condone 1 multiplication error, addition not necessary. **OR** M1 for a complete grid. Condone 1 multiplication error, addition not necessary. **OR** M1 for sight of a complete partitioning method, condone 1 multiplication error. Final addition not necessary. A1 for digits 27616 A1 ft (dep on M1) for correct placement of decimal point after addition (of appropriate values) (SC: B1 for attempting to add 32 lots of 8.63)
23	(a)		30	2	M1 for 25 ÷ 10 **or** 2.5 seen **or** 10 ÷ 25 **or** 0.4 seen **or** 12+12+6 oe **or** a complete method eg. 25×12÷10 oe A1 cao
	(b)	1000 ÷ 200 × 12	60	2	M1 for 500 ÷ 50 **or** 1000 ÷ 200 **or** 500 ÷ 10 **or** correct scale factor clearly linked with one ingredient eg 10 with sugar or 5 with butter or flour or 50 with milk **or** an answer of 120 or 600 A1 cao

PMT

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
24		Acton after 24, 48, 72, 96, .. Barton after 20, 40, 60, 80, .. LCM of 20 and 24 is 120 9:00 am + 120 minutes **OR** Acton after 24, 48, 1h 12 min... Barton after 20, 40, 1 h LCM is 2 hours 9:00 am + 2 hours **OR** Times from 9:00 am when each service leaves the bus station Acton at 9:24, 9:48, 10:12.. Barton at 9:20, 9:40, 10:00.. **OR** 20 = 2 × 2 × 5 24 = 2 × 2 × 2 × 3 2 × 2 × 2 × 3 × 5 = 120	11:00 am	3	M1 for listing multiples of 20 and 24 with at least 3 numbers in each list ; multiples could be given in minutes or in hours and minutes (condone one addition error in total in first 3 numbers in lists) A1 identify 120 (mins) **or** 2 (hours) as LCM A1 for 11:00 (am) **or** 11(am) **or** 11 o'clock **OR** M1 for listing times after 9am when each bus leaves the bus station, with at least 3 times in each list (condone one addition error in total in first 3 times after 9am in lists) A1 for correct times in each list up to and including 11:00 A1 for 11:00 (am) **or** 11(am) **or** 11 o'clock **OR** M1 for correct method to write 20 and 24 in terms of their prime factors 2, 2, 5 and 2, 2, 2, 3 (condone one error) A1 identify 120 as LCM A1 for 11:00 (am) **or** 11(am) **or** 11 o'clock

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
25	(a)		$6y - 15$	1	B1 cao
	(b)		$4x(2x + y)$	2	B2 cao (B1 for $x(8x + 4y)$ **or** $2x(4x + 2y)$ **or** $4(2x^2 + xy)$ **or** $4x(ax + by)$ where a, b are positive integers **or** $ax(2x + y)$ where a is a positive integer **or** $4x(2x \quad y)$)
	(c)	$10t = gh$ $h = \frac{10t}{g}$	$\frac{10t}{g}$	2	M1 for clear intention to multiply both sides of the equation by 10 (eg. ×10 seen on both sides of equation) **or** clear intention to divide both sides of the equation by g (eg. ÷ g seen on both sides of equation) **or** $10t = gh$ **or** $\frac{t}{g} = \frac{h}{10}$ **or** fully correct reverse flow diagram eg. ← ×10 ← ÷g ← A1 for $\frac{10t}{g}$ oe

1MA0_1F					
Question		**Working**	**Answer**	**Mark**	**Notes**
26	(a)	$2 \times 5 \times 2 = 20$ $300 \div 20 =$	15	3	M2 for $300 \div (2 \times 5 \times 2)$ oe (M1 for $2 \times 5 \times 2$ **or** 20 seen **or** $300 \div (2 \times 5)$ **or** 30 seen A1 cao
	(b)	$c = \frac{30 \times 40}{150}$ =	8	2	M1 for $\frac{30 \times 40}{150}$ **or** 1200 seen A1 cao
27		$3x - 15 = 2x + 24$ $x = 39$ **OR** $2x + 3x - 15 + 2x + 2x + 24 = 360$ $9x + 9 = 360$ $9x = 351$ $x = 39$ **OR** $2x + 2x + 24 = 180$ $4x + 24 = 180$ $4x = 156$ $x = 39$ **OR** $2x + 3x - 15 = 180$ $5x - 15 = 180$ $5x = 195$ $x = 39$	39	3	M1 for forming an appropriate equation eg $3x - 15 = 2x + 24$ **or** $2x + 3x - 15 + 2x + 2x + 24 = 360$ oe **or** $2x + 2x + 24 = 180$ oe **or** $2x + 3x - 15 = 180$ oe **or** $2x + 3x - 15 = 2x + 2x + 24$ M1 (dep) for correct operation(s) to isolate x and non-x terms in an equation to get $ax = b$ A1 cao **OR** M2 for $\frac{351}{9}$ **or** $\frac{195}{5}$ **or** $\frac{156}{4}$ oe A1 cao

PMT

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481
Email publication.orders@edexcel.com

Order Code UG032619 Summer 2012

For more information on Edexcel qualifications, please visit our website
www.edexcel.com

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual
Llywodraeth Cynulliad Cymru
Welsh Assembly Government
Rewarding Learning

PMT

edexcel

Mark Scheme (Results)

November 2015

Pearson Edexcel GCSE
In Mathematics B (2MB01)
Higher (Calculator) Unit 1

ALWAYS LEARNING PEARSON

PMT

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

November 2015
Publications Code UG042904
All the material in this publication is copyright
© Pearson Education Ltd 2015

NOTES ON MARKING PRINCIPLES

1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.

2 Mark schemes should be applied positively.

3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Note that in some cases a correct answer alone will not score marks unless supported by working; these situations are made clear in the mark scheme. Examiners should be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.

5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

6 Mark schemes will award marks for the quality of written communication (QWC).
The strands are as follows:

i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labelling conventions.

ii) *select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.

iii) *organise information clearly and coherently, using specialist vocabulary when appropriate.*
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

PMT

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.
If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.
If there is no answer on the answer line then check the working for an obvious answer.
Partial answers shown (usually indicated in the ms by brackets) can be awarded the method mark associated with it (implied).
Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks; transcription errors may also gain some credit. Send any such responses to review for the Team Leader to consider.
If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.
Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect cancelling of a fraction that would otherwise be correct
It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.

10 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).
Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.
If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.
If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded (embedded answers).

12 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers

Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

14 The detailed notes in the mark scheme, and in practice/training material for examiners, should be taken as precedents over the above notes.

Guidance on the use of codes within this mark scheme
M1 – method mark for appropriate method in the context of the question A1 – accuracy mark B1 – Working mark C1 – communication mark QWC – quality of written communication oe – or equivalent cao – correct answer only ft – follow through sc – special case dep – dependent (on a previous mark or conclusion) indep – independent isw – ignore subsequent working

PMT

5MB1H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
1.	(a)		0.2	2	M1 for 1 – (0.15 + 0.41 + 0.24) A1 cao
	(b)		12	2	M1 for 50 × 0.24 oe A1 cao
2.	(a)		(146, 41)	1	B1 for point plotted
	(b)		Description	1	B1 for description of dynamic relationship or statement of positive correlation
	(c)		155 – 158	2	M1 for a single straight line segment with positive gradient that could be used as a line of best fit or a horizontal line from 47.5 A1 answer in the range 155 – 158 or ft from line

PMT

<table>
<tr><th colspan="5">5MB1H_01 November 2015</th></tr>
<tr><th colspan="2">Question</th><th>Working</th><th>Answer</th><th>Mark</th><th>Notes</th></tr>
<tr><td>3.</td><td></td><td><table><tr><td></td><td>Bus</td><td>Pre</td><td>Ec</td><td>Tot</td></tr><tr><td>M</td><td>(30)</td><td>28</td><td>34</td><td>(92)</td></tr><tr><td>F</td><td>14</td><td>32</td><td>(62)</td><td>108</td></tr><tr><td>Total</td><td>(44)</td><td>(60)</td><td>96</td><td>(200)</td></tr></table>( ) value given</td><td>28</td><td>4</td><td>M1 for total female students 200 – 92 or 108 seen; or for total Economy passengers 200 – 44 – 60 or 96 seen.
M1 for total male passengers in Economy “96”-62 or 34 seen; or for total female Premium “108” – 62 – (44-30) or 32 seen
M1 for 92 – 30 – “34” or for 60 – “32”
A1 cao

OR

Answers may appear in a two-way table with no other method seen
B1 for Female total 108 or Total Economy 96
M1 for “96” – 62 or 34 seen in Male Economy;
or “108” – 62 – (44 –30) or 32 seen in Female Premium
M1 for 92 – 30 – “34” or for 60 – “32”
A1 cao</td></tr>
<tr><td>4.</td><td>(a)</td><td></td><td>Out: 1023/1148
Return: 1737/1918
in Colwyn 5h 49min</td><td>4</td><td>B1 for selecting 1023 train (cheapest out)
B1 for selecting 1737
B1 for 5h 49min (spent in Colwyn)
C1 (dep on at least B1) clearly stating departure and arrival times of both trains used and correct time in Colwyn for their trains</td></tr>
<tr><td></td><td>(b)</td><td></td><td>133.40</td><td>3</td><td>M1 for 30% of any adult fare
M1 for 2 × adult fare + 3 × child fare for at least one of their chosen trains
A1 (accept 133.4)

(SC B2 for £188.60)</td></tr>
</table>

PMT

5MB1H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
5.			Reasons	2	B1 for one reason eg no quantity (units), no time frame, overlapping intervals, not exhaustive B1 for second (different) reason
6.			$\frac{5x + 4}{3}$	2	M1 $x + x + 4 + 3x$ $(= 5x + 4)$ A1 for $(x + x + 4 + 3x) \div 3$ oe
7.		2500 + 2500 × 3.5÷100 = 2500 + 87.50 = 2587.50 2587.50 + 2587.50 × 3.5 ÷ 100 = 2587.50 + 90.5625	2678.06	3	M1 for 2500 × 1.035 or 2500 + 2500 × 0.035 oe or for 2587.5(0) or 87.5(0) or 8750 or 2412.5(0) M1 (dep) for "2587.5" × 1.035 or for "2587.5" + "2587.5" × 0.035 or for "2578.5" + "90.56(25)" or for 2678 or 2678.1(0) or 2678.07 or 2678.06… A1 cao NB: if correct answer seen then ignore any extra years Alternative method: M2 for 2500 × 1.035^n where $n \geq 2$ or for 2678 or 2678.07 or 2678.06… A1 cao

PMT

5MB1H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
8.	(a)		$10 < n \leq 20$	2	M1 for use of cf to find the 20.5^{th} or 21st value A1 cao
	(b)	$(5 \times 14) + (15 \times 13) + (25 \times 8)$ $+ (35 \times 4) + (45 \times 2)$ $70 + 195 + 200 + 140 + 90$ $695 \div 41$	16.95	4	M1 for $\sum fx$ with consistent values within interval (including end points); accept one error M1 for $\sum fx$ with correct midpoint values M1 (dep M1) for "$\sum fx$" $\div 41$ or "695" $\div 41$ A1 for 16.95(12….)
9.			Box plot	3	M1 for UQ = 1.7 + 2.7 (= 4.4) or 4.4 shown on box plot B1 for box plot with at least 3 pieces of information correctly plotted B1 for fully correct box plot NB: there must be a box
10.	(a)		5, 23, 35, 39, 40	1	B1 for all correct
	(b)		cf graph	2	B1 for 4 or 5 "points" plotted consistently dep on sensible table (condone 1 addition error) B1 correct graph
	(c)		3	2	M1 for evidence of reading off from 187 (eg vertical line) on cf graph or $35 + \frac{2}{5} \times 4$ A1 ft from cf graph

5MB1H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
11.			20 shown	5	B1 for 3 combinations (1 + 8, 5 + 4, 7 + 2) spacing amended M1 for partial working $\frac{3}{20}\times 80$ or $\frac{3}{20}\times 3$ oe or 80 × 3 (= 240) M1 for complete working $\frac{3}{20}\times 80\times 3$ oe M1 (income) 80 × 70 (= 5600) or 80 × 0.7 (= 56) C1 for "56 – 36=20" clearly stated
12.	(a)		3	2	M1 for $\frac{50}{600}\times 40$ oe or for 3.33… oe A1 cao
	(b)		75	2	M1 for $\frac{n}{600}\times 40$ or for $\frac{5\times 600}{40}$ where '5' can be 4.5 to 5.5 A1 for 75; accept answers in the range 68 – 82 supported by working.

PMT

5MB1H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
13.			400 and correct assumption	4	M1 for partial working eg $\frac{60}{12}$ oe or 20% or $\frac{1}{5}$ seen or 80 ÷ 12 (= 6.66..) or $\frac{12}{80}$ oe M1 for complete method eg $\frac{80\times 60}{12}$ or 80 × 5 or 6.66..× 60 or $\frac{12}{60}=\frac{80}{n}$ oe or 80 ÷ 0.2 oe A1 cao C1 for a correct mathematical assumption eg population has not changed overnight or mark which does not wear off or sample is random etc

PMT

5MB1H_01 November 2015					
Question		**Working**	**Answer**	**Mark**	**Notes**
14.			$\frac{56}{110}$	4	M1 for use of 10 as denominator of second probability M1 for a correct method to find the probability of at least one possible combination, $\frac{4}{11}\times\frac{7}{10}$ or $\frac{7}{11}\times\frac{4}{10}$ M1 for complete and correct method shown, eg $\frac{4}{11}\times\frac{7}{10}+\frac{7}{11}\times\frac{4}{10}$ A1 for $\frac{56}{110}$ oe **Alternative scheme for replacement** M1 for $\frac{4}{11}\times\frac{7}{11}$ or $\frac{7}{11}\times\frac{4}{11}$ M1 for $\frac{4}{11}\times\frac{7}{11}+\frac{7}{11}\times\frac{4}{11}$ OR B2 for $\frac{56}{121}$
15.		Frequency densities of 8 ÷ 10 = 0.8 16 ÷ 10 = 1.6 15 ÷ 5 = 3 12 ÷ 5 = 2.4 6 ÷ 20 = 0.3	Fully labelled histogram	3	B3 for fully correct histogram (B2 for 4 correct blocks) (B1 for 3 correct blocks) (If B0 then SC B1 for correct key eg $1cm^2$=2 birds or frequency ÷ class interval for at least 3 frequencies) NB apply the same mark scheme if a different frequency density is used

Modifications to the mark scheme for Modified Large Print (MLP) papers.

Only mark scheme amendments are shown where the enlargement or modification of the paper requires a change in the mark scheme.

The following tolerances should be accepted on marking MLP papers, unless otherwise stated below:
Angles: ±5°
Measurements of length: ±5 mm

PAPER: 5MB1H_01			
Question		**Modification**	**Notes**
Q02	(a)	Grid enlarged. Crosses changed to filled in circles. 146 cm changed to 150 cm. 41 kg changed to 40 kg.	
Q04		Crewe to Colwyn – First 2 columns removed. Colwyn to Crewe – First and last columns removed.	
Q06		MLP only: x → f.	
Q09		Weights changed to 1.5 kg, 6.5 kg, 2.5 kg, 1.5 kg, 3.5 kg. Grid enlarged.	
Q10	(b) (c)	Frequencies changed to 5, 15, 10, 5, 5. Grid enlarged. 187 cm changed to 182.5 cm.	
Q15		Frequencies changed to 5, 15, 15, 10, 10. Grid enlarged.	

PMT

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom

Question 21

Though a significant number of students did not attempt this question, a good proportion of those students who did attempt it scored full marks for a complete and correct solution. The most common errors made by those students who followed a correct method, that of finding the total number of onions weighing less than 60g or more than 120g, usually involved obtaining an incorrect frequency for the number of onions weighing between 120g and 135g or for the number of onions weighing between 135g and 180g, or both. This usually arose because of using an incorrect class interval . Students who used the strategy of finding that there were 420 onions weighing between 60g and 120g were generally more successful. Some students opted to work with percentages, either giving their final answer as a percentage, or rounding a recurring decimal to 1decimal place and hence introducing a rounding error. In these cases the final accuracy mark was usually lost. There were a significant number of students who used the heights of the bars to represent frequencies. This approach could not be awarded any credit.

Question 18

Only a few gained marks on this question. Some were able to find at least one frequency but then failed to proceed. In general, those who got to 32 were able to draw the bar at 3.2 Incorrect answers came from just adding the heights of the bars or misreading the scales when drawing the bar.

Question 17

It was relatively rare to see a fully correct solution to this question with the consequent award of 4 marks. However, many more students were able to access the two marks for correct processes to work out the frequencies represented by each bar. These marks were given provided that these frequencies were linked to the method leading to the student's final answer. As is often the case with a question involving a histogram, weaker students often used the heights of the bars as frequencies and so no marks could be awarded.

Question 17

This histogram question was pleasing to mark, with a surprising number of students gaining some credit. Many were able to gain the first mark for one (or more) correct frequencies, and of these, a significant number going on to gain three marks for the correct process to find the total and then find 20%. The last mark proved extremely challenging but gaining 3 marks at this level showed good performance for a number of students. It was interesting to see that almost all worked with frequencies rather than areas, which shows a good understanding of histograms and frequency density.

Question 17

In part (a) of this question, lower attaining students often drew a bar chart and did not recognise the need to find and use frequency densities. Some students used the midpoints of the class intervals erroneously together with the frequencies to get values which they then used to draw a diagram. Having said that, there was a good proportion of students who did successfully calculate frequency densities and use these to draw a histogram. It was interesting to note that a significant number of students drew the first four bars correctly but then made an error drawing the bar representing the interval $3.0 < d \leq 5.0$

Part (b) of the question was less well done with only a small proportion of students able to use the ratio of the areas of the two bars concerned to find a correct expression. Examiners did award one mark to a significant number of students who showed that they needed to compare the heights, widths and areas of the two bars concerned.

Question 14

The students who worked out frequency densities and used these to draw a histogram often gained all three marks. Some students drew the first three bars correctly but then made an error drawing the bar representing the interval $90 < w \leq 110$ and gained two of the three marks. A few students with the correct frequency densities did not draw the bars in the correct intervals and gained one mark only. Weaker students often did not recognise the need to find frequency densities and drew a bar chart. Some students used the midpoints of the class intervals and the frequencies to get values which they then used to draw a frequency polygon.

Question 16

It was clear that histograms is a topic that many candidates continue to find difficult, and so many were unable to draw the missing bar correctly. Very few worked with frequency density and the fact so many achieved a single mark was down to the bar ranging from 20 to 30 hours having the same class width as the one going from 30 to 40, and candidates found the correct frequency for that bar using the difference in heights. Very few written calculations were seen in part (a). Students should be encouraged to write working down such as how to calculate the scale. Values were given in the question, so a simple calculation of 28 divided by the class width would have got a mark. These errors continued into part (b) where again we saw few fully correct responses, although some candidates gained 1 mark for summing 3 out of 4 correct frequencies.

Question 15

This question, whilst relatively standard, especially parts (a) and (b), was answered very poorly by the majority of candidates. The first two parts required students to fill in a frequency table from a histogram, and then to complete the graph. Part (c) asked for an estimate of the median. In fact, students were more likely to score a mark in part (c), normally for finding at least 2 areas, but it was very rare to see any further correct working.

Question 13

In part (a) the students who worked out the frequency densities and used these to draw a histogram often drew five correct bars. This alone was not sufficient for full marks because the two axes needed to be correctly scaled and labelled. Some arithmetic errors such as $22 \div 5 = 4.5$ were made when working out the frequency densities. Students should be encouraged to show the method they use to work out each frequency density so that examiners can see if incorrect values come from a correct method of dividing frequency by class width. Some students made an error drawing one of the bars but still gained two of the three marks. A few students with the correct frequency densities did not draw the bars in the correct intervals and gained one mark only. Weaker students often did not recognise the need to find frequency densities and drew a bar chart or a frequency polygon or a cumulative frequency graph.

A similar proportion of students gained full marks in part (b) for working out an estimate for the fraction of the 150 people who were in the shop for between 20 minutes and 40 minutes. As this can be done by using the table or by using the histogram some students who had gained no marks in part (a) were able to give a correct fraction in part (b) by using the table. The marks were also awarded in part (b) for answers which followed through correctly from incorrect histograms. A common mistake was to give the answer as 67, not as a fraction, and this gained one mark only.

Question 16

This histogram question pulled together several different skills. Firstly, candidates had to find the frequencies for two completed bars on the histogram. They then had to use these to find the frequencies of the missing bars, then their frequency densities and finally draw the missing bars in.
Those who made a successful start to the question did so by finding at least one of the missing frequencies correctly. The second mark was then gained by subtracting the frequencies from 60 and halving to find the frequency for the two missing bars. Although a good number made a single error in reading the frequency density of the given bars, they were able to still gain the next two method marks. The final method mark was for a method to find the frequency density for one of the missing bars, or for drawing one bar correctly. We did see cases of correctly drawing the first bar, but then simply drawing the second at the same height.

Question 15

Many responses showed that they knew the frequency can be found by multiplying the frequency density by the class width and gained the first mark for one (or more) correct frequencies. Frequencies were often written on the bars of the histogram. Candidates should be encouraged to show the method they use to work out each frequency so that examiners can see if incorrect values come from a correct method. Many responses went on to gain all four marks for working out a correct estimate for the probability that the biscuit will have a weight between 20 grams and 40 grams. Sometimes the accuracy mark was lost because of poor addition and 30 + 100 + 120 + 150 = 300 was a common mistake when finding the total frequency. It was not uncommon for responses to correctly find the number of biscuits with a weight between 20 grams and 40 grams (130) and then not find the total number of biscuits. These responses gained two of the four marks. Weaker responses often used the heights of the bars as frequencies and no marks could be awarded.